1    Nanoparticle-mediated transformation expands horizon of organism engineering
2
3

4    Katherine E. French*[1]
5
6    [1] Department of Plant and Microbiology, University of California Berkeley, 111 Koshland Hall, Berkeley,
7    CA
8
9    Correspondence: katherine.french@berkeley.edu

10
11

12   **Our ability to engineer organisms will unlock new discoveries in how cellular processes work and enable new**
13   **advances in biotechnology. However, many organisms are recalcitrant to traditional transformation methods.**
14   **Here, we describe a new nanoparticle-based method of transformation which can be used to transform bacteria,**
15   **plants, algae and diatoms.**

16
17
18
Our ability to engineer organisms will unlock new discoveries in how cellular processes work and enable

new advances in biotechnology (Khalil and Collins 2010). This process is known as 'transformation' and

relies on electric currents, heat shock, or chemicals to create tiny pores in organism cell membranes that

allow foreign genetic material to enter the host. However, many organisms are recalcitrant to these

methods (Brenner et al. 2008). Recent advances in nanoparticle technology have illustrated that

quantum dots can deliver siRNA (typically of 20-22 bp) and other molecules to human and mice cells,

but these methods do not work on bacteria or organisms with cell walls (Kim et al. 2017; Pierrat et al.

2015). Here, we describe a new carbon-nanoparticle based transformation system. Nanoparticle

composition, polymer coating, pH, and plasmid DNA (p-DNA) concentration were key factors in

transformation success. Pilot experiments with eukaryotic, prokaryotic, and difficult-to-transform

species suggest any organism that can take-up the nanoparticles we designed can be transformed using

this method.

We prepared 15 different nanoparticles to assess the following properties (1) organism uptake; (2)

cellular localization; (3) transformation ability; and (4) cellular toxicity. We replicated two nanoparticle

types from previous studies; the rest of the nanoparticles are unique to this study. For nanoparticles

with desired optical and functional properties, we experimented with different polymer coatings (bPEI

and PEG), polymer concentrations (0.0125-0.05 g/mL), plasmid concentrations (1 ng-1 ug/ul), and

nanoparticle-plasmid incubation time (10 min. to 24 h). The particles best suited for general (non-

organelle) transformation were made of citric acid or glucose, passivated with low concentrations of

bPEI (0.01-0.02 g/mL) and plasmid (1-4ng/ul), and with a short nanoparticle-plasmid conjugation time

(15-20 min.). Particles made with threonine localized to chloroplasts and those made with arginine,

lysine, valine and serine localized to mitochondria (**SI Table 1**). Nanoparticles made with specific amino

acid combinations may mimic the role of terminal sequences in guiding proteins targeted to specific

cellular compartments (Carrie et al. 2009; de Castro et al. 1996; Lodish et al. 2000).

Cellular uptake studies were carried out in the following organisms: plants/mosses (*Arabidopsis*

*thaliana, Nicotiana benthamiana, Physcomitrella patens* and *Selaginella kraussiana*), protists (*Physarum*

*polycephalum*), bacteria (*Pseudomonas flourescens*), cyanobacteria (*Synecoccochus*), diatoms

(*Thalassiosira oceanica*), algae (*Penium margaritaceum, Spyrogira* sp.), fungi (*Trichoderma reesei),* and

coccolithophores (*Pleurochrysis certerae, Gephyrocapsa oceanica*). This is the first study to show algae,

diatoms and coccolithophores will take up nanoparticles.   Nanoparticles entered cells in two ways: (1)

through membrane pores or (2) invagination of cell membranes.  For example, in diatoms nanoparticles

appear to enter the cell through external pores in the frustule; in algae such as *Penium*, in-folding of the

cell membrane around nanoparticle aggregates was observed and these vesicles seemed to transport

nanoparticles in and out of the cell (**Fig. 1 A and B**). The latter correlates with Giraldo et al. 2014, who

proposed that single-walled carbon nanotubes may enter the double-membranes of chloroplasts via

diffusion, becoming coated with glycerolipids in the process. Experiments with vascular plants,

filamentous fungi, and *P. polycephalum* indicate nanoparticles can be transported long distances (either

within the vascular system or within vesicles) (**SI Fig. 1**). Based on material composition, nanoparticles

localized to six places: the cell membrane, cytoplasm, vacuole, nucleus, mitochondria, and chloroplasts.

In general, carbon-based nanoparticles are non-toxic to cells. This is in contrast to nanoparticles made

out of heavy metals (e.g. 'quantum dots') which can kill cells and may cause environmental harm

(Hardman 2006; Valizadeh et al. 2012). Study organisms did show signs of stress when exposed to

nanoparticles with high bPEI (0.05-0.1 g/mL) and plasmid concentrations (300 ng-1 ug/ul) or with low pH

(< pH 4). Signs of stress included retraction of the cytoplasm from the cell wall in vascular plants; cell

rupture; and loss of innate chloroplast structure (specifically in algae). Once nanoparticles were

optimized to decrease cell toxicity, organisms could survive on media (agar gel plates or liquid growth

media) containing nanoparticles for over 30 days.

The following organisms were successfully transformed using glucose and citric acid based nanoparticles

coated with bPEI and conjugated with a Venus NLS plasmid, a MCherry NLS plasmid, or a GFP HDEL

plasmid: *A. thaliana*, *N. benthamiana*, *S. kraussiana*, *P. margaritaceum* and *T. oceanica* (**Fig. 1 A, C-F**; **SI**

**Fig. 2**). Organisms were exposed to the following treatments: control (no treatment), bPEI, bPEI + pDNA,

CND, CND + pDNA. No transformation was seen using plasmids bound to bPEI, a transfection method

commonly used in mammalian cells (Kim et al. 2017; Pierrat et al. 2015), and exposure to bPEI alone

resulted in cell death for plants and algae. Successful transformation using plasmids bound to CNDs was

indicated by fluorescence of the nuclear envelope and the endoplasmic reticulum. To further confirm

transformation, we transformed *P. margaritaceum* with a 35S ubi-GUS plasmid. Non-transformed cells

appear clear or yellowish; transformed cells appear blue (**Fig. 1 E**). This system can be used for transient

and stable expression of foreign genes in both chassis organisms and difficult to transform species. For

example, the alga *P. margaritaceum* showed nuclear YFP and MCherry labelling eight days after

transfection, suggesting the plasmid DNA was inserted into the hosts' genomes. Other images show cells

dividing, with both cells exhibiting nuclear fluorescence, suggesting the DNA is inherited (**Fig. 1 F**).

Carbon nanoparticle-based transformation presents a promising new method of altering the genetic

make-up of cells. This is the first study to show the transport and expression of large plasmids (>10 kbp)

into recalcitrant organisms using carbon nanoparticles. Future studies should look at the metabolic

impact of CND exposure, the degradation of CNDs in cells and the environment, plasmid size limitations

which may affect transformation, and whether organelle-targeting nanoparticles can be used to

genetically modify chloroplasts and mitochondria. Further development of this technology will lead to

new discoveries across the life sciences and commercial gains in the growing biotech sector.

92
93

94
95 **Fig. 1**. **Organism uptake of carbon nanoparticles and transformation**. A. Expression on VNLS in *T. oceanica*. The
96 image shows dorsal and lateral views of several diatoms in different stages of life. B. Uptake of CNDs in *P.*
97 *margaritaceum*. Arrows point to vesicular structures filled with nanoparticles. C. Transformation of *N.*
98 *benthamiana* with VNLS. D. Expression of VNLS in *S. kraussiana* 16 hours after exposure to CND-pDNA complexes;

boxes show magnified cells with fluorescent nuclei. E. GUS-staining of *P. margaritaceum*. F. Dividing *P. margaritaceum* eight days after exposure to CND-pDNA complexes. N= nucleus. Confocal images (A, B, C, F) taken using excitation 405/488/543nm. Images are composed of superimposed z-sections. Images D and E were taken with a Leica light microscope using a YFP filter and an LED light respectively.

Author contributions

KEF designed the nanoparticles, performed the nanoparticle-uptake and transformation experiments, imaged the experiments, and wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

**References**

Brenner, K., L. You and F. Arnold. 2008. Engineering microbial consortia: a new frontier in synthetic biology. *Trends in Biotechnology* 26.9: 483-489.

Carrie, C., E. Giraud and J. Whelan. 2009. Protein transport in organelles: Dual targeting of proteins to mitochondria and chloroplasts. *FEBS Journal* 276: 1187–1195.

de Castro Silva Filho, M., F. Chaumont, S. Leterme, M. Boutry. 1996. Mitochondrial and chloroplast targeting sequences in tandem modify protein import specificity in plant organelles. *Plant Molecular Biology* 30: 769-780.

Giraldo, J.P., M. P. Landry, S. M. Faltermeier, T. P. McNicholas, N. M. Iverson, A. A. Boghossian, N. F. Reuel, A. J. Hilmer, F. Sen, J. A. Brew and M. S. Strano. 2014. Plant nanobionics approach to augment photosynthesis and biochemical sensing. Nature Materials 13: 400-410. DOI: 10.1038/NMAT3890.

Hardman, R. 2006. A Toxicologic Review of Quantum Dots: Toxicity Depends on Physicochemical and Environmental Factors. *Environmental Health Perspectives* 114.2: 165–172.

Kim, S., Y. Choi, G. Park, C. Won, Y. Park, Y. Lee. 2017. Highly efficient gene silencing and bioimaging based on fluorescent carbon dots *in vitro* and *in vivo*. *NanoResearch* 10.2: 503–519.

Khalil, A. and J. Collins. 2010. Synthetic biology: applications come of age. *Nature Reviews Genetics* 11: 367-379.

*Molecular Cell Biology*. 4th edition. Lodish ,H., A. Berk, S.L. Zipursky, et al. New York: W. H. Freeman. 2000.

Pierrat, P., R. Wang, D. Kereselidze, M. Lux, P. Didier, A. Kichler, F. Pons, L. Lebeau. 2015. Efficient in vitro and in vivo pulmonary delivery of nucleic acid by carbon dot-based nanocarriers. *Biomaterials* 51: 290e302.

Valizadeh, A., H. Mikaeili, M. Samiei, S. M. Farkhani, N. Zarghami, M. Kouhi, A. Akbarzadeh, S. Davaran. 2012. Quantum dots: synthesis, bioapplications, and toxicity. *Nanoscale Research Letters* 7.1: 480.