Quantifying the efficiency and biases of forest *Saccharomyces* sampling strategies

Primrose J. Boynton*[1], Vienna Kowallik[2], Doreen Landermann[1], Eva H.

Stukenbrock[1,3]

1) Max-Planck Institute for Evolutionary Biology, Environmental Genomics Research

Group, Plön, Germany

2) Okinawa Institute of Science and Technology, Okinawa, Japan

3) Christian-Albrechts Universität, Botanisches Institut, Kiel, Germany

* To whom correspondence should be addressed:

pboynton@evolbio.mpg.de

August-Thienemann-Str. 2, 24306 Plön, Germany

Keywords: *Saccharomyces paradoxus*, *Saccharomyces cerevisiae*, forest, enrichment

culture, soil, environmental isolates

Abstract

Saccharomyces yeasts are emerging as model organisms for ecology and

evolution, and researchers need environmental *Saccharomyces* isolates to test

ecological and evolutionary hypotheses. However, methods for isolating

Saccharomyces from nature have not been standardized and isolation methods can

influence the genotypes and phenotypes of studied strains. We developed a direct

isolation method for forest floor *Saccharomyces* and compared its success and

phenotypic biases to a previously published enrichment-based isolation method. In a

European forest, direct isolation was more successful at isolating *S. paradoxus*, but

also more labor intensive, than enrichment culturing. Average growth rates of *S.*

paradoxus isolates collected using the two methods did not differ at the enrichment

isolation temperature, but variances in growth rates did: direct isolation produced a

collection of *S. paradoxus* isolates with less variation in growth rates than enrichment

culturing. In other words, enrichment culturing sampled more phenotypic diversity

than direct isolation. Enrichment culturing also sampled more *Saccharomyces* species

diversity than direct isolation, including our only isolations of rare *S. cerevisiae*.

Enrichment culturing may sample higher *Saccharomyces* phenotypic and species

diversity because of variations in interactions between yeasts and the other microbes

that were present in the soil and leaf litter samples. We recommend direct culturing

for researchers interested in randomly sampling their study habitats and enrichment

culturing for researchers interested in discovering new *Saccharomyces* phenotypes or

rare *Saccharomyces* species from natural environments. We include step-by-step

sampling protocols in the supplemental materials.

Introduction

Naturally-occurring *Saccharomyces* populations are models for ecology and

evolution (Boynton & Greig, 2014). Use of these models has led to exciting

discoveries about the ecology and evolution of microbial phenotypes; for example,

adaptation to climate can lead to speciation (Leducq et al., 2014), domesticated *S.*

cerevisiae is more phenotypically diverse than wild *S. paradoxus* (Warringer et al.,

2011), and interspecific hybrids can have high fitnesses in stressful environments

(Bernardes, Stelkens, & Greig, 2017; Stelkens, Brockhurst, Hurst, Miller, & Greig,

2014). These studies made inferences based on the phenotypes and genotypes of

isolates collected from wild and domesticated substrates. And *Saccharomyces*

substrates are diverse: wild substrates include tree bark, insect guts, leaf litter, soil,

fruits, and parasitic *Cyttaria* galls, (Kowallik & Greig, 2016; Libkind et al., 2011;

Mortimer & Polsinelli, 1999; Sampaio & Goncalves, 2008; Stefanini et al., 2012), and

domesticated substrates include wine, beer, bread, kimchi, kombucha, palm wine, and

pulque, among many other substrates (Boynton & Greig, 2016; Carbonetto,

Ramsayer, Nidelet, Legrand, & Sicard, 2018; Estrada-Godina et al., 2001; Ezeronye

& Okerentugba, 2001; Gallone et al., 2016; Greenwalt, Steinkraus, & Ledford, 2000;

Jeong, Jung, Lee, Jin, & Jeon, 2013). One challenge of environmental yeast sampling

is to minimize sampling biases so researchers can assure that differences among

source environments, not sampling techniques, are responsible for observed

phenotypic patterns.

Enrichment culturing is a reliable and frequently-used method for isolating

difficult-to-culture microbes, including *Saccharomyces*, from natural environments

(Schlegel & Jannasch, 1967; Sniegowski, Dombrowski, & Fingerman, 2002) (Figure

1A). Microbiologists have been relying on enrichment cultures for over a century

Figure 1: Schematic illustration of sampling strategies used to isolate *Saccharomyces* for this project. A) Enrichment culturing B) Direct plating. Photo: Doreen Landermann

(Beijernick, 1961), and have used enrichment culturing to isolate many of the model

Saccharomyces strains commonly used in laboratory studies (Johnson et al., 2004;

Liti et al., 2009; Sniegowski et al., 2002). To isolate a microbe using enrichment

culturing, a researcher adds a small amount of natural material to a growth medium

designed to be hospitable to the target microbe and inhospitable to other microbes

(Liti, Warringer, & Blomberg, 2017; Schlegel & Jannasch, 1967). If the enrichment

medium is well-designed, the target microbe is expected to grow in abundance, and

after some incubation time, this enrichment culture can be streaked to solid media and

colonies of the target microbe can be easily isolated. An alternative to enrichment

culturing is to spread a microbial substrate directly onto a selective solid medium,

with or without dilution, and to pick colonies which morphologically resemble the

target microbe (Stefanini et al., 2012) (Figure 1B).

Because it can be difficult to isolate *Saccharomyces* from natural substrates,

many investigations of wild *Saccharomyces* rely on enrichment culturing, usually in

high-sugar, acidic media (Charron, Leducq, Bertin, Dube, & Landry, 2014; Robinson,

Pinharanda, & Bensasson, 2016; Sniegowski et al., 2002; Sweeney, Kuehne, &

Sniegowski, 2004). Comparative studies of *Saccharomyces* genomes have been

carried out using *Saccharomyces* strains isolated using disparate strategies, including

both enrichment and direct culturing (Liti et al., 2009; Peter et al., 2018). However,

isolation strategy can influence the genotypes and phenotypes of isolated microbes

(Stefani et al., 2015), and we were concerned about the biases that might be

introduced during enrichment culturing of *Saccharomyces* yeasts. Specifically, we

were concerned that isolation method might bias results in studies of *Saccharomyces*

phenotypes. For example, enrichment culturing might select for individuals with high

relative fitness in the enrichment medium. Such potential biases in sampled yeast

phenotypes are likely to lead to biases in sampled genotypes because genetic

information is responsible for expressed phenotypes. Isolation biases have also been

suggested as potential explanations for differences between results of culture-

dependent and culture-independent studies of environmental *Saccharomyces*

(Alsammar et al., 2018).

This study's goals were to compare isolation success between enrichment

culturing and a direct culturing strategy, and to quantify biases in *Saccharomyces*

phenotypes (and therefore genotypes) that might be introduced when sampling a

forest environment. We tested the assumption that it is easier to sample

Saccharomyces from forest substrates using enrichment cultures than direct plating.

We also compared growth rates between *S. paradoxus* isolated using enrichment and

direct strategies. Enrichment culturing might decrease or increase sampled *S.*

paradoxus phenotypic diversity compared to direct plating, thereby decreasing or

increasing variance among *S. paradoxus* growth rates. For example, variance among

growth rates would be low (and average growth rates high) among *S. paradoxus*

isolated using enrichment cultures if the enrichment temperature and media select for

the fastest growing *S. paradoxus* present in every sample. Conversely, variance

among growth rates would be high for *S. paradoxus* isolated using enrichment

cultures if diversity in the non-*Saccharomyces* microbial communities present on

sampled substrates select for diverse *S. pardoxus* among samples.

111   To test these predictions, we compared *Saccharomyces* sampling success and

isolates' growth rates among soil and leaf litter samples from a well-studied northern

German forest (Kowallik & Greig, 2016; Kowallik, Miller, & Greig, 2015). A

previous study showed that *S. paradoxus*, the wild sister species of the lab model *S.*

cerevisiae, is readily isolated using enrichment cultures from oak leaf litter in this

forest (Kowallik & Greig, 2016). We were also previously able to isolate *S.*

paradoxus directly from these forest substrates without enrichment (Kowallik, 2015).

Methods

Field sampling and yeast isolation

121   All isolates were sampled from a mixed hardwood and conifer forest in

Nehmten, Schleswig-Holstein, northern Germany (Nehmtener Forst). We sampled

leaf litter and soil material from close to the bases of ten oak trees at four sampling

dates (Table 1), although not all trees were sampled at every date. Trees were between

12 and 744 m from one another. At each date, samples were collected from leaf litter

and the top organic layer of soil within one meter of the base of each tree. Paired leaf

litter and soil samples were collected on the north, south, east, and west side of each

tree at all collection days except 7 April, when samples were collected at an arbitrary

two of the four cardinal directions.

130   Material was collected simultaneously for the **direct plating** and **enrichment**

131   collections at each sampling point (Figure 1). First, leaf litter was collected by

132 Table 1: Number of sampling points from each tree at each timepoint (each sampling
133 point includes one enrichment and one direct plating sample)

	21 March 2017		7 April 2017		12 June 2017		10 July 2017	
	soil	litter	soil	litter	soil	litter	soil	litter
Tree 1	4	4	2	2	4	4	4	4
Tree 2	4	4	2	2	4	4	4	4
Tree 3	4	4	2	2	4	4	4	4
Tree 4	4	4	2	2	4	4	4	4
Tree 5	4	4					4	4
Tree 6			2	2	4	4	4	4
Tree 7			2	2	4	4	4	4
Tree 8			2	2	4	4	4	4
Tree 9					4	4	4	4
Tree 10							4	4

aseptically transferring litter into sterile collection tubes: approximately 5 ml of

compressed leaf litter was collected for the direct plating method and approximately

2 ml for the enrichment method. Then, the remaining leaf litter was removed from the

soil surface and the top approximate 2 cm of soil (mostly composed of soil organic

layer) were aseptically transferred into sterile collection tubes. As for leaf litter,

approximately 5 ml of compressed soil was collected for the direct plating method

and approximately 2 ml for the enrichment method. Instruments were sterilized

between samples using 70% ethanol. Samples were transported between the field and

lab at ambient temperature and processed within four hours of collection.

For **direct plating** (Figure 1A), material was mixed with 20 ml sterile water in

a sterile 50 ml tube, the mixture was vigorously mixed for at least 10 seconds with a

vortex mixer on its highest setting, and 0.2 ml of the resulting dirty liquid was

pipetted on each of two plates containing solid modified selective media PIM1 (3 g

yeast extract, 5 g peptone, 10 g sucrose, 3 g malt extract, 1 mg chloramphenicol, 80

ml ethanol, 5.2 ml 1 M HCl, and 20 g agar per liter) (Kowallik & Greig, 2016;

Sniegowski et al., 2002). Liquid was spread on plates using sterile glass beads, and

plates were left open in a laminar flow hood until dry. Plates were incubated for three

days at 30 ºC before colonies were picked.

For **enrichments** (Figure 1B), material was mixed with 10 ml liquid selective

media PIM1 (composition as for solid PIM1 but without agar) in a 15-ml sterile tube,

mixtures were inverted, and tubes were incubated, slightly open and without shaking,

at 30 ºC. After 10 days, a sterile wooden stick was inserted into each enrichment tube

and a small amount of liquid was streaked onto a single plate with solid selective

media PIM2 (20 g Methyl-(alpha)-D-glucopyranoside, 1 ml 5% Antifoam Y-30

emulsion, 6.7 g Yeast Nitrogen Base without amino acids, 4 ml 1M HCl, and 20 g

agar per liter) (Kowallik & Greig, 2016; Sniegowski et al., 2002), and plates were

incubated 4 days at 30ºC before colonies were picked.

We include these procedures as step-by-step protocols for the convenience of

future researchers in the supplementary materials (Supplemental File 1).

<u>Yeast identification</u>

After incubation, we streaked colonies with yeast-like morphology to fresh

YPD media (10 g yeast extract, 20 g peptone, 20 g dextrose, and 25 g agar per liter).

For each method, up to 6 (March and April sampling days) or 12 (June and July

sampling days) colonies per sample were selected. After one day of growth on YPD at

30 ºC, cultures were frozen at -80 ºC in 20% glycerol and a small amount of each

culture was transferred to sporulation media (20 g potassium acetate, 2.2g yeast

extract, 0.5 g dextrose, 870 mg complete amino acid mixture, and 25 g agar per liter).

Any cultures with bacteria-like morphology on YPD media (slimy culture and/or cells

smaller than 1 micron across) were not frozen and were discarded. Sporulation

cultures were incubated for at least three days at room temperature before being

screened under a compound microscope for *Saccharomyces*-like asci (tetrads).

177   All cultures producing tetrads were identified using sequencing of the internal

transcribed sequence (ITS), a region neighboring rRNA-coding DNA (Schoch et al.,

2012). We sequenced every strain using the ITS1/ITS4 primer pair (White, Bruns,

Lee, & Taylor, 1990). PCR mixes were 7-15 µl in volume and contained one yeast

colony, 0.5 µM each primer, and either 50% Phusion® High-Fidelity PCR master mix

with HF buffer or 1x HF-buffer, 100 µM dNTP mix, 3% DMSO, and 1 U/50 µl

Phusion DNA polymerase. PCR reactions were cycled at 98 ºC for 30 s, then 35

cycles of 98 ºC for 5 s, 62 ºC for 20 s, and 72 ºC for 30 s, plus a 10 min terminal

extension at 72 ºC. PCR products were cleaned using illustra™ ExoProStar™

according to the manufacturer's instructions, and sequenced on an ABI 3130xl

sequencer.

188   ITS sequences were compared to sequences from the type or neotype strains of

S. paradoxus, *S. cerevisiae*, *S. kudriavzevii*, and *S. mikatae* (Genbank accession

numbers NR_138272.1, NR_111007.1, KY105195.1, and KY105198.1). If a

sequence did not align with *Saccharomyces* sequences, we compared the sequence

with all sequences in the NCBI database from type strains using BLAST (Zhang,

Schwartz, Wagner, & Miller, 2000). If the sequence aligned with *Saccharomyces*

sequences but had more than one base pair different from its closest match, we

supplemented ITS sequences with sequences from the gene for translation elongation

factor 1 using primers EF1-983F and EF1-2212R (Rehner & Buckley, 2005) using the

protocols above, but with a PCR annealing temperature of 57 ºC. In some cases,

cultures originating from apparent single colonies were in fact mixtures of two yeast

species. We counted these colonies as *Saccharomyces* if sequences from one of the

species was *Saccharomyces*.

<u>Growth rates</u>

We compared the distributions of maximum growth rates between two groups

of *S. paradoxus* strains: strains collected using enrichment culturing and strains

collected using direct plating. To avoid confounding effects of environmental source

(*i.e.*, combination of substrate, date collected, and tree), we compared growth rates for

pairs of *S. paradoxus* strains originating from the same environmental source. In other

words, we collected a dataset of *S. paradoxus* growth rates from two groups of strains

with equal representations of combinations of substrate, date collected, and tree, and

differing only in the method used to isolate the strains. To ensure that all isolates were

pure *S. paradoxus* cultures, we streaked all isolates used for growth rate

measurements to single-colony cultures a second time and reidentified these cultures

by mating them with a *S. paradoxus* tester strain (NCYC 3708, α, ura3::KANMX,

ho::HYGMX). In total, 110 isolates (55 from each sampling method) were measured.

Growth rates were measured using an Epoch 2 microplate reader (Biotek

Instrument, Inc., Winooski, VT, USA) and calculated using the included Gen5

software version 3.03.14 (Biotek Instrument, Inc., Winooski, VT, USA). We first

inoculated strains in 0.2 ml liquid YPD media (composition as for solid YPD, but

without agar) in a 96-well microplate and incubated cultures without shaking or

measurement in the microplate reader at 30 ºC for 24 hours to condition strains to

microplate reader conditions. We then transferred 2 μl from each culture to 198 μl

fresh liquid YEPD in a new microplate and incubated the new microplate under the

same conditions for 20-24 hours. OD_{660} was measured during the second incubation

every ten minutes, and maximum growth rate (mOD_{660}/min) was calculated from the

maximum slope of each growth curve over four points (30 min total) using Gen5

software. Reported growth rates for each isolate are means of three replicates.

Statistical analyses

We compared sampling success across substrates (leaf litter or soil) and

methods (direct plating or enrichment) using a generalized linear mixed-effects model

with probability of isolating *Saccharomyces* (including *S. paradoxus* and *S.*

cerevisiae) as the response variable, substrate and method as fixed effects, and tree

and date as random effects. We selected the best model using a top-down strategy,

comparing Akaike's Information Criteria (AIC) after removing predictors from a full

model one by one.

We compared growth rate distributions by first comparing variances using

Levene's test (Levene, 1960) for homogeneity of variance, and then comparing

medians using a paired Wilcoxon signed rank test. Statistics were computed using R

version 3.3.1 (R Development Core Team, 2016) and the car and lme4 packages

(Bates, Machler, Bolker, & Walker, 2015; R Development Core Team, 2016).

Graphics were produced using the ggplot2 package (Wickham, 2016).

Results

Influence of sampling method on success isolating *Saccharomyces*

Direct plating was more successful than enrichment culturing for isolating

Saccharomyces from natural substrates ($z = 6.1$, $p < .001$) (Tables 2, 3, Figure 2). We

found *Saccharomyces* isolates in 45% of direct plating cultures and 19% of

enrichment cultures. However, enrichment culturing produced the only *S. cerevisiae*

found in this study: we found six *S. cerevisiae* isolates from a single enrichment

culture from tree 3 in March of 2017. All other *Saccharomyces* isolates found in this

study were *S. paradoxus*. Other detected yeast species included *Saccharomycodes*

ludwigii, *Torulaspora delbrueckii*, *Pichia membranifaciens*, and *Hanseniaspora*

osmophila, all of which have previously been found alongside *Saccharomyces* yeasts

in beverage fermentations (Domizio et al., 2011; Gschaedler, 2017).

Table 2: Model selection (Mixed-effects generalized linear model)

model	fixed effects	random effects	AIC	compared to	better model
1	method + substrate + method:substrate	1|tree + 1|month	447.20		
2	method + substrate + method:substrate	1|month	470.94	model 1	1
3	method + substrate + method:substrate	1|tree	451.99	model 1	1
4	**method + substrate**	**1|tree + 1|month**	**446.76**	**model 1**	**4**
5	method	1|tree + 1|month	480.34	model 4	4
6	substrate	1|tree + 1|month	485.71	model 4	4

257 Table 3: Model summary table (model 4)

	Estimate	Std. Error	z	p
intercept	-2.5324	0.4599	-5.506	< .001
method (plating)	1.5494	0.2559	6.054	< .001
substrate (soil)	1.4445	0.2542	5.683	< .001

Figure 2: Percentages of samples in which *Saccharomyces* could be detected using A) enrichment cultures or B) direct plating. Bars represent all samples for each category of sampling, and colors represent *Saccharomyces* species. *S. paradoxus* and *S. cerevisiae* were the only detected *Saccharomyces* species.

While the direct plating method was more successful than the enrichment

method, it was also more labor-intensive (Table 4). We screened 3.4 times as many

colonies for tetrads when using the direct plating method than we did using the

enrichment method. Only 32% of the processed direct plating colonies were *S.*

paradoxus, compared to 74% of enrichment colonies.

Table 4: Colonies processed and sampling success for each method

method	total colonies picked	sequenced colonies with *Saccharomyces*-like ascus morphology	*Saccharomyces* isolates
enrichment	284	246 (87%)	211 (74%)
plating	969	344 (35%)	307 (32%)

Both methods isolated *Saccharomyces* colonies from both substrates, most

trees, and all timepoints (Figure 2). We had significantly more sampling success on

soil than leaf litter substrates (z = 5.7, p < .001, Table 3), but other relationships

among sampling success, sampling method, and sampling environments were

idiosyncratic. For example, direct plating did not produce any *Saccharomyces*

isolates from tree 6, while three enrichment samples from this tree isolated *S.*

paradoxus, and enrichments produced more *Saccharomyces* isolates in March than

direct plating did (Figure 2). Because our sampling effort was not the same for all

trees at all months, we did not model tree habitat or sampling month as fixed effects;

instead, we modeled these parameters as random effects, and found that models

including tree and month fit the data better than models without tree and month (Table

2). A list of sequenced yeasts from each sample is included in the supplemental

materials (Supplemental File 2).

Phenotypes of sampled *S. paradoxus*

282   Variances in growth rate, but not median growth rates, were different between

283 strains isolated using the two methods (Figure 3). Growth rates of *S. paradoxus*

284 isolated using enrichment culturing had a larger variance than growth rates of *S.*

285 *paradoxus* isolated using direct plating (Levine's test $F_{1,108}$ = 5.42, p = .02). Median

286 growth rates for the two groups of *S. paradoxus* strains did not differ (Wilcoxon

287 signed rank test V = 6320, p = .39). Measured growth rates of each strain are included

288 in the supplemental materials (Supplemental File 2).

Figure 3: Histograms representing distributions of growth rates for *S. paradoxus* clones isolated using A) enrichment culturing or B) direct plating.

290 **<u>Discussion</u>**

291 <u>Direct plating detects *S. paradoxus* more frequently than enrichment culturing</u>

Enrichment culturing did not increase *Saccharomyces* sampling success from

forest leaf litter and soil over direct plating, in spite of researchers' long history of

using enrichment culturing to isolate *Saccharomyces* from forest environments

(Kowallik & Greig, 2016; Naumov, Naumova, & Sniegowski, 1998; Sniegowski et

al., 2002). We expect reliable *Saccharomyces* isolation from this forest using direct

plating to be a result of high *S. paradoxus* abundance on forest floor substrates.

Indeed, it is common to find hundreds to tens of thousands of *S. paradoxus* cells per

gram of leaf litter near the bases of oak trees in this forest (Kowallik & Greig, 2016).

We expect direct plating to be less successful in environments in which

Saccharomyces are rarer, and note that enrichment culturing is frequently used to

isolate *Saccharomyces* from tree bark, which may be a habitat with lower

Saccharomyces density than the forest floor habitats we sampled (Kowallik et al.,

2015; Sniegowski et al., 2002).

<u>Isolation using enrichment culturing samples more phenotypic diversity than direct</u>

<u>plating</u>

Conditions in enrichment cultures resulted in isolating a different (albeit

higher) phenotypic diversity than random colony selection on plate cultures did

(Figure 3). There are several potential methodological and ecological explanations for

the high phenotypic diversity in enrichment isolates. But we expect that interactions

with microbes *during the enrichment culturing* are most likely to be responsible for

the increased phenotypic diversity among enrichment-sourced *S. paradoxus* isolates.

Microbes that potentially interact with *S. paradoxus* were doubtless present in

our enrichment cultures, and it is realistic to expect these microbes to influence the

phenotypes of the *S. paradoxus* that were ultimately recovered. Our enrichment

cultures contained all of the microbes that were present in the soil and leaf litter

samples, and soil and leaf litter include a wide diversity of bacterial and fungal taxa

that can interact with fungi (Curd, Martiny, Li, & Smith, 2018; Glassman et al., 2018;

Santonja et al., 2018). For example, some agricultural soils contain *Bacillus* and

Pseudomonas species that secrete compounds toxic to phytopathogenic fungi (Islam,

Jeong, Lee, & Song, 2012; Petatán-Sagahón et al., 2011). Conversely, some soil

bacteria promote mycorrhizal fungal growth and host plant colonization (Xie et al.,

2018).

We expect a variety of similar inhibition and facilitation interactions to

determine the identities of the *Saccharomyces* strains that ultimately reached high

frequencies in enrichment cultures. Bacteria and fungi co-occurring with *S. paradoxus*

on oak bark can both inhibit and facilitate *S. paradoxus* growth (Kowallik et al.,

2015). These effects are dependent on temperature and interacting microbe identities,

and likely also depend on other environmental conditions and *S. paradoxus* genotype.

The *Saccharomyces* strains recovered at the end of our enrichment cultures were

probably the fittest *Saccharomyces* strains present in the cultures, but this high

relative fitness was as likely to be a result of interactions with co-occurring microbes

that occur by chance in the same enrichment cultures as it was to be the result of

intrinsic growth rate. Microbial diversity during enrichment may similarly explain our

idiosyncratic sampling success across months and trees (Figure 2). For example, it is

possible that a bacterium that facilitates rare *S. paradoxus* growth in the enrichment

medium was more common in spring than summer months, resulting in higher

enrichment sampling success in spring than summer. Unfortunately, we did not

measure the microbial diversity of the enrichment cultures and do not know which

microbes may have interacted with *Saccharomyces*.

It is also possible, although in our opinion less likely, that methodological

biases resulted in relatively low phenotypic diversity among directly plated *S.*

paradoxus compared to enrichment *S. paradoxus*. While we aimed to randomly select

colonies with *Saccharomyces*-like morphologies from direct culture plates, biases in

colony picking could have selected for low variance in *S. paradoxus* phenotypes. For

example, if a *S. paradoxus* colony had an unusual morphology, it might have been

mistaken for a bacterial colony and not isolated. However, direct culturing isolated

more non-*Saccharomyces* yeast isolates than enrichment culturing did, and we expect

interspecific morphological variation to be higher than intraspecific morphological

variation. We therefore consider it unlikely that biases in colony picking decreased

phenotypic variance among *S. paradoxus* isolates.

We continue to expect plate culturing to be a more random sampling of

Saccharomyces diversity than enrichment culturing. Colonies are physically isolated

on a plate and unlikely to influence each other's growth, and plated colonies come

from a well-mixed mixture of environmental substrate and sterile water. We therefore

expect *Saccharomyces* colonies on plates to accurately reflect the diversity of

Saccharomyces strains present in nature, as long as all strains are able to grow on the

selective media chosen. Biases in sample *success* can still be introduced to direct

plating samples by the presence of morphologically similar yeast species. On petri

dishes, a high density of non-*Saccharomyces* yeasts with morphologies similar to

Saccharomyces could have prevented us from detecting *Saccharomyces*. This issue

can be mitigated by designing a more selective isolation medium, picking more

colonies per plate (we picked up to 6 or 12 colonies across two plates), or replica

plating selective plates to media containing a color indicator for the target yeast (*e.g.,*

Wallerstein media) (Hall, 1971).

<u>Recommendations for future yeast sampling</u>

Researchers should consider both resources available for sampling and study

goals when choosing a *Saccharomyces* field sampling strategy. Our results identified

a tradeoff between resources spent on sampling and resources spent on sequencing:

enrichment culturing was less successful than direct plating at finding *Saccharomyces*

per sample collected, but more successful per ITS region sequenced (Figure 2, Table

4). Researchers with a few precious samples are therefore better off isolating

Saccharomyces using direct plating than enrichments, especially if *Saccharomyces* is

common on their substrates. Conversely, if samples are easy to get but funds available

for sequencing are limited, researchers may prefer to use enrichment culturing.

Researchers with limited time or freezer space who would like assurances that most

picked colonies are *Saccharomyces* may also prefer enrichment culturing.

However, the scientific question to be answered by environmental

Saccharomyces samples may be more important than sampling and sequencing

resource limitations, especially as sequencing becomes cheaper. Direct plating

samples a more random collection of *Saccharomyces* cells from environmental

samples than enrichment culturing, and we recommend that researchers who need

random environmental samples to answer ecological and evolutionary questions rely

on direct plating. Researchers targeting phenotypic diversity, especially for applied

yeast biology (*e.g.,* food microbiology, drug discovery) may uncover more diversity

by isolating environmental *Saccharomyces* using enrichment cultures. Researchers

interested in detecting rare *Saccharomyces* species in an environment (*e.g., S.*

cerevisiae from our study forest, *S. mikatae* and *S. eubayanus* from European forests)

(Alsammar et al., 2018) may also have more success using enrichment culturing or a

combination of enrichment and direct plating strategies.

Conclusions

Isolation protocols do indeed influence characteristics of isolated microbes. As

researchers continue to develop *Saccharomyces* yeasts as model organisms for

ecology and evolution, they must also consider how isolation history of environmental

strains can influence the ecological stories the strains can tell. The results of this study

highlight the need for consistent sampling within studies, and are a warning to

researchers comparing phenotypes (and perhaps also genotypes) among

Saccharomyces strains from different sources. Much of what we know about

Saccharomyces evolution has come from culture collections (Strope et al., 2015;

Warringer et al., 2011)—these collections are invaluable, as are the data and

conclusions they enable, but sampling information about deposited strains is not

always available. As technology improves and both genome sequencing and

phenotype assays become cheaper and more accessible (Porter & Hajibabaei, 2018;

Stewart et al., 2018; van Dijk, Auger, Jaszczyszyn, & Thermes, 2014; Zackrisson et

al., 2016), environmental sampling may emerge as a limiting step to studying

Saccharomyces ecology and evolution. Researchers should carefully consider the

consequences of their chosen sampling strategies as they conduct this exciting

microbiological research.

Acknowledgements

We would like to thank Danielle Stevens and Tjorben Nawroth for help in the field,

Jenna Gallie and the Gallie lab for help with growth curves, and Amine Hassani for

helpful conversations on microbial diversity in enrichment cultures. Thank you to

Christoph Freiherr von Fürstenberg-Plessen for permission to work in the Nehmten

forest. This work was supported through a Max Planck Fellowship to Eva H.

Stukenbrock.

Data Accessibility

All data for this project are included in Supplementary File 2.

Figure Legends

Figure 1: Schematic illustration of sampling strategies used to isolate *Saccharomyces*

for this project. A) Enrichment culturing B) Direct plating. Photo: Doreen

Landermann.

Figure 2: Percentages of samples in which *Saccharomyces* could be detected using A)

enrichment cultures or B) direct plating. Bars represent all samples for each category

of sampling, and colors represent *Saccharomyces* species. *S. paradoxus* and *S.*

cerevisiae were the only detected *Saccharomyces* species.

Figure 3: Histograms representing distributions of growth rates for *S. paradoxus*

clones isolated using A) enrichment culturing or B) direct plating.

Citations

Alsammar, H. F., Naseeb, S., Brancia, L. B., Gilman, R. T., Wang, P., & Delneri, D.
(2018). Targeted metagenomics approach to capture the biodiversity of
Saccharomyces genus in wild environments. *Environmental Microbiology*
Reports. doi:10.1111/1758-2229.12724
Bates, D., Machler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting Linear Mixed-
Effects Models Using lme4. *Journal of Statistical Software, 67*(1), 1-48.

Beijernick, M. W. (1961). Enrichment culture studies with urea bacteria (T. D. Brock,
Trans.). In T. D. Brock (Ed.), *Milestones in Microbiology*. London, United
Kingdom: Prentice-Hall International.
Bernardes, J. P., Stelkens, R. B., & Greig, D. (2017). Heterosis in hybrids within and
between yeast species. *Journal of Evolutionary Biology, 30*(3), 538-548.
doi:10.1111/jeb.13023
Boynton, P. J., & Greig, D. (2014). The ecology and evolution of non-domesticated
Saccharomyces species. *Yeast, 31*(12), 449-462. doi:10.1002/yea.3040
Boynton, P. J., & Greig, D. (2016). Species richness influences wine ecosystem
function through a dominant species. *Fungal Ecology, 22*, 61-72.
doi:10.1016/j.funeco.2016.04.008
Carbonetto, B., Ramsayer, J., Nidelet, T., Legrand, J., & Sicard, D. (2018). Bakery
yeasts, a new model for studies in ecology and evolution. *Yeast, 35*(11), 591-
603. doi:10.1002/yea.3350
Charron, G., Leducq, J. B., Bertin, C., Dube, A. K., & Landry, C. R. (2014).
Exploring the northern limit of the distribution of *Saccharomyces cerevisiae*
and *Saccharomyces paradoxus* in North America. *FEMS Yeast Research,*
14(2), 281-288. doi:10.1111/1567-1364.12100
Curd, E. E., Martiny, J. B. H., Li, H. Y., & Smith, T. B. (2018). Bacterial diversity is
positively correlated with soil heterogeneity. *Ecosphere, 9*(1), 16.
doi:10.1002/ecs2.2079
Domizio, P., Romani, C., Comitini, F., Gobbi, M., Lencioni, L., Mannazzu, I., &
Ciani, M. (2011). Potential spoilage non-Saccharomyces yeasts in mixed
cultures with *Saccharomyces cerevisiae*. *Annals of Microbiology, 61*(1), 137-
144. doi:10.1007/s13213-010-0125-1
Estrada-Godina, A. R., Cruz-Guerrero, A. E., Lappe, P., Ulloa, M., Garcia-Garibay,
M., & Gomez-Ruiz, L. (2001). Isolation and identification of killer yeasts
from Agave sap (aguamiel) and pulque. *World Journal of Microbiology &*
Biotechnology, 17(6), 557-560. doi:10.1023/a:1012210106203
Ezeronye, O. U., & Okerentugba, P. O. (2001). Genetic and physiological variants of
yeast selected from palm wine. *Mycopathologia, 152*(2), 85-89.
doi:10.1023/a:1012323721012
Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., . . .
Verstrepen, K. J. (2016). Domestication and Divergence of *Saccharomyces*
cerevisiae Beer Yeasts. *Cell, 166*(6), 1397-+. doi:10.1016/j.cell.2016.08.020
Glassman, S. I., Weihe, C., Li, J. H., Albright, M. B. N., Looby, C. I., Martiny, A. C.,
. . . Martiny, J. B. H. (2018). Decomposition responses to climate depend on
microbial community composition. *Proceedings of the National Academy of*
Sciences of the United States of America, 115(47), 11994-11999.
doi:10.1073/pnas.1811269115
Greenwalt, C. J., Steinkraus, K. H., & Ledford, R. A. (2000). Kombucha, the
fermented tea: Microbiology, composition, and claimed health effects. *Journal*
of Food Protection, 63(7), 976-981. doi:10.4315/0362-028x-63.7.976
Gschaedler, A. (2017). Contribution of non-conventional yeasts in alcoholic
beverages. *Current Opinion in Food Science, 13*, 73-77.
doi:10.1016/j.cofs.2017.02.004
Hall, J. F. (1971). Detection of wild yeasts in the brewery. *Journal of the Institute of*
Brewing, 77(6), 513-516. doi:10.1002/j.2050-0416.1971.tb03413.x
Islam, M. R., Jeong, Y. T., Lee, Y. S., & Song, C. H. (2012). Isolation and
Identification of Antifungal Compounds from *Bacillus subtilis* C9 Inhibiting

494  the Growth of Plant Pathogenic Fungi. *Mycobiology, 40*(1), 59-66.
495  doi:10.5941/myco.2012.40.1.059
496 Jeong, S. H., Jung, J. Y., Lee, S. H., Jin, H. M., & Jeon, C. O. (2013). Microbial
497  succession and metabolite changes during fermentation of dongchimi,
498  traditional Korean watery kimchi. *International Journal of Food*
499  *Microbiology, 164*(1), 46-53. doi:10.1016/j.ijfoodmicro.2013.03.016
500 Johnson, L. J., Koufopanou, V., Goddard, M. R., Hetherington, R., Schafer, S. M., &
501  Burt, A. (2004). Population genetics of the wild yeast *Saccharomyces*
502  *paradoxus*. *Genetics, 166*(1), 43-52. doi:10.1534/genetics.166.1.43
503 Kowallik, V. (2015). *The natural ecology of Saccharomyces yeasts.* (Dr. rer. nat.),
504  Christian Albrechts University of Kiel,
505 Kowallik, V., & Greig, D. (2016). A systematic forest survey showing an association
506  of *Saccharomyces paradoxus* with oak leaf litter. *Environmental Microbiology*
507  *Reports, 8*(5), 833-841. doi:10.1111/1758-2229.12446
508 Kowallik, V., Miller, E., & Greig, D. (2015). The interaction of *Saccharomyces*
509  *paradoxus* with its natural competitors on oak bark. *Molecular Ecology, 24*(7),
510  1596-1610. doi:10.1111/mec.13120
511 Leducq, J. B., Charron, G., Samani, P., Dube, A. K., Sylvester, K., James, B., . . .
512  Landry, C. R. (2014). Local climatic adaptation in a widespread
513  microorganism. *Proceedings of the Royal Society B-Biological Sciences,*
514  *281*(1777), 9. doi:10.1098/rspb.2013.2472
515 Levene, H. (1960). Robust tests for equality of variances. In I. Olkin & H. Hotelling
516  (Eds.), *Contributions to Probability and Statistics: Essays in Honor of Harold*
517  *Hotelling* (pp. 278-292): Stanford University Press.
518 Libkind, D., Hittinger, C. T., Valerio, E., Goncalves, C., Dover, J., Johnston, M., . . .
519  Sampaio, J. P. (2011). Microbe domestication and the identification of the
520  wild genetic stock of lager-brewing yeast. *Proceedings of the National*
521  *Academy of Sciences of the United States of America, 108*(35), 14539-14544.
522  doi:10.1073/pnas.1105430108
523 Liti, G., Carter, D. M., Moses, A. M., Warringer, J., Parts, L., James, S. A., . . . Louis,
524  E. J. (2009). Population genomics of domestic and wild yeasts. *Nature,*
525  *458*(7236), 337-341. doi:10.1038/nature07743
526 Liti, G., Warringer, J., & Blomberg, A. (2017). Isolation and Laboratory
527  Domestication of Natural Yeast Strains. *Cold Spring Harb Protoc, 2017*(8),
528  pdb.prot089052. doi:10.1101/pdb.prot089052
529 Mortimer, R., & Polsinelli, M. (1999). On the origins of wine yeast. *Research in*
530  *Microbiology, 150*(3), 199-204. doi:10.1016/s0923-2508(99)80036-9
531 Naumov, G. I., Naumova, E. S., & Sniegowski, P. D. (1998). *Saccharomyces*
532  *paradoxus* and *Saccharomyces cerevisiae* are associated with exudates of
533  North American oaks. *Canadian Journal of Microbiology, 44*(11), 1045-1050.
534  doi:10.1139/cjm-44-11-1045
535 Petatán-Sagahón, I., Anducho-Reyes, M. A., Silva-Rojas, H. V., Arana-Cuenca, A.,
536  Tellez-Jurado, A., Cárdenas-Álvarez, I. O., & Mercado-Flores, Y. (2011).
537  Isolation of Bacteria with Antifungal Activity against the Phytopathogenic
538  Fungi *Stenocarpella maydis* and *Stenocarpella macrospora*. *International*
539  *Journal of Molecular Sciences, 12*(9), 5522-5537. doi:10.3390/ijms12095522
540 Peter, J., De Chiara, M., Friedrich, A., Yue, J. X., Pflieger, D., Bergstrom, A., . . .
541  Schacherer, J. (2018). Genome evolution across 1,011 *Saccharomyces*
542  *cerevisiae* isolates. *Nature, 556*(7701), 339-+. doi:10.1038/s41586-018-0030-
543  5

Porter, T. M., & Hajibabaei, M. (2018). Scaling up: A guide to high-throughput
genomic approaches for biodiversity analysis. *Molecular Ecology, 27*(2), 313-
338. doi:10.1111/mec.14478
R Development Core Team. (2016). R: A language and Environment for Statistical
Computing. Vienna, Austria: R Foundation for Statistical Computing.
Rehner, S. A., & Buckley, E. (2005). A *Beauveria* phylogeny inferred from nuclear
ITS and EF1-alpha sequences: evidence for cryptic diversification and links to
Cordyceps teleomorphs. Mycologia, 97(1), 84-98.
doi:10.3852/mycologia.97.1.84
Robinson, H. A., Pinharanda, A., & Bensasson, D. (2016). Summer temperature can
predict the distribution of wild yeast populations. *Ecology and Evolution, 6*(4),
1236-1250. doi:10.1002/ece3.1919
Sampaio, J. P., & Goncalves, P. (2008). Natural populations of *Saccharomyces*
kudriavzevii in Portugal are associated with oak bark and are sympatric with *S.*
cerevisiae and *S. paradoxus. Applied and Environmental Microbiology, 74*(7),
2144-2152. doi:10.1128/aem.02396-07
Santonja, M., Foucault, Q., Rancon, A., Gauquelin, T., Fernandez, C., Baldy, V., &
Mirleau, P. (2018). Contrasting responses of bacterial and fungal communities
to plant litter diversity in a Mediterranean oak forest. *Soil Biology &*
Biochemistry, 125, 27-36. doi:10.1016/j.soilbio.2018.06.020
Schlegel, H. G., & Jannasch, H. W. (1967). Enrichment cultures. *Annual Reviews*
Microbiology, 21, 49-70. doi:10.1146/annurev.mi.21.100167.000405
Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C.
A., . . . Fungal Barcoding, C. (2012). Nuclear ribosomal internal transcribed
spacer (ITS) region as a universal DNA barcode marker for Fungi.
Proceedings of the National Academy of Sciences of the United States of
America, 109(16), 6241-6246. doi:10.1073/pnas.1117018109
Sniegowski, P. D., Dombrowski, P. G., & Fingerman, E. (2002). *Saccharomyces*
cerevisiae and *Saccharomyces paradoxus* coexist in a natural woodland site in
North America and display different levels of reproductive isolation from
European conspecifics. *FEMS Yeast Research, 1*(4), 299-306.
doi:10.1111/j.1567-1364.2002.tb00048.x
Stefani, F. O. P., Bell, T. H., Marchand, C., de la Providencia, I. E., El Yassimi, A.,
St-Arnaud, M., & Hijri, M. (2015). Culture-Dependent and -Independent
Methods Capture Different Microbial Community Fractions in Hydrocarbon-
Contaminated Soils. *PLoS One, 10*(6), 16. doi:10.1371/journal.pone.0128272
Stefanini, I., Dapporto, L., Legras, J. L., Calabretta, A., Di Paola, M., De Filippo, C., .
. . Cavalieri, D. (2012). Role of social wasps in *Saccharomyces cerevisiae*
ecology and evolution. *Proceedings of the National Academy of Sciences of*
the United States of America, 109(33), 13398-13403.
doi:10.1073/pnas.1208362109
Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., Miller, E. L., & Greig, D. (2014).
The effect of hybrid transgression on environmental tolerance in experimental
yeast crosses. *Journal of Evolutionary Biology, 27*(11), 2507-2519.
doi:10.1111/jeb.12494
Stewart, E. L., Croll, D., Lendenmann, M. H., Sanchez-Vallet, A., Hartmann, F. E.,
Palma-Guerrero, J., . . . McDonald, B. A. (2018). Quantitative trait locus
mapping reveals complex genetic architecture of quantitative virulence in the
wheat pathogen *Zymoseptoria tritici. Molecular Plant Pathology, 19*(1), 201-
216. doi:10.1111/mpp.12515

Strope, P. K., Skelly, D. A., Kozmin, S. G., Mahadevan, G., Stone, E. A., Magwene,
P. M., . . . McCusker, J. H. (2015). The 100-genomes strains, an *S. cerevisiae*
resource that illuminates its natural phenotypic and genotypic variation and
emergence as an opportunistic pathogen. *Genome Research, 25*(5), 762-774.
doi:10.1101/gr.185538.114
Sweeney, J. Y., Kuehne, H. A., & Sniegowski, P. D. (2004). Sympatric natural
Saccharomyces cerevisiae and *S. paradoxus* populations have different
thermal growth profiles. *FEMS Yeast Research, 4*(4-5), 521-525.
doi:10.1016/s1567-1356(03)00171-5
van Dijk, E. L., Auger, H., Jaszczyszyn, Y., & Thermes, C. (2014). Ten years of next-
generation sequencing technology. *Trends in Genetics, 30*(9), 418-426.
doi:10.1016/j.tig.2014.07.001
Warringer, J., Zorgo, E., Cubillos, F. A., Zia, A., Gjuvsland, A., Simpson, J. T., . . .
Blomberg, A. (2011). Trait Variation in Yeast Is Defined by Population
History. *Plos Genetics, 7*(6), 30. doi:10.1371/journal.pgen.1002111
White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and Direct
Sequencing fo Fungal Ribosomal RNA Genes for Phylogenetics. In M. A.
Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR Protocols: a*
Guide to Methods and Applications (pp. 315-322): Academic Press.
Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*: Springer.
Xie, L., Lehvavirta, S., Timonen, S., Kasurinen, J., Niemikapee, J., & Valkonen, J. P.
T. (2018). Species-specific synergistic effects of two plant growth-Promoting
microbes on green roof plant biomass and photosynthetic efficiency. *PLoS*
One, 13(12), 16. doi:10.1371/journal.pone.0209432
Zackrisson, M., Hallin, J., Ottosson, L. G., Dahl, P., Fernandez-Parada, E.,
Landstrom, E., . . . Blomberg, A. (2016). Scan-o-matic: High-Resolution
Microbial Phenomics at a Massive Scale. *G3-Genes Genomes Genetics, 6*(9),
3003-3014. doi:10.1534/g3.116.032342
Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for
aligning DNA sequences. *Journal of Computational Biology, 7*(1-2), 203-214.
doi:10.1089/10665270050081478
625