**Anosognosia for Hemiplegia as a tripartite disconnection syndrome**

Pacella V. [a,b*], Foulon C. [c,d,e], Jenkinson P.M. [f], Scandola M. [b], Bertagnoli S. [b], Avesani R. [g], Fotopoulou A. [h,§], Moro V. [b,§] & Thiebaut De Schotten M. [c,d,i,§*]

[a] *Social and Cognitive Neuroscience Laboratory, Department of Psychology, Sapienza University of Rome, Rome, Italy.*

[b] *NPSY.Lab-VR, Department of Human Sciences, University of Verona, Italy.*

[c] *Brain Connectivity and Behaviour Laboratory, Sorbonne Universities, Paris, France.*

[d] *Frontlab, Institut du Cerveau et de la Moelle épinière (ICM), UPMC UMRS 1127, Inserm U 1127, CNRS UMR 7225, Paris, France.*

[e] *Computational Neuroimaging Laboratory, Department of Diagnostic Medicine, The University of Texas at Austin Dell Medical School*

[f] *School of Life and Medical Sciences, University of Hertfordshire, UK.*

[g] *Department of Rehabilitation, Sacro Cuore-Don Calabria Hospital, Negrar, Verona, Italy.*

[h] *Clinical, Educational and Health Psychology, Division of Psychology and Language Sciences, University College London, UK.*

[i] *Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives-UMR 5293, CNRS, CEA University of Bordeaux, Bordeaux, France.*

[§]*Equal contribution.*

[*]*Corresponding Author: valentina.pacella.90@gmail.com & michel.thiebaut@gmail.com*

**Abstract**

The rare syndrome of Anosognosia for Hemiplegia (AHP) can provide unique insights into

the neurocognitive processes of motor awareness. Yet, prior studies have only explored

predominately discreet lesions. Using advanced structural neuroimaging methods in 174 patients with

a right-hemisphere stroke, we were able to identify three neural networks that contribute to AHP,

when disconnected: the (1) premotor loop (2) limbic system, and (3) ventral attention network. Our

results suggest that human motor awareness is contingent on the joint contribution of these three

systems.

Motor awareness allows individuals to have insight into their motor performance, a
fundamental aspect of self-awareness. However, following damage to the right hemisphere, patients
with left paralysis may show delusions of intact motor ability, or anosognosia for hemiplegia (AHP,
1). Hence, studying AHP offers unique opportunities to explore the neurocognitive mechanisms of
motor awareness.

While early studies regarded AHP as secondary to concomitant spatial deficits such as hemineglect
2 caused by parietal lesions, more recent experimental and voxel-based, lesion-symptom mapping
(VLSM) results suggest that AHP is an independent syndrome. These earlier studies address AHP as
an impairment of action and body monitoring, with lesions to the lateral premotor cortex and the
anterior insula (3,4), affecting patients' ability to detect discrepancies between feed-forward motor
predictions and sensorimotor feedback. However, these hypotheses are insufficient to explain all the
AHP symptoms, such as patients' inability to update their beliefs based on social feedback or more
general difficulties experienced in their daily living (5,6). Indeed, others have suggested that AHP
can be caused by a functional disconnection between regions processing top-down beliefs about the
self and those processing bottom-up errors regarding the current state of the body (5,7). Nevertheless,
to date the brain disconnection hypothesis could not be explored due to the relatively small sample
size and the standard methodology of previous studies, which favours the implication of discreet
lesion locations in the pathogenesis of AHP.

Here, to overcome this gap, we took advantage of (1) the largest cohort of AHP patients to date (N =
174; 95 AHP patients diagnosed by (8) and 79 hemiplegic controls) and (2) an advanced lesion
analysis method (BCBtoolkit, 9). This method generates a probabilistic map of disconnections from
each patient's brain lesion to identify the disconnections that are associated with given
neuropsychological deficits at the group level. Previous use of this connectivity approach has already
proven fruitful in the study of neuropsychological deficits (10–12).

We predicted that AHP would be associated not only with focal grey matter lesions, but also with
long-range disconnections due to the white matter damage, in particular to tracts associated with
sensorimotor monitoring and self-reflection. Specifically, we anticipated the possibility that motor
awareness emerges from the integrated activation of separated networks (13,14), whose contributions
feed into the multifaceted expression of the syndrome.

## Results

To test these predictions, we first conducted anatomical investigations to identify lesion sites and created probability maps of white matter tracts' disconnection. These results were statistically analysed by means of regression analyses, to identify the contribution of grey and white matter structures in AHP, taking into account differences in age, lesion size, lesion onset-assessment interval and critical motor and neuropsychological deficits (i.e. covariates of non-interest). Considering our sample size and a power of 95%, t values above 2 correspond to a medium effect size (cohen d > 0.5) and t values above 3.6 correspond to a large effect size (cohen d > 0.8).

The regression computed on the lesion sites (**Figure 1a**) indicated the involvement of grey matter structures previously associated with AHP (15), such as the insula (anterior long gyrus, t = 4.89; p = 0.002), the temporal pole (t = 4.77; p = 0.003), and the striatum (t = 4.68; p = 0.003) as well as a very large involvement of white matter (t = 4.98; p = 0.002). The second regression on white matter maps of disconnection (**Figure 1b**) revealed a significant contribution of the cingulum (t = 3.85; p = 0.008), the third branch of the superior longitudinal fasciculus (SLF III; t = 4.30; p = 0.003), and

connections to the pre-supplementary motor area (preSMA; t = 3.37; p = 0.013), such as the frontal aslant and the fronto-striatal connections.

**Figure 1**: On the top half, statistical mapping of the lesioned areas in AHP. a) right hemisphere b) striatum c) insula d) axial sections. Pal: pallidum; put: putamen; ALg: anterior long gyrus; PSg: posterior short gyrus; MSg: middle short gyrus; Tp: temporal pole. On the bottom half, statistical mapping of the brain disconnections in AHP. e) right hemisphere lateral view; f) right hemisphere medial view; g) axial sections. TPJ: temporo-parietal junction; VPF: ventral prefrontal cortex; preSMA: pre-supplementary area; H:

hippocampus; Cing: cingulum; SLF III: third (ventral) branch of the superior longitudinal fasciculus;

PreSMA: pre-supplementary motor area.

To test whether AHP emerges from the disconnection of each of these networks independently or together as a whole, we investigated the contribution pattern of the tracts' disconnection to AHP (individual or integrated), by means of Bayesian computation of generalised linear multilevel models. All of the possible binomial models (n=95) were computed, starting from the null model (with only the covariates of non-interest) to the full model, with all the covariates of non-interest, the tracts, and all the interactions among them (16) (see Methods section). The results confirmed that the disconnection of each tract is critical to AHP (Cingulum, $BF_{10}$ = 270.98; FST, $BF_{10}$ = 180.48; FAT, $BF_{10}$ = 367.61; SLF III, $BF_{10}$ = 571.49). However, results indicate that the model that best fits with our data (99% of probability) includes the contribution of all the four tracts (**Figure 2**).

**Figure 2**: Motor awareness network a) right hemisphere medial view; b) right hemisphere lateral view c) Bayes Factors for the four models, each one representing the hypothesis that the presence of disconnection in a tract is necessary to explain AHP, against the null model (i.e. no disconnection is necessary to explain the presence of AHP). The dashed line represents $BF_{10}$ = 150, the boundary suggested by (17) as the decision criteria for very strong support of hypotheses. The plot shows that each tract contributes to AHP symptoms. d) proportion of AHP over non-AHP patients, normalised to its maximum (20). Darker the line, the greater the presence of AHP compared to non-AHP. The disconnections of tracts are divided into four categories: 25% - (0, 25]; 50%: (25, 50]; 75% - (50, 75] and 100% - (75, 100]. In each subplot, the y-axis shows the percentage of FAT disconnection and the x-axis shows the percentage of Cingulum disconnection. The subplots are divided horizontally by the percentage of SLF III disconnection, and vertically by

the percentage of FST disconnection. In the all squares, binomial tests were computed to check if the
number between AHP and non-AHP is different. The only significant difference (p=.01, the number
of AHP is greater) is when the probability of disconnection for each tract is: SLFIII = 100%, FST =
100%, Cingulum = 100% and FAT = 100% (the black square on the extreme top-right).

These results, derived from the largest lesion mapping study on AHP to date, show that white matter
disconnections in three networks contribute to AHP: (1) posterior parts of the limbic network (i.e.
connections between the amygdala, the hippocampus and the cingulate gyrus); (2) the ventral
attentional network (i.e. connections between temporo-parietal junction and ventral frontal cortex),
through the SLF III; and (3) the premotor loop (i.e. connections between the striatum, the preSMA
and the inferior frontal gyrus).

## Discussion

Previous lesion mapping studies in AHP have highlighted the role of discrete cortical lesions in areas such as the lateral premotor cortex or the insula, and suggested corresponding theories of motor and body awareness (3,4 and 5,6, for a critical review). By contrast, our results suggest that AHP is a tripartite disconnection syndrome involving disruptions in networks that include, but also extend beyond, sensorimotor circuits. Correspondingly, motor awareness should be regarded as the collaborative effort (i.e. integration of a number of cognitive processes, rather than a purely motor monitoring function. Indeed, this interpretation is consistent with the delusional features of AHP (see 18) and a variety of experimental findings in AHP, such as the fact that awareness can be influenced by mood (19,20), or perspective-taking (21,22).

Specifically, the cingulum connects limbic system structures that have been previously associated with emotional and memory processing, and is part of the default mode network (23)–a pattern of intrinsic connectivity observed during self-referential, introspective states, including autobiographical retrieval, future imaging and mentalisation. These abilities relate to well-documented deficits in AHP patients' general awareness ("why are you in hospital?"), anticipatory awareness ("are you able to reach the table with your left hand?", 24) and mentalisation (25; "the doctors think there is some paralysis, do you agree?", 26).

The ventral attentional network (i.e. SLF III connections between temporo-parietal junction and ventral frontal cortex, as well as lesions in the insula and temporal pole) reorients attention towards relevant stimuli (27). This disconnection prevents the possibility to appreciate the bottom-up stimuli referring to one's own paralysis and (along with the limbic system) to update beliefs regarding the current body's condition ("I can walk as I have always done"; "I have just clapped my hands", 21). The insula is also crucial in these processes due to its important role in integrating external sensory information with internal emotional and bodily state signals (28).

Finally, the observed disconnections of the pre-motor network (pre-SMA, striatum and inferior frontal gyrus) suggest difficulties in monitoring motor signals and learning from action failures ("did you execute the action? Yes, I have").

Crucially, none of these networks (i.e. the limbic system, the ventral attentional network, and the premotor system) alone can fully explain AHP (**Figure 2c**). It thus appears that bottom up deficits in interoceptive and motor salience monitoring need to be combined with higher-order deficits, collectively leading to a multifaceted syndrome in which premorbid beliefs and emotions about the non-paralysed self dominates current cognition about the paralysed body.

These results open up interesting hypotheses on the hierarchical or parallel relations between the three

networks, in terms of temporal activations (either serial, parallel or recurrent) that remain to be

explored in future studies.

The main limitation of the study is related to manual lesion delineation and registration methods

(29,30) and the sensitivity level of neuroimaging techniques that do not depict the full extent of

damage produced by stroke lesions (31). However, these limitations mainly apply to small sample

studies, while here, the large number of patients investigated reduces these risks.

In conclusion, on the basis of a large (N = 174) and advanced lesion-mapping study, we demonstrate

a tripartite contribution of disconnections to the pre-motor network, the limbic system, and the ventral

attentional network to motor unawareness. We thus suggest that motor awareness is not limited to

sensorimotor attention and monitoring but also requires the joint contribution of higher-order

cognitive components.

**Acknowledgements**

We thank Lauren Sakuma for useful discussion and edits to the manuscript. This project has received

funding from the European Research Council (ERC) under the European Union's Horizon 2020

research and innovation programme (grant agreement No. 818521). This study was supported by a

European Research Council Starting Investigator Award (ERC-2012-STG GA313755, to A.F.);

Fondation pour la Recherche Médicale (FRM DEQ20150331725); by MIUR Italy (PRIN

20159CZFJK), and University of Verona (Bando di Ateneo per la Ricerca di Base 2015 project

MOTOS) (to V.M.).

## Materials and Methods

### 1. Design and Statistical Analysis

The aim of the study was to explore the white matter disconnections involved in AHP. To this end, we investigated the neural systems that contribute to the symptoms of AHP. To the best of our knowledge, this approach has never been applied to the study of AHP and it can shed light on the theoretical and phenomenological complexity of the disease, by integrating and going beyond existing findings gained through classic lesion studies (3,4,15,18,32,33).

For this purpose, we collected neuroimaging and clinical data from a large sample of right hemisphere stroke patients. To compute lesion sites and map of disconnections that were strictly related to the AHP pathology, we compared our target group of AHP patients with a group of stroke patients with hemiplegia but without AHP. Patients' map of lesions and disconnections were statistically compared between the two groups and adjusted for covariates of non-interest. Variance related to patients' demographic variables (age and education level) was removed. As previous lesion studies (15,32) found some differences in neuronal correlates of AHP in acute and chronic stages, the interval between lesion onset and neuropsychological assessment was considered as a covariate of non-interest as well. We also included the lesions' size of our sample, taking into account the number of voxels of each lesion as a nuisance variable.

Finally, when computing the AHP map of lesions and disconnections we controlled for the clinical (onset-assessment interval, motor deficits) and neuropsychological (personal, external neglect and memory impairments) symptoms that are often associated with AHP but are related to different patterns of disconnection.

The tracts emerging from this analysis were further analysed by means of Bayesian models to confirm the individual involvement of each tract and test their joint contribution to AHP (see details below).

### 2. Patients

Data from 195 stroke patients with unilateral right hemisphere damage were collected from two collaborating centers based in Italy and the United Kingdom over a period of 10 years.

Patients' inclusion criteria were: (i) unilateral right hemisphere damage, secondary to a first-ever stroke, as confirmed by clinical neuroimaging; (ii) severe plegia of their contralateral upper limb (AHP left arm, MRC $\leq$ 2), as clinically assessed (MRC scale). Exclusion criteria were: (i) previous history of neurological or psychiatric illness; (ii) medication with severe cognitive or mood side-effects; (iii) severe language, general cognitive impairment, or mood disturbance that precluded completion of the study assessments.

The MRI or CT neuroimaging data were available for 174 out of 195 patients. They were divided into two groups according to the presence/absence of AHP (see below for AHP assessment details), resulting in a group of 95 AHP patients and 79 non-AHP, hemiplegic control (HP) subjects. Among these, clinical and anatomical data of 40 AHP patients and 27 controls has been described in a previous study (15). Groups were balanced for demographic data (age, education, interval period between lesion and assessments) and lesion size. As the data were collected from different stroke recovery units, we took into account the neurological and neuropsychological tests that were most commonly administrated to all the patients across the different centers (**Table 1**).

All patients gave written, informed consent and the research was conducted in accordance with the guidelines of the Declaration of Helsinki (2013) and approved by the Local Ethical Committees of each center.

### 3. *Neurological and Neuropsychological Assessment*

Patients were identified as anosognosic or control according to their score in the Bisiach scale (8). This investigates the explicit form of awareness related to one's limb paralysis. During the scale administration, patients were required to verbally answer a 4-point interview about their current condition: a '0' score indicates a spared consciousness of the disease (= the disorder is spontaneously reported or mentioned by the patient following a general question about his/her complains), a '1' score is assigned when patients refer to their disability only after specific questions about the strength of their left limbs, while patients scoring '2' or '3' are considered anosognosic for their awareness of the disease emerging only after a demonstration through a routine technique of neurological examination (score 2) or not emerging at all (score 3).

Plegia of the contralesional upper limb was assessed through the Medical Research Council 5-point scale (34), ranging from 5 (normal functioning) to 0 (no movement).

Personal neglect was assessed by means of the 'Comb' test, from the 'Comb/Razor test' (35). We referred to each patient's score on the line cancellation subtest of the BIT as our measure of extra-personal neglect (Behavioral Inattention Test, 36). Finally, we used the digit/word span (37,38) to assess working memory. The 3-nearest neighbour computation replaced the missing data from the demographic and clinical variables (education: 4.9%; lesion-assessment interval: 0.05%; motricity index: 1.9%; personal neglect: 1.7%; extra-personal neglect: 2.5%; memory span: 6.2%).

In order to compare results expressed in different scoring ranges, all the scores from neuropsychological tests were transformed to z-scores, with higher scores corresponding to better performances.

*4. Lesions drawing*

Patients' neuroimaging data was acquired via Computerized Tomography (CT) and Magnetic

Resonance (MRI) and lesions were segmented and co-registered using the manual procedure already

described by Moro and colleagues (15).

The lesion drawing was performed blindly and independently by two of the authors (VM, SB), prior

(blind) to the group classification. In cases of disagreement on a lesion drawing, a third anatomist's

opinion was consulted (<10%).

Scans were registered on the ICBM152 template of the Montreal Neurological Institute, furnished

with the MRIcron software (ch2, http://www.mccauslandcenter.sc.edu/mricro/mricron/). First the

standard template was rotated on the three plans (size: 181 x 217 x 181 mm, voxel resolution: 1 mm$^2$)

in order to match the orientation of patient's MRI or CT scan. Lesions were outlined on the axial

slices of the rotated template. The resulting lesion volumes were then rotated back into the canonical

orientation, as to align the lesion volumes of each patient to the same stereotaxic space. Finally, in

order to remove voxels of lesions outside the white and grey matter brain tissue, lesion volumes were

filtered by means of custom masks based on the ICBM152 template.

*5. Disconnectome Maps*

Disconnectome maps were computed with the 'disconnectome map' tool of the BCBToolkit software

(9). The first step of the procedure is the tracking of white matter fibres passing through each patient's

lesion, by means of the registration of lesions on the diffusion weighted imaging dataset of 10 healthy

controls (39). This produces a percentage overlap map that takes into account the inter-individual

variability of tractography in healthy controls' dataset (40). Therefore, in the resulting disconnectome

maps computed for each lesion, voxels show the probability of disconnection from 0 to 100% (11).

These disconnection probabilities of each patient are then used for statistical analyses.

*6. Statistical analysis producing the sites of lesion and tract disconnection.*

We ran 2 separate regression analyses for lesion sites and tract disconnections, using the same

procedure. We used the tool "randomize" (41), part of FSL package (http://www.fmrib.ox.ac.uk/fsl/,

version 5.0), which performs nonparametric statistics on neuroimaging data. Lesion drawings or

disconnectome maps were taken into account as dependent variables within the general linear model

implemented in 'randomize', in order to test the difference between the two groups in terms of

disconnected brain regions. Demographic (age, education), clinical (lesion size, lesion onset-

assessment interval, motor deficit) and neuropsychological (personal and extrapersonal neglect and

memory impairment) data were considered in the model as covariates of non-interest. Threshold-Free

Clusters Enhancement option was applied as to boost cluster-like structures of voxels and results that
survived 5000 permutations testing were controlled for family-wise error rate (p>0.95).

*7. Comparison of disconnection results with a brain atlas*

In order to confirm the matching between white matter disconnection emerging from regression and
the anatomy of each single tract, we used an atlas of human brain connections (42).

Maps of this atlas were first thresholded at 90% (i.e. the tract position in at least 90% of healthy
population) and binarized to produce masks representing the cingulum, the frontal aslant (FAT) and
the fronto-striatal tracts (FST) as well as the third branch of the superior longitudinal fasciculus (SFL
III).

Then, these masks were used to extract the probabilities of disconnection for each tract from each
patient's disconnectome map. These probabilities were used to investigate the contribution of each
tract and their disconnection co-occurrence.

*8. Integration among tracts*

To confirm the individual involvement of each tract and test their joint contribution to AHP, statistical
analyses were conducted by using Bayesian models (R software, 43; brms package, 44) and
generalised linear multilevel models were computed (Stan, 45).

The presence of AHP (1) or its absence (0) was used as the dependent variable, while keeping
demographic, clinical and neuropsychological variables as covariates of non-interest. As covariates
of interest, we used the probability of each tract disconnection, ranged between 0 (=no lesion) to 1
(=full lesion).

Then, we fitted 95 binomial models, starting from the null model (i.e., with only the covariates of
non-interest) to the full model (i.e. with all the covariates of non-interest, the tracts, and all the
interactions among them). The posterior samples were obtained by 4 chains, with 2500 burn-in and
2500 sampling iterations, resulting in a total of 10000 iterations for each posterior sample.

As a first step we tested whether the single tracts can explain the presence of AHP better than the null
model. For this, we used the Bayes Factor ($BF_{10}$, 17). A $BF_{10}$ greater than 3 shows positive support
for the hypothesis that the tract is necessary, a $BF_{10}$ greater than 150 shows very strong support (17).

Then, all the models were compared among them by means of their marginal likelihood, showing
which is the winning model with a probability from 0 (=no representative) to 1 (=the best model).

*9. Data availability*

The raw data used for this research (lesions) as well as the dependent variable and covariates are

provided in full as supplementary data.

| | AHP | HP |
|---|---|---|
| | (N=95) | (N=79) |
| **Demographic and clinical** | | |
| Age (years) | 68.48 ± 12.54 | 63,01 ± 13.49 |
| Education (years) | 9.46 ± 3.74 | 11 ± 3.77 |
| Interval (days) | 35.74 ± 40.58 | 44.42 ± 46.7 |
| Lesion Size (voxels) | 134327.74 ± 113196.17 | 113082.73 ± 120844.22 |
| **Anosognosia** | | |
| Bisiach score | 2.46 ± 0.6 | 0 ± 0 |
| **Personal neglect** | | |
| Comb ($\frac{left-right\ strokes}{left+ambiguous+right\ strokes}$) | -0.3 ± 0.4 | -0.06 ± 0.47 |
| **Extra-personal neglect** | | |
| Line cancellation (number of items cancelled) | 19.26 ± 11.9 | 28.35 ± 10.77 |
| **Memory Span** | | |
| Digit/verbal span (number of items recalled) | 5.65 ± 2.14 | 6.83 ± 2.46 |
| **Motor index** | | |
| MRC (UUL) | 0.15 ± 0.42 | 0.6 ± 0.99 |

294 **Table 1.** For AHP and control groups, mean and (± standard deviation) of demographic and clinical
295 variables, neurological and neuropsychological assessments are reported.

**References**

1.    J. Babinski, Contribution to the study of mental disorders in organic cerebral hemiplegia
(anosognosia). *Rev. Neurol*. **27**, 845-888 (1914).

2.    E. Bisiach, "The neuropsychological approach to consciousness" in *Handbook of clinical and*
*experimental neuropsychology*, G. Denes, & L. Pizzamiglio, Eds. (Psychology Press, Hove,
UK, 1999), pp 671-685.

3.    A. Berti, G. Bottini, M. Gandola, L. Pia, N. Smania, A. Stracciari, … E. Paulesu, Shared
Cortical Anatomy for Motor Awareness and Motor Control. *Science* **309**, 488-491 (2005).
https://doi.org/10.1126/science.1110625

4.    H.-O. Karnath, B. Baier & T. Nägele, Awareness of the Functioning of One's Own Limbs
Mediated by the Insular Cortex? *J. Neurosci.* **25,** 7134–7138 (2005).
https://doi.org/10.1523/JNEUROSCI.1590-05.2005

5.    A. Fotopoulou, Time to get rid of the 'Modular' in neuropsychology: A unified theory of
anosognosia as aberrant predictive coding. *J. Neuropsychol.* **8,** 1–19 (2014).
https://doi.org/10.1111/jnp.12010

6.    P. Vuilleumier, Anosognosia: The Neurology of Beliefs and Uncertainties. *Cortex* **40,** 9–17
(2004). https://doi.org/10.1016/S0010-9452(08)70918-3

7.    D. C. Mograbi & R. G. Morris, Implicit awareness in anosognosia: Clinical observations,
experimental evidence, and theoretical implications. *Cogn. Neurosci.* **4,** 181–197 (2013).
https://doi.org/10.1080/17588928.2013.833899

8.    E. Bisiach, G. Vallar, D. Perani, C. Papagno & A. Berti, Unawareness of disease following
lesions of the right hemisphere: anosognosia for hemiplegia and anosognosia for hemianopia.
*Neuropsychologia* **24,** 471–82 (1986). https://doi.org/10.1016/0028-3932(86)90092-8

9.    C. Foulon, L. Cerliani, S. Kinkingnéhun, R. Levy, C. Rosso, M. Urbanski, … M. Thiebaut de
Schotten, Advanced lesion symptom mapping analyses and implementation as BCBtoolkit.
*Gigascience* **7,** 1–17 (2018). https://doi.org/10.1093/gigascience/giy004

10.   M. Thiebaut De Schotten, F. Tomaiuolo, M. Aiello, S. Merola, M. Silvetti, F. Lecce, … F.
Doricchi, Damage to white matter pathways in subacute and chronic spatial neglect: A group
study and 2 single-case studies with complete virtual 'in vivo' tractography dissection.
*Cereb. Cortex* **24,** 691–706 (2014). https://doi.org/10.1093/cercor/bhs351

11.   M. Thiebaut De Schotten, F. Dell'Acqua, P. Ratiu, A. Leslie, H. Howells, E. Cabanis, … M.

Catani, From phineas gage and monsieur leborgne to H.M.: Revisiting disconnection
syndromes. *Cereb. Cortex* **25,** 4812–4827 (2015). https://doi.org/10.1093/cercor/bhv173

12.   M. D. Fox, Mapping Symptoms to Brain Networks with the Human Connectome. *N. Engl. J.*
*Med.* **379,** 2237–2245 (2018). https://doi.org/10.1056/NEJMra1706158

13.   A. R. Luria, *Higher Cortical Functions in Man*. Eds. (Tavistock, London, UK, 1966).

14.   J. M. Shine, M. Breakspear, P. T. Bell, K. Ehgoetz Martens, R. Shine, O. Koyejo, … R.
Poldrack, Human cognition involves the dynamic integration of neural activity and
neuromodulatory systems. *Nat. Neurosci.* 1 (2019). https://doi.org/10.1038/s41593-018-
0312-0

15.   V. Moro, S. Pernigo, M. Tsakiris, R. Avesani, N. M. J. Edelstyn, P. M. Jenkinson, & A.
Fotopoulou, Motor versus body awareness: Voxel-based lesion analysis in anosognosia for
hemiplegia and somatoparaphrenia following right hemisphere stroke. *Cortex* **83,** 62–77
(2016). https://doi.org/10.1016/j.cortex.2016.07.001

16.   A. Gelman & J. Hill, *Data Analysis Using Regression and Multilevel/Hierarchical Models*.
(Cambridge University Press, 2006). https://doi.org/10.1017/CBO9780511790942

17.   A. E. Raftery, "Baesyan model selection in social research" in *Sociological methodology,* P.
V. Marsden, (Cambridge: Blackwells, 1995).

18.   A. Fotopoulou, S. Pernigo, R. Maeda, A. Rudd & M. A. Kopelman, Implicit awareness in
anosognosia for hemiplegia: unconscious interference without conscious re-representation.
*Brain* **133,** 3564–3577 (2010). https://doi.org/10.1093/brain/awq233

19.   S. Besharati, , S. J. Forkel, M. Kopelman, M. Solms, P. M. Jenkinson, & A. Fotopoulou, The
affective modulation of motor awareness in anosognosia for hemiplegia: Behavioural and
lesion evidence. *Cortex* **61,** 127–140 (2014). https://doi.org/10.1016/j.cortex.2014.08.016

20.   D. D'Imperio, C. Bulgarelli, S. Bertagnoli, R. Avesani & V. Moro, Modulating anosognosia
for hemiplegia: The role of dangerous actions in emergent awareness. *Cortex* **92,** 187–203
(2017). https://doi.org/10.1016/j.cortex.2017.04.009

21.   A. J. Marcel, R. Tegnér & I. Nimmo-Smith, Anosognosia for Plegia: Specificity, Extension,
Partiality and Disunity of Bodily Unawareness. *Cortex* **40,** 19–40 (2004).
https://doi.org/10.1016/S0010-9452(08)70919-5

22.   A. Fotopoulou, A. Rudd, P. Holmes & M. Kopelman, Self-observation reinstates motor
awareness in anosognosia for hemiplegia. *Neuropsychologia* **47,** 1256–1260 (2009).

https://doi.org/10.1016/j.neuropsychologia.2009.01.018

23.    M. D. Greicius, K. Supekar, V. Menon & R.F. Dougherty, Resting-State Functional

Connectivity Reflects Structural Connectivity in the Default Mode Network. *Cereb. Cortex*

**19,** 72–78 (2009). https://doi.org/10.1093/cercor/bhn059

24.    V. Moro, M. Scandola, C. Bulgarelli, R. Avesani & A. Fotopoulou, Error-based training and

emergent awareness in anosognosia for hemiplegia. *Neuropsychol. Rehabil.* **25,** 593–616

(2015). https://doi.org/10.1080/09602011.2014.951659

25.    S. Besharati, S. J. Forkel, M. Kopelman, M. Solms, P. M. Jenkinson, & A. Fotopoulou,

Mentalizing the body: spatial and social cognition in anosognosia for hemiplegia. *Brain* **139,**

971–85 (2016). https://doi.org/10.1080/09602011.2014.951659

26.    T. E. Feinberg, D. M. Roane, & J. Ali, Illusory limb movements in anosognosia for

hemiplegia. *J. Neurol. Neurosurg. Psychiatry* **68,** 511–3 (2000).

https://doi.org/10.1136/JNNP.68.4.511

27.    M. Corbetta, G. Patel & G. L. Shulman, The Reorienting System of the Human Brain: From

Environment to Theory of Mind. *Neuron* **58,** 306–324 (2008).

https://doi.org/10.1016/j.neuron.2008.04.017

28.    A. D. Craig, How do you feel — now? The anterior insula and human awareness. *Nat. Rev.*

*Neurosci.* **10,** 59–70 (2009). https://doi.org/10.1038/nrn2555

29.    B. de Haan & H.-O. Karnath, A hitchhiker's guide to lesion-behaviour mapping.

*Neuropsychologia* **115,** 5–16 (2018). https://doi.org/10.1016/j.neuropsychologia.2017.10.021

30.    C. Rorden & H.-O. Karnath, Using human brain lesions to infer function: a relic from a past

era in the fMRI age? *Nat. Rev. Neurosci.* **5,** 813–9 (2004). https://doi.org/10.1038/nrn1521

31.    A. E. Hillis, P. B. Barker, N. J. Beauchamp, B. Gordon & R. J. Wityk, MR perfusion

imaging reveals regions of hypoperfusion associated with aphasia and neglect. *Neurology* **55,**

782–8 (2000). https://doi.org/10.1212/WNL.55.6.782

32.    R. Vocat, F. Staub, T. Stroppini & P. Vuilleumier, Anosognosia for hemiplegia: a clinical-

anatomical prospective study. *Brain* **133,** 3578–3597 (2010).

https://doi.org/10.1093/brain/awq297

33.    V. Moro, S. Pernigo, P. Zapparoli, Z. Cordioli & S. M. Aglioti, Phenomenology and neural

correlates of implicit and emergent motor awareness in patients with anosognosia for

hemiplegia. *Behav. Brain Res.* **225,** 259–269 (2011).

https://doi.org/10.1016/j.bbr.2011.07.010

34.    M. B. Matthews, *Aids to the examination of the peripheral nervous system* (Her Majesty's
Stationary Office, London, 1976).

35.    R. D. McIntosh, E. E. Brodie, N. Beschin & I. H. Robertson, I. H. Improving the Clinical
Diagnosis of Personal Neglect: A Reformulated Comb and Razor Test. *Cortex* **36,** 289–292
(2000). https://doi.org/10.1016/S0010-9452(08)70530-6

36.    B. Wilson, J. Cockburn & P. Halligan, Development of a behavioral test of visuospatial
neglect. *Arch. Phys. Med. Rehabil.* **68,** 98–102 (1987).

37.    A. D. Baddeley, N. Thomson & M. Buchanan, Word length and the structure of short-term
memory. *J. Verbal Learning Verbal Behav.* **14,** 575–589 (1975).
https://doi.org/10.1016/S0022-5371(75)80045-4

38.    A. Baddeley, Exploring the Central Executive. *Q. J. Exp. Psychol. Sect. A* **49,** 5–28 (1996).
https://doi.org/10.1080/713755608

39.    M. Thiebaut de Schotten, M. Urbanski, B. Batrancourt, R. Levy, B. Dubois, L. Cerliani, & E.
Volle, Rostro-caudal Architecture of the Frontal Lobes in Humans. *Cereb. Cortex* **27,** 4033–
4047 (2016). https://doi.org/10.1093/cercor/bhw215

40.    P. L. Croxson, S. J. Forkel, L. Cerliani & M. Thiebaut de Schotten, Structural Variability
Across the Primate Brain: A Cross-Species Comparison. *Cereb. Cortex* **28,** 3829–3841
(2018). https://doi.org/10.1093/cercor/bhx244

41.    A. M. Winkler, G. R. Ridgway, M. A. Webster, S. M. Smith & T. E. Nichols, Permutation
inference for the general linear model. *Neuroimage* **92,** 381–397 (2014).
https://doi.org/10.1016/j.neuroimage.2014.01.060

42.    K. Rojkova, E. Volle, M. Urbanski, F. Humbert, F. Dell'Acqua, & M. Thiebaut de Schotten,
Atlasing the frontal lobe connections and their variability due to age and education: a
spherical deconvolution tractography study. *Brain Struct. Funct.* **221,** 1751–1766 (2016).
https://doi.org/10.1007/s00429-015-1001-3

43.    R Core Team, R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna (2018). https://www.R-project.org

44.    P.-C. Bürkner, **brms** : An *R* Package for Bayesian Multilevel Models Using *Stan*. *J. Stat.*
*Softw.* **80,** 1–28 (2017). https://doi.org/10.18637/jss.v080.i01

45.    B. Carpenter, A. Gelman, M. D. Hoffman, D. Lee, B. Goodrich, M. Betancourt, … A.

Riddell, Stan: A Probabilistic Programming Language. *J. Stat. Softw.* **76,** 1–32 (2017).

https://doi.org/10.18637/jss.v076.i01