Nonparametric expression analysis using inferential replicate counts

Anqi Zhu[1], Avi Srivastava[2], Joseph G. Ibrahim[1], Rob Patro[2], and Michael I. Love [*1,3]

[1]Department of Biostatistics, University of North Carolina-Chapel Hill
[2]Department of Computer Science, Stony Brook University
[3]Department of Genetics, University of North Carolina-Chapel Hill

February 25, 2019

Abstract

A primary challenge in the analysis of RNA-seq data is to identify differentially expressed genes or transcripts while controlling for technical biases present in the observations. Ideally, a statistical testing procedure should incorporate information about the inherent uncertainty of the abundance estimates, whether at the gene or transcript level, that arise from quantification of abundance. Most popular methods for RNA-seq differential expression analysis fit a parametric model to the counts or scaled counts for each gene or transcript, and a subset of methods can incorporate information about the uncertainty of the counts. Previous work has shown that nonparametric models for RNA-seq differential expression may in some cases have better control of the false discovery rate, and adapt well to new data types without requiring reformulation of a parametric model. Existing nonparametric models do not take into account the inferential uncertainty of the observations, leading to an inflated false discovery rate, in particular at the transcript level. Here we propose a nonparametric model for differential expression analysis using inferential replicate counts, extending the existing *SAMseq* method to account for inferential uncertainty, batch effects, and sample pairing. We compare our method, "SAMseq With Inferential Samples Helps", or *Swish*, with popular differential expression analysis methods. *Swish* has improved control of the false discovery rate, in particular for transcripts with high inferential uncertainty. We apply *Swish* to a single-cell RNA-seq dataset, assessing sensitivity to recover DE genes between sub-populations of cells, and compare its performance to the Wilcoxon rank sum test.

Introduction

Quantification uncertainty in RNA-seq arises from fragments that are consistent with expression of one or more transcripts within a gene, or fragments which map to multiple genes. Abundance estimation algorithms probabilistically assign fragments to transcripts, and may simultaneously estimate technical bias parameters, providing a point estimate of transcript abundance. Transcript abundance can be compared across samples directly, or transcript abundances can be summarized within a gene to provide gene-level comparisons across samples. However, incorporating the uncertainty of the quantification step into statistical testing of abundance differences across samples is challenging, but nevertheless critical, as the inferential uncertainty is not equal across transcripts or genes, but is higher for genes with many similar transcripts (transcript-level uncertainty) or genes that belong to large families with high level of sequence homology (gene-level uncertainty).

Methods designed for transcript-level differential expression (DE) analysis include *BitSeq* [1], *EBSeq* [2], *Cuffdiff2* [3], *IsoDE* [4], and *sleuth* [5]. *BitSeq* uses MCMC samples from the posterior distribution during abundance estimation of each sample, to build a model of technical noise when comparing abundance across samples. A parametric Bayesian model is specified with the distribution for the abundance in a single sample being conjugate, such that the posterior inference comparing across samples is exact. *EBSeq* takes into account the number of transcripts within a gene when performing dispersion estimation, while *Cuffdiff2* models biological variability across replicates through over-dispersion, as well as the

---

*michaelisaiahlove@gmail.com

uncertainty of assigning counts to transcripts within a gene, using the beta negative binomial mixture distribution. *IsoDE* and *sleuth* leverage a measurement of quantification uncertainty from bootstrapping reads when performing DE analysis. These methods therefore may offer better control of FDR for transcripts or genes with high uncertainty of the abundance. *BitSeq*, *EBSeq*, *Cuffdiff2*, and *sleuth* incorporate biological variance and inferential uncertainty into a parametric model of gene or transcript expression, while *IsoDE* focuses on comparisons of bootstrap distributions for two samples at a time, aggregating over random pairs or all pairs of samples in two groups in the case of biological replicates. Another method, *RATs* [6], provides a test for differential transcript usage (DTU). *RATs* takes inferential uncertainty into account by repeating the statistical testing procedure across inferential replicates (posterior samples or bootstrap replicates), and then offers a filter on the fraction of replicate analyses which achieved statistical significance. In this study, we focus on differential transcript expression (DTE) and differential gene expression (DGE), the latter defined as a change in the total expression of a gene across all of its transcripts.

BitSeq, *EBSeq*, *Cuffdiff2*, and *sleuth* rely on parametric models of the transcript or gene abundance, assuming underlying distributions for the data and sometimes also for the parameters in the context of a Bayesian model. The distributions are typically chosen for being robust across many experiments and for efficient inference. In a recent simulation benchmark of DTE and DGE, *EBSeq* and *sleuth* did have improved performance for detecting DTE compared to methods designed for gene-level analysis, for *EBSeq* when sample size was large ($n \geq 6$) and for *sleuth* when the sample size was low to moderate ($n \in [3, 6]$) [7], lending support for these models specialized for transcript-level inference. In this simulation benchmark, *EBSeq* and *sleuth* performed similarly to methods designed for detecting DGE, where uncertainty in abundance estimation is greatly reduced as described by Soneson et al. [8].

Previous studies have shown that nonparametric methods for differential expression analysis may be more robust compared to parametric models when the data for certain genes or transcripts deviates from distributional assumptions of the model [9]. Li and Tibshirani [9] proposed *SAMseq*, which has high sensitivity and good control of FDR at the gene-level, when the sample size is 4 per group or higher [7, 10, 11]. *SAMseq* uses a multiple re-sampling procedure to account for differences in library size across samples, and then averages a rank-based test statistic over the re-sampled datasets to produce a single score per gene. We considered whether the *SAMseq* model could be extended to account for inferential uncertainty during abundance estimation, as well as batch effects and sample pairing.

Further, we investigated whether a nonparametric method for comparison of counts, which accounts for inferential uncertainty, would provide benefit for new types of data exhibiting different distributions compared to bulk RNA-seq experiments. Single cell RNA-seq (scRNA-seq) provides researchers with high-dimensional transcriptomic profiles of single cells [12], revealing inter-cell heterogeneity and in particular sub-populations of cells that may be obscured in bulk datasets. Due to the limited amount of RNA material in a single cell and the low capture efficiency, all current protocols for scRNA-seq experiment use amplification techniques. To reduce the amplification bias, it is common to use Unique Molecular Identifiers (UMIs), short random sequences that are attached to the cDNA molecules per cell [13]. Ideally, by counting the total number of distinct UMIs per gene and per cell, one obtains a direct count of cDNA molecules, reducing bias from amplification.

Among the multiple different types of assays available for UMI-tagged scRNA-seq, droplet-based protocols have been of particular importance because of their relatively low cost, higher capture rate, robustness and accuracy in gene-count estimation. However, current dscRNA-seq (droplet-based single-cell RNA sequencing) protocols are limited in their resolution, as they only sequence the 3' end of the mRNA molecule. This complicates full transcript-level quantification, and in fact such quantification is not possible for collections of transcripts that are not disambiguated within a sequenced fragment's length of the 3' end. Moreover, since dscRNA-seq protocols typically rely on considerable amplification of genomic material, the UMI sequences designed to help identify duplicate fragments are often subject to errors during amplification and sequencing. Most dscRNA-seq quantification pipelines [14–17] thus include some method for de-duplicating similar UMIs (i.e. UMIs within some small edit distance) attached to reads that align to the same gene. These dscRNA-seq quantification pipelines [14–17] simplify the problem of gene-level count estimation by discarding the gene-ambiguous reads, which represent a considerable portion of the data (12-23% as reported by Srivastava et al. [18]). *Alevin* [18] is the first tool to propose a method to account for sequencing and amplification errors in cellular barcodes (CB) and UMIs while also taking into account gene multi-mapping reads during quantification. To resolve gene ambiguity, *Alevin* first uses the Parsimonious UMI Graph (PUG) resolution algorithm. In the case that parsimony fails to distinguish a single best gene, abundance is estimated by utilizing the proportion of uniquely-mapping reads per gene to distribute ambiguous counts by an expectation-maximization procedure. *Alevin* also can produce inferential replicates in the form of multiple matrices of estimated counts,

via bootstrapping of the counts in their PUG model. We assessed how incorporation of this inferential uncertainty using *Swish* resulted in different sets of DE genes compared to those called by a rank-based test ignoring uncertainty, in the context of calling DE genes between sub-populations of cells detected in scRNA-seq.

We describe a nonparametric method for gene- or transcript-level differential expression analysis, "SAMseq With Inferential Samples Helps", or *Swish*, that propagates quantification uncertainty from Gibbs posterior samples generated by the *Salmon* [19] method for transcript quantification. In addition to the incorporation of uncertainty in the counts, *Swish* also extends the two-group comparison in *SAMseq* by handling experiments with discrete batches or sample pairing. We show through simulation studies that the proposed method has better control of FDR and better sensitivity for DTE, compared to existing methods designed for transcript-level or gene-level analysis. For DGE, *Swish* tends to perform as well as existing methods designed for gene-level analysis, with better control of FDR than *SAMseq* for genes with high inferential uncertainty. We also compare *Swish*, leveraging estimation uncertainty, with a Wilcoxon test on gene expression in scRNA-seq sub-populations of cells in the developing mouse brain.

1 Materials and Methods

1.1 Median-Ratio Scaled Counts

In order to apply nonparametric rank-based methods to estimated counts per gene or transcript, we first remove biases on the counts due to library size differences or changes in the effective length of the gene or transcript across samples [19]. For 3' tagged datasets, where we do not expect counts to be proportional to length, we correct only for library size but not for effective length, which is not computed. *scaledTPM* counts are generated which are roughly on the scale of original estimated counts, but are adjusted to account for technical biases with respect to effective transcript or gene length [8]. We perform median-ratio normalization [20] of the *scaledTPM*, to remove any residual differences in *scaledTPM* across samples due to library size differences. The resulting scaled counts can then be directly compared across samples.

Suppose a matrix Y^0 of estimated counts from *Salmon*. The rows of the matrix represents genes or transcripts, $(g = 1, \ldots, G)$, and columns represent samples, $(i = 1, \ldots, m)$. Let Y_{gi}^0 denote the count of RNA-seq fragments assigned to gene g in sample i. The estimated counts Y_{gi}^0 are divided by a bias term accounting for the sample-specific effective length of the gene or transcript, denoted as l_{gi} [8, 21]. We first scale the l_{gi} by the geometric mean over i, giving a bias correction term b_{gi} centered around 1:

$$b_{gi} = l_{gi} / \left(\prod_{j=1}^{m} l_{gj} \right)^{\frac{1}{m}}.$$

Then we divide the b_{gi} from the estimated counts:

$$Y_{gi}^* = \frac{Y_{gi}^0}{b_{gi}}.$$

Again, for 3' tagged data, we set $l_{gi} = b_{gi} = 1$ in the above. The Y_{gi}^* are scaled to the geometric mean of sequencing depth, by

$$Y_{gi}^{**} = \frac{Y_{gi}^*}{\sum_{g=1}^{G} Y_{gi}^*} \times \left(\prod_{j=1}^{m} \sum_{g=1}^{G} Y_{gj}^0 \right)^{\frac{1}{m}}.$$

For each sample i a median-ratio size factor is computed as

$$\hat{s}_i = \text{median}_g \frac{Y_{gi}^{**}}{\left(\prod_{j=1}^{m} Y_{gj}^{**} \right)^{\frac{1}{m}}}, \tag{1}$$

where the median is taken over g where the numerator and denominator are both greater than zero. For scRNA-seq data, an alternative size factor estimate is used, as described in a later section. Lastly, the scaled count used for nonparametric methods is given by:

$$Y_{gi} = \frac{Y_{gi}^{**}}{\hat{s}_i}. \tag{2}$$

After scaling, we filter features (genes or transcripts) with low scaled counts and therefore low power: for bulk RNA-seq, we recommend keeping features with a estimated count of at least 10 across a minimal

number of samples, and here we set the minimal number of samples to 3. For UMI de-duplicated scRNA-seq data, we lowered the minimum count to 3 and require at least 5 cells, as described in a later section.

We scale all of *Salmon*'s Gibbs sampled estimated count matrices using the above procedure. For posterior sampling, *Salmon* adopts a similar approach to *MMSEQ* [22] to perform Gibbs sampling on abundance estimates, alternating between (1) Multinomial sampling of fragment allocation to equivalence classes of transcripts, and (2) conditioning on these fragment allocations, Gamma sampling of the transcript abundance, as described in further detail in the Supplementary Note 2 of Patro et al. [19]. The Gibbs samples are thinned every 16th iteration to reduce auto-correlation (*Salmon*'s default setting), and 20 thinned Gibbs samples are used as the posterior samples in the following methods.

1.2 Inferential Relative Variance

For each gene (or transcript) from each sample, S Gibbs sampled estimated counts are generated using *Salmon*. Estimated counts are then scaled and filtered as described above.

We define the *inferential relative variance* (InfRV), per gene and per sample, as

$$\text{InfRV} = \frac{\max(s^2 - \mu, 0)}{\mu},$$

where s^2 is the sample variance of scaled counts over Gibbs samples, and μ is the mean scaled count over Gibbs samples. In practice, we also add a pseudocount of 5 to the denominator and add 0.01 overall, in order to stabilize the statistic and make it strictly positive for log transformation. InfRV is therefore a measure of the uncertainty of the estimated counts, which is roughly stabilized with respect to μ. The larger the InfRV, the higher the uncertainty in quantification for that gene or transcript in that sample relative to other genes with the same mean. Note that InfRV is not used in the *Swish* statistical testing procedure, but only used in this work for visualization purpose, to categorize genes or transcripts by their inferential uncertainty, in a way that is roughly independent of the range of counts for the feature.

1.3 Test Statistic over Gibbs Samples

In *Swish*, we consider experiments with samples across two conditions of interest. We additionally allow for two or more batches to be adjusted for, or comparisons of paired data, described below. We first consider an experiment with two conditions and no batches. Let m_k denote the sample size in condition k, $k = 1, 2$, and $m_1 + m_2 = m$. Let C_k denote the collection of samples that are from condition k.

For the following, consider a single matrix Y_{gi}^s of scaled counts from one of the Gibbs samples, $(s = 1, \cdots, S)$. For gene g, the scaled counts Y_{gi}^s estimated from (2) are ranked over i. To break ties in Y_{gi}^s over the m biological samples, a small random value is draw from Uniform$(0, 0.1)$ and added before computing the ranks, as in Li and Tibshirani [9]. Let R_{gi}^s denote the rank of sample i among m biological samples for gene g. Also following Li and Tibshirani [9] we use the Mann-Whitney Wilcoxon test statistic [23] for gene g:

$$W_g^s = \sum_{i \in C_2} R_{gi}^s - \frac{m_2(m+1)}{2}. \tag{3}$$

Under the null hypothesis that gene g is not differentially expressed, W_g^s has expectation zero.

For every gene $g = 1, \cdots, G$, we calculate W_g^s for all Gibbs sampled scaled count matrices ($s = 1, \cdots, S$). While *SAMseq* computes the mean over imputed count matrices, we use the 25% quantile of the S test statistics when $\text{median}_s(W_g^s) \geq 0$,

$$T_g = Q_1\{W_g^1, \cdots, W_g^S\}, \tag{4}$$

or the 75% quantile when $\text{median}_s(W_g^s) < 0$. Finally, if $|T_g| > |\text{median}_s(W_g^s)|$, we take the median as the test statistic.

We use the 25% (or 75%) quantile of the W_g^s instead of the mean, as we found that the lower (or upper) quantile had more exposure to the quantification uncertainty than the mean; often for genes with high InfRV, the mean of W_g^s was nearly the same as the Wilcoxon test statistic of the point estimate of scaled counts, i.e. the test statistic ignoring inferential uncertainty. By using 25% (or 75%) quantile of the test statistics, we penalize those genes or transcripts with higher quantification uncertainty.

1.4 Stratified Test Statistic

In experiments with two or more batches, the above test statistic can be generalized to a stratified test statistic as described in Van Elteren [24]. We use the following to produce a summarized test statistic:

1. Stratify the samples by the nuisance covariate (e.g. batch). Suppose we have J strata. In stratum j the sample size for condition k is $m_{j,k}$ and the total sample size is m_j, i.e. $m_{j,1} + m_{j,2} = m_j$.

2. For stratum j, compute the test statistic T_g^j.

3. The final test statistic is calculated as the weighted combination of T_g^j over J strata

$$T_g^{\text{strat}} = \sum_{j=1}^{J} \omega_j T_g^j, \tag{5}$$

where $\omega_j = \dfrac{1}{m_j + 1}$ is the weight assigned to strata j.

When the difference in expression for gene g is consistent across batches, the weighted test statistic maximizes the power of test [25].

1.5 Signed-Rank Statistic for Paired Data

In experiments with paired samples, e.g. before and after treatment, we use a Wilcoxon signed-rank test statistic [23] in (4) in lieu of the Mann-Whitney Wilcoxon statistic (3). For pair index $p = 1, \ldots, m/2$:

$$W_g^s = \sum_p \text{sign}(\Delta_{gp}^s) R_{gp}^s \tag{6}$$

where Δ_{gp}^s is the difference in Y_{gi}^s for the two samples in pair p, and R_{gp}^s is the rank over pairs of $|\Delta_{gp}^s|$.

Unlike the Mann-Whitney Wilcoxon and stratified version described previously, the signed-rank statistic is not necessarily identical after transformation of the data by a monotonic function, e.g. by the square root or shifted logarithm. We choose to keep the data as scaled counts when computing the signed-rank statistic, and not to perform the square root or other transformation, so as to promote the association of higher ranks to matched samples with higher scaled counts for a given gene or transcript.

1.6 Permutation for Estimation of FDR

Under the null hypothesis that gene g is not differentially expressed, the distribution of test statistic T_g in (4)-(6) is not known. Instead, following the approach in *SAMseq* [9] we use permutation tests to construct the null distribution for T_g. In the case of stratified data, we perform permutations independently within strata and calculate a weighted test statistic as in (5). For paired data, we randomly permute condition labels within pairs. Finally we use the permutation plug-in method [26] to estimate the q-value for each gene g, leveraging functions in the R package *samr*. We provide the option to estimate π_0, the proportion of nulls, from the observed test statistics and the permutation null distribution, but by default we set $\hat{\pi}_0 = 1$. In practice, 4 samples per group is sufficient for the permutation method to have sensitivity to detect differential expression [9, 11].

1.7 Simulation

In order to evaluate *Swish* and other methods, we make use of a previously published simulated RNA-seq dataset [7], which we briefly describe here. *polyester* [27] was used to simulate paired-end RNA-seq reads across four groups of samples: two condition groups with various differential expression patterns described below, which were balanced across two batches. One batch demonstrated realistic fragment GC bias, and one batch demonstrated approximately uniform coverage. The fragment GC bias was derived via *alpine* [28] from GEUVADIS samples [29]. Likewise, the baseline transcript expression was derived from a GEUVADIS sample. Each condition group had 12 samples in total, equally split among the two batches, giving 24 samples in all. Fragments were simulated from 46,933 expressed transcripts belonging to 15,017 genes (Gencode [30] release 28 human reference transcripts).

70% of the genes were simulated with no differential expression across condition, while 10% of the genes had all of their transcripts differentially expressed with the same fold change, 10% of the genes had a single transcript differentially expressed, and 10% of the genes had shifts in expression among two of

their transcript isoforms which resulted in no change in total expression. For the purpose of gene-level differential expression, 20% of the genes were non-null due to changes in total expression. For transcript-level differential expression, all transcripts expressed in the first category were differentially expressed, as well as one transcript per gene from the second category, and two transcripts per gene from the third category.

Transcripts were quantified using *Salmon* v0.11.3 and *kallisto* [31] v0.44.0, with bias correction enabled for both methods, and generating 20 Gibbs samples for *Salmon* and 20 bootstrap samples for *kallisto*. The scripts used to generate the simulation are provided in the Code Availability section, and links to the raw reads and quantification files are listed in the Data Availability section.

1.8   Simulation Evaluation

We evaluated methods across four settings:

1. DGE analysis with one batch of samples (fragment GC bias), 6 samples in each of the two conditions,

2. DGE analysis with two balanced batches of samples, 12 samples in each of the two conditions,

3. DTE analysis with one batch of samples (fragment GC bias), 6 samples in each of the two conditions,

4. DTE analysis with two balanced batches of samples, 12 samples in each of the two conditions.

To evaluate the performance of methods on DGE and DTE, we use the iCOBRA package [32] to construct plots of the true positive rate (TPR) over the false discovery rate (FDR) at three nominal FDR thresholds: 1%, 5%, and 10%. The methods compared include *Swish*, *DESeq2* [11], *EBSeq* [2], *limma* with *voom* transformation [33], *SAMseq* [9], and *sleuth* [5]. We chose a set a methods to represent popular choices among classes of approaches. We note that *edgeR* [34] and *edgeR* with quasi-likelihood [35] performed well in a previous benchmark of this dataset [7], with both performing similarly to *limma*. *EBSeq* and *sleuth* are designed for DTE or DGE analysis taking the inferential uncertainty of quantification into account, while *SAMseq*, *DESeq2*, and *limma* are designed for gene-level analysis, but have been shown nevertheless to be able to recover DTE when supplied with transcript-level estimated counts [7]. We provided minimal filtering across all methods, removing transcript or genes without an estimated count of 10 or more across at least 3 samples. We also used the default recommended filtering functions `filterByExpr` and `sleuth_prep` for *limma* and *sleuth*, respectively.

For *DESeq2*, *EBSeq*, *limma*, and *SAMseq*, we provided the methods with counts on the gene- or transcript-level generated by the *tximport* package [8], using the *lengthScaledTPM* method. For the two balanced batches experiment, we provided all methods with the batch variable, except *SAMseq* which does not have such an option. For *SAMseq*, for the two batch experiment, we first adjusted the counts using the known batch variable and the `removeBatchEffect` function in the *limma* [36] package, which improved *SAMseq*'s performance. For *sleuth* we directly provided *kallisto* quantification files to the method, and for detection of DGE, changes in the total expression of all the transcripts of a gene across condition, we used the count summarization option.

1.9   Single Cell RNA-seq of Mouse Neurons

We downloaded a dataset of scRNA-seq of the developing mouse brain in order to evaluate the performance of *Swish* on RNA-seq counts with a different distribution and with more gene-level inferential uncertainty than bulk RNA-seq. In this dataset, cells from the cortex, hippocampus and ventricular zone of a embryonic (E18) mouse were dissociated and sequenced on the 10x chromium pipeline [37]. scRNA-seq data was quantified by *Alevin* [18] v0.13.0 with the default command line parameters along with `--numCellBootstraps 30` to generate 30 bootstrap estimated count matrices. The total number of cells was 931, and the number of genes was 52,325 (Gencode [30] release M16 mouse reference transcripts).

In a comprehensive benchmark, Soneson and Robinson [38] compared DE methods designed for both scRNA-seq and bulk RNA-seq, and found that the Mann-Whitney Wilcoxon test [23] had favorable performance compared to numerous other methods in detection of DE genes across cells. However, the Wilcoxon test was listed as having the downside of not accommodating complex designs (for example, covariate control, sample pairing, or interaction terms). We therefore compared *Swish* to the Wilcoxon test statistic in detection of genes differentially expressed across clusters.

Seurat [39] was used to cluster the cells in the dataset. First, cells with low quality were removed, based on quality control plots (Supplementary Figure 1). Those cells having less than 1,500 or greater than 6,500 genes expressed were removed based on the quality control plots. Additionally, those cells with greater than 4% percent of mitochondrial genes detected were removed. These filters resulted in

835 cells remaining in the dataset. For *Seurat* clustering, gene counts were then normalized by their total count, multiplied by 10,000, and the shifted logarithm was applied, with a shift of 1 to preserve zeros.

Only highly variable genes were used for clustering by *Seurat*, which was determined by calculating the mean expression of scaled counts and dispersion for each gene, binning genes by mean expression into 20 bins, and calculating z-scores for s^2/μ, (the sample variance over the sample mean) within each bin. The highly variable genes were defined as those with mean expression between 0.0125 and 3 on the natural log scale and z-score of 0.5 or larger. A linear model was fit to the log scaled counts, to further remove any technical artifacts associated with the total number of UMIs detected per cell and with the percent of mitochondrial genes detected. The top 10 principal components of residuals of highly variable genes were used to cluster the cells, using *Seurat*'s graph-based unsupervised clustering method.

After defining clusters using the highly variable genes, we returned to the matrix of estimated counts and attempted to find genes which were differentially expressed across two specific clusters. The clusters were chosen based on expression of neural differentiation marker genes reported by a previous study [40]. We applied both *Swish* and a standard two-sample Wilcoxon test to the scaled counts of cells in the two clusters. We scaled the counts using size factors estimated with the setting `type="poscounts"` in the *DESeq2* package [11], as recommended for scRNA-seq differential expression analysis [41]. Genes without a count of 3 or more in 5 or more cells were removed prior to the DE analysis across clusters. *Swish* produces a q-value, while the p-values from the Wilcoxon test were adjusted using the Benjamini-Hochberg [42] method to produce nominal FDR bounded sets.

2 Results

2.1 Simulation

We evaluated the InfRV in the simulated bulk RNA-seq dataset at the gene and transcript level. At the transcript level, there are two main sources of abundance uncertainty: 1) among different transcripts within a gene, and 2) due to homologous sequence of transcripts across genes. At the gene level, we observed substantially less abundance uncertainty, as expected (Figure 1). At the gene level the uncertainty among transcripts within a gene has been collapsed, and only the latter source of uncertainty remains, also noted by Soneson et al. [8].

At the gene level, where there was less quantification uncertainty, *Swish* had similar performance as DGE methods overall (Figure 2). For the experiment with a single batch, *Swish* had similar TPR and FDR to *EBSeq*, *limma*, *SAMseq*, and *sleuth*, and better control of FDR compared to *DESeq2*. *Swish* had improved FDR control compared to *SAMseq*, but only slightly so, and this minor performance difference was expected as there was relatively low InfRV compared to transcript-level analysis. For the experiment with two batches, *limma* and *EBSeq* showed the best performance, with good sensitivity and control of FDR. *Swish* exceeded the FDR (e.g. observed FDR of 13% for nominal 10% target overall), but had improved control relative to *SAMseq* for genes with the highest inferential uncertainty (Supplementary Figure 2). *sleuth* exhibited loss of control of the FDR for the 5% and 10% target in the two batch experiment (Supplementary Figure 3).

At the transcript level, *Swish* had consistently high sensitivity and good control of FDR, where other methods either had low sensitivity or loss of FDR control in some setting. *Swish* controlled FDR for all nominal FDR targets (1%, 5%, and 10%), while *SAMseq* tended to exceed the 5% and 10% target for transcripts with medium-to-high InfRV, both for the single batch experiment (Figure 3), and the two batch experiment (Supplementary Figure 4). *limma* had reduced sensitivity for all InfRV categories, for both experiments. *EBSeq* had high sensitivity but loss of FDR control for the transcripts with high InfRV. *sleuth* performed well in the single batch experiment, but lost control of FDR for the 5% and 10% target in the two batch experiment (Supplementary Figure 5).

Overall, *Swish* performed well at the gene level, although not better than *limma* and *EBSeq* for the two batch experiment. *Swish* out-performed other methods at the transcript level, both for the single and two batch experiments, where other methods either had reduced sensitivity or loss of FDR in some setting.

2.2 scRNA-seq DE comparison of clusters

We evaluated the extent to which inclusion of uncertainty in the gene-level abundance would affect differential expression in an scRNA-seq dataset. As described in the Methods, *Seurat* was used to identify clusters in a dataset of 835 cells from embryonic mouse brain. *Seurat* identified a total of nine clusters,

visualized in Supplementary Figure 6. Expression level of known cell-type markers for developing mouse brain, as reported by Loo et al. [40], was overlaid on the t-Distributed Stochastic Neighbor Embedding (tSNE) [43] plot. *Eomes*, a gene expressed in neural progenitor cells, was highly expressed in cluster 6 and 7, while *Mki67*, a proliferation marker, was expressed mostly in cluster 7 (Supplementary Figure 7). This pattern suggested that cluster 6 and 7 might represent populations of non-cycling and cycling neural progenitor cells, respectively.

We compared the test results of a two-sample Mann-Whitney Wilcoxon [23] test to *Swish* across clusters 6 and 7. We defined sets of genes which were called by the Wilcoxon test at a 1% FDR threshold (n=316) or by *Swish* at a 1% FDR threshold (n=480). Examining the intersection of genes called by either Wilcoxon test or *Swish* revealed that, when genes were called exclusively by Wilcoxon test and not by *Swish*, these genes tended to have higher InfRV, compared to genes called only by *Swish* and to genes called by both methods (Supplementary Figure 8).

We additionally performed gene set enrichment analysis with *GOseq* [44], which can account for potential selection bias due to gene length if it exists. In this dataset no systematic enrichment for longer genes was detected by *GOseq*, which was expected as the 3' tagged data does not exhibit length bias. The GO categories [45] detected by *GOseq*, using either significant gene sets derived from Wilcoxon tests or from *Swish* were mostly overlapping, with genes relating to cell cycle and cell division making up a large proportion of the DE genes between cluster 6 and 7 (Supplementary Tables 1-2). We hypothesized therefore that cluster 7 represents active, cycling neural progenitor cells, while cluster 6 represents non-cycling neural progenitor cells.

In order to assess whether we would see a similar pattern of higher InfRV for genes called DE only by Wilcoxon test, we performed another comparison of clusters 4 and 6, which appeared distinct in the tSNE plot of cells (Supplementary Figure 7). Again, a similar pattern was found, where those genes called exclusively by Wilcoxon test tended to have higher InfRV relative to those found only by *Swish* or by both methods (Supplementary Figure 9). We performed gene set enrichment analysis with *GOseq* on the gene sets found by either method, and the results were highly overlapping, with many genes related to neurogenesis and neuron differentiation found by both methods.

3   Discussion

Many parametric models have been proposed to include quantification uncertainty into statistical testing procedures when evaluating differential transcript or gene expression. We extended an existing nonparametric framework, *SAMseq*, through the incorporation of inferential replicate count matrices, so that the test statistic becomes exposed to the uncertainty of the counts. We show that this extension recovers control of FDR in particular for transcripts with high inferential relative variance. We used posterior samples generated by a Gibbs sampler with thinning to reduce autocorrelation, although bootstrapping of reads to generate inferential replicates is also compatible with our proposed method. *Swish* additionally extends *SAMseq* by allowing for control of discrete batches using the stratified Wilcoxon test, and allowing for analysis of paired samples using the Wilcoxon signed-rank test.

Swish has the greatest advantage over other competing methods at the transcript level, while it performs similar to other methods at the gene level. We additionally demonstrated the use of *Swish* on scRNA-seq counts generated by the *Alevin* method, making use of bootstrap inferential replicates at the gene level. We found that *Swish* detected more differentially expressed genes across sub-populations of cells compared to a Wilcoxon test. *Swish* was more conservative for some genes with elevated inferential relative variance, compared to the Wilcoxon test, which was desired behavior as the 3' tagged scRNA-seq datasets exhibit higher uncertainty on the gene-level compared to bulk RNA-seq. In this scRNA-seq analysis, we did not take into account that we first clustered the data based on highly variable genes, and then performed differential analysis using those and other genes, which may generate anti-conservative inference as noted recently by Zhang et al. [46]. We are interested in extensions which may alleviate this problem and in general help to avoid comparisons across unstable clusters [47–50].

Some of the limitations of our proposed work include the types of experiments that it can be used to analyze. Because it relies on permutation of samples to generate a null distribution over all transcripts or genes, *SAMseq* and *Swish* in practice do not have sufficient power to detect differential expression when the sample size is less than 4 per group. Additionally, certain terms which can be included in linear models, such as interaction terms between categorical or continuous variables, cannot be easily incorporated into the nonparametric framework. We focused on inference across a discrete nuisance covariate, for example sample preparation batches, which motivated the stratified test statistic. We cannot yet accommodate more continuous measures of batch, such as those estimated by *RUVSeq* [51]

or *svaseq* [52], unless such factors of unwanted variation could be discretized without substantial loss of information. Currently, *Swish* supports two group comparisons, with discrete covariate correction and sample pairing, although we could extend to other analysis types already supported by *SAMseq*, including multi-group comparisons via the Kruskal-Wallis statistic, quantitative association via the Spearman correlation, or survival analysis via the Cox proportional hazards model.

4   Code Availability

The `swish` function is available in the *fishpond* R/Bioconductor package at `https://github.com/mikelove/fishpond`. The package has been submitted to Bioconductor [53]. The implementation is efficient, with vectorized code for all row-wise ranking and quantile calculations via the *matrixStats* package [54]. The software runs in comparable time to *DESeq2*, taking 2-3 seconds per 1,000 genes for small datasets, e.g. total $n \in [10 - 20]$. For larger datasets, *Swish* scales better than *DESeq2* (Supplementary Figure 10).

The package contains a detailed vignette with live code examples, showing how to import *Salmon* quantification files using the *tximeta* package, perform differential transcript or gene expression analysis with `swish`, and visualize results for each transcript or gene. The package vignette includes an example of a paired RNA-seq analysis, using a subset of the RNA-seq data from a recent publication on human macrophage immune response [55] (Supplementary Figure 11).

The scripts used to generate the Love et al. [7] simulation data are available at the following link: `https://dx.doi.org/10.5281/zenodo.1410443`. The scripts used in evaluating the methods here, and in analyzing the scRNA-seq dataset have been posted at the following link: `https://github.com/azhu513/swishPaper`.

The versions of software used in the paper are: *fishpond* (*Swish* method): 0.99.5, *samr* (*SAMseq* method): 2.0, *Salmon*: 0.11.3, *Alevin*: 0.13.0, *tximport*: 1.10.1, *kallisto*: 0.44.0, *sleuth*: 0.30.0, *DESeq2*: 1.20.0, *EBSeq*: 1.22.0, *limma*: 3.38.3, *Seurat*: 2.3.4, *GOseq*: 1.34.1

5   Data Availability

The simulated data used to evaluate the methods is published in Love et al. [7]. The raw read files can be accessed at the following links: `https://doi.org/10.5281/zenodo.1291375`, `https://doi.org/10.5281/zenodo.1291404`, `https://doi.org/10.5281/zenodo.2564176`, `https://doi.org/10.5281/zenodo.2564261`. The processed simulation quantification files can be accessed at the following link: `https://doi.org/10.5281/zenodo.2564115`. The mouse neurons dataset is available under the heading *Chromium Demonstration (v2 Chemistry) > Cell Ranger 2.1.0 > 1k Brain Cells from an E18 Mouse*, from `https://support.10xgenomics.com/single-cell-gene-expression/datasets`.

6   Acknowledgments

We thank Naim Rashid, Yun Li, Jason Stein, and Scott Van Buren for useful discussions.

7   Funding

MIL is supported by R01 HG009937, R01 MH118349, P01 CA142538, and P30 ES010126. JGI and AZ are supported by R01 GM070335 and P01 CA142538. AS and RP are supported by R01 HG009937. This work was supported by National Science Foundation award CCF-1750472 and by grant number 2018-182752 from the Chan Zuckerberg Initiative DAF, an advised fund of Silicon Valley Community Foundation.

Figure 1: Mean of InfRV over samples for genes and transcripts in the simulation studies (log10 scale). InfRV is binned by increments of 0.1, so that 10 bins span an order of magnitude. (A) Gene level for the single batch experiment. (B) Transcript level for the single batch experiment.

Figure 2: True positive rate (y-axis) over false discovery rate (x-axis) for DGE analysis with only one batch of samples, with 6 samples in each of the two conditions. The first panel depicts overall performance, while the subsequent three panels depict the genes stratified into thirds by InfRV averaged over samples.

Figure 3: True positive rate (y-axis) over false discovery rate (x-axis) for DTE analysis with only one batch of samples, with 6 samples in each of the two conditions. As in Figure 2, the first panel depicts overall performance, while the subsequent three panels depict the transcripts stratified into thirds by InfRV averaged over samples.

References

[1] Peter Glaus, Antti Honkela, and Magnus Rattray. Identifying differentially expressed transcripts from RNA-seq data with biological variation. *Bioinformatics*, 28(13), 2012.

[2] Ning Leng, John A Dawson, James A Thomson, Victor Ruotti, Anna I Rissman, Bart MG Smits, Jill D Haag, Michael N Gould, Ron M Stewart, and Christina Kendziorski. EBSeq: an empirical Bayes hierarchical model for inference in RNA-seq experiments. *Bioinformatics*, 29(8):1035–1043, 2013.

[3] Cole Trapnell, Adam Roberts, Loyal Goff, Geo Pertea, Daehwan Kim, David R Kelley, Harold Pimentel, Steven L Salzberg, John L Rinn, and Lior Pachter. Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nature Protocols*, 7(3):562, 2012.

[4] Sahar Al Seesi, Yvette Temate-Tiagueu, Alex Zelikovsky, and Ion I Măndoiu. Bootstrap-based differential gene expression analysis for RNA-Seq data with and without replicates. *BMC Genomics*, 15(Suppl 8), 2014.

[5] Harold Pimentel, Nicolas L Bray, Suzette Puente, P Melsted, and Lior Pachter. Differential analysis of RNA-seq incorporating quantification uncertainty. *Nature Methods*, 14(7):687–690, 07 2017.

[6] Kimon Froussios, Kira Mourão, Gordon Simpson, Geoff Barton, and Nicholas Schurch. Relative Abundance of Transcripts (RATs): Identifying differential isoform abundance from RNA-seq [version 1; referees: awaiting peer review]. *F1000Research*, 8(213), 2019.

[7] Michael I Love, Charlotte Soneson, and Rob Patro. Swimming downstream: statistical analysis of differential transcript usage following Salmon quantification [version 3; referees: 3 approved] . *F1000Research*, 7(952), 2018.

[8] Charlotte Soneson, Michael I Love, and Mark D Robinson. Differential analyses for RNA-seq: transcript-level estimates improve gene-level inferences [version 2; referees: 2 approved]. *F1000Research*, 4(1521), 2016.

[9] Jun Li and Robert Tibshirani. Finding consistent patterns: a nonparametric approach for identifying differential expression in RNA-Seq data. *Statistical Methods in Medical Research*, 22(5):519–536, 2013.

[10] Charlotte Soneson and Mauro Delorenzi. A comparison of methods for differential expression analysis of RNA-seq data. *BMC Bioinformatics*, 14(1):91, 2013.

[11] Michael I Love, Wolfgang Huber, and Simon Anders. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, 15(12):550, 2014.

[12] Aleksandra A Kolodziejczyk, Jong Kyoung Kim, Valentine Svensson, John C Marioni, and Sarah A Teichmann. The technology and biology of single-cell RNA sequencing. *Molecular Cell*, 58(4): 610–620, 2015.

[13] Saiful Islam, Amit Zeisel, Simon Joost, Gioele La Manno, Pawel Zajac, Maria Kasper, Peter Lönnerberg, and Sten Linnarsson. Quantitative single-cell RNA-seq with unique molecular identifiers. *Nature Methods*, 11(2):163, 2014.

[14] Grace XY Zheng, Jessica M Terry, Phillip Belgrader, Paul Ryvkin, Zachary W Bent, Ryan Wilson, Solongo B Ziraldo, Tobias D Wheeler, Geoff P McDermott, Junjie Zhu, et al. Massively parallel digital transcriptional profiling of single cells. *Nature Communications*, 8(14049), 2017.

[15] Evan Z Macosko, Anindita Basu, Rahul Satija, James Nemesh, Karthik Shekhar, Melissa Goldman, Itay Tirosh, Allison R Bialas, Nolan Kamitaki, Emily M Martersteck, et al. Highly parallel genome-wide expression profiling of individual cells using nanoliter droplets. *Cell*, 161(5):1202–1214, 2015.

[16] Allon M Klein, Linas Mazutis, Ilke Akartuna, Naren Tallapragada, Adrian Veres, Victor Li, Leonid Peshkin, David A Weitz, and Marc W Kirschner. Droplet barcoding for single-cell transcriptomics applied to embryonic stem cells. *Cell*, 161(5):1187–1201, 2015.

[17] Tom Smith, Andreas Heger, and Ian Sudbery. UMI-tools: modeling sequencing errors in unique molecular identifiers to improve quantification accuracy. *Genome Research*, 27(3):491–499, 2017.

[18] Avi Srivastava, Tom Sean Smith, Ian Sudbery, and Rob Patro. Alevin: An integrated method for dscRNA-seq quantification. *bioRxiv*, 2018.

[19] Rob Patro, Geet Duggal, Michael I Love, Rafael A Irizarry, and Carl Kingsford. Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*, 14:417–419, 2017.

[20] Simon Anders and Wolfgang Huber. Differential expression analysis for sequence count data. *Genome Biology*, 11(10):R106, 2010.

[21] Cole Trapnell, Brian A Williams, Geo Pertea, Ali Mortazavi, Gordon Kwan, Marijke J Van Baren, Steven L Salzberg, Barbara J Wold, and Lior Pachter. Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nature Biotechnology*, 28(5):511, 2010.

[22] Ernest Turro, Shu-Yi Su, Ângela Gonçalves, Lachlan JM Coin, Sylvia Richardson, and Alex Lewin. Haplotype and isoform specific expression estimation using multi-mapping RNA-seq reads. *Genome Biology*, 12(2):R13, 2011.

[23] Frank Wilcoxon. Individual comparisons by ranking methods. *Biometrics Bulletin*, 1:80–83, 1945.

[24] PH Van Elteren. On the Combination of Independent Two-Sample Tests of Wilcoxon. *Bulletin of the International Statistical Institute*, 37:351–361, 1960.

[25] Devan V Mehrotra, Xiaomin Lu, and Xiaoming Li. Rank-based analyses of stratified experiments: alternatives to the van elteren test. *The American Statistician*, 64(2):121–130, 2010.

[26] John D Storey and Robert Tibshirani. Statistical significance for genomewide studies. *Proceedings of the National Academy of Sciences*, 100(16):9440–9445, 2003.

[27] Alyssa C Frazee, Andrew E Jaffe, Ben Langmead, and Jeffrey T Leek. Polyester: simulating RNA-seq datasets with differential transcript expression. *Bioinformatics*, 31(17):2778–2784, 2015.

[28] Michael I Love, John B Hogenesch, and Rafael A Irizarry. Modeling of RNA-seq fragment sequence bias reduces systematic errors in transcript abundance estimation. *Nature Biotechnology*, 34:1287–1291, 2016.

[29] Tuuli Lappalainen, Michael Sammeth, Marc R Friedländer, Peter AC 't Hoen, Jean Monlong, Manuel A Rivas, Mar Gonzalez-Porta, Natalja Kurbatova, Thasso Griebel, Pedro G Ferreira, et al. Transcriptome and genome sequencing uncovers functional variation in humans. *Nature*, 501(7468):506, 2013.

[30] Adam Frankish, Alexandra Bignell, Andrew Berry, Andrew Yates, Anne Parker, Bianca M Schmitt, Bronwen Aken, Carlos GarcaGirn, Daniel Zerbino, Eloise Stapleton, Fergal J Martin, Fiona Cunningham, If Barnes, Irina Sycheva, Jane Loveland, Jonathan M Mudge, Jose Manuel Gonzalez, Magali Ruffier, Marie-Marthe Suner, Matthew Hardy, Osagie G Izuogu, Sarah Donaldson, Shamika Mohanan, Thibaut Hourlier, Tiago Grego, Toby Hunt, Paul Flicek, James Wright, Jyoti S Choudhary, Julien Lagarde, Silvia CarbonellSala, Roderic Guig, Fernando Pozo, Laura Martnez, Michael L Tress, Toms DiDomenico, Paul Muir, Barbara Uszczynska-Ratajczak, Benedict Paten, Ian T Fiddes, Joel Armstrong, Mark Diekhans, Tim JP Hubbard, Alexandre Reymond, Anne-Maud Ferreira, Jacqueline Chrast, Rory Johnson, Irwin Jungreis, Manolis Kellis, Baikang Pei, Fabio CP Navarro, Jinuri Xu, Yan Zhang, Mark Gerstein, and Cristina Sisu. GENCODE reference annotation for the human and mouse genomes. *Nucleic Acids Research*, 47(D1):D766–D773, 2018.

[31] Nicolas L Bray, Harold Pimentel, Páll Melsted, and Lior Pachter. Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, 34(5):525, 2016.

[32] Charlotte Soneson and Mark D Robinson. iCOBRA: open, reproducible, standardized and live method benchmarking. *Nature Methods*, 13(283), 2016.

[33] Charity W Law, Yunshun Chen, Wei Shi, and Gordon K Smyth. Voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biology*, 15(2):R29, 2014.

[34] Davis J McCarthy, Yunshun Chen, and Gordon K Smyth. Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation. *Nucleic Acids Research*, 40(10): 4288–4297, 2012.

[35] Steven P Lund, Dan Nettleton, Davis J McCarthy, and Gordon K Smyth. Detecting differential expression in RNA-sequence data using quasi-likelihood with shrunken dispersion estimates. *Statistical Applications in Genetics and Molecular Biology*, 11(5), 2012.

[36] Gordon K Smyth. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, 3(1):1–25, 2004.

[37] 10x Genomics. Single cell gene expression datasets, 2019. URL `https://support.10xgenomics.com/single-cell-gene-expression/datasets`.

[38] Charlotte Soneson and Mark D Robinson. Bias, robustness and scalability in single-cell differential expression analysis. *Nature Methods*, 15(4):255, 2018.

[39] Andrew Butler, Paul Hoffman, Peter Smibert, Efthymia Papalexi, and Rahul Satija. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nature Biotechnology*, 36(5):411, 2018.

[40] Lipin Loo, Jeremy M Simon, Lei Xing, Eric S McCoy, Jesse K Niehaus, Jiami Guo, ES Anton, and Mark J Zylka. Single-cell transcriptomic analysis of mouse neocortical development. *Nature Communications*, 10(1):134, 2019.

[41] Koen Van den Berge, Fanny Perraudeau, Charlotte Soneson, Michael I Love, Davide Risso, Jean-Philippe Vert, Mark D Robinson, Sandrine Dudoit, and Lieven Clement. Observation weights unlock bulk rna-seq tools for zero inflation and single-cell applications. *Genome Biology*, 19(1):24, 2018.

[42] Yoav Benjamini and Yosef Hochberg. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, 57 (1):289–300, 1995. ISSN 00359246.

[43] Laurens van der Maaten and Geoffrey Hinton. Visualizing data using t-SNE. *The Journal of Machine Learning Research*, 9:2579–2605, 2008.

[44] Matthew D Young, Matthew J Wakefield, Gordon K Smyth, and Alicia Oshlack. Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biology*, 11(2):R14, 2010.

[45] The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing strong. *Nucleic Acids Research*, 47(D1):D330–D338, 2018.

[46] Jesse Min Zhang, Govinda M Kamath, and David N Tse. Towards a post-clustering test for differential expression. *bioRxiv*, 2018.

[47] Vladimir Yu Kiselev, Kristina Kirschner, Michael T Schaub, Tallulah Andrews, Andrew Yiu, Tamir Chandra, Kedar N Natarajan, Wolf Reik, Mauricio Barahona, Anthony R Green, et al. SC3: consensus clustering of single-cell RNA-seq data. *Nature Methods*, 14(5):483, 2017.

[48] Peijie Lin, Michael Troup, and Joshua WK Ho. CIDR: Ultrafast and accurate clustering through imputation for single-cell RNA-seq data. *Genome Biology*, 18(1):59, 2017.

[49] S Freytag, L Tian, I Lönnstedt, M Ng, and M Bahlo. Comparison of clustering tools in R for medium-sized 10x Genomics single-cell RNA-sequencing data [version 2; referees: 3 approved]. *F1000Research*, 7(1297), 2018.

[50] Yuchen Yang, Ruth Huh, Houston W Culpepper, Yuan Lin, Michael I Love, and Yun Li. SAFE-clustering: Single-cell Aggregated (from Ensemble) clustering for single-cell RNA-seq data. *Bioinformatics*, 2018. doi: 10.1093/bioinformatics/bty793.

[51] Davide Risso, John Ngai, Terence P Speed, and Sandrine Dudoit. Normalization of RNA-seq data using factor analysis of control genes or samples. *Nature Biotechnology*, 32:896 EP–, 08 2014.

[52] Jeffrey T Leek. svaseq: removing batch effects and other unwanted noise from sequencing data. *Nucleic Acids Research*, 42(21):e161–e161, 2014.

[53] Wolfgang Huber, Vincent J Carey, Robert Gentleman, Simon Anders, Marc Carlson, Benilton S Carvalho, Hector Corrada Bravo, Sean Davis, Laurent Gatto, Thomas Girke, Raphael Gottardo, Florian Hahne, Kasper D Hansen, Rafael A Irizarry, Michael Lawrence, Michael I Love, James Mac-Donald, Valerie Obenchain, Andrzej K Oles, Hervé Pagès, Alejandro Reyes, Paul Shannon, Gordon K Smyth, Dan Tenenbaum, Levi Waldron, and Martin Morgan. Orchestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*, 12:115 EP –, 2015.

[54] Henrik Bengtsson. matrixStats: Functions that Apply to Rows and Columns of Matrices (and to Vectors). *CRAN R package*, 2017.

[55] Kaur Alasoo, Julia Rodrigues, Subhankar Mukhopadhyay, Andrew J Knights, Alice L Mann, Kousik Kundu, Christine Hale, Gordon Dougan, Daniel J Gaffney, and HIPSCI Consortium. Shared genetic effects on chromatin and gene expression indicate a role for enhancer priming in immune response. *Nature Genetics*, 50(3):424–431, 2018.