The great tit HapMap project: a continental-scale analysis of genomic variation in a songbird

Lewis G. Spurgin[1,2]*, Mirte Bosse[3,4,5], Frank Adriaensen[6], Tamer Albayrak[7], Christos Barboutis[8], Eduardo Belda[9], Andrey Bushuev[10], Jacopo G. Cecere[11], Anne Charmantier[12], Mariusz Cichon[13], Niels J. Dingemanse[14], Blandine Doligez[15,16], Tapio Eeva[17], Kjell Einar Erikstad[18], Vyacheslav Fedorov[10]**, Matteo Griggio[19]**, Dieter Heylen[6,20,21], Sabine Hille[22], Camilla A. Hinde[23], Elena Ivankina[24], Bart Kempenaers[25], Anvar Kerimov[10], Milos Krist[26], Laura Kvist[27], Veronika N. Laine[3,28], Raivo Mänd[29], Erik Matthysen[6], Ruedi Nager[30], Boris P. Nikolov[31], A. Claudia Norte[32], Markku Orell[27], Jenny Ouyang[25,33], Gergana Petrova-Dinkova[31], Heinz Richner[34], Diego Rubolini[35], Tore Slagsvold[36], Vallo Tilgar[29], János Török[37], Barbara Tschirren[38], Csongor I. Vágási[39,40], Teru Yuta[41], Martien A.M. Groenen[4], Marcel E. Visser[3,4,42], Kees van Oers[3], Ben C. Sheldon[2] and Jon Slate[43]*

1. School of Biological Sciences, University of East Anglia, Norwich Research Park, United Kingdom
2. Edward Grey Institute, University of Oxford, United Kingdom
3. Department of Animal Ecology, Netherlands Institute of Ecology (NIOO-KNAW), Wageningen, the Netherlands
4. Animal Breeding and Genomics Centre, Wageningen University, the Netherlands
5. Department of Ecological Science, Animal Ecology Group, Vrije Universiteit Amsterdam, Amsterdam, The Netherlands
6. Evolutionary Ecology Group, Department of Biology, University of Antwerp, Universiteitsplein 1, 2610 Antwerp, Belgium
7. Department of Biology, Mehmet Akif Ersoy University, Science and Art Faculty, Ortulu, Burdur, Turkey

8. Hellenic Ornithological Society / BirdLife Greece, Themistokleous 80, GR-10681 Athens

9. Institut d'Investigació per a la Gestió Integrada de Zones Costaneres, Campus de Gandia, Universitat Politècnica de València, Carrer Paranimf 1, E-46730 Grau de Gandia (València), Spain

10. Faculty of Biology, Lomonosov Moscow State University, Moscow 119234, Russia

11. Area Avifauna Migratrice, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), via Ca' Fornacetta 9, I-40064, Ozzano Emilia, (BO), Italy

12. CEFE-CNRS, UMR 5175, 1919, route de Mende, F34293 Montpellier Cedex 5, France

13. Institute of Environmental Sciences, Jagiellonian University, Gronostajowa 7, 30-387 Kraków, Poland

14. Behavioural Ecology, Department of Biology, Ludwig Maximilians University of Munich, Planegg-Martinsried, Germany

15. UMR CNRS 5558—LBBE, Biométrie et Biologie Évolutive, UCB Lyon 1 - Bât. Grégor Mendel, 43 bd du 11 novembre 1918, 69622 VILLEURBANNE cedex, France

16. Department of Ecology and Evolution, Animal Ecology, Evolutionary Biology Centre, Uppsala University, Sweden

17. Department of Biology, University of Turku, Turku 20014, Finland

18. Norwegian Institute for Nature Research, FRAM-High North Research Centre for Climate and the Environment, 9296 Tromsø, Norway

19. Department of Biology, University of Padova, Via U. Bassi 58/B, I-35131, Padova, Italy

20. Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ, United States of America

21. Interuniversity Institute for Biostatistics and statistical Bioinformatics, Hasselt University, Diepenbeek, Belgium

22. Institute of Wildlife Biology and Game Management, University of Natural Resources and Life Science, A-1180 Vienna, Austria

23. Behavioural Ecology Group, Department of Life Sciences, Anglia Ruskin University, East Road Cambridge, Cambridgeshire, CB1 1PT, UK.

24. Zvenigorod Biological Station of Lomonosov Moscow State University, P.O. Box Shikhovo, Odintsovo District, Moscow 143036, Russia

25. Max Planck Institute for Ornithology, Department of Behavioural Ecology & Evolutionary Genetics, Eberhard-Gwinner-Straße, House 5, 82319 Seewiesen (Starnberg), Germany

26. Department of Zoology and Laboratory of Ornithology, Faculty of Science, Palacký University, Olomouc 77147, Czech Republic

27. Department of Ecology and Genetics, P.O.Box 3000, 90014-University of Oulu, Finland

28. Finnish Museum of Natural History, University of Helsinki, FI-00014, Finland

29. Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu, Vanemuise 46, Tartu 51014, Estonia

30. Institute of Biodiversity, Animal Health and Comparative Medicine, University of Glasgow, Glasgow G12 8QQ, United Kingdom

31. Bulgarian Ornithological Centre, Institute of Biodiversity and Ecosystem Research, Bulgarian Academy of Sciences, 1 Tsar Osvoboditel Blvd, 1000 Sofia, Bulgaria

32. MARE - Marine and Environmental Sciences Centre, Department of Life Sciences, Faculty of Sciences and Technology, University of Coimbra, Portugal

33. University of Nevada, Reno, NV, United States of America

34. Evolutionary Ecology Lab, Institute of Ecology and Evolution, University of Bern, Bern 3012, Switzerland

35. Dipartimento di Scienze e Politiche Ambientali, Università degli Studi di Milano, via Celoria 26, I-20133, Milano, Italy

36. Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, P.O. Box 1066, Blindern, 0316 Oslo, Norway

37. Behavioural Ecology Group, Department of Systematic Zoology and Ecology, Eötvös Loránd University, Budapest H-1117, Hungary

38. University of Exeter, Centre for Ecology and Conservation, Treliever Road, Penryn, TR10 9FE, United Kingdom

39. Evolutionary Ecology Group, Hungarian Department of Biology and Ecology, Babeş-Bolyai University, Cluj-Napoca, Romania

40. Behavioural Ecology Research Group, Department of Evolutionary Zoology, University of Debrecen, Debrecen, Hungary

41. Graduate School of Environmental Science, Hokkaido University, N10 W5 Sapporo, Hokkaido 060-0810, Japan

42. Groningen Institute for Evolutionary Life Sciences (GELIFES), University of Groningen,
Groningen, the Netherlands

43. Department of Animal and Plant Sciences, University of Sheffield, United Kingdom

* **Correspondence**: l.spurgin@uea.ac.uk; j.slate@sheffield.ac.uk

** Deceased

Abstract

A major aim of evolutionary biology is to understand why patterns of genomic diversity

vary among populations and species. Large-scale genomic studies of widespread species

are useful for studying how the environment and demographic history shape patterns of

genomic divergence, and with the continually decreasing cost of sequencing and

genotyping, such studies are now becoming feasible. Here, we carry out one of the most

geographically comprehensive surveys of genomic variation in a wild vertebrate to date;

the great tit (*Parus major*) HapMap project. We screened *ca* 500,000 SNP markers across

647 individuals from 29 populations, spanning almost the entire geographic range of the

European great tit subspecies. We found that genome-wide variation was consistent with a

recent colonisation across Europe from a single refugium in South-East Europe, with

bottlenecks and reduced genetic diversity in island populations. Differentiation across the

genome was highly heterogeneous, with clear "islands of differentiation" even among

populations with very low levels of genome-wide differentiation. Low local recombination

rate in the genome was a strong predictor of high local genomic differentiation (F_{ST}),

especially in island and peripheral mainland populations, suggesting that the interplay

between genetic drift and recombination is a key driver of highly heterogeneous

differentiation landscapes. We also detected genomic outlier regions that were confined to

one or more peripheral great tit populations, most likely as a result of recent directional

selection at the range edges of this species. Haplotype-based measures of selection were

also related to recombination rate, albeit less strongly, and highlighted population-specific

sweeps that likely resulted from positive selection. These regions under positive selection

contained candidate genes associated with morphology, thermal adaptation and

colouration, providing promising avenues for future investigation. Our study highlights

how comprehensive screens of genomic variation in wild organisms can provide unique

insights into evolution.

Introduction

Since the first studies of allozyme variation in humans [1] and *Drosophila* [2,3], there has

been great interest in explaining how evolutionary and ecological processes shape the

patterns of genetic variation observed within and among natural populations. One focus of

research and debate in this area has been on quantifying the roles of adaptive and neutral

processes in explaining observed levels of genetic variation [4]. However, adaptation does

not occur in isolation, but acts on genetic variation that is also shaped by mutation,

recombination, gene flow, and genetic drift [5]. More recently there has been increased

effort in understanding how these fundamental evolutionary forces operate in concert to

generate and maintain the levels of genetic diversity commonly observed in natural

populations [6,7].

The increasing feasibility of high-throughput sequencing and genotyping, alongside

subsequent characterisation of genome-wide variation across large numbers of individuals,

has revealed that at the genomic level, patterns of variation and divergence among natural

populations and species are highly heterogeneous [8]. A key feature of these "genomic

landscapes" of divergence that has received particular attention is the presence of so-called

"islands of differentiation": outlier regions of the genome with high levels of divergence

estimated from statistics such as F_{ST} or d_{xy} that consistently emerge at the same loci in

different comparisons between populations or related species [8–11]. Initially these

regions were termed "islands of speciation", and were thought to arise as a result of

reduced gene flow in genomic regions associated with reproductive isolation [8,12].

Subsequent research has revealed that highly heterogeneous patterns of genomic

divergence can occur even in the complete absence of gene flow, as a result of

recombination rate variation and linked selection [13,14]. In genomic regions of low

recombination, selection for beneficial mutations (positive selection), or against

deleterious mutations (background selection), will impact relatively large genomic regions

as a result of high levels of linkage disequilibrium (LD) among sites. Selection within these

regions reduces diversity within populations, and increases levels of differentiation among

them, resulting in "islands" of increased differentiation that persist over evolutionary time

[14,15]. Another, less well explored reason for which islands of divergence can arise is due

to the differential effects of genetic drift in response to variation in effective population size

across different genomic regions; something that may be particularly important in recently

colonised populations [16,17]. These circumstances promote fixation of haplotypes and

therefore result in either reduced or inflated local differentiation.

Comparing patterns of genomic differentiation among sets of populations or species at

different stages of the divergence/speciation continuum is a powerful way of disentangling

the forces that shape variation among populations. Martin et al. [18] showed that, across

multiple *Heliconius* butterfly populations and species, patterns of genomic variation were

shaped by a combination of gene flow and selection, particularly in genomic regions

harbouring genes involved in wing patterning. In contrast, Renaut et al. [10] showed that in

Helianthus sunflowers, genomic architecture was the main driver of genomic

differentiation across sets of populations. Similarly, recent research in birds has revealed

that differentiation landscapes are conserved across populations, species and even across

avian families, with the same islands of differentiation arising among populations of

distantly related species [19–21]. This latter pattern appears to have arisen, at least in part,

as a result of a highly conserved synteny and recombination landscape in birds [22], with

background selection in regions of low recombination producing recurrent islands of

differentiation [23].

It is now clear that the recombination landscape and linked selection are key drivers of

genomic variation within and among populations. However, we are only just beginning to

understand how this linked selection interacts with other evolutionary forces to shape

patterns of differentiation across natural populations and species [23–28]. A recent, large-

scale analysis of threespine sticklebacks (*Gasterosteus aculeatus*) showed that islands of

differentiation were more likely to arise in low recombination regions when gene flow

occurred between populations [29]. There is also a significant impact of divergence time; in

recently separated populations the differentiation landscape is most likely to reflect

selective sweeps. Then, as divergence accumulates, genomic architecture is expected to

play an increasingly important role in generating these genomic islands [23].

Widespread continental species are excellent models for studying how demography and

the environment shape genetic and phenotypic variation among populations, due to their

large effective population sizes and ecologically varied ranges. Insight into the evolutionary

history of such species can be gained if genetic variation is characterized across much of its

geographical range. Cross-population comparisons of genetic variation can then be utilised

to make inferences about phylogeography, levels of gene flow between populations and

how adaptation to different environmental and ecological conditions occurs [30]. The first

large-scale studies were performed in humans - i.e. the HapMap Projects [31–33] which

characterized human genetic variation on different continents, with a view to determining

the feasibility of association mapping studies. Similar studies have been conducted in

domesticated species and their wild ancestors [34–36], and in model organisms [37,38].

More recently, there is a growing appreciation that HapMap-type studies are useful for

studying signatures of selection and adaptation in natural populations of species with large

effective population sizes and high levels of gene flow [39–42].

The European great tit (*Parus major major*) is an excellent model for ecological and

evolutionary studies [43]. As is the case with several avian species which are amenable to

long-term study [44], a wealth of ecological data exists across multiple great tit populations

[45–47], enabling informed hypotheses about selection to be tested in this system.

Phyleogeographic research using mitochondrial DNA suggests that this species has recently

expanded across its European range, possibly from a single refugium in South-East Europe

[48]. Contemporary populations are characterised by large effective population sizes and

low levels of genetic differentiation [49,50]. However, these previous cross-population

molecular studies have relied on a modest number of microsatellite loci and mitochondrial

DNA, making the detection of genomic regions under selection impossible. The genome of

the great tit has recently been sequenced [51], and a high density panel of *ca* 500,000 SNP

markers has been developed [52]. A recent study of two European populations using this

marker panel suggests that rapid adaptation has occurred at the genomic and phenotypic

levels, with pronounced selection on morphology [53].

Here, we perform a HapMap study of 647 unrelated individuals across 29 populations (Fig.

1), to examine how genomic architecture, natural selection and population history have

shaped patterns of genomic variation across recently colonised European great tit

populations. Using a large SNP panel typed across all individuals, we first characterise

genome-wide patterns of variation within and among populations, in order to infer

population history. We then examine how variation is partitioned across the genome, and

test the hypothesis that highly divergent genomic regions have arisen in genomic regions of

low recombination [13,14]. Finally, we examine how genomic divergence accumulates

along the colonisation route of this species, with the aim of inferring how recent natural

selection and demography drive variation across the genome in the wild.

Figure 1 Sampling locations of great tit populations. Population names and sample sizes are given in Table S1, and numbers on the map correspond to the "code" column in Table S1.

Results and Discussion

Genetic diversity and population history

Sampling locations and sample sizes for each population are given in Table S1. Levels of

genetic diversity (π_{SNP}) were generally high, but we observed substantial differences

among populations (Fig. 2A). Similarly, LD declined rapidly with genomic distance in all

populations, reaching baseline levels within ~5kb in all populations, but also varied among

populations (Fig. S1). Highest levels of LD (and lowest levels of genetic diversity) were

observed in the Mediterranean island populations of Crete (Greece) and Sardinia (Italy),

with lowest levels of LD in central and western Europe (Fig. S1). This is consistent with

reduced effective population size in these island populations, either as result of the

colonisation process or more recent bottlenecks, along with low levels of subsequent gene

flow from the continent to the islands [54,55].

Mean genome-wide F_{ST} between all pairs of European great tit populations was 0.015, with

no significant pattern of isolation-by-distance (Mantel test; r = 0.13, p = 0.18; Fig. S2).

Instead, the highest levels of F_{ST} were found in comparisons involving the Mediterranean

island populations of Corsica (France), Sardinia and Crete (Fig. S2). Admixture analysis was

consistent with this pattern (Fig. S3); the K = 2 analysis assigned individuals in Sardinia

and Corsica to one genetic cluster, and the remaining populations to the second. Thus, it is

likely that much of the genetic structure between European great tit populations is a result

of genetic drift in these small island populations. Admixture analysis also revealed some

structure between (mainly peripheral) mainland and larger island populations. At K = 3

(the model that best fitted the great tit data; Fig. S4), Spain was separated from the rest of

mainland Europe. Increasing values of K resulted in the separation of populations in

Scotland (K = 4), Sardinia (from Corsica; K = 5), southern France (K = 6), Crete (K = 7) and

England (K = 8). The Admixture output at K = 8 is displayed in Fig. 2C as this gives the most

detailed picture of genetic structure among European great tit populations. Further

increases in K did not generate patterns of structure that corresponded to geographical

variation (Fig. S3), and were increasingly less well supported (Fig. S4). Thus, even with

hundreds of thousands of markers Admixture was unable to separate many of the

European populations, confirming that levels of divergence are extremely low [51]. PCA

largely corroborated the Admixture results, with PC1 separating Corsica and Sardinia from

the remaining populations, PC2 separating Spain, while PC3 and PC4 separated Scotland,

England, Corsica, Sardinia and Crete (Fig. S5).

Maximum likelihood analyses implemented in TreeMix showed that a model with no

migration explained 97.8% of variance in relatedness between populations [56]; increasing

the number of migration events substantially improved the percentage of relatedness

explained, up to 99.7% when 10 migration events were fitted (Fig. S6). In Figure 3 we

display the maximum likelihood tree with three migration events, after which the variance

in relatedness explained plateaued when more migration events were added (Fig. S6). The

tree was generally characterised by short branch lengths, with the exception of the island

populations of Sardinia and Crete, which were grouped with the population from mainland

Italy (Fig. 3). Thus, the TreeMix analysis is consistent with large populations and low

overall genomic divergence, with the exception of the Mediterranean island populations.

However, much (though not all) of the grouping that did occur among continental

populations made geographical sense, with populations from Finland and Estonia grouped

together, as were some populations from South-East Europe, and populations from

England and Scotland (Fig. 3). Interestingly, TreeMix grouped the Spanish and Corsican

populations, which is consistent with previous subspecies descriptions of European great

tits [57]. The three fitted migration edges all involved Sardinia, with migration from

eastern Europe to Sardinia, and from Sardinia to Corsica and Spain (Fig. 3).

We next tested the hypothesis that great tits colonised Europe from a single refugium in

South-East Europe. This scenario has been suggested before [48], but due to the low

number of genetic markers available there has been limited power with which to test this

hypothesis. Using our genome-wide panel of SNP markers, we compared genetic and

geographic distance between each population and the proposed refugial populations.

Because of the elevated structure in Corsica, Sardinia and Crete (Fig. S2), we excluded

comparisons involving these populations. We found that F_{ST} was significantly related to

distance from Turkey (r = 0.81, p < 0.001; Fig. 2B) and Romania/Bulgaria (r = 0.44, p =

0.001). The same relationship was not found for alternative potential refugial populations

[58] in Spain (r = -0.09, p = 0.55), or southern Italy (r = 0.04, p = 0.77). Our results

therefore lend empirical support to the hypothesis [48] that great tits colonised Europe

primarily from a single refugium in the south-east. Clearly, although our sampling was

extensive, it is not exhaustive, and more fine-scaled sampling in eastern Europe would be

required to determine the extact location and extent of refugial great tit populations.

Sampling in North Africa would also be useful to determine whether further refugia exist,

and to quantify the extent of admixture between European and African great tit

populations.

Figure 2 Genetic diversity and structure in European Great tit populations. **A** Nucleotide
diversity within each population. **B** Pairwise F_{ST} in relation to geographic distance from the
Turkey, only including comparisons involving Turkey. **C** Output from Admixture analysis at
$K = 8$. Population details can be found in Table S1.

Figure 3 Maximum likelihood tree inferred by TreeMix, allowing three migration events. The two migration events (arrows) are coloured according to their weight (red = higher migration), and horizontal branch lengths are proportional to the amount of genetic drift that has occurred along the branch. A population of the great tit's sister species, *Parus minor*, was used as an outgroup. Population details are given in Table S1.

Genomic landscapes of differentiation

It is likely that many, and perhaps the majority, of wild populations are characterised by

highly heterogeneous patterns of differentiation across the genome [27]. This is the case in

European great tits - despite the extremely low average F_{ST}, we found genomic regions

with very high levels of genomic structure (maximum F_{ST} for 10kb and 500kb windows

was 0.98 and 0.07, respectively). To examine how landscapes of genomic divergence have

formed along the colonisation route of European great tits, we calculated windowed F_{ST} in

500kb bins between each population and the proposed refugial population in Turkey. We

found that F_{ST} varied markedly across the genome in all comparisons (Fig. 4; Fig. S7).

Outlier regions (windows with standardized F_{ST},hereafter zF_{ST}, > 10) were found in all

comparisons apart from Crete and Sardinia, in which overall levels of divergence were

highest, with some outlier regions found across multiple comparisons (Fig. 4). Our results

suggest, therefore, that genomic islands of differentiation can and do arise even among

recently separated populations.

Considering all populations, F_{ST} calculated in 500kb windows was strongly negatively

correlated with local variation in recombination rate (Spearman correlation, r = -0.50, p <

0.001). Although recombination rate and gene density were positively correlated, (r = 0.16,

p < 0.001), F_{ST} was only moderately correlated with gene density in 500kb windows (r = -

0.09, p < 0.001), and this correlation became weak when calculated in 10kb windows (r = -

0.01, p < 0.001). Thus it appears that the negative relationship between F_{ST} and

recombination rate is not driven by gene density. Examining how the relationship between

genomic differentiation and recombination varied among populations revealed that F_{ST}

was negatively related to recombination rate in almost all comparisons with Turkey (Fig.

5). The relationship between F_{ST} and recombination rate was generally weak, but in a

handful of populations this relationship was substantially stronger - most notably in the

island populations of Corsica, Sardinia and Crete, and in the peripheral mainland

populations of England, Scotland, Spain, Finland and France, which are the populations that

are likely most susceptible to drift (Fig. 5).

As a further examination of how allele frequency variation and haplotype structure may be

shaped by neutral and adaptive processes, we compared F_{ST} and recombination rate with

Rsb - a measure which aims to detect regions of the genome under positive selection by

comparing extended haplotype homozygosity profiles between populations, and is

expected to be less sensitive to variation in recombination rate because extended

haplotypes should be present in both populations when recombination frequency is lower

[59]. Specifically, we compared the distributions of these statistics in 500kb windows using

three populations in the Netherlands, Finland and Spain, with the aim of exploring how *Rsb*

varies with recombination rate and with F_{ST}. We found that, across the genome,

correlations between absolute *Rsb* and recombination rate were only slightly weaker

(Spearman rank r between -0.25 and -0.20) compared to those between F_{ST} and

recombination for the same pairwise comparisons (r between -0.36 and -0.20), and that the

regions of high F_{ST} in these comparisons tended to be in these same regions of low

recombination and high *Rsb* (Fig. S8). Notably, though, there were several cases where

windows with extreme *Rsb* values occurred in regions of high recombination, and most

high F_{ST} regions in high recombination regions were also *Rsb* outliers (Fig. S8). Below we

explore the possibility that such regions are strong candidates for recent adaptive

evolution.

Background selection is a key driver of genomic differentiation in birds [20,21], and other

organisms [60], but is not expected to play a major role in driving islands of differentiation

over the timescales (tens of thousands of years) relevant to this study [23,29]. On the other

hand, differential effects of drift operating across the genome could contribute

substantially to the highly heterogeneous patterns of differentiation observed in European

great tits. Our finding that the relationship between recombination rate and F_{ST} is strongest

in populations experiencing the greatest levels of drift (e.g. islands; Fig. 5) is consistent

with this - low recombining regions have reduced effective population size and increased

rates of lineage sorting due to drift. Drift has been suggested to be a driver of

heterogeneous genomic landscapes in other systems [16,17], and our study suggests that it

may play a key role in shaping genomic structure in widespread, continental species.

Outlier regions of very high differentiation (zF_{ST} > 10) mostly occurred in areas of the

genome with low recombination rates (Fig. 5) and the recombination rate of outlier regions

was (just) significantly lower than the genome-wide average (Wilcoxon test, P = 0.048). Of

the 11 outlier regions, nine were found in only one or two comparisons, while the other

two were found in 12 and 10 comparisons, respectively (Table S2). We hereafter refer to

outlier regions found in one or two comparisons as "population-specific" outlier regions,

and to those found in more than two comparisons as "shared" regions.

Regions of high differentiation that are not shared among populations could potentially

arise as a result of recent drift, but are also candidate regions for recent positive selection

[23]. Several lines of evidence suggest that this may be the case in European great tits.

Firstly, population-specific regions tended to occur in regions of higher recombination than

shared outliers, although low sample size prevents a formal statistical analysis of this.

Second, population-specific outliers tended to be found in the most peripheral European

great tit populations, with three found in Scotland, and two in England, Spain and Finland;

the remaining outlier regions were found in comparisons involving the Czech republic,

Russia, Vlieland (Netherlands) and Belgium (Table S2). Notably, these populations are not

the lowest in genetic diversity and therefore drift effects are not necessarily stronger in

these peripheral populations. Third, the Rsb tests, which indicate which population has

undergone a selective sweep, also tend to find evidence of positive selection in the more

peripheral population, as expected under an adaptive evolution scenario (Fig. S8). This

directionality cannot be assessed with F_{ST}, because it is a combined measure of allele

frequency differences. Of course, across the genome, both positive selection and linked

background selection are likely to be operating. Observational and experimental research

shows that adaptation at range edges is a key feature shaping divergence among recently

colonised and expanding populations [61–63]. Population-specific regions appeared less

likely to be in regions of low recombination than shared regions (Fig. 5), although the small

number of regions precluded testing this hypothesis formally. Thus, it is likely that genomic

architecture plays a key role in determining how both selection and drift have shaped

genomic variation across the recent evolutionary history of European great tits.

To further explore how selection may have shaped variation in F_{ST} outlier regions, we

estimated levels of nucleotide and haplotype diversity within these regions. Nucleotide

diversity (π_{SNP}) in outlier regions varied from 0.21 to 0.48, and diversity in these regions

was significantly lower than the genome-wide average (Wilcoxon test, P = 0.018; Fig. S9A).

However, there appeared to be little difference in nucleotide diversity between shared and

population-specific regions (Fig. S9A). Haplotype diversity varied substantially among

regions, with haplotype richness ranging from 72 to 1033. Both haplotype richness and

marker density in shared regions tended to be lower than those in population-specific

regions (Fig. S9B), consistent with lower recombination frequency. A detailed examination

of haplotype structure in one shared and one population-specific region is displayed in

Figure S10. The population-specific outlier region (to Finland, situated on chromosome 1A)

was characterised by a complex structure, with a single haplotype at high frequency in

Finland compared to other populations, indicating a population-specific selective sweep

and high background diversity (Fig. S10). In contrast, the shared region on chromosome 2

was much less complex, demonstrating high frequency haplotypes that are found across a

range of populations. Our data therefore suggest that examining patterns of haplotype

diversity in outlier regions may help to separate recent episodes of positive selection from

drift and background selection (Figs S9, S10).

Potential candidate genes found within shared and population-specific outlier regions are

displayed in Table S2. Perhaps most notable among these is *COL4A5*, a gene found to be

associated with bill length, and under selection between populations in England and the

Netherlands, in a recent great tit study [53]. Here we found that the *COL4A5* region is an F_{ST}

outlier in England and Scotland, but not in any other European populations (Table S2). UK

great tits have been described as a separate subspecies based on beak shape [64], and our

results here, combined with previous results, suggest that this divergence is the result of

recent natural selection in the UK [53]. Another notable candidate gene potentially

involved in beak morphology, and previously found to be under selection in UK great tits is

BMPR1A, which plays a key role in palate development [65] and in this study was found in

an outlier region in Scotland. Other candidate morphology and obesity genes in the

population-specific outlier regions in the UK included *PPP1CB*, which may play a role in

adipogenesis [66] and *GHITM*, which appears to have been subject to natural selection in

human pygmy populations [67]. Thus, morphological traits may frequently be involved in

adaptation in great tits.

In addition to morphological candidates in the UK, we found outlier regions specific to cold

populations in Scotland, Finland and Russia (Table S2), containing at least one candidate

gene for thermal stress (*CDKN1B*) [68]. Other genomic outlier regions contained potential

candidate genes associated with malaria infection (*MRPL33*) [69] and colour variation

(*SOX10*) [70]. This is thus far an exploratory analysis, and we are therefore reluctant to

speculate whether these candidate genes are genuine targets for natural selection, and

more reluctant still to speculate as to how selection might be driving variation at these

regions. Regardless, these candidates will provide useful starting points for future genomic

and ecological investigation.

HapMap style projects have been hugely informative in shaping our understanding of how

natural selection operates in humans and other model species [31,38]. This study is one of

the largest to date of genomic variation in a wild vertebrate, which has helped to reveal the

evolutionary history of great tits, and to identify candidate genes and traits that may have

been involved in adaptation during and/or after postglacial recolonisation. Further, this

work will form the foundation of many future analyses. Clearly, we have only touched on

haplotype-based methods to infer adaptation here, and this will be the subject of future

work. Environmental association approaches are also highly suited to detecting adaptation

in widespread continental species [71,72], and further work will test how variation in the

environment has shaped patterns of genomic variation in great tits [73]. This combination

of environmental and genomic data in species such as great tits, in which a wealth of

ecological and genomic resources are available, is likely to generate interesting insights

into the the genetic and phenotypic basis of natural selection.

Figure 4 Landscapes of relative genomic differentiation in European great tit populations.

zF_{ST} across the genome is averaged in 500kb windows, with each panel displaying a

pairwise comparison with the proposed refugial population in Turkey. Red lines represent

F_{ST} outliers (windows with mean F_{ST} values at least 10 standard deviations greater than

the global mean for that comparison) shared across more than two comparisons (solid red

lines), or specific to one or two comparisons (dashed red lines). Mean, untransformed F_{ST}

values are given in the top-right of each panel, and are fully displayed in Fig. S7.

Figure 5 Genomic differentiation and recombination rate variation in European great tit populations. Each point is the mean of a 500kb window, with each panel displaying a pairwise comparison with the proposed refugial population in Turkey. Coloured points represent F_{ST} outliers (mean standardized F_{ST} values of $zF_{ST} > 10$) shared across more than two comparisons (red), or specific to one or two comparisons (dark blue). Solid and dotted lines represent median and 99% quantiles of F_{ST} windows from bins of 0.1 log cM/Mbp.

Materials and Methods

Sampling and molecular methods

Samples were collected from 29 populations from 22 regions across Europe (Fig. 1; Table S1). Samples were pooled into regions either based on geographical proximity (e.g. Cambridge and Wytham woods), or based on sample size (e.g. Romania and Bulgaria). An exploratory analysis considering all sampled populations separately yielded virtually identical results to those shown here, and in no cases did we observe substructure within pooled populations in our Admixture analyses (Fig. S3).

Birds were trapped from nest boxes, or using mist nets, and ringed with a uniquely numbered aluminium ring. Blood was taken via brachial or tarsal venipuncture, and stored in either 1 ml Cell Lysis Solution (Gentra Puregene Kit, Qiagen, USA), Queen's buffer, or absolute ethanol. All samples were genotyped using a custom-made Affymetrix® great tit 650K SNP chip at Edinburgh Genomics (Edinburgh, United Kingdom), following the approaches outlined in [52]

Analyses

Unless stated otherwise, all population genetic statistics were calculated in PLINK version 1.9 [74], and downstream analysis and plotting was carried out in R version 3.3 [75]. For population genetic analyses, we used the filtering approaches outlined in [53]. Briefly, we randomly removed individuals from pairs with relatedness values > 0.4, and for demographic analyses we used a LD pruned dataset (based on VIF > 0.2), with SNPs

associated with an inversion on chromosome 1A [**???**] removed. After filtering, a total of

647 samples typed at 483888 SNPs were retained for analysis.

In each population, we estimated LD (R^2) for each pair of markers within 50kb on the same

chromosome, and compared this to physical distance between marker pairs. We calculated

observed heterozygosity for each SNP and population using a reduced SNP dataset, which

was pruned based on LD to remove all markers with R^2 > 0.1, then thinned with a

probability of retaining each variant of 0.25. We calculated genome-wide (mean) F_{ST}

between each pair of populations using the pruned and thinned dataset described above.

Pairwise F_{ST} was compared to geographic distance between populations using Mantel tests,

implemented in the Ecodist package in R [76]. We tested whether genetic structure was

related to distance from candidate refugial populations (in Romania/Bulgaria, Turkey,

Spain and Italy), using Pearson correlations. We also estimated population structure using

Admixture version 1.3, with default settings [77]. We varied values of K from one to ten; by

which point increasing values of K provided no informative information about population

structure (see results). Model support for each value of K was estimated by calculating 5-

fold cross-validation error. Finally, we visualised the evolutionary history among European

great tit populations by generating a maximum likelihood tree in TreeMix version 1.13

[56]. We rooted the tree using a sample of *P. minor* individuals sampled from Amur, Russia

[52]. We fitted models allowing for range of migration events (0-10), and used a window

size of 500 SNPs [56]. To assess model fit, we calculated the proportion of variance in

relatedness between populations explained by each model [56].

Recombination rates at each locus were estimated by comparing the location of SNPs on

the genome assembly (v1.1) with their location on the great tit linkage map [78]. Previous

linkage mapping, using a lower density SNP chip, was independently carried out in UK and

Netherlands great tit populations and the two maps were almost identical [78]. For the

purposes of this analysis, we used SNPs and marker intervals from the UK comprehensive

map. A total of 2706 SNPs were located on both the genome assembly and the linkage map.

Thus a positon in Mb and cM of each of these SNPs is known. All other SNPs on the HD chip

have a physical (Mb) position but no known linkage map position. The great tit genome

v1.1 is 1.02 Gb long, so the average physical interval between mapped SNPs is ~376Kb. The

linkage map position of each unmapped SNP was estimated by interpolation; by taking the

closest mapped SNP in either direction, and assuming a constant recombination rate in the

interval between those SNPs. For example, an unmapped SNP with physical position 1.4Mb,

flanked by mapped SNPs at 1.0Mb/0.5cM and 2.0Mb/1.0cM, would be estimated to be

located at 0.5 + (1.4-1.0)/(2.0-1.0) * (1.0-0.5) = 0.7cM. Having interpolated cM position of

every SNP, the local recombination rate was calculated as the cM interval spanned by the

nearest neighbouring SNPs, divided by the physical distance (bp) spanned by those same

neighbouring SNPs. In other words, for the ith SNP, the recombination rate is estimated as

the linkage distance between the i-1th and i+1th SNP, divided by the physical distance

between the i-1th and i+1th SNP. For downstream analyses, local recombination rates were

estimated by averaging across all SNPs in each 500kb window. We calculated gene density

in 10kb and 500kb windows using custom R scripts and the annotated great tit genome (v

1.1) [51].

We examined the genomic landscape of differentiation across European great tit

populations by calculating F_{ST} in 10kb and 500kb bins, using python scripts obtained from

Github (https://github.com/simonhmartin/genomics_general). We did not estimate d_{xy}, as

this parameter is difficult to estimate accurately from single SNP loci [14]. We also

calculated standardised F_{ST} (zF_{ST}) by mean-centring windowed values and dividing them

by the standard deviation among windows. We defined outlier regions as 500kb bins with

zF_{ST} values greater than ten. Finally, Rsb [59] was calculated for three pairwise

comparisons (in Spain, Finland and the Netherlands), using the R package Rehh [79], and

this averaged into 500kb windows. Unlike F_{ST}, the Rsb statistic gives an indication of which

population an adaptively important haplotype is under positive selection in, and is thought

to be less sensitive to local recombination rates [59]. Rsb estimation requires phased

genotype data, so phasing was performed using shapeIT2 v 2.r837 [80]. The –duohmm

argument was used to ensure that family information, where available, was used to

improve the accuracy of phasing. The –effective-size parameter was set at 500,000

reflecting the large effective population size of European great tit populations [51]. Local

recombination rates (measured in cM/Mbp) were used in the map files. Comparisons

between F_{ST}, Rsb, recombination rate and gene density were carried out using Spearman

rank correlation (as all statistics were highly skewed).

Ethics Statement

Great tit DNA samples were provided by researchers studying hole-nesting great tit

populations with permissions and licenses to study and sample birds granted by the

following organisations: Bundesministerium für Wissenschaft und Forschung (Ref. II/3b

Gentechnik und Tierversuche), Czech Republic Ministry of Education Youth and Sports

(MSMT-56147/2012-310), Dutch legal entity Dier Experimenten Comissie (NIOO 14.11,

NIOO 07.04), Dutch Central Authority for Scientific Procedures on Animals (AVD-

801002017831), Estonian Ministry of Agriculture - Animal Procedures Committee (License

108), Ethical Committee of the Canton of Bern, Ethics Committee for animal

experimentation of Languedoc Roussillon CEEA-LR (APAFIS#8608-2017012011062214),

Ethics Committee of Lomonosov Moscow State University (Approval numbers 26ch, 37zh,

49zh, 59zh, 88zh), Federal Office for the Environment - Switzerland (License 234), Flemish

Ministry - Agency for Nature and Forest, Generalitat Valenciana - Conselleria de Medi

Ambient, Hellenic Ministry of Environment and Energy, Istituto Superiore per la Protezione

e la Ricerca Ambientale - ISPRA (Law 157/1992 Art. 4 (1) and Art. 7 (5)), Mehmet Akif

Ersoy University - Local Ethical Committee on Animal Experiments (93773921-9), Middle

Danube Valley Inspectorate for Environmental Protection, Nature Conservation and Water

Management (43355-1/2008), Regierung Oberbayern (55.2-1-54-2532-140-11),

Regierungspräsidium Freiburg - Departments for Animal Welfare and Nature Conservation,

Regional State Administrative Agency of Southern Finland (ESAVI/1650/04.10.03/2012,

ESAVI-1804/04.10.03/2011), Romanian Academy of Sciences (Permit number 2257),

Swedish Committee for Experiments on Animals (Licence number C 108/7), UK Home

Office (License PPL 30/2409), University of Oxford - Department of Zoology Local Ethical

Review Committee, University of Glasgow AWERB committee, Veterinary Office of the

Canton of Zurich (License 195/2010)

Samples were imported to the UK under DEFRA general import license IMP/GEN/2011/01,

according to regulations outlined by the UK Animal Health and Veterinary Laboratories

Agency.

Acknowledgements

This work was supported by grants from the Natural Environment Research Council (grant

NE/J012599/1 to J.S. and B.C.S) and the European Research Council (grant 202487 to J.S.

and grant 339092 – E-Response to MEV) L.G.S was supported by fellowships from the

Edward Grey Institute for Ornithology and the BBSRC (BB/N011759/1). We thank Claire

Bloor, Geoff Scopes and Alessandro Davassi of Affymetrix for their help during the chip

design and genotyping calling processes. Richard Talbot and Alison Downing of Edinburgh

Genomics provided the genotyping service.

Data Availability

All data and code to reproduce the results are available on Github:

https://github.com/lgs85/SpurginBosse_Hapmap

References

1. Harris H. Enzyme polymorphisms in man. Proceedings of the Royal Society B: Biological

Sciences. 1966;164: 298–310. doi:10.1098/rspb.1966.0032

2. Lewontin RC, Hubby JL. A molecular approach to the study of genic heterozygosity in

natural populations. II. Amount of variation and degree of heterozygosity in natural

populations of <i>Drosophila pseudoobscura</i>. Genetics. 1966;54: 595–609.

doi:10.1111/j.1601-5223.1973.tb01163.x

3. Hubby JL, Lewontin RC. A molecular approach to the study of genic heterozygosity in

natural populations. I. The number of alleles at different loci in Drosophila pseudoobscura.

Genetics. 1966;54: 577–594.

4. Nei M. Selectionism and Neutralism in Molecular Evolution. Molecular Biology and

Evolution. 2005;22: 2318–2342. doi:10.1093/molbev/msi242

5. Hartl DL, Clark AG. Principles of population genetics. Sinauer associates Sunderland, MA;

1997.

6. Feder JL, Egan SP, Nosil P. The genomics of speciation-with-gene-flow. Trends in

Genetics. 2012;28: 342–350. doi:10.1016/J.TIG.2012.03.009

7. Lanfear R, Kokko H, Eyre-Walker A. Population size and the rate of evolution. Trends in

Ecology & Evolution. 2014;29: 33–41. doi:10.1016/J.TREE.2013.09.009

8. Turner TL, Hahn MW, Nuzhdin SV, Mitchison G, Spencer F. Genomic Islands of Speciation

in Anopheles gambiae. PLoS Biology. 2005;3: e285. doi:10.1371/journal.pbio.0030285

9. Nadeau NJ, Whibley A, Jones RT, Davey JW, Dasmahapatra KK, Baxter SW, et al. Genomic

islands of divergence in hybridizing Heliconius butterflies identified by large-scale targeted

sequencing. Philosophical Transactions of the Royal Society of London B: Biological

Sciences. 2011;367. Available: doi: 10.1098/rstb.2011.0198

10. Renaut S, Grassa CJ, Yeaman S, Moyers BT, Lai Z, Kane NC, et al. Genomic islands of

divergence are not affected by geography of speciation in sunflowers. Nature

Communications. 2013;4: 1827. doi:10.1038/ncomms2833

11. Poelstra JW, Vijay N, Bossu CM, Lantz H, Ryll B, Muller I, et al. The genomic landscape

underlying phenotypic integrity in the face of gene flow in crows. Science. 2014;344: 1410–

1414. doi:10.1126/science.1253226

12. Wolf JBW, Ellegren H. Making sense of genomic islands of differentiation in light of

speciation. Nature Reviews Genetics. 2017;18: 87–100. doi:10.1038/nrg.2016.133

13. Noor MAF, Bennett SM. Islands of speciation or mirages in the desert? Examining the

role of restricted recombination in maintaining species. Heredity. 2009;103: 439–444.

doi:10.1038/hdy.2009.151

14. Cruickshank TE, Hahn MW. Reanalysis suggests that genomic islands of speciation are

due to reduced diversity, not reduced gene flow. Molecular Ecology. 2014;23: 3133–3157.

doi:10.1111/mec.12796

15. Turner TL, Hahn MW. Genomic islands of speciation or genomic islands and speciation?

Molecular Ecology. 2010;19: 848–850. doi:10.1111/j.1365-294X.2010.04532.x

16. Ma T, Wang K, Hu Q, Xi Z, Wan D, Wang Q, et al. Ancient polymorphisms and divergence

hitchhiking contribute to genomic islands of divergence within a poplar species complex.

Proceedings of the National Academy of Sciences of the United States of America. National

Academy of Sciences; 2018;115: E236–E243. doi:10.1073/pnas.1713288114

17. Campagna L, Gronau I, Silveira LF, Siepel A, Lovette IJ. Distinguishing noise from signal

in patterns of genomic divergence in a highly polymorphic avian radiation. Molecular

Ecology. 2015;24: 4238–51. doi:10.1111/mec.13314

18. Martin SH, Dasmahapatra KK, Nadeau NJ, Salazar C, Walters JR, Simpson F, et al.

Genome-wide evidence for speciation with gene flow in Heliconius butterflies. Genome

Research. 2013;23: 1817–1828. doi:10.1101/gr.159426.113

19. Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, et al. Linked selection and

recombination rate variation drive the evolution of the genomic landscape of

differentiation across the speciation continuum of Ficedula flycatchers. Genome research.

2015;25: 1656–1665. doi:10.1101/gr.196485.115

20. Vijay N, Weissensteiner M, Burri R, Kawakami T, Ellegren H, Wolf JBW. Genomewide

patterns of variation in genetic diversity are shared among populations, species and

higher-order taxa. Molecular Ecology. 2017;26: 4284–4295. doi:10.1111/mec.14195

21. Van Doren BM, Campagna L, Helm B, Illera JC, Lovette IJ, Liedvogel M. Correlated

patterns of genetic diversity and differentiation across an avian family. Molecular Ecology.

2017;26: 3982–3997. doi:10.1111/mec.14083

22. Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, et al. Comparative genomics reveals insights

into avian genome evolution and adaptation. Science. 2014;346: 1311–1320.

23. Burri R. Dissecting differentiation landscapes: a linked selection's perspective. Journal

of Evolutionary Biology. 2017;30: 1501–1505. doi:10.1111/jeb.13108

24. Ellegren H, Wolf JBW. Parallelism in genomic landscapes of differentiation, conserved

genomic features and the role of linked selection. Journal of Evolutionary Biology. 2017;30:

1516–1518. doi:10.1111/jeb.13113

25. Jiggins CD, Martin SH. Glittering gold and the quest for Isla de Muerta. Journal of

Evolutionary Biology. 2017;30: 1509–1511. doi:10.1111/jeb.13110

26. Lohse K. Come on feel the noise - from metaphors to null models. Journal of

Evolutionary Biology. 2017;30: 1506–1508. doi:10.1111/jeb.13109

27. Ravinet M, Faria R, Butlin RK, Galindo J, Bierne N, Rafajlović M, et al. Interpreting the

genomic landscape of speciation: a road map for finding barriers to gene flow. Journal of

Evolutionary Biology. 2017;30: 1450–1477. doi:10.1111/jeb.13047

28. Perrier C, Charmantier A. On the importance of time scales when studying adaptive

evolution. Evolution Letters. John Wiley & Sons, Ltd; 2019;3: 240–247.

29. Samuk K, Owens GL, Delmore KE, Miller SE, Rennison DJ, Schluter D. Gene flow and

selection interact to promote adaptive divergence in regions of low recombination.

Molecular Ecology. 2017;26: 4378–4390. doi:10.1111/mec.14226

30. Perrier C, Rougemont Q, Charmantier A. Demographic history and genomics of local

adaptation in blue tit populations. Evol Appl. John Wiley & Sons, Ltd; 2020;13: 1145–1165.

31. Altshuler D, Donnelly P. A haplotype map of the human genome. Nature. 2005;437:

1299–1320. doi:10.1038/nature04226

32. Frazer KA, Ballinger DG, Cox DR, Hinds DA, Stuve LL, Gibbs RA, et al. A second

generation human haplotype map of over 3.1 million SNPs. Nature. 2007;449: 851–861.

doi:10.1038/nature06258

33. Altshuler DM, Gibbs RA, Peltonen L, Altshuler DM, Gibbs RA, Peltonen L, et al.

Integrating common and rare genetic variation in diverse human populations. Nature.

2010;467: 52–8. doi:10.1038/nature09298

34. Kijas JW, Lenstra JA, Hayes B, Boitard S, Porto Neto LR, San Cristobal M, et al. Genome-

wide analysis of the world's sheep breeds reveals high levels of historic mixture and strong

recent selection. PLoS Biology. 2012;10: e1001258. doi:10.1371/journal.pbio.1001258

35. Gibbs RA, Taylor JF, Van Tassell CP, Barendse W, Eversole KA, Gill CA, et al. Genome-

wide survey of SNP variation uncovers the genetic structure of cattle breeds. Science.

2009;324: 528–532. doi:10.1126/science.1167936

36. Chia J-M, Song C, Bradbury PJ, Costich D, Leon N de, Doebley J, et al. Maize HapMap2

identifies extant variation from a genome in flux. Nature Genetics. 2012;44: 803–807.

doi:10.1038/ng.2313

37. Lindblad-Toh K, Wade CM, Mikkelsen TS, Karlsson EK, Jaffe DB, Kamal M, et al. Genome

sequence, comparative analysis and haplotype structure of the domestic dog. Nature.

2005;438: 803–819. doi:10.1038/nature04338

38. Kirby A, Kang HM, Wade CM, Cotsapas C, Kostem E, Han B, et al. Fine mapping in 94

inbred mouse strains using a high-density haplotype resource. Genetics. 2010;185: 1081–

95. doi:10.1534/genetics.110.115014

39. Horton MW, Hancock AM, Huang YS, Toomajian C, Atwell S, Auton A, et al. Genome-wide

patterns of genetic variation in worldwide Arabidopsis thaliana accessions from the

RegMap panel. Nature Genetics. 2012;44: 212–216. doi:10.1038/ng.1042

40. Alonso-Blanco C, Andrade J, Becker C, Bemm F, Bergelson J, Borgwardt KM, et al. 1,135

Genomes Reveal the Global Pattern of Polymorphism in Arabidopsis thaliana. Cell.

2016;166: 481–491. doi:10.1016/j.cell.2016.05.063

41. Williamson RJ, Josephs EB, Platts AE, Hazzouri KM, Haudry A, Blanchette M, et al.

Evidence for widespread positive and negative selection in coding and conserved

noncoding regions of Capsella grandiflora. PLoS Genetics. 2014;10: e1004622.

doi:10.1371/journal.pgen.1004622

42. Mueller JC, Kuhl H, Boerno S, Tella JL, Carrete M, Kempenaers B. Evolution of genomic

variation in the burrowing owl in response to recent colonization of urban areas.

Proceedings of the Royal Society B: Biological Sciences. The Royal Society; 2018;285:

20180206. doi:10.1098/rspb.2018.0206

43. Gosler A. The great tit. Hamlyn Species Guides; 1993.

44. Culina A, Adriaensen F, Bailey LD, Burgess MD, Charmantier A, Cole EF, et al. Connecting

the data landscape of long-term ecological studies: The SPI-Birds data hub. J Anim Ecol.

John Wiley & Sons, Ltd; 2020;n/a.

45. Visser ME, Noordwijk AJv, Tinbergen JM, Lessells CM. Warmer springs lead to mistimed

reproduction in great tits (Parus major). Proceedings of the Royal Society B: Biological

Sciences. 1998;265: 1867–1870. doi:10.1098/rspb.1998.0514

46. Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LEB, Sheldon BC. Adaptive

phenotypic plasticity in response to climate change in a wild bird population. Science.

2008;320: 800–803. doi:10.1126/science.1157174

47. Dingemanse NJ, Bouwman KM, Pol M van de, Overveld T van, Patrick SC, Matthysen E, et

al. Variation in personality and behavioural plasticity across four populations of the great

tit Parus major. Journal of Animal Ecology. 2012;81: 116–126. doi:10.1111/j.1365-

2656.2011.01877.x

48. Kvist L, Ruokonen M, Lumme J, Orell M. The colonization history and present-day

population structure of the European great tit (Parus major major). Heredity. 1999;82:

495–502. doi:10.1038/sj.hdy.6885130

49. Kvist L, Martens J, Higuchi H, Nazarenko AA, Valchuk OP, Orell M. Evolution and genetic

structure of the great tit (Parus major) complex. Proceedings of the Royal Society B:

Biological Sciences. 2003;270: 1447–1454. doi:10.1098/rspb.2002.2321

50. Lemoine M, Lucek K, Perrier C, Saladin V, Adriaensen F, Barba E, et al. Low but

contrasting neutral genetic differentiation shaped by winter temperature in European

great tits. Biological Journal of the Linnean Society. 2016;118: 668–685.

doi:10.1111/bij.12745

51. Laine V, Gossmann T, Schachtschneider K, Garroway C, Madsen O, Verhoeven K, et al.

Evolutionary signals of selection on cognition from the great tit genome and methylome.

Nature Communications. 2016;7: 10474. doi:10.1038/ncomms10474

52. Kim J-M, Santure AW, Barton HJ, Quinn JL, Cole EF, Great Tit HapMap Consortium ME, et

al. A high-density SNP chip for genotyping great tit (Parus major) populations and its

application to studying the genetic architecture of exploration behaviour. Molecular

Ecology Resources. 2018;18: 877–891. doi:10.1111/1755-0998.12778

53. Bosse M, Spurgin L, Laine V, Cole E, Firth J, Gienapp P, et al. Recent natural selection

causes adaptive evolution of an avian polygenic trait. Science. 2017;358: 365–368.

doi:10.1126/science.aal3298

54. Postma E, Noordwijk AJ van. Gene flow maintains a large genetic difference in clutch

size at a small spatial scale. Nature. 2005;433: 65–68. doi:10.1038/nature03083

55. James JE, Lanfear R, Eyre-Walker A. Molecular Evolutionary Consequences of Island

Colonization. Genome Biology and Evolution. 2016;8: 1876–1888.

doi:10.1093/gbe/evw120

56. Pickrell JK, Pritchard JK. Inference of Population Splits and Mixtures from Genome-

Wide Allele Frequency Data. PLoS Genetics. 2012;8: e1002967.

doi:10.1371/journal.pgen.1002967

57. Clements J. The Clements checklist of the birds of the world. New York: Cornell

University Press; 2007.

58. Hewitt GM. Postglacial re-colonisation of European biota. Biological Journal of the

Linnean Society. 1999;68: 87–112.

59. Tang K, Thornton KR, Stoneking M. A new approach for using genome scans to detect

recent positive selection in the human genome. PLoS Biology. 2007;5: 1587–1602.

doi:10.1371/journal.pbio.0050171

60. Comeron JM. Background selection as null hypothesis in population genomics: insights

and challenges from Drosophila studies. Philosophical Transactions of the Royal Society B:

Biological sciences. The Royal Society; 2017;372. doi:10.1098/rstb.2016.0471

61. Sexton JP, McIntyre PJ, Angert AL, Rice KJ. Evolution and Ecology of Species Range

Limits. Annual Review of Ecology, Evolution, and Systematics. 2009;40: 415–436.

doi:10.1146/annurev.ecolsys.110308.120317

62. Hill JK, Griffiths HM, Thomas CD. Climate Change and Evolutionary Adaptations at

Species' Range Margins. Annual Review of Entomology. 2011;56: 143–159.

doi:10.1146/annurev-ento-120709-144746

63. Weiss-Lehman C, Hufbauer RA, Melbourne BA. Rapid trait evolution drives increased

speed and variance in experimental range expansions. Nature Communications. 2017;8:

14303. doi:10.1038/ncomms14303

64. Gosler AG. A comment on the validity of the British Great Tit Parus major newtoni.

Bulletin of The British Ornithologists' Club. 1999;119: 47–55. Available:

http://biostor.org/reference/111964

65. Baek J-A, Lan Y, Liu H, Maltby KM, Mishina Y, Jiang R. Bmpr1a signaling plays critical

roles in palatal shelf growth and palatal bone formation. Developmental Biology. 2011;350:

520–531. doi:10.1016/J.YDBIO.2010.12.028

66. Cho Y-L, Min J-K, Roh KM, Kim WK, Han BS, Bae K-H, et al. Phosphoprotein phosphatase

1CB (PPP1CB), a novel adipogenic activator, promotes 3T3-L1 adipogenesis. Biochemical

and Biophysical Research Communications. 2015;467: 211–217.

doi:10.1016/J.BBRC.2015.10.004

67. Migliano AB, Romero IG, Metspalu M, Leavesley M, Pagani L, Antao T, et al. Evolution of

the Pygmy Phenotype: Evidence of Positive Selection from Genome-wide Scans in African,

Asian, and Melanesian Pygmies. Human Biology. 2013;85: 251–284.

doi:10.3378/027.085.0313

68. Logan CA, Somero GN. Effects of thermal acclimation on transcriptional responses to

acute heat stress in the eurythermal fish Gillichthys mirabilis (Cooper). American Journal of

Physiology-Regulatory, Integrative and Comparative Physiology. 2011;300: R1373–R1383.

doi:10.1152/ajpregu.00689.2010

69. Videvall E, Cornwallis CK, Palinauskas V, Valkiūnas G, Hellgren O. The Avian

Transcriptome Response to Malaria Infection. Molecular Biology and Evolution. 2015;32:

1255–1267. doi:10.1093/molbev/msv016

70. Gunnarsson U, Kerje S, Bed'hom B, Sahlqvist A-S, Ekwall O, Tixier-Boichard M, et al. The

Dark brown plumage color in chickens is caused by an 8.3-kb deletion upstream of SOX10.

Pigment Cell & Melanoma Research. 2011;24: 268–274. doi:10.1111/j.1755-

148X.2011.00825.x

71. Coop G, Witonsky D, Di Rienzo A, Pritchard JK. Using environmental correlations to

identify loci underlying local adaptation. Genetics. 2010;185: 1411–23.

doi:10.1534/genetics.110.114819

72. Frichot E, Schoville SD, Bouchard G, François O. Testing for associations between loci

and environmental gradients using latent factor mixed models. Molecular Biology and

Evolution. 2013;30: 1687–1699. doi:10.1093/molbev/mst063

73. Salmón P, Jacobs A, Ahrén D, Biard C, Dingemanse NJ, Dominoni DM, et al. Repeated

genomic signatures of adaptation to urbanisation in a songbird across europe. bioRxiv.

2020. p. 2020.05.05.078568.

74. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MA, Bender D, et al. PLINK: A Tool

Set for Whole-Genome Association and Population-Based Linkage Analyses. The American

Journal of Human Genetics. 2007;81: 559–575. doi:10.1086/519795

75. R Development Core Team. R: A Language and Environment for Statistical Computing

[Internet]. Team RDC, editor. 2011. doi:10.1007/978-3-540-74686-7

76. Goslee SC, Urban DL. The ecodist Package for Dissimilarity-based Analysis of Ecological

Data. Journal of Statistical Software. 2007;22: 1–19. doi:10.18637/jss.v022.i07

77. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in

unrelated individuals. Genome Research. 2009;19: 1655–1664. doi:10.1101/gr.094052.109

78. Oers K van, Santure AW, De Cauwer I, Bers NE van, Crooijmans RP, Sheldon BC, et al.

Replicated high-density genetic maps of two great tit populations reveal fine-scale genomic

departures from sex-equal recombination rates. Heredity. 2014;112: 307–316.

doi:10.1038/hdy.2013.107

79. Gautier M, Vitalis R. Rehh: An R package to detect footprints of selection in genome-

wide SNP data from haplotype structure. Bioinformatics. 2012;28: 1176–1177.

80. O'Connell J, Gurdasani D, Delaneau O, Pirastu N, Ulivi S, Cocca M, et al. A general

approach for haplotype phasing across the full spectrum of relatedness. PLoS Genet.

2014;10: e1004234.