BIOLOGICAL SCIENCES

Density-dependent selection mediates harvest-induced evolution

Alix Bouffet-Halle[1], Jacques Mériguet[2,3], David Carmignac[1], Simon Agostini[2], Alexis Millot[2], Samuel

Perret[2,4], Eric Motard[1], Beatriz Decenciere[2], Eric Edeline[1,5]*

1: Sorbonne Université, Université Paris Diderot, UPEC, CNRS, INRA, IRD, Institut d'Ecologie et des

Sciences de l'Environnement de Paris (iEES-Paris), F-75252 Paris, France.

2: CEREEP Ecotron Île-de-France, UMS CNRS ENS 3194, 78 rue du Château, 77140 Saint Pierre-lès-

Nemours, France.

3: Institut de Biologie de l'Ecole Normale Supérieure, CNRS, INSERM, PSL Research University,

Paris, France.

4: Centre d'Ecologie Fonctionnelle et Evolutive CEFE, UMR 5175, Campus CNRS, Université de

Montpellier, Université Paul-Valéry Montpellier, Montpellier Cedex 5, France.

5: INRA, Ecology and Ecosystem Health, 65 rue de Saint-Brieuc, 35042 Rennes Cedex, France.

* Corresponding author: Eric.Edeline@inra.fr

Key words: Age and size at maturation, Anthropogenic change, Body size, Fisheries, Harvesting

yields, Rapid evolution, Size-dependent selection, *K*-selection, Life-history evolution, Somatic growth

rate.

Abstract. Harvesting has been demonstrated to cause rapid, yield-decreasing trait change towards

slower somatic growth and earlier maturation in wild populations. These changes are largely

considered to result from direct, density-*in*dependent harvest selection on traits. Here, we show that

exact same trait changes may also indirectly result from a harvest-induced relaxation of density-

dependent (K) natural selection for faster growth and delayed maturation. We exposed 12 pond

populations of medaka fish (*Oryzias latipes*) to contrasted size-selective harvesting during 5 years, and

show that harvesting effectively changed juvenile natural mortality from density-dependent to density-

*in*dependent. We then laboratory-reared medaka progeny under contrasted food levels mimicking the

environmental effects of a harvest-induced density gradient. Interaction between past harvest regime

and present food environment on progeny traits revealed that harvest-induced trait changes in medaka

resulted from selection in a low-food environment only, i.e., were driven by relaxed K-selection only,

not by direct harvest selection. Feeding trials further demonstrated that trait changes were associated

with reorganizations in rates of food acquisition, assimilation and allocation that were contingent upon

the food environments. This is the first study to demonstrate that harvesting can induce undesirable

distortions of natural selection that impair productivity traits. We conclude that sustaining harvesting

yields over extended time scales requires a preservation of high population densities.

Significance statement: Fisheries management often opposes a density-dependent approach which

prioritize the preservation of high population densities, and an evolutionary approach which consider

that alleviating change towards smaller body sizes is paramount to the sustainability of harvesting. The

evolutionary approach consider harvest-induced body downsizing to be density-*in*dependent, i.e., to

result only from direct harvest selection against large-bodied individuals. Here, we show instead that

harvest-induced body downsizing may be density-dependent because, by decreasing population density,

fishing relaxes natural, density-dependent selection for large-bodied individuals. Therefore, preserving

population numbers and alleviating body downsizing in harvested populations are not independent lines

of management, but are in fact two necessary and complementary routes to reaching the same

management objectives.

Introduction

Harvesting potentially creates a mixture of selective pressures acting in parallel both directly and

indirectly on life-history traits. In particular, size-selective harvesting directly selects against an old

age, thus favoring early-maturing genotypes, and against large-bodied individuals at a given age, thus

favoring slow-growing genotypes (1). This direct, "brute-force" warping of naturally-selected fitness

landscapes is currently the prevailing model to explain harvest-induced evolution in wild populations

(1–3). However, in parallel harvesting also lowers population densities and is thus susceptible to

indirectly warp the naturally-selected fitness landscape through relaxing the strength of density-

dependent natural selection (4), also known as *K*-selection (5–7). So far, however, this density-

dependent pathway to harvest-induced selection remains unexplored empirically or experimentally. To

bridge this gap in our knowledge, we conducted a 5-year size-selective harvesting experiment on 12

populations of medaka fish (*Oryzias latipes*) maintained in outdoor ponds under natural conditions with

no artificial feeding, followed by a 1-generation common garden experiment in the laboratory. The 12

founder medaka populations originated from parents wild-caught in Kiyosu (Japan).

In ponds, our experimental size-selective fishery removed 81% of the catch (catch rate = 98%) by

specifically targeting large-sized individuals (Fig. 1a), and thus successfully reproduced a typical direct

harvest selection pattern. In parallel, our experimental fishery relaxed negative density-dependence in

medaka populations. Pond medaka populations followed Ricker stock-recruitment dynamics (Fig. 1b),

a population dynamics model used in many fisheries management schemes (8). Fishing consistently

decreased stock (population size in March) density below ca. 50 individuals (red squares in Fig. 1b), a

density region in which increasing stock size had a positive effect on the number of summer-born

juveniles (recruitment, black curves, Fig. 1b), indicating demographic "undercompensation" due to

density-*in*dependence of vital rates (9). In contrast, unharvested populations had stock sizes above ca.

50 individuals (blue triangles in Fig. 1b), a density region where increasing stock size had a negative

effect on recruitment, indicating demographic "overcompensation" due negative density-dependence of

vital rates (black curves, Fig. 1b).

Overcompensating recruitment may operate through decreased fecundity and/or through increased

mortality. To discriminate between the two mechanisms, we counted newborn larvae hiding in artificial

vegetation in each pond during 3 years. In harvested populations, newborn medaka larvae were on

average less numerous than in unharvested populations (P-value = 0.003, Fig. 2, Table S1), but average

recruit numbers were similar among unharvested and harvested populations (85 vs. 73 recruits

respectively, Fig. 1b, non statistically-significant difference), indicating that overcompensating

recruitment was mediated by increased post-larval mortality in medaka populations.

Density-dependent post-larval mortality is expected to select for a larger body size and delayed

reproductive investment. Experiments with *Drosophila* demonstrated that resource competition under a

high density favors the evolution of increased food intake and/or conversion efficiency, ultimately

resulting in faster somatic growth rates under standardized food conditions (10, 11). In fish,

cannibalism is a further source of density-dependent mortality also predicted to favor faster somatic

growth to larger body sizes (12, 13). Finally, high density and food limitation are expected to select for

a delayed reproduction at a larger body size (6, 14, 15), a prediction that was validated in *Drosophila*

(15). Therefore, we predicted that, exactly like direct harvest selection, harvest-induced relaxation of

K-selection should have favored slower somatic growth rates and earlier maturation in medaka.

In a previous laboratory selection experiment, Kiyosu medaka were unable to respond to selection for a

smaller body size but were able to respond to selection for a larger body size (16). This previous result

suggests that any harvest-induced change in somatic growth or maturation evolved by pond medaka in

the present experiment would more likely result from *K*-selection than from direct fishery selection.

However, a further, efficient way to discriminate between the direct *vs*. density-mediated effects of

harvesting is through the measurement of interactions between harvest treatments and food levels on

trait expression (17–20). This is because the genes that control a given trait are often environment-

specific (18–22). Consequently, trait differences measured under different standardized environments

may be used to infer the direction of selection in each environment (17, 19, 20).

For instance, mice selected for a fast (slow) somatic growth in a high-food environment grow faster

(slower) than unselected mice, but only in a high-food environment (18, 19). In contrast, mice selected

on somatic growth in a low-food environment showed a phenotypic response to selection under both a

low- and high-food environments, suggesting that selection on somatic growth in a low-food

environment tends to erase the sensitivity of growth to food variation (18–20). Following this rationale,

we predicted that somatic growth response to direct, density-*in*dependent harvest selection in medaka

should be manifest in a high-food environment, while somatic growth response to *K*-selection should

be manifest in any food environment (19, 20).

Evolution of maturation is also expected to be contingent upon the food environment. For instance,

predation-induced evolution towards earlier maturation in guppies *Poecilia reticulata* is more

pronounced under a high-food environment because predators decrease guppy density and thus select

in a high-food environment (17). Hence, we further predicted in medaka that maturation response to

direct, density-*in*dependent harvest selection should be more pronounced in a high-food environment,

while maturation response to *K*-selection should be more pronounced under a low-food environment

(17).

Results

To test these predictions we measured in the laboratory the somatic growth of F_1 progeny from pond-

sampled parents. We applied a low-, medium- and high-food regimes intended to mimic the

environmental effects of an increasing harvest intensity from feeding the progeny once every second

day to feeding twice daily.

Under all three food environments, harvested medaka grew significantly slower than unharvested

medaka (low food P-value = 0.008, medium food P-value < 0.001, high food P-value = 0.002, Fig. 3a).

Accordingly, a deviance analysis shows that there was no significant harvest by food interaction (P-

value = 0.2650, Table 1), indicating that the amplitude of harvest-induced decrease in somatic growth

was food-independent. This result suggests that medaka responded to selection for fast-growth in a

low-food environment (19, 20), i.e., responded to *K*-selection for faster somatic growth, but not to

direct harvest selection for slower somatic growth in a high-food environment. This result is further in

line with our previous finding that medaka from Kiyosu are unable to respond to selection for slower

somatic growth under laboratory conditions but that they do respond to selection for faster somatic

growth (see above).

Supporting our second prediction, a deviance analysis shows that the effect of harvesting on the age-

dependency of maturation was significantly food-dependent (Age $\times$ Harvesting $\times$ Food

interaction, Table 1). Specifically, harvesting changed the size-corrected effect of age on maturation

probability from significantly positive (P-value = 0.025, Table S1) to significantly negative (P-value =

0.020, Table S1), reflecting that harvesting induced earlier maturation only in a low-food environment

(Fig. 3b). These results suggest that medaka responded to selection for delayed maturation in a low-

food environment (17), i.e., responded to *K*-selection, but not to direct harvest selection for earlier

maturation in a high-food environment. In line with this result, we previously found that Kiyosu

medaka are unable to respond to selection for earlier maturation in the laboratory (16).

K-selected changes in somatic growth and maturation may be mediated by combined changes in energy

acquisition, assimilation or allocation rates. To gain insights into these regulatory pathways we

measured acquisition rates through individual feeding trials on laboratory-born F_1 individuals. We

starved fish overnight, presented them with 20 prey (nauplii of *Artemia salina*), and counted the

number of prey eaten during 5 minutes (repeated 3 times per individual).

Progeny from harvested populations ate significantly less prey than progeny from unharvested

populations, but only in a medium-food environment (P-value = 0.011, Fig. 3c). This result suggest that

changes occurred in all three pathways of energy acquisition, assimilation and allocation, but that the

respective contributions of these pathways to the expression of life-history change was environment-

specific. In a low-food environment, slower somatic growth (Fig. 3a), earlier maturation (Fig. 3b) but

unchanged energy acquisition (Fig. 3c, P-value = 0.523) in harvested medaka together suggest energy

re-allocation from growth to reproduction. In a medium-food environment, the slower somatic growth

(Fig. 3a) of in harvested medaka was apparently mediated by decreased energy acquisition (Fig. 3c),

but unchanged maturation (Fig. 3b) also suggests energy re-allocation from growth to reproduction.

Finally in a high-food environment, slower somatic growth (Fig. 3a) but unchanged rates of maturation

(Fig. 3b) and energy acquisition (Fig. 3c, P-value = 0.424) together suggest decreased energy

assimilation rates in harvested medaka. These results are consistent with previous studies showing that

evolution towards slower somatic growth in fish may be underlaid by decreases in food consumption

rate and conversion efficiency (23).

Discussion

Our results demonstrate that harvesting caused evolution towards slower somatic growth and earlier

maturation in medaka through relaxed *K*-selection. However, in ponds the body size of 0+ juvenile

medaka did not show any statistically significant temporal trend in harvested or unharvested

populations (MCMC P-values = 0.365 and 0.262, respectively, Fig. 4). Phenotypic stasis despite known

evolutionary change (i.e., cryptic evolution) is typical of responses to environmental deterioration

where decreased environment quality selects for higher competitive ability but, as highly competitive

genotypes spread in the population the environment further deteriorates, resulting in no detectable

effects on phenotypes (24, 25). Such cryptic "Red Queen" evolutionary dynamics are expected in all

density-dependent populations and are thus probably commonplace in harvested systems. However,

their detection requires using common garden experiments or specific quantitative genetic methods (24,

25), and studies of harvest-induced trait change based on field data published so far thus maybe

underestimate potential for harvest-induced evolution.

The direct and density-dependent pathways to harvest-induced selection act in the same direction on

life-history traits, but have different implications for management. Phenotypic changes from direct

harvest selection may be alleviated by moulding the shape of artificial selection onto the shape of

natural selection through, for instance, adjusting gear selectivity. In contrast, the consequences of

density-dependent harvest selection can be alleviated only by relaxing the harvest effort (4).

Additionally, post-moratorium phenotypic recovery from direct harvest selection is expectedly slow

because the strength of natural selection is predicted to be constant and modest relative to the strength

of harvest selection. In contrast, recovery from density-dependent harvest selection should be rapid

because natural selection strengthens when fishing is relaxed (13).

Recent studies have shown that predator-induced life-history evolution may be, at least partly, mediated

by relaxed K-selection (26) and by an associated adaptation to increased food availability (17). Our

study reinforces and extends these previous results by experimentally demonstrating that harvest-

induced trait changes previously ascribed to direct, density-independent selection in the literature may,

in fact, have also emerged through a relaxation of *K*-selection. Hence, the more ecologically sustainable

harvesting strategies also produce smaller evolutionary changes (4), and the next-generation harvest

management methods should thus converge towards an integration of the reciprocal effects between

ecological dynamics and rapid evolutionary change.

183   **Materials and Methods**

Pond medaka populations

Origin and maintenance

Our start medaka populations descended from 100 wild medaka caught in Kiyosu (27) (Toyohashi,

Aichi Prefecture, Japan) in June 2011. These 100 Japanese breeders were maintained in five 20L

aquariums and their eggs were collected daily from July to September 2011. Hatched larvae were

stocked in 12 circular outdoor ponds (3.57 m diameter, 1.2 m deep).

Prior to medaka introduction, the 12 ponds were bottom-coated with a 5 cm layer of Loire River sand,

filled with tap water and mildly enriched with a plant fertilizer. After a few weeks of algal

development, tanks were seeded with a diverse community of zooplankton collected from surrounding

water bodies. Medaka introduction was performed after ponds had reached a clear-water state

indicating algal control by zooplankton. After introduction, two pairs of floating plastic brushes were

placed in each tank to provide fish with a spawning substrate and shelter for larvae. Each pond was

covered with a net to prevent avian predation, and was outlet-secured with a stainless steel filter to

prevent any fish or egg escapement. No food was added to the ponds which thus represented natural,

replicated ecosystems.

Medaka harvesting and phenotyping in ponds

From 2012 to 2016, each of the 12 pond populations was sampled in March before medaka

reproduction (pre-recruitment) and in November after medaka reproduction (post-recruitment). Fish

were concentrated using a seine net and then fished using handnets (catchability = 98 ± 0.6% SD

estimated using removal sampling). All sampled fish were individually weighted to the nearest mg and

estimated for standard body length (from the tip of the snout to the base of the caudal fin) using a body

mass-length relationship (R^2 = 0.98 on a log-log scale, n = 2722). In March in the 6 harvested

populations all the fish that were too large to pass through a 2 mm-wide screen were removed, while in

unharvested populations all fish were released after phenotyping. In November, all fish from both

harvested and unharvested populations were released after phenotyping.

Larvae counts

We visually counted the number of newly-hatched larvae hiding in each pair of floating plastic brushes

(summed for the two brush pairs) from one to three times per day at irregular intervals during the 2014,

2015 and 2016 spawning periods (April to September).

Medaka F_1 in the laboratory

Parental fish

In November 2016, between 6 and 10 individuals were randomly kept from each of the 12 pond

populations to serve as parents for a F_1 generation in the laboratory. These parental fish were

maintained in a greenhouse at air temperature in 12, 150L tanks with live food. In January 2017,

parental fish were weighted to the nearest mg, measured for standard body length with ImageJ, and

grouped to form 3 breeding pairs per population (except one harvested population that had only one

female). Each of the resultant 36 pairs was transferred to the laboratory in a 3.5L aquarium and induced

to spawn by progressively raising temperature to 27.0 ± 0.3°C and setting a 15-h light:9h dark

photoperiod. Dry food (Skretting Gemma Micro) was provided twice per day and live nauplii of

Artemia salina once per day. After initiation of spawning by all breeding pairs, eggs from each

breeding pair were collected daily during a 4-day period, enumerated and incubated in separate jars so

as to keep track of individual parental identity (but not spawning day). We found no significant effect

of the harvest treatment on parental body size, body condition, fertility or fecundity.

F_1 progeny phenotyping and food environments

We collected F_1 larvae born from the 7th to the 10th day after the weighted average date of spawning.

Larvae hatched from the same breeding pair on the same day were transferred to 1.5L aquariums by

groups of 3 larvae, and were maintained under the same temperature and light regime as their parents.

We kept 1-4 groups of F_1 larvae per breeding pair (average 2.9 groups per breeding pair). At 15 days

post hatch (dph), all F_1 individuals were weighted and measured as described above and only one

individual per aquarium was randomly kept for subsequent phenotyping, making it possible to track

individual developmental trajectories. Individual phenotyping was repeated at 30 dph, 40 dph and then

once per week until 90 dph (11 individual measurements). From 40 dph onwards, phenotyping further

included detection of the maturity status from the presence of secondary sexual characters (28).

Specifically, the maturity criteria were first appearance of a round-shaped anal papilla in females, and

of the papillar process on the anal fin in males. Additionally, at around 48, 56 and 63 dph, each

individual F_1 medaka was measured for feeding rate. We counted the number of live prey (nauplii of

Artemia salina) eaten when the medaka was placed alone with 20 prey during 5 minutes in a 80 mL

container. Medaka were starved overnight prior to each behavioural test.

From 15 dph onwards, we varied resource levels by applying three food environments to F_1 progeny.

We chose feeding regimes so has to mimic a high-density, scarce-food environment in which predators

are not able to daily catch a prey, a low-density, food-rich environment in which predators are replete

with prey, and an intermediate environment. In the low-food environment, individuals were fed with

2mL of a solution containing nauplii of *Artemia salina* at a standard concentration on day 1, nothing on

day 2, dry food (see below) on day 3, nothing on day 4 and so on. In the high-food environment,

medaka were fed twice daily, once with nauplii and once with dry food. Finally, in the medium-food

environment, medaka were fed once daily alternating nauplii and dry food.

Volume of dry food doses and pellet size were increased during fish development to fit with the

ontogenetic increase in energy needs and prey size. From 0 to 40 dph, 40 to 60 dph, and 60 dph

onwards, medaka received daily 4, 6 and 14µL of food, respectively. From 0 to 20 dph, 20 to 40 dph,

and 40 dph onwards, dry food was made from 100% 150 µm pellets, 50% mixture of 150-300 µm

pellets and 100% 300 µm pellets, respectively.

Statistical analyses

We below provide a short summary of the statistical analyses. A full description is given in the SI

Appendix.

Analysis of pond data

Medaka age was inferred by fitting a mixture of two Gaussian distributions to individual standard body

lengths measurements (n = 17908). We further estimated temporal trends in body size of November 0+

recruits (n = 9688 individuals) using a version of the Gaussian mixture model that was modified to

include a harvest treatment-specific (n = 2 treatments) hierarchical regression of mean recruit standard

body length on year of sampling (n = 5 years). We estimated the relationship between individual

standard body length and probability to survive through the fishery in March (n = 3970 individuals)

using a mixed effects Bernoulli GLM with a logit link function. The Gaussian mixture model described

above allowed us to estimate the number of November 0+ recruits in each pond and year. We then

visualized the strength of negative density-dependence in pond medaka populations by plotting Ricker

"stock-recruitment" relationships (Fig. 1b). Finally, larvae counts in ponds were modelled using a

mixed-effects zero-inflated negative binomial model, which parameter estimates are provided in Table

S1.

Analysis of laboratory data

We estimated the effects of harvesting and food environments on the growth trajectories of F_1 progeny

in the laboratory using a second order polynomial regression of standard body length on age (parameter

estimates provided in Table S1). We fitted probabilistic maturation reaction norms (PMRNs) to medaka

maturation data using the "direct estimation" method for PMRNs (29) in a mixed-effects Bernoulli

GLM with a logit link function (parameter estimates provided in Table S1). Counts of the number of

nauplii larvae eaten by individual medaka were modelled using a mixed-effects zero-inflated negative

binomial model (parameter estimates provided in Table S1).

Acknowledgements. We are grateful to Prof. Kiyoshi Naruse (NIBB, Okazaki, Japan) for his support

in obtaining and maintaining wild medaka from Kiyosu.

Funding. This work has benefited from technical and human resources provided by CEREEP-Ecotron

IleDeFrance (CNRS/ENS UMS 3194) as well as financial support from the Regional Council of Ile-de-

France under the DIM Program R2DS bearing the references I-05-098/R and 2015-1657. It has

received a support under the program "Investissements d'Avenir" launched by the French government

and implemented by ANR with the references ANR-10-EQPX-13-01 Planaqua and ANR-11-INBS-

0001 AnaEE France, and from Pépinière interdisciplinaire CNRS de site PSL (Paris-Sciences et

Lettres) "Eco-Evo-Devo". EE also acknowledges support from the Research Council of Norway

(projects EvoSize RCN 251307/F20 and REEF RCN 255601/E40).

Author contributions. ABH performed the laboratory F_1 experiment, contributed to data analysis,

wrote the first draft of the manuscript and contributed to subsequent versions. EE designed the study,

contributed to the pond experiment, performed data analysis, and led manuscript writing from the

second version. JM, DC, SA, AM, SP, EM and BD contributed to the pond and laboratory experiments.

Competing interests. The authors declare no competing interests.

Data archiving statement. All data and codes used in this paper will be archived.

Ethical statement. The protocols used in this study were designed to minimize discomfort, distress and

pain of animals, and were approved by the Darwin Ethical committee (case file #Ce5/2010/041).

Supplementary Materials

Table S1: MCMC parameter estimates for models 4-7.

References

1. Heino M, Díaz Pauli B, Dieckmann U (2015) Fisheries-induced evolution. *Annu Rev Ecol Evol Syst* 46(1):461–480.

2. Edeline E (2016) Life history evolution, human impacts on. *The Encyclopedia of Evolutionary Biology*, ed Kliman R (Academic Press, Oxford), pp 335–342. 1st Ed.

3. Carlson SM, et al. (2007) Four decades of opposing natural and human-induced artificial selection acting on Windermere pike (Esox lucius). *Ecol Lett* 10(6):512–521.

4. Engen S, Lande R, Sæther B-E, Associate Editor: Jürgen Groeneveld, Editor: Troy Day (2014) Evolutionary consequences of nonselective harvesting in density-dependent populations. *Am Nat* 184(6):714–726.

5. MacArthur RH, Wilson EO (1967) *The theory of island biogeography* (Princeton University Press, Princeton). 1st Ed.

6. Pianka ER (1970) On r- and K-Selection. *Am Nat* 104(940):592–597.

7. Reznick D, Bryant MJ, Bashey F (2002) r- and K-selection revisited: the role of population regulation in life-history evolution. *Ecology* 83(6):1509–1520.

8. Hilborn R, Walters C (1992) *Quantitative fisheries stock assessment: choice, dynamics and uncertainty* (Springer US). 1st Ed.

9. Bellows TS (1981) The descriptive properties of some models for density dependence. *J Anim Ecol* 50(1):139–156.

10. Mueller LD (1988) Evolution of competitive ability in Drosophila by density-dependent natural selection. *Proc Natl Acad Sci U S A* 85(12):4383–4386.

11. Sarangi M, Nagarajan A, Dey S, Bose J, Joshi A (2016) Evolution of increased larval competitive ability in Drosophila melanogaster without increased larval feeding rate. *J Genet*:1–13.

12. Claessen D, de Roos AM, Persson L (2000) Dwarfs and giants: cannibalism and competition in size   structured populations. *Am Nat* 155(2):219–237.

13. Edeline E, et al. (2007) Trait changes in a harvested population are driven by a dynamic tug-of-war between natural and harvest selection. *Proc Natl Acad Sci* 104(40):15799–15804.

14. Holliday R (1989) Food, reproduction and longevity: is the extended lifespan of calorie-restricted animals an evolutionary adaptation? *BioEssays News Rev Mol Cell Dev Biol* 10(4):125–127.

15. Sgrò CM, Partridge L (2000) Evolutionary responses of the life history of wild caught Drosophila melanogaster to two standard methods of laboratory culture. *Am Nat* 156(4):341–353.

16. Renneville C, et al. (2018) Anthropogenic selection along directions of most evolutionary resistance. *bioRxiv*:498683.

17. Walsh MR, Reznick DN (2008) Interactions between the direct and indirect effects of predators determine life history evolution in a killifish. *Proc Natl Acad Sci* 105(2):594.

18. Falconer DS (1960) Selection of mice for growth on high and low planes of nutrition. *Genet Res* 1(1):91–113.

19. Falconer DS, Latyszewski M (1952) The environment in relation to selection for size in mice. *J Genet* 51(1):67.

20. Falconer DS (1990) Selection in different environments: effects on environmental sensitivity (reaction norm) and on mean performance. *Genet Res* 56(1):57–70.

21. Jinks JL, Connolly V (1973) Selection for specific and general response to environmental differences. *Heredity* 30:33.

22. Falconer DS, Mackay TFC (1996) *Introduction to quantitative genetics* (Longman, Harlow, Essex, UK). 4th Ed.

23. Walsh MR, Munch SB, Chiba S, Conover DO (2006) Maladaptive changes in multiple traits caused by fishing: impediments to population recovery. *Ecol Lett* 9(2):142–148.

24. Wolf JB (2003) Genetic architecture and evolutionary constraint when the environment contains genes. *Proc Natl Acad Sci U S A* 100(8):4655–4660.

25. Hadfield JD, Wilson AJ, Kruuk LEB (2011) Cryptic evolution: does environmental deterioration have a genetic basis? *Genetics* 187(4):1099–1113.

26. Bassar RD, Lopez-Sepulcre A, Reznick DN, Travis J (2013) Experimental evidence for density-dependent regulation and selection on Trinidadian guppy life histories. *Am Nat* 181(1):25–38.

27. Spivakov M, et al. (2014) Genomic and phenotypic characterization of a wild medaka population: towards the establishment of an isogenic population genetic resource in fish. *G3 GenesGenomesGenetics* 4(3):433–445.

28. Yamamoto T (1975) *Medaka (killifish): biology and strains* (Keigaku Pub. Co, Tokyo). 1st Ed.

29. Heino M, Dieckmann U (2008) Detecting fisheries-induced life-history evolution: an overview of the reaction-norm approach. *Bull Mar Sci* 83(1):69–93.

30. Stearns SC, Koella JC (1986) The evolution of phenotypic plasticity in life-history traits: predictions of reaction norms for age and size at maturity. *Evolution* 40(5):893–913.

31. Heino M, Dieckmann U, Godø OR (2002) Measuring probabilistic reaction norms for age and size at maturation. *Evolution* 56(4):669–678.

300 **Table 1. Analysis of deviance table for GLMs testing for the harvest by food interaction on life-**

301 **history traits in laboratory-born F$_1$ medaka progeny.** The "Deviance" column gives the reduction in

302 the residual deviance as each predictor is added in turn into the model. The P-values compare the

303 reduction in deviance to the residual deviance in an F test.

Trait	Distribution	Link	Predictor	Df	Deviance	Resid. DF	Resid. Dev	F	P-val
Body length	Gaussian	Identity	Age	1	18100	1130	2651	17144	**<0.0001**
			Age^2	1	468	1129	2183	443	**<0.0001**
			Harvesting	1	130	1128	2053	123	**<0.0001**
			Age x Harvesting	1	24	1127	2029	22	**<0.0001**
			Age x Food	2	841	1125	1188	398	**<0.0001**
			Age x Harvesting x Food	2	3	1123	1186	1	0.2650
Maturation	Bernoulli	Logit	Age*	1	96	589	432	164	**<0.0001**
			Length*	1	97	588	335	166	**<0.0001**
			Harvesting	1	2	587	333	3	0.064
			Food	2	3	585	329	3	0.0513
			Age* x Harvesting	1	8	584	321	14	**0.0002**
			Length* x Harvesting	1	1	583	320	2	0.1745
			Age* x Food	2	12	581	309	10	**<0.0001**
			Length* x Food	2	1	579	307	1	0.3263
			Age* x Harvesting x Food	2	10	577	297	9	**0.0002**
			Length* x Harvesting x Food	2	2	575	295	2	0.2027

Fig. 1. Direct and density-mediated harvest-selection in ponds. a: Size- and age-dependent harvest selection. Light grey bars represent raw standard length data in harvested populations. Superimposed Gaussians represent mean MCMC estimates for the density of 0+ juveniles (short-dashed curve) and 1+ and older adults (long-dashed curve) individuals. The magenta logistic curve shows the mean relationship between exploitation rate by the fishery and standard body length. **b: Stock-recruitment relationships.** Points show mean MCMC recruitment estimates with 95% credible intervals for unharvested (blue triangles) and harvested (red squares) populations. Black curves show year-specific Ricker functions fitted to mean estimates using maximum likelihood.

312 **Fig.2. Larvae count seasonal dynamics in ponds.** Thick curves represent mean MCMC estimates for

313 daily counts of newly-hatched larvae for unharvested (dot-dashed, blue curve) and harvested (dashed,

314 red curve) populations. Thin curves show 95% credible intervals around mean MCMC estimates.

Fig. 3. Individually-raised F$_1$ progeny in the laboratory. a: Mean growth trajectories. MCMC mean growth curves for individuals originating from unharvested (dot-dashed blue curves) and harvested (dashed red curves) populations in a low-, medium- or high-food environments. Grey dots show the raw data. **b: Probabilistic maturation reaction norms (PMRNs).** PMRNs show the combination of age and lengths at which maturation probability equals 0.5. They account for the plastic effect of growth on maturation, and a shift in PMRNs is thus suggestive of a non-plastic, evolutionary change in maturation schedules (30, 31). Coloured lines show MCMC mean estimates with 95%

credible intervals for PMRNs of medaka originating from unharvested (dot-dashed blue line) and

harvested (dashed red line) populations. Thin grey curves in the background show raw growth

trajectories for medaka originating from unharvested (dot-dashed) and harvested (dashed) populations.

Solid black lines show the mean growth trajectories in a low-, medium- or high-food environment

(averaged across harvesting treatments). **c: Feeding rates.** Coloured, open points symbols show mean

MCMC estimates with 95% credible intervals for the number of prey eaten by medaka originating from

unharvested control (blue triangles) and harvested (red squares) populations and maintained in a low-,

medium- or high- food environment. Grey, filled symbols show the raw data.

Fig. 4. Body length time series estimates for November 0+ recruits in pond medaka populations.

Points show mean MCMC recruitment estimates with 95% credible intervals for unharvested (blue

triangles) and harvested (red squares) populations.

Supplementary Information for

Density-dependent selection mediates harvest-induced evolution

Alix Bouffet-Halle, Jacques Mériguet, David Carmignac, Simon Agostini, Alexis Millot,

Samuel Perret, Eric Motard, Beatriz Decenciere, Eric Edeline*

* Corresponding author: eric.edeline@inra.fr

This PDF file includes:
Supplementary methods: statistical analyses.
Table S1.

341   **Statistical analyses**

342   *Medaka aging in ponds*

343   Medaka juveniles are too small to be tagged and, unlike in Japan (*1, 2*), no winter check was deposited

344   in medaka otoliths in our experimental populations. We therefore relied on analysis of length-frequency

345   distributions to infer medaka age. We fitted a mixture of two Gaussian distributions to individual

346   standard body lengths Sdl_i :

$$Sdl_i \sim \sum_{j=1}^{J} \sum_{k=1}^{K} \pi_{j,k} N\left(\mu_{j,k}, \sigma_j^2\right)$$
347    $$\mu_{2,k} \sim N\left(\mu_{H[k]}, \sigma^2\right) \qquad (1a),$$
$$\mu_{1,k} = \delta_k \mu_{2,k}$$
$$\delta_k \sim U(0,1)$$

where i indexes individuals (n = 17908), j indexes age groups (0+ *vs.* 1+ and older such that

J = 2), k indexes a sampling event, i.e., indexes one population in a particular year and month (

K = 109 sampling events), N is the normal distribution, and U is the uniform distribution.

$H[k]$ indexes the harvest treatment (harvested *vs.* non harvested) associated with sampling event

k . $\pi_{j,k}$ is the proportion of age j individuals at each sampling event k such as for each

k :

354
$$\pi_j \geq 0, \sum_{j=1}^{J} \pi_j = 1 \qquad \text{(1b)}.$$

Indexes in line 1 in Eq. 1a show that our model estimated a mean standard body length separately for

each age group at each sampling event, while body length variance was assumed to vary only with age.

Line 2 in Eq. 1a shows that we assumed the mean standard body length of age 1+ and older medaka at

each sampling event $\mu_{2,k}$ to be a normally-distributed random variable with mean specific to each

harvest treatment, because harvesting was expected to restrict the maximum age and size of medaka.

Lines 3-4 in Eq. 1a show that mean standard body length of 0+ medaka at each sampling event $\mu_{1,k}$

was estimated as proportional to $\mu_{2,k}$ with a proportionality constant δ_k following a uniform

distribution U between 0 and 1. Model 1 provided us with MCMC age samples for each individual

fish in the dataset, allowing us to compute age-specific survival rates through the fishery.

We estimated temporal trends in mean standard body length of November 0+ recruits using a modified

version of model 1 that included a hierarchical regression:

$$Sdl_i \sim \sum_{j=1}^{J} \sum_{k=1}^{K} \pi_{j,k} N(\mu_{j,k}, \sigma_j^2)$$

$$\mu_{2,k} \sim N(\mu_{H[k]}, \sigma_2^{\ 2}) \qquad (1c)$$

$$\mu_{1,k} \sim N(\hat{\mu}_{1,k}, \sigma_1^{\ 2})$$

$$\hat{\mu}_{1,k} = \alpha_{H[k]} + \beta_{H[k]} Year_k$$

where i indexes November-sampled fish (n = 9688 individuals, K = 60), $\alpha_{H[k]}$ and $\beta_{H[k]}$

are harvest treatment-specific temporal regression parameters, and *Year* was scaled to 0 mean.

Other variables and subscripts are as described above.

Fishery exploitation rate and selection in ponds

We estimated the relationship between individual standard body length and probability to survive

through the fishery using a Bernoulli GLM with a logit link function:

$$y_i \sim Bern(p_i)$$

$$\ln\left(\frac{p_i}{1-pi}\right) = \alpha_0 + \alpha_{j[i]} + (\beta_0 + \beta_{j[i]}) Sdl_i \qquad (2),$$

$$\begin{pmatrix} \alpha_j \\ \beta_j \end{pmatrix} \sim N\left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma_\alpha & \rho\,\sigma_\alpha\sigma_\beta \\ \rho\,\sigma_\beta\sigma_\alpha & \sigma_\beta \end{pmatrix} \right)$$

where subscripts i and j index individuals (n = 3970) and groups, respectively, to which

individuals belong. There was n = 6 fished populations and n = 5 sampling years, yielding

$j=1,2\ldots30$ groups. Finally, *Bern* is the Bernoulli distribution, and $\ln$ is the natural

logarithm.

Eq. 2 indicates that we modelled the intercept and slope of the survival-mass relationship as normally-

varying among groups j, including a correlation parameter ρ between intercept and slope.

Parameter estimates α_0 and β_0 from Eq. 2 define a mean size-dependent survival function

$s(Sdl)=1/(1+\exp(-(\alpha_0+\beta_0 Sdl)))$ plotted in Fig. 1a.

Stock-recruitment relationship in ponds

Model (1) described above allowed us to estimate the number R_k of 0+ medaka (recruits) at each

November sampling event k (n = 60 November sampling events). We then visualized the strength of

negative density-dependence in pond medaka populations by plotting (Fig. 1b) Ricker (*3*) "stock-

recruitment" relationships between R_k and the number S_k of fish released in March (stock of

spawners):

388
$$R_k \sim P(\lambda_k)$$
$$\ln(\lambda_k)=\ln(S_k)+\alpha_{Year[k]}+\beta_{Year[k]}S_k \quad (3),$$

389 where P is the Poisson distribution and $Year[k]$ indexes indicate that one Ricker curve was

390 fitted for each year from 2012 to 2016.

391 *Larvae counts*

392 Larvae counts L followed a zero-inflated negative binomial distribution and were modelled as (*4*):

393
$$L_i \sim NB(\phi_i, r_i)$$
$$\phi_i=\frac{r_i}{r_i+\lambda_i(1-\theta_i)}$$
$$r_i=\gamma_{H[i]} \quad (4a),$$
$$\ln(\lambda_i)=\alpha_{Year[i],Pond[i]}+\beta_{H[i]}+\delta_{H[i]}Day_i$$
$$\alpha_{Year[i],Pond[i]}\sim N(0,\sigma_\alpha^2)$$

where subscript i indexes sampling events corresponding to a given observer in a given pond on a

given sampling day (n = 2,004 sampling events), NB is the negative binomial distribution with

success probability ϕ and number of failures r. Lines 4 and 5 in Eq. (4a) show that we modelled

positive (non-zero) counts λ as harvest treatment-specific linear regressions of the day of year

(scaled to 0 mean), with a normally-distributed random effect of the year and pond combination (n = 36

groups).

The θ latent variable for absence of larvae was modelled as a Bernoulli process having a linear

dependency on the day of year:

402
$$\theta_i \sim B(\psi_i)$$
$$\ln\left(\frac{\psi_i}{1-\psi_i}\right) = \zeta + \omega \, Day_i \quad \text{(4b),}$$

where B is the Bernoulli distribution with probability for absence of larvae ψ.

Line 3 in Eq. 4A shows that we allowed for r, which enters in the computation of the variance of

the distribution (*4*), to be different among the two harvest treatments H. Harvest treatment-specific

mean larvae count is given by $E(L_H) = \bar{\lambda}_H(1-\bar{\theta})$ and variance by

$var(L_H) = \bar{\lambda}_H(1-\bar{\theta})(\bar{\lambda}_H(1-\bar{\theta}) + \gamma_H)$, and we computed the dispersion index (*4*) in each harvest

treatment as $DI_H = E(L_H)/var(L_H)$ (Table S1).

Somatic growth trajectories of F_1 progeny in the laboratory

We estimated the effects of harvesting and food environments on medaka growth trajectories using a

second order polynomial regression of standard body length on age:

412
$$Sdl_i \sim N(\mu_i, \sigma_i^2)$$
$$\mu_i = \alpha_{P[i]} + \beta_{H[i]} + (\gamma_{H[i],F[i]} + \delta_{P[i]}) * Age_i + \eta\, Age_i^2$$
$$\alpha_{P[i]} \sim N(0, \sigma_\alpha^2)$$
$$\delta_{P[i]} \sim N(0, \sigma_\delta^2)$$
$$\ln(\sigma_i^2) = A_{H[i],F[i]} + B_{H[i],F[i]}\, Age_i$$

(5),

where i indexes observations (n = 1144 observations from 104 individuals), $H[i]$ indexes the

harvest treatment associated with observation i, $H[i], F[i]$ indexes the interaction of harvest

treatment and food environment (n = 2 * 3 = 6 groups), and $P[i]$ indexes the parental breeding pair

associated with observation i (n = 36 pairs), treated as a normally-distributed random effect on both

size-at-hatch α and linear somatic growth rate δ (lines 3 and 4 in Eq. 5, respectively).

In this model, we assumed both linear somatic growth rate and the regression of (ln-transformed)

residuals variance on age to be different among harvest treatments and food environments (lines 2 and

5 in Eq. 5, respectively). In contrast, size-at-hatch $\beta_{H[i]}$ was allowed to vary only due to harvest

treatment because food environments were applied only starting from 15 dph.

Probabilistic maturation reaction norms of F_1 progeny in the laboratory

Probabilistic maturation reaction norms (PMRNs) describe the probability that an immature individual

at a given age and size will mature during a given interval of time (*5*). Provided that plasticity in the

maturation process is captured by growth trajectories, PMRNs separate the effects of evolution from

plasticity on maturation. PMRNs have been extensively used to explore genetic effects of exploitation

on the maturation process in wild populations (*6, 7*). Following the "direct estimation" method for

PMRNs (*7*), we fitted a Bernouilli model to individual medaka maturity (0 or 1) data y_i , truncated

so as to keep only the first maturity event for each individual:

430
$$
\begin{gathered}
y_i \sim B(\mathrm{M}_i) \\
\ln\!\left(\frac{\mathrm{M}_i}{1-\mathrm{M}_i}\right)=\alpha_{P[i]}+\beta_{H[i]}+\gamma_{H[i]}Age_i+\delta_{H[i]}Sdl_i \qquad (6), \\
\alpha_{P[i]}\sim N\left(0,\sigma_\alpha^2\right)
\end{gathered}
$$

where M is maturity probability. Other subscripts or variables are as described above. Eq. 6 shows

that we allowed harvest-specific intercept and slopes of age and standard body length effects on

maturation probability. Harvest-specific PMRNs corresponding to length at 50% maturation probability

for each age in each treatment group H was then computed as $Sdl\,50_H=-(\beta_H+\gamma_H\,Age)/\delta_H$.

Predatory behaviour of F₁ progeny in the laboratory

Counts C_i of number of prey eaten by individual medaka followed a zero-inflated negative binomial

distribution and were modelled similarly as larvae counts in model 4 above:

438
$$
\begin{aligned}
&C_i \sim NB(\phi_i, r_i) \\
&\phi_i = \frac{r_i}{r_i+\lambda_i(1-\theta_i)} \\
&r_i = \gamma_{H[i],F[i]} \qquad\qquad (7a), \\
&\ln(\lambda_i)=\alpha_{I[i]}+\beta_{H[i],F[i]} \\
&\alpha_{I[i]}\sim N(0,\sigma_\alpha^2)
\end{aligned}
$$

439   where number of failures r and positive (non-zero) counts λ were both modelled as being

440   different among harvest treatments H in each food environment F, while $\alpha_{I[i]}$ was a

441   normally-distributed random individual effect on λ (n = 3 counts per individual). The θ latent

442   variable was modelled as:

$$
\begin{aligned}
\theta_i &\sim B(\psi_i) \\
\ln\left(\frac{\psi_i}{1-\psi_i}\right) &= \gamma + \delta_{I[i]} \quad \text{(7b)}, \\
\delta_{I[i]} &\sim N(0, \sigma_\delta^2)
\end{aligned}
$$

444   where δ_I is a normally-distributed random individual effect.

445   *Analysis of deviance*

446   We tested for the overall statistical significance of harvest by food interaction on somatic growth and

447   maturation in the laboratory using analyses of deviance. Specifically, we fitted the following models:

$$
\begin{aligned}
Sdl_i &\sim N(\mu_i, \sigma_i^2) \\
\mu_i &= \alpha_{H[i]} + \left(\beta_{H[i]} + \gamma_{F[i]} + \delta_{H[i],F[i]}\right) Age_i + \zeta\, Age_i^2
\end{aligned}
\quad \text{(8), and}
$$

$$
\begin{aligned}
y_i &\sim B(M_i) \\
\ln\left(\frac{M_i}{1-M_i}\right) &= \alpha_{H[i]} + \beta_{F[i]} + \left(\gamma_{H[i]} + \delta_{F[i]} + \zeta_{H[i],F[i]}\right) Age_i + \left(\eta_{H[i]} + \theta_{F[i]} + \iota_{H[i],F[i]}\right) Sdl_i
\end{aligned}
\quad \text{(9),}
$$

450   where variables are as in models (5) and (6). We then used an F test to evaluate the significance of each

451   predictor separately (Table 1).

Parameter estimation

Models 3, 8 and 9 were fitted using maximum likelihood (glm function) in R 3.4.4 (*8*). Other models

were fitted by Markov chain Monte Carlo (MCMC) in JAGS 4.2.0 (*9*), through the jagsUI package

(*10*). To ease model convergence and avoid slope-intercept correlations, all numerical predictors were

scaled to zero mean. For each model, we ran three independent MCMC chains thinned at a period of 5

iterations until parameter convergence was reached, as assessed using the Gelman–Rubin statistic (*11*).

Parameter estimates for models 4-7 are provided in Table S1. Statistical significance of harvest- and

food-treatment effects reported in the main text was assessed from the posterior distributions of

parameter differences in a test equivalent to a bilateral *t* test. In these tests, the MCMC P-value was

twice the proportion of the posterior for which the sign was opposite to that of the mean posterior

value. For instance, in Eq. 4a the posterior differences $\beta_{H=1}-\beta_{H=0}$ and $\delta_{H=1}-\delta_{H=0}$ measure the

effect of harvest treatment (H = 0 for unharvested, H = 1 for harvested) on intercept and slope of day

effect for $\ln(\lambda)$, respectively.

Priors were chosen to be weakly informative. In model 1 we used a Dirichlet prior for the $\pi_{j,k}$ and

prevented label switching by assigning age class 0+ to fish shorter than 8 mm and age class 1+ and

older to fish longer than 35 mm (*12*).

We assessed goodness of fit of our models by using a Bayesian P-value (*13*). Briefly, we computed

residuals for the actual data as well as for synthetic data simulated from estimated model parameters

(i.e., residuals from fitting the model to "ideal" data). The Bayesian P-value is the proportion of

simulations in which ideal residuals are larger than true residuals. If the model fits the data well, the

Bayesian P-value is close to 0.5. Bayesian P values for our models ranged from 0.47 to 0.57 and were

on average 0.51, indicating excellent model fit to the data.

References

1. O. Terao, thesis, Tokyo (1985).

2. E. Edeline, O. Terao, K. Naruse, Empirical evidence for competition-driven semelparity in wild medaka. *Popul. Ecol.* **58**, 371–383 (2016).

3. W. E. Ricker, Stock and recruitment. *J. Fish. Res. Board Can.* **11**, 559–623 (1954).

4. I. Ntzoufras, *Bayesian modeling using WinBUGS* (Wiley, Hoboken (NJ), ed. 1, 2009).

5. M. Heino, U. Dieckmann, O. R. Godø, Measuring probabilistic reaction norms for age and size at maturation. *Evolution*. **56**, 669–678 (2002).

6. E. M. Olsen *et al.*, Maturation trends indicative of rapid evolution preceded the collapse of northern cod. *Nature*. **428**, 932–935 (2004).

7. M. Heino, U. Dieckmann, Detecting fisheries-induced life-history evolution: an overview of the reaction-norm approach. *Bull. Mar. Sci.* **83**, 69–93 (2008).

8. R Core Team, *R: a language and environment for statistical computing* (R Foundation for Statistical Computing, Vienna, Austria, 2018; https://www.R-project.org/).

9. M. Plummer, (Vienna, Austria, 2003).

10. K. Kellner, *jagsUI: a wrapper around "rjags" to streamline "JAGS" analyses* (2017; https://CRAN.R-project.org/package=jagsUI).

11. A. Gelman, D. B. Rubin, Inference from iterative simulation using multiple sequences. *Stat. Sci*, 457–472 (1992).

12. H. Chung, E. Loken, J. L. Schafer, Difficulties in drawing inferences with finite-mixture models: a simple example with a simple solution. *Am. Stat.* **58**, 152–158 (2004).

13. A. Gelman, X. L. Meng, H. Stern, Posterior predictive assessment of model fitness via realized discrepancies. *Stat. Sin.* **6**, 733–807 (1996).

475 **Table S1. Structure and MCMC parameter estimates for models 4-7.** The MCMC P-value is twice

476 the proportion of the posterior for which the sign was opposite to that of the mean posterior value. The

477 MCMC P-values is not relevant for variance parameters that are constrained to be non-zero.

Model	Response	N	Distribution	Link	Effect	Mean estimate	SD of the estimate	MCMC P-value
4	Larvae count	2004	Bernoulli in ZINB	logit	Int.	-8.309	0.942	0.000
					Slope of day	-0.189	0.021	0.000
			Negative binomial in ZINB	ln	Int. no-harvest	2.081	0.251	0.000
					Int. harvest	1.006	0.244	0.000
					Slope of day no-harvest	0.001	0.001	0.380
					Slope of day harvest	0.016	0.002	0.000
					Dispersion index no-harvest	8.702	2.089	
					Dispersion index harvest	2.417	0.359	
					SD of year by pond effect (random)	0.998	0.137	
5	Standard body length	1144	Gaussian	Identity	Int. no-harvest	4.410	0.106	0.000
					Int. harvest	4.548	0.099	0.000
					Slope of age no-harvest low food	0.224	0.005	0.000
					Slope of age harvest low food	0.210	0.005	0.000
					Slope of age no-harvest medium food	0.250	0.005	0.000
					Slope of age harvest medium food	0.231	0.005	0.000
					Slope of age no-harvest high food	0.263	0.005	0.000
					Slope of age harvest high food	0.248	0.004	0.000
					Slope of age squared	-0.001	0.000	0.000
					Int. residual variance no-harvest low food	-0.021	0.157	0.854
					Int. residual variance harvest low food	-0.549	0.127	0.001
					Int. residual variance no-harvest medium food	-0.597	0.155	0.002
					Int. residual variance harvest medium food	-0.384	0.130	0.004
					Int. residual variance no-harvest high food	-0.520	0.129	0.000
					Int. residual variance harvest high food	-0.295	0.145	0.043
					Slope of age residual variance no-harvest low food	-0.005	0.003	0.079
					Slope of age residual variance harvest low food	0.011	0.002	0.000
					Slope of age residual variance no-harvest medium f	0.000	0.004	0.923
					Slope of age residual variance harvest medium food	0.010	0.002	0.000
					Slope of age residual variance no-harvest high food	0.011	0.002	0.000
					Slope of age residual variance harvest high food	-0.011	0.003	0.001
					SD of parental pair effect on int. (random)			
					SD of parental pair on slope of Age effect (random)			
6	Maturation probability	591	Bernoulli	logit	Int. no-harvest	-4.138	0.698	0.000
					Int. harvest	-4.762	0.771	0.000
					Slope of age no-harvest	-0.054	0.025	0.020
					Slope of age harvest	0.055	0.024	0.025
					Slope of length no-harvest	1.662	0.286	0.000
					Slope of length harvest	1.521	0.271	0.000
					SD of parental pair effect on int. (random)	1.470	0.356	
7	Prey count	311	Bernoulli in ZINB	logit	Int.	-1.960	0.541	0.000
					SD of individual effect (random)	0.903	0.534	
			Negative binomial in ZINB	ln	Int. no-harvest, low food	2.035	0.208	0.000
					Int. harvest, low food	1.848	0.231	0.000
					Int. no-harvest, medium food	1.928	0.245	0.000
					Int. harvest, medium food	0.986	0.286	0.001
					Int. no-harvest, high food	0.357	0.270	0.188
					Int. harvest, high food	0.672	0.309	0.025
					Dispersion index no-harvest, low food	2.388	0.722	
					Dispersion index harvest, low food	5.994	2.141	
					Dispersion index no-harvest, medium food	6.509	3.857	
					Dispersion index harvest, medium food	5.012	2.357	
					Dispersion index no-harvest, high food	2.033	0.710	
					Dispersion index harvest, high food	5.708	2.642	
					SD of individual effect (random)	0.681	0.136	