Analysis of movement recursions to detect reproductive events and estimate their fate in central
place foragers

Simona Picardi[1*], Brian J. Smith[2], Matthew E. Boone[1], Peter C. Frederick[3], Jacopo G. Cecere[4],

Diego Rubolini[5], Lorenzo Serra[4], Simone Pirrello[4], Rena R. Borkhataria[6], Mathieu Basille[1]

[1]Department of Wildlife Ecology and Conservation, Fort Lauderdale Research and Education

Center, University of Florida, 3205 College Ave, Davie, FL 33314, USA

[2]Deparmtent of Wildland Resources, Ecology Center, Utah State University, Logan, UT 84322,

USA

[3]Department of Wildlife Ecology and Conservation, University of Florida, 368 Newins-Ziegler

Hall, Gainesville, FL 32611, USA

[4]Area Avifauna Migratrice, Istituto Superiore per la Protezione e la Ricerca Ambientale

(ISPRA), via Cà Fornacetta 9, I-40064 Ozzano Emilia (BO), Italy

[5]Dipartimento di Scienze e Politiche Ambientali, Università degli Studi di Milano, via Celoria

26, I'20133 Milan, Italy

[6]Department of Wildlife Ecology and Conservation, Everglades Research and Education Center,

University of Florida, 3200 E Palm Beach Rd, Belle Glade, FL 33430, USA

[*]Corresponding author – simona.picardi@ufl.edu

Abstract

   Recursive movement patterns have been used to detect behavioral structure within individual movement trajectories in the context of foraging ecology, home-ranging behavior, and predator avoidance. Some animals exhibit movement recursions to locations that are tied to reproductive functions, including nests and dens; while existing literature recognizes that, no method is currently available to explicitly target different types of revisited locations. Moreover, the temporal persistence of recursive movements to a breeding location can carry information regarding the fate of breeding attempts, but it has never been used as a metric to quantify recursive movement patterns. Here, we introduce a method to locate breeding attempts and estimate their fate from GPS-tracking data of central place foragers. We tested the performance of our method in three bird species differing in breeding ecology (wood stork (*Mycteria americana),* lesser kestrel (*Falco naumanni),* Mediterranean gull (*Ichthyaetus melanocephalus*)) and implemented it in the R package 'nestR'. We identified breeding sites based on the analysis of recursive movements within individual tracks. Using trajectories with known breeding attempts, we estimated a set of species-specific criteria for the identification of nest sites, which we further validated using non-reproductive individuals as controls. We then estimated individual nest survival as a binary measure of reproductive fate (success, corresponding to fledging of at least one chick, or failure) from nest-site revisitation histories during breeding attempts, using a Bayesian hierarchical modeling approach that accounted for temporally variable revisitation patterns, probability of visit detection, and missing data. Across the three species, positive predictive value of the nest-site detection algorithm varied between 87-100% and sensitivity between 88-92%, and we correctly estimated the fate of 86-100% breeding attempts. By providing a method to formally distinguish among revisited locations that serve different ecological functions and introducing a probabilistic framework to quantify temporal persistence of movement recursions, we demonstrated how the analysis of recursive movement patterns can be applied to estimate reproduction in central place foragers. Beyond avian species, the principles of our method can be applied to other central place foraging breeders such as denning mammals. Our method estimates a component of individual fitness from movement data and will help bridge the gap between movement behavior, environmental factors, and their fitness consequences.

   Keywords

   breeding-site detection – fitness – GPS telemetry – Bayesian hierarchical models – nest survival – nestR – R package – recursive movement patterns

Background

A major goal of movement ecology is to uncover behaviors underlying movement trajectories (Gurarie et al. 2016). Knowing what animals are doing when moving a certain way can improve our understanding of the links between movement and resource dynamics, species interactions, distribution, and individual fitness (Mueller and Fagan 2008, Schick et al. 2008, Morales et al. 2010). In the past decade, the availability of animal tracking data has increased exponentially and, in parallel, methodological approaches to infer behavior from movement have proliferated (Wilmers et al. 2015).

One way to detect behavioral structure in movement trajectories is to analyze movement recursions (Berger-Tal and Bar-David 2015). Recursive movement patterns are common across taxa, as animals exhibit periodic returns to places of ecological significance such as nests, dens, foraging patches, etc. (Riotte-Lambert et al. 2013, Berger-Tal and Bar-David 2015, Bracis et al. 2018). By analyzing recursive movement patterns, researchers can identify the location of ecologically relevant places and unlock the signature of underlying ecological processes, such as responses to temporally variable resources (Bar-David et al. 2009, Berger-Tal and Bar-David 2015). In this paper, we demonstrate the use of movement recursions to detect reproductive events and estimate their fate from the trajectories of central place foragers.

Berger-Tal and Bar-David (2015) reviewed the literature on recursive movement patterns and identified three main lines of research: traplining behavior, path recursions, and the ecology of fear. While these lines of research focus on different spatial scales and have traditionally focused on different taxa, the common thread among them is the analysis of the relationship between movement recursions and temporal variation in resource availability, alone or in trade-off with predator avoidance (Berger-Tal and Bar-David 2015). For example, Garrison and Gass (1999) showed how the long-billed hermit (*Phaethornis longirostris*), a traplining hummingbird, adjusts the rate of revisitation to feeders according to changes in food delivery rates; Bar-David et al. (2009) analyzed path recursions to identify individual responses to cycles of vegetation depletion and replenishment in African buffalos (*Syncerus caffer*), and English et al. (2014) in forest elephants (*Elephas maximus borneensis*); Riotte-Lambert et al. (2013) linked recursive movement patterns of an impala (*Aepyceros melampus*) to open areas with temporally varying predation risk. As Berger-Tal and Bar-David's (2015) review illustrates, most of the existing research on recursive movement patterns is deeply rooted in foraging ecology, with ties to optimal foraging theory, home-ranging behavior, and predator-prey interactions.

From a technical standpoint, methods to identify places of ecological relevance along individual

trajectories are often based on the use of spatial buffers around visited locations (Barraquand and

Benhamou 2008, Bracis et al. 2018). Other approaches that do not entail the use of spatial buffers exist,

such as detecting recursions as closed paths within a trajectory (Bar-David et al. 2009) or using

utilization-distribution based methods (Benhamou and Riotte-Lambert 2012). The advantage of buffer-

based approaches is that they allow simple computation of metrics describing the temporal patterns of

site revisitation (Bracis et al. 2018).

Temporal patterns of revisitation are used to uncover behavioral signatures underlying

movement recursions (Barraquand and Benhamou 2008, Riotte-Lambert et al. 2013, Bracis et al. 2018).

Different temporal metrics have proven useful to quantify recursive movement patterns for different

objectives. Residence time, i.e., the cumulative time an animal spends at a certain location or area, has

been used to quantify the profitability of resource patches, with high-quality patches attended for

longer stretches of time (Barraquand and Benhamou 2008). Similarly, the frequency of visits has also

been used to quantify the intensity of use of different areas within animals' home ranges, with

individuals visiting profitable patches more often (Benhamou and Riotte-Lambert 2012). Conversely,

shorter times between visits (or time-to-return) characterize preferred areas within an animal's home

range (Van Moorter et al. 2016), as well as lower speeds (Nelson et al. 2015). The periodicity of

movement recursions, i.e., the temporal repeatability of visits to a certain location, detected using

Fourier- and wavelet-based methodologies, has been used to quantify responses of animals to cyclical

variation in resource availability and/or predation risk (Riotte-Lambert et al. 2013). More complex

analyses of movement recursions involve quantifying repetitiveness in the sequence of returns to

multiple locations (Riotte-Lambert et al. 2017). Recent software developments, such as the 'recurse' R

package (Bracis et al. 2018), have made it easier for researchers to quantify spatio-temporal patterns of

movement recursions along individual tracks by identifying recurrently visited locations and

calculating metrics such as number of visits, visit duration, intervals between visits, and residence time.

Important gaps remain in our current approach to the study of movement recursions. First, from

a conceptual standpoint, most of the existing work has used movement recursions to gain insight on

foraging ecology, home-ranging behavior, and responses to resource dynamics (Berger-Tal and Bar-

David 2015). However, recursive movement patterns may lock away the signatures of other behavioral

processes as well, including reproduction; yet, this application remains unexplored. For example, in

central place foragers (Orians and Pearson 1979), the central return location is often associated with

reproductive activities, like in the case of nests for birds (Andersson 1981, Kacelnik 1984, Burke and

Montevecchi 2009) or dens for some mammals (Frame et al. 2004, Castillo et al. 2011, Olson et al.

2011). The potential for recursive movement patterns to provide information on reproduction has been

so far neglected. Current literature acknowledges that, besides foraging patches, recurrently visited

locations also include nests and dens (Bracis et al. 2018), but it makes no formal effort to provide ways

to explicitly distinguish between different types of revisited locations. Without this ability, current

approaches cannot take advantage of recursive movement patterns to specifically detect reproductive

events, which is nonetheless a relevant application in the case of central place foragers. Other studies

have analyzed movement tracks with the objective of detecting reproductive events along them in

ungulates (caribou (*Rangifer tarandus),* DeMars et al. 2013, Bonar et al. 2018; moose (*Alces alces*)

Nicholson et al. 2019). However, these species do not necessarily exhibit movement recursions because

of reproductive behavior; rather, the signature of reproductive events consists in a slow-down of

movements due to limited motion capacity of calves, and thus the appropriate diagnostic metric used to

detect reproduction in these species was movement rate (DeMars et al. 2013, Bonar et al. 2018,

Nicholson et al. 2019). Second, while researchers have devoted effort to calculating and interpreting

temporal patterns of movement recursions in terms of metrics such as residence time and periodicity

(e.g., Barraquand and Benhamou 2008, Riotte-Lambert et al. 2013), little or no attention has been given

to another temporal metric that can prove informative: the persistence of movement recursions, i.e., the

time until the recursive movement behavior to a certain location ceases. In the case of reproduction in

central place foragers, the temporal persistence of returns to a nest or den may inform us of the fate of

reproductive attempts. For example, many bird species act as central place foragers while breeding,

leaving the nest to go forage and returning to incubate their eggs and bring back food for their offspring

(Kacelnik 1984, Alonso et al. 1994, Burke and Montevecchi 2009). Altricial bird species keep returning

to the nest (often with varying frequency throughout the development of nestlings, e.g., Wiley and

Wiley 1980) until their young are independent (i.e., until fledging or for some time after that; Grüebler

and Naef-Daenzer 2008, Tarwater and Brawn 2010). If their attempt fails, i.e., the eggs are lost or

nestlings die before fledging, altricial bird parents abandon the nest and stop visiting its location

(Calder 1973, Stake and Cimprich 2003, Zangmeister et al. 2009); in some cases, nest abandonment

itself can be the cause of failure (Nelson and Hamer 1995, Sabine et al. 2006, Becker et al. 2010).

Thus, the persistence of recursive movements around the nest site can inform us of the fate of breeding

attempts in altricial bird species, based on whether returns to the nest site persist for a duration

compatible with the full developmental cycle of nestlings or cease before it is completed. The challenge

with using the persistence of recursive movement patterns to pinpoint the cessation of an underlying

behavior lies in the fact that movement data are collected as discrete locations rather than continuously,

and thus visits to a location of interest may be missed if they happen between fixes. Moreover, in the

case of breeding birds, nest attendance often decreases as the nestlings grow older (Clark 1980, Cadiou

and Monnat 1996, Weimerskirch et al. 2001), which may make the daily detectability of nest visits

lower later in the attempt. To account for imperfect detection of visits, the persistence of movement

recursions needs to be conceptualized in a probabilistic framework.

In this paper, we present a method to apply the analysis of movement recursions to 1) detect

reproductive events along the trajectories of central place foragers and 2) estimate their fate. The first

step is necessary for the second, but the method serves both purposes. Unlike other available methods

for the identification of revisited sites (e.g., Bracis et al. 2018), our method allows to distinguish

between different types of revisited locations and isolate those of interest, thus addressing an important

shortcoming of current approaches. Moreover, our method introduces the persistence of movement

recursions as a useful metric to detect the cessation of behaviors of interest and allow inference on

latent processes. Specifically, we use the persistence of movement recursions to estimate the fate of

reproductive events in central place foragers, accounting for imperfect detection of visits to the central

location. Our method thus demonstrates the use of recursive movement patterns to infer a component of

individual fitness. Given the central role of fitness in ecological and evolutionary processes, estimating

components of it from movement data is a long-sought goal in movement ecology (Morales et al. 2004,

Singh and Ericsson 2014). As an illustration, we apply our method to identify nest sites and estimate

the fate of individual breeding attempts using real GPS-tracking data for three avian species differing in

their breeding habitat and ecology: wood storks (*Mycteria americana*), lesser kestrels (*Falco*

naumanni), and Mediterranean gulls (*Ichthyaetus melanocephalus*).

Methods

Method description

Our method is composed of two parts: first, the detection of breeding sites, and second, the

estimation of reproductive fate (Fig. 1). The workflow is implemented in the R package 'nestR', which

was designed with avian species in mind (https://github.com/picardis/nestR; Picardi et al. 2019).

Although the purpose of this paper is to introduce the method *per se*, we included a synthetic

description of the package structure and functions in Additional file 1; Additional file 2 provides the

complete package documentation; moreover, the 'nestR' package vignette includes worked examples of

our study cases (Additional file 3).

Nest-site detection – The objective of Part I is to identify nest sites (and thus, breeding events)

along individual trajectories (Fig. 1). Nest sites are identified as repeatedly visited locations along

individual trajectories (Fig. 1, "Find recurrently visited locations"). Returns to a location are defined as

returns to a circular area of a user-defined radius buffering each point of the trajectory (Fig. 2A). Using

buffers accounts for the spatial scattering of GPS points around the actual central location due to both

behavior and GPS error (Frair et al. 2010). The buffer size sets the spatial scale at which revisitation

patterns will be calculated and should be chosen according to the expected scale of movements, which

should be small in the case of a nest (compared, for example, to returns to a same foraging area but not

exact location; Bracis et al. 2018). We recommend using this rationale to choose a reasonable buffer

size for the initial round of analysis; then, we recommend performing a sensitivity analysis on the

effect of buffer size on nest-detection performance and refining the spatial scale of the analysis to

optimize performance, if necessary (Additional file 4).

For the purpose of distinguishing between different types of revisited locations, we describe

revisitation patterns using the following set of temporal parameters: the maximum number of

consecutive days a location is visited; the percentage of days it is visited between the first and last visit;

and the percent fixes at a location on the day with longest attendance (we later refer to this as the "top

day", in short). We calculate these parameters for every revisited location that we initially detect, each

of which is a potential nest site. Parameter values are then used as diagnostic features to filter actual

nest sites among revisited locations (Fig. 1, "Find nests among revisited locations"), based on the

rationale that revisitation patterns differ between nest and non-nest sites (i.e., other types of revisited

locations such as foraging patches, roosts, etc.). Nest sites are often visited for longer stretches of

consecutive days, on more days, and more frequently or for longer within a day than other types of

revisited locations; the parameters we use to describe revisitation patterns are meant to capture these

diagnostic behaviors. The method can be tailored to different case studies by restricting the analysis

within the breeding season for a given species and accounting for data sampling rate and fix failure rate

(Additional files 2 and 3).

Several approaches are possible to identify sets of parameter values to distinguish nest from

non-nest sites. The simplest option is to use prior knowledge of the behavior of the focal species or

expert opinion (Fig. 1, "Prior information on revisitation patterns of nests?"). If such information is

available, users can directly skip to the final step of the first part of the workflow (Fig. 1, "Find nests

among revisited locations"). Alternatively, we propose a data-driven approach to agnostically identify

optimal parameter values based on Classification And Regression Trees (CART; De'ath and Fabricius

2000). CART can be used to find the optimal set of parameter values that discriminate between nest

sites and other types of revisited locations, based on the comparison of known nest and non-nest sites

(Fig. 1, "Find revisitation patterns that discriminate between nests and non-nests"). If the true location

of nests is known for a subset of the data (Fig. 1, "Coordinates known for some nests?"), researchers

can use these known nests and compare their revisitation patterns with those of other locations that are

not nests. If on-ground data are not directly available, an alternative is to visually explore the data and

identify trusted nest sites, where possible (Fig. 1, "Visualize data to find trusted nests"). For example,

likely nest sites may be recognized in some species based on habitat features or proximity to known

breeding colonies. Once known or trusted nest sites are identified, non-nest sites can be selected based

on a criterion of temporal overlap (Fig. 1, "Select nests and non-nests to compare"); locations revisited

simultaneously with a breeding attempt can be considered non-nest sites, assuming birds cannot breed

in two places at the same time (which may not be true in all study systems). Once a dataset of known

nest and non-nest sites is built, users can apply CART to this dataset and prune the tree to the optimal

number of nodes based on a minimum relative error criterion (De'ath and Fabricius 2000). More

sophisticated classification tools, such as random forests (Breiman 2001), may also be appropriate for

this task, but CART has the advantage of providing outputs that are easy to interpret biologically, as it

returns numeric cut-offs for each of the input parameters that most efficiently split the data between the

groups of interest.

The (one or more) sets of parameters identified by CART as best discriminating between nest

and non-nest sites can be manually applied to the complete set of revisited locations to identify nest

sites among them (Fig. 1, "Find nests among revisited locations"; Fig 2B). In case multiple nests that

satisfy the criteria are identified for one individual, if any of them temporally overlap with each other

(and again assuming birds cannot breed in two places at the same time), the recommended option is to

pick the most visited candidate and discard those that temporally overlap with it.

Because subsequent analyses rely on accurate nest-detection results, it is of paramount

importance that users verify that performance of the nest-detection algorithm is satisfactory before

proceeding with the estimation of reproductive fate. Both failing to identify some nests or identifying

spurious ones might result in mis-estimating reproductive fate down the line. On one hand, failure to identify a nest in an individual's track would lead to concluding that that individual did not attempt to breed; the consequences of this wrong conclusion might weigh more or less according to, first, whether the missed attempt was successful or not, and second, whether we are interested in assessing net reproductive success or relative to effort. For instance, if we are not interested in measuring reproductive fate relative to effort, concluding that an individual did not attempt to breed or attempted and failed does not make a difference in terms of net output, but it does when considered in light of return per energy investment. Both behavior and data quality might affect the probability to miss a nest: for example, the risk of failing to detect a nest is higher if data resolution is low or if nest visits are infrequent. In most cases, we would expect successful attempts to not be missed except in case of persistent malfunctioning of a GPS tag; thus, missed detections are likely to be biased towards failed attempts – especially early failures, where the nest was not visited for long enough to be picked up by the algorithm. On the other hand, identifying as a nest a revisited location which is actually not a nest might lead to concluding that an individual bred when it actually did not or failed at breeding when it actually did not attempt. This type of error is affected by behavior more than data characteristics, and it may be more likely for some species than others: for example, the signature of recursive movements around a nest may be more likely to be confounded with non-breeding movements in perch-and-wait predators, who spend long and frequent stretches of time on the same perch, than in seabirds, who spend most of their time wandering at sea when not breeding. The gravity of detection errors depends on specific study objectives (e.g., measuring net fate vs. relative to effort) and committing one type of error might be more undesirable than committing the other; we recommend that researchers think critically about how much they care about minimizing each and interpret performance metrics accordingly.

We propose four metrics to quantify nest-detection performance: positive predictive value, i.e., the proportion of true nest sites among those that are found; sensitivity, i.e., the proportion of true nest sites that are successfully identified; false negative rate, i.e., the proportion of true nest sites that the algorithm failed to identify; and false positive rate, i.e., the proportion of non-nest sites that are identified as nests. The first three metrics can be evaluated using data on known or trusted nest-site locations; the fourth metric can be assessed using non-breeder trajectories, for example outside of the breeding season or for individuals under reproductive age.

Reproductive fate estimation – Part II of the workflow addresses the second objective of our

method: estimating the fate of the reproductive events identified in Part I (Fig. 1). The fate of breeding

attempts is estimated based on the temporal persistence of returns to the nest: essentially, we are asking

whether the tracked individual kept revisiting the nest for long enough to carry the attempt to term,

based on how long the developmental cycle is in the focal species. A breeding attempt is considered

successful if the nest site was visited until the end of a complete breeding cycle for the focal species,

which includes nest-building, egg-laying, incubation, and chick-rearing until the nestlings reach

autonomy and no longer receive parental care. Thus, we define success of a breeding attempt as

survival of at least one nestling until fledging and failure to none. First, for each nest site identified in

Part I, we compile a history of nest revisitation (Fig. 1, "Format nest revisitation histories"), by

converting the movement track to a presence/absence time series at the nest (0/1 for each GPS point).

This presence/absence time series will then be treated as capture-recapture data and used to estimate

the fate of each breeding attempt using a survival analysis in a Bayesian hierarchical modeling

framework (Fig. 1, "Run nest survival models"). Modeling persistence of nest revisitation with a

hierarchical modeling approach allows us to take into account imperfect detection of nest visits and

missing GPS fixes, as described in the following paragraph.

The nest survival model specification includes two processes: the survival process, which is not

directly observable, and the observation process, which is the revisitation history. Much like a Bayesian

implementation of a Cormack-Jolly-Seber capture-mark-recapture model (Lebreton et al. 1992, Schaub

and Royle 2014), the latent nest survival variable, z, is modeled as a Bernoulli variable at the daily

scale as a function of survival status and daily survival probability, φ, at the previous time-step:

$$z_t \sim Bern(z_{t-1} \times \phi_{t-1})$$

288   Observed visits on a given day are modeled as a binomial variable as a function of current nest

289 survival status, probability of visit detection, p, and number of GPS fixes available, N, on day t:

$$Y_t \sim Bin(N_t, z_t \times p_t)$$

290   Where the probability of detection is conditional to N and to the nest being alive on that day:

$$p_t = Pr(visit detected \lor z_t = 1, N_t)$$

291   Reproductive fate is defined as the probability P that the nest was still surviving on the last day

292 of the theoretical duration of a complete breeding attempt, T:

$$P = Pr(z_T = 1)$$

293   Both survival and detection probability are modeled using a binomial Generalized Linear Model

294   (GLM) as a function of the day of the attempt:

$$logit(\phi_t) = \beta_{\phi_0} + \beta_{\phi_1} \times t$$
$$logit(p_t) = \beta_{p_0} + \beta_{p_1} \times t$$

The model is fully specified by using uninformative priors on the β parameters, in this case a

normal distribution with a mean of 0 and precision of 1e-5. In the current implementation, daily

survival and detection are assumed to be the same for all nests in the population. The model outputs

daily estimates of survival and detection probability at the population level, as well as daily survival

estimates for each breeding attempt along with credible intervals. From these values, results can be

summarized in different ways: for instance, they could be expressed in terms of value of P on the final

day of the breeding cycle for each attempt, or the last day when P was greater than a chosen threshold

(e.g., 90%), and so on. The MCMC (Markov Chain Monte Carlo) algorithm is implemented in JAGS

(Just Another Gibbs Sampler; Plummer 2003) via the R package 'rjags' (Plummer 2018).

Assumptions underlying the nest survival model include: 1) birds are tracked for the entire

duration T of a complete nesting attempt (if birds were tagged part-way through an attempt, T needs to

be adjusted by subtracting the age of the nest (in days) at tagging); 2) the GPS tag does not

permanently fail before the end of the attempt; 3) parents visit the nest until fledging, or nestling

mortality is negligible between the time when parental care is interrupted and fledging; 4) parents stop

visiting a nest after failing.

<u>Study cases</u>

We applied our method to GPS-tracking data for 148 individual-years for wood storks

(henceforth storks), 56 for lesser kestrels (henceforth kestrels) and 29 for Mediterranean gulls

(henceforth gulls; Table 1). All tags were solar-powered and recorded fixes primarily during daytime.

Details about devices, settings, harnesses, and study areas regarding storks and kestrels can be found in

Borkhataria et al. (2008) and Cecere et al. (2018), respectively. To find nest sites, we restricted the

analysis to the breeding season only for each species (Table 1). While both kestrels and gulls have a

well-defined breeding season between April and August in our study areas (Snow et al. 1997), storks in

the southeastern U.S. can breed at slightly different times of the year depending on latitude (Coulter et

al. 1999; Table 1). In this case, we used a conservative approach and only excluded the period where no

breeding activities were expected to occur anywhere in the range.

Given the spatial resolution of the GPS data (Table 1) and the expected scale of movements

around the nest site for all three species, we initially ran the nest-detection algorithm using a buffer of

40 m around each GPS position. We then ran a sensitivity analysis on buffer size using values between

10 and 100 m (Additional file 4), picked the radius that maximized performance for each species, and

re-ran the analysis to refine results.

We initially screened trajectories for any revisited locations using non-constraining values for

parameters describing revisitation patterns (thus not applying any filtering). We then used on-ground

data on known nest locations to select true nests and non-nest sites from the revisited locations.

Kestrels and gulls were captured at the nest site (Table 1), so the location of the nest was known. For

storks, on-ground data on nest locations was available for 10 individual-years (Bear D., unpublished

data). We explored the remaining stork trajectories and identified those for which the top visited

location was at a known breeding colony (data from U.S. Fish and Wildlife Service 2018). We marked

these as trusted and treated them as known nest sites for the rest of the analysis.

We used CART to compare revisitation patterns between nest and non-nest sites, and used the

resulting sets of parameter values to filter nest sites among revisited locations in the trajectories of

breeding individuals. We only retained individual-years where data exceeded the minimum number of

consecutive days visited indicated by CART (Table 1). Even when CART did not suggest that the

number of consecutive days visited was an important predictor of true nest sites, we chose a reasonable

value to use as a threshold for this parameter, as we did not expect to have enough power to discern

nest from non-nest sites for attempts that failed in the first handful of days. We only retained the

candidate with the most visits among any sets of breeding attempts that were temporally overlapping.

We used non-breeder trajectories (sub-adults in the case of storks, non-breeding season data in the case

of kestrels and gulls) to validate our results against false positives. We calculated positive predictive

value of our algorithm as the percentage of nest sites that were known among those we found for each

species; sensitivity as the percentage of known nest sites that were identified; false negative rate as the

percentage of known nest sites that we failed to identify; and false positive rate as the percentage of

non-breeding individual-years for which we erroneously identified a nest site.

We fit the nest survival model described above to estimate the fate of identified breeding

attempts, using only individual-years for which the tag was active throughout the attempt to meet

model assumptions (Table 1). Since kestrels and gulls were captured after they had already started

breeding (immediately after hatching and in late incubation, respectively, although the exact age of the

nest at tagging was unknown), the initial part of every breeding attempt was missing from the

movement data. To account for this, we subtracted the theoretical number of days until hatching (for

kestrels, 25 days) and late incubation (for gulls, 20 days) from the value of T (Table 1). We evaluated

performance of the method by comparing survival estimates to known fates.

Results

The initial screening with no filtering identified 9871 revisited locations (i.e., potential nest

sites) for storks, 511 for kestrels, and 1379 for gulls. Results from CART showed that the optimal set of

parameter values to discriminate nest from non-nest sites was 14 minimum consecutive days visited

and 79% minimum nest attendance on the top day for storks, 7 minimum consecutive days visited for

kestrels, and 26% minimum attendance on the top day for gulls (Fig. 3). Because CART did not

indicate a minimum number of consecutive days visited for gulls, we added a reasonable constraint for

this value by exploring the data and determining which value would allow us to rule out most non-nest

sites while retaining most nest sites (8 days). Based on results of the sensitivity analysis, we used

buffers of 30, 60, and 20 m for storks, kestrels, and gulls, respectively (Additional file 4). By filtering

revisited locations using these optimal buffer sizes and the parameter values indicated by CART, we

identified 108 nest sites for storks, 46 for kestrels, and 24 for gulls, which closely matched the number

of nest sites we were expecting to find (Table 1). As a consequence, the positive predictive value of the

algorithm ranged between 87-100%, the sensitivity between 88-92%, and the false negative rate

between 8-12% for the three species (Table 2). The false positive rate was 2% for storks and 0% for

gulls but reached 50% for kestrels (Table 2). The probability of detecting nest visits decreased

throughout the breeding attempt for all three species, while survival remained constant (Fig. 4). We

correctly estimated the fate of 100% of breeding attempts for gulls and 86% for kestrels (1 failure and 2

successes incorrectly estimated; Fig. 5). No data on true fates were available for storks, therefore we

were unable to verify survival estimates for this species.

Discussion

We presented a method to detect reproductive events and estimate their fate from recursive

movement patterns of central place foragers. As an illustration, we applied our method to identify nest-

site locations and estimate the fate of breeding attempts from GPS-tracking data for three avian species

differing in breeding behavior and ecology (Fasola and Canova 1992, Coulter et al. 1999, Gustin et al.

2017). To our knowledge, this is the first time the analysis of movement recursions is applied to infer

reproduction; more broadly, ours is also among the first attempts to infer the reproductive component

of fitness from telemetry data (but see DeMars et al. 2013, Bonar et al. 2018, Nicholson et al. 2019),

and to our knowledge, the first applied to birds. Our method broadens the applicability of current

approaches and tools for the analysis of movement recursions, opening the possibility of inferring

reproduction in addition to traditionally targeted latent processes such as foraging behavior and home-

ranging (Berger-Tal and Bar-David 2015). The application of recursive movement analysis to

estimating the fate of reproduction is made possible by introducing persistence of movement recursions

as a metric to detect the cessation of latent behaviors. Conservation and management applications may

both benefit from the availability of our method and its implementation in an open-access, user-friendly

R package, 'nestR'. Knowledge of the biology and ecology of the target species and careful

consideration of data characteristics and limitations are critical for successful use of the tools we

presented.

Performance of nest-site detection – Our nest-site detection method performed well for all three

species, allowing us to correctly identify most or all known nest sites from movement trajectories of

breeding individuals. We achieved high positive predictive value (87-100%) and sensitivity (88-92%)

for all species. Importantly, the positive predictive value quantifies how many of the nest sites we

found were known, which does not necessarily imply that the remaining were non-nest sites: it is

possible that those we were unable to confirm for storks and gulls included second attempts (true but

unknown nest sites) in addition to non-nest sites, as both species may attempt to breed again at a

different location if their first clutch fails early in the season (Rodgers et al. 1987, Cecere pers. obs.). In

support of this possibility, all unknown nest sites we found for gulls were from birds whose known

attempt failed early on, and they were thus plausible second attempts. The same might be true for

storks, although we did not have on-ground data to confirm it. False negative rates were low for all

species (8-12%) and mostly associated with early failures: 2 out of 2 nest sites that we failed to identify

for gulls and 2 out of 5 for kestrels corresponded to attempts that failed before the enforced limit of

consecutive days visited (as early as the day after tagging in the case of gulls). This may be true for

storks as well, where the breeding attempts we were unable to identify might have failed before the 14-

day mark. Not identifying breeding attempts whose duration does not exceed the minimum constraint

applied is a logical implication of the approach rather than a failure of the algorithm. The remaining 3

nest sites that we were unable to identify for kestrels did not fail within the first week, but were never

visited for 7 consecutive days. False positives were none or negligible for gulls and storks (0% and 2%

respectively), but reached 50% for kestrels. This is likely explained by species-specific behavior:

kestrels spend long stretches of time and consecutive days on a perch while scanning for prey or resting

(Hernández-Pliego et al. 2017). Distinguishing these patterns of attendance and revisitation from those

of a nest might be challenging without applying restrictions based on seasonality and geographical area

(e.g., breeding versus wintering range).

Error rates for nest-site identification vary in importance depending on the study objectives. If

the objective is to estimate reproductive fate (especially if regardless of effort), ensuring that attempts

are not missed should receive priority over avoiding the selection of non-nest sites. Any revisited

location that gets erroneously identified as a nest site would likely be classified as a failed attempt

eventually anyway. In this case, we suggest that researchers may want to focus on minimizing false

negatives. Conversely, if the objective of a study is, for instance, to analyze factors associated with

nest-site selection, minimizing false positives should be the priority.

Once on-ground data on nest locations are used to identify parameter values to distinguish nests

among revisited locations, these parameter values can then be applied to new individuals of the same

species for which on-ground information is not available, assuming other data characteristics are the

same. If CART is the tool of choice to inform the choice of parameter values, we recommend that

parameter values suggested by the algorithm should be critically evaluated for their biological

significance before use, and that adjustments should be made as needed based on knowledge of the

species biology. Future efforts to improve our method for the identification of nest locations will

include incorporating uncertainty in our estimates of nest sites, allowing us to interpret classification

results in a probabilistic framework.

Performance of reproductive fate estimation – We correctly estimated reproductive fate of

100% of breeding attempts for gulls and 86% for kestrels, with probability of success estimated as $P >$

0.97 for true successes and as $P = 0$ for true failures. The remaining attempts were two successes that

we estimated as failures ($P <= 0.3$) and one failure that we estimated as a success ($P = 0.98$). The two

attempts that we erroneously estimated as failures corresponded to one male and one female kestrel

whose original clutch included four eggs and was partially lost, leading to two and one fledglings,

respectively. Partial brood loss may lead to a faster completion of the breeding cycle (Stenning 1996),

which may have compromised our ability to detect these attempts as successful as they did not reach

the benchmark $T = 60$. Specifically, one of the two attempts was completed within 27 days of tagging,

which corresponds to $T = 52$. However, the other attempt was completed within 33 days of tagging,

corresponding to $T = 58$, which is a similar duration to other successful attempts that we estimated

correctly. In this case, our inability to recognize the attempt as successful might have depended on

behavioral differences between parents, whereby the male we were tracking might have interrupted

parental care before the female did. This result highlights the importance of taking into account sex

differences in breeding behavior, where that applies (Hernández-Pliego et al. 2017). For example, in

species exhibiting uniparental care, inference should only be based on the sex that carries out parental

care (e.g., Birks 1997). The failed attempt that we erroneously estimated as successful corresponded to

a male that occasionally visited the nest site after failing, thus violating one of the assumptions of our

model. Unfortunately, this was the only failed attempt for kestrels in our dataset, which makes it

difficult to generalize our ability to estimate nest failures for this species. Overall, the three instances of

incorrect estimation might suggest that model assumptions, such as interruption of nest visits after

failure, might not always hold across species; or that the duration of a complete breeding cycle may be

too variable to lend itself to generalizations in some species; or that not knowing the exact age of the

nest at tagging might have reduced our power to distinguish late failures from successful attempts that

were completed in less-than-average time.

We did not have on-ground data to validate our estimation of reproductive fate for storks;

however, most attempts were estimated as either $P = 1$ or $P = 0$, while intermediate values (between

0.25 and 0.75) were relatively rare (14 out of 109). This is an important result given that data for storks

were at lower temporal resolution compared to kestrels and gulls (Table 1). Low temporal resolution of

data in combination with decreasing frequency of nest visits can, in principle, increase the uncertainty

of fate estimation by reducing probability of visit detection especially towards the end of a breeding

attempt (Fig. 4). Thus, the higher proportion of intermediate values for estimates of breeding success

probabilities we observed in storks compared to kestrels and gulls was to be expected, but results were

still rather polarized, suggesting that the method is largely able to distinguish between successes and

failures at this temporal resolution, given the frequency of nest visits in storks.

Innovations in the analysis of movement recursions – Our method introduces key innovations

compared to current approaches to the analysis of recursive movement patterns. First, compared to

other point-based methods, our method provides the ability to distinguish between locations that are

revisited for different purposes and are thus functionally different from one another from an ecological

point of view. Bracis et al. (2018) acknowledge that revisited locations can include foraging patches,

roosts, nests, dens, etc. and they speculate post-hoc on what type of revisited locations are those they identify for the vulture in their study, but their method does not provide a way to explicitly tell them apart. Similarly, Riotte-Lambert et al. (2017) recognize that foraging patches are not the only type of revisited location, but they do not attempt to distinguish between these types *a-priori*, rather they interpret the ecological function of recursive movements *post-hoc*. Other approaches (e.g., Bar-David et al. 2009, English et al. 2014) focus on quantifying individual foraging behavior while implicitly assuming that any revisited location is a foraging patch (which may be a valid assumption in species that do not exhibit recursive movements when reproducing). Finally, approaches that aim to quantify the intensity of use of different areas within individual home ranges usually do not attempt to discriminate between areas used for different functions (e.g., Barraquand and Benhamou 2008, Benhamou and Riotte-Lambert 2012, Van Moorter et al. 2016). Second, our method draws attention to a neglected temporal metric, i.e., the persistence of movement recursions. Previous studies focused on inferring latent processes from recursive movement patterns use residence time (Barraquand and Benhamou 2008), time-to-return (Moorter et al. 2016), periodicity (Riotte-Lambert et al. 2013), or repeatability of visitation sequences (Riotte-Lambert et al. 2017) as temporal metrics, but not the persistence of movement recursions. We introduced a probabilistic approach to quantify the persistence of recursions to a central location (nest site) and to estimate the cessation of a latent behavior (administering parental care to nestlings) using a survival analysis in a Bayesian hierarchical modeling framework. This approach allowed us to account for imperfect detection of visits to the central location, thus incorporating uncertainty into our estimates of temporal persistence. Combined, the two elements of novelty that we described unlocked the possibility of using recursive movement patterns to infer reproduction from movement data in central place foragers.

Among birds, important exceptions to the applicability of our method are precocial species and nest parasites, where parental care is limited (Starck et al. 1998) or absent (Lyon and Eadie 1991). Another limitation of our approach is that it does not provide estimation of reproductive success in terms of number of offspring, but only in terms of overall success or failure. Under this aspect, our method does not compare to the level of detail obtainable with conventional field methods (e.g., Coulson and Porter 1985, Mayer and Ryan 1991, Bruant et al. 2019).

We demonstrated an application of our method to three avian species, but the method can in principle be adapted to other central place foragers too. For example, some carnivores such as wolves (*Canis lupus*, Frame et al. 2004), Canada lynx (*Lynx canadensis*; Olson et al. 2011), and Pampas foxes

(*Lycalopex gymnocercus*, Castillo et al. 2011) also act as central place foragers when rearing their
young, performing trips to and from the den until the offspring are weaned. Some denning mammals,
including Canada lynx, use multiple dens sequentially within a single breeding season (e.g., Olson et al.
2011), so the method may need to be adjusted to sequentially identify these, and the persistence of
movement recursions would then be analyzed in relation to multiple locations. The persistence of
returns to the den (or dens) could be used like in the present study to estimate whether the young
survived until weaning. More broadly, the principle of distinguishing types of revisited locations that
serve different ecological functions can be applied across species and contexts.

Synthesis and significance – Our method can appeal to researchers with different objectives.
First, it can be used by researchers who aim to detect reproductive events from movement data in
central place foragers, such as any non-parasitic altricial bird species. Researchers may want to identify
reproductive events for analyses of reproductive-site fidelity or selection or to discover the geographic
location of unknown breeding sites. Researchers interested in this application of our method can stop at
the first part of the workflow. Second, the method can serve researchers who aim to estimate whether
individual reproductive attempts were successful or not. This application of our method can be
especially useful when collecting field data on reproductive fate is impractical or risky, for example
because of inaccessible locations or high risk of disturbing reproductive activities (Götmark 1992,
Mayer-Gross et al. 1997, Etterson et al. 2011, Wilmers et al. 2015). Researchers interested in this
application of our method will use both parts of our method sequentially. Finally, having pinpointed the
location of reproductive sites and estimated the fate of reproductive events, researchers can pair this
information with environmental conditions experienced by breeding individuals at locations visited
away from the reproductive site (Cagnacci et al. 2010, Pettorelli et al. 2014), which are usually
unknown when data on reproduction is collected in the field and individuals are not tracked.
Knowledge of sites visited by breeding animals when not at the reproductive site can be used, for
example, to test for foraging conditions that may enhance or hamper reproductive success. Our method
can be applied both in situations of opportunistic use of historical tracking data or in cases where the
study is explicitly designed with these objectives in mind. Our work pioneers the use of recursive
movement analysis to obtain estimates of reproductive events from tracking data in central place
foragers. Reproductive fate is an important component of fitness, and estimating it from tracking data
will help establish the long-sought bridge between movement and fitness at the individual level
(Nathan et al. 2008, Morales et al. 2010).

Conclusions

We presented a method to locate breeding sites and estimate the fate of breeding events from the movement trajectories of central place foragers. Our approach complements the existing literature on recursive movement patterns by allowing the distinction of functionally different types of revisited locations, and by introducing the temporal persistence of recursions as a metric to detect the cessation of latent behaviors. In our case, the focal type of revisited location are breeding sites and the latent behavior of interest is the administration of parental care, which informs us of the fate of breeding events. By estimating reproductive fate from movement trajectories, our method helps bridge the gap between movement and individual fitness.

List of abbreviations

GPS: Global Positioning System

CART: Classification And Regression Trees

MCMC: Markov Chain Monte Carlo

JAGS: Just Another Gibbs Sampler

GLM: Generalized Linear Model

Declarations

Ethics approval and consent to participate – Not applicable.

Consent for publication – Not applicable.

Availability of data and material – The GPS-tracking datasets generated and analyzed during the current study will be made publicly available on MoveBank. The datasets on nest locations and fates are available from the corresponding author on reasonable request.

Competing interests – The authors declare that they have no competing interests.

Funding – This work was supported by the USDA National Institute of Food and Agriculture (Hatch project 1009101). The wood stork tracking project was funded by the U. S. Geological Survey's Southeast Region, the U. S. Fish and Wildlife Service, the Army Corps of Engineers, the National Park Service, the Environmental Protection Agency (STAR Fellowship to R.B.), and the Everglades Foundation (ForEverglades Scholarship to S.P.). The Mediterranean gull tracking project was funded by the Ministry for Environment, Land and Sea Protection of Italy. Lesser kestrel tracking project

activities were carried out within the framework of the LIFE+Natura project "Un falco per amico"
(LIFE11/NAT/IT000068).

Authors' contributions – S. Picardi conceived the idea. S. Picardi., B. J. S., and M. E. B.
implemented the analyses. S. Picardi wrote the paper. M. B. and P. C. F. supervised research. J. G. C.,
D. R., L. S., S. Pirrello, and R. R. B. collected the data. All authors contributed to revisions.

Acknowledgments – We thank D. Bear for providing data on known nest locations for wood
storks in northern Florida. We thank anonymous reviewers, as well as J. Hightower and C. Poli, for
their constructive criticism that greatly helped to improve previous drafts of the manuscript. The
Mediterranean gull tracking project was carried out with the support of Ente Parco Delta del Po Emilia-
Romagna, Reparto Carabinieri per la Biodiversità di Punta Marina, R. Nardelli, A. Andreotti, G.
Meneghini, F. Spina. Lesser kestrel tracking project activities were carried out with the support of S.
Podofillini, E. Fulco, P. Giglio, S. C. Pellegrino, M. Lorusso (Comune di Altamura), F. Parisi (Comune
di Gravina in Puglia), and D. Ciampanella (project manager). Any use of trade, firm, or product names
is for descriptive purposes only and does not imply endorsement by the U. S. Government.

References

Alonso, J. C. et al. 1994. Flock Size and Foraging Decisions in Central Place Foraging White Storks,
     Ciconia Ciconia. - Behaviour 129: 279–292.

Andersson, M. 1981. Central Place Foraging in the Whinchat, Saxicola Rubetra. - Ecology 62: 538–
     544.

Bar-David, S. et al. 2009. Methods for assessing movement path recursion with application to African
     buffalo in South Africa. - Ecology 90: 2467–2479.

Barraquand, F. and Benhamou, S. 2008. Animal Movements in Heterogeneous Landscapes: Identifying
     Profitable Places and Homogeneous Movement Bouts. - Ecology 89: 3336–3348.

Becker, C. et al. 2010. Nest success and parental investment in the Critically Endangered Maui
     parrotbill Pseudonestor xanthophrys with implications for recovery. - Endangered Species
     Research 11: 189–194.

Benhamou, S. and Riotte-Lambert, L. 2012. Beyond the Utilization Distribution: Identifying home range areas that are intensively exploited or repeatedly visited. - Ecological Modelling 227: 112–116.

Berger-Tal, O. and Bar-David, S. 2015. Recursive movement patterns: review and synthesis across species. - Ecosphere 6: 1–12.

Birks, S. M. 1997. Paternity in the Australian brush-turkey, Alectura lathami, a megapode bird with uniparental male care. - Behav Ecol 8: 560–568.

Bonar, M. et al. 2018. Implementing a novel movement-based approach to inferring parturition and neonate caribou calf survival (DB Hardy, Ed.). - PLOS ONE 13: e0192204.

Borkhataria, R. R. et al. 2008. A Preliminary Model of Wood Stork Population Dynamics in the Southeastern United States. - Waterbirds 31: 42–49.

Bracis, C. et al. 2018. Revisitation analysis uncovers spatio-temporal patterns in animal movement data. - Ecography 41: 1801–1811.

Breiman, L. 2001. Random Forests. - Machine Learning 45: 5–32.

Bruant, A. et al. 2019. Effect of Parental Care on Reproductive Success in Wood Storks (Mycteria Americana). - bioRxiv: 592840.

Burke, C. M. and Montevecchi, W. A. 2009. The foraging decisions of a central place foraging seabird in response to fluctuations in local prey conditions. - Journal of Zoology 278: 354–361.

Cadiou, B. and Monnat, J.-Y. 1996. Parental attendance and squatting in the Kittiwake Rissa tridactyla during the rearing period. - Bird Study 43: 164–171.

Cagnacci, F. et al. 2010. Animal ecology meets GPS-based radiotelemetry: a perfect storm of opportunities and challenges. - Philosophical Transactions of the Royal Society B: Biological Sciences 365: 2157–2162.

Calder, W. A. 1973. The Timing of Maternal Behavior of the Broad-Tailed Hummingbird Preceding Nest Failure. - The Wilson Bulletin 85: 283–290.

Castillo, D. F. et al. 2011. Diet of Adults and Cubs of Lycalopex gymnocercus in Pampas Grassland: A Validation of the Optimal Foraging Theory? - anzf 48: 251–256.

Cecere, J. G. et al. 2018. Spatial segregation of home ranges between neighbouring colonies in a diurnal raptor. - Scientific Reports 8: 11762.

Clark, E. S. 1980. The Attentiveness and Time Budget of a Pair of Nesting Wood Storks. - Proceedings of the Colonial Waterbird Group 3: 204–215.

Coulson, J. C. and Porter, J. M. 1985. Reproductive success of the Kittiwake Rissa tridactyla: the roles of clutch size, chick growth rates and parental quality. - Ibis 127: 450–466.

Coulter, M. C. et al. 1999. Wood Stork(Mycteria americana). - The Birds of North America: 24.

De'ath, G. and Fabricius, K. E. 2000. Classification and Regression Trees: A Powerful yet Simple Technique for Ecological Data Analysis. - Ecology 81: 3178–3192.

DeMars, C. A. et al. 2013. Inferring parturition and neonate survival from movement patterns of female ungulates: a case study using woodland caribou. - Ecology and Evolution 3: 4149–4160.

English, M. et al. 2014. Foraging site recursion by forest elephants Elephas maximus borneensis. - Curr Zool 60: 551–559.

Etterson, M. A. et al. 2011. Modeling fecundity in birds: Conceptual overview, current models, and considerations for future developments. - Ecological Modelling 222: 2178–2190.

Fasola, M. and Canova, L. 1992. Nest Habitat Selection by Eight Syntopic Species of Mediterranean Gulls and Terns. - Colonial Waterbirds 15: 169–178.

Frair, J. L. et al. 2010. Resolving issues of imprecise and habitat-biased locations in ecological analyses using GPS telemetry data. - Philosophical Transactions of the Royal Society B: Biological Sciences 365: 2187–2200.

Frame, P. F. et al. 2004. Long Foraging Movement of a Denning Tundra Wolf. - Arctic 57: 196–203.

Garrison, J. S. E. and Gass, C. L. 1999. Response of a traplining hummingbird to changes in nectar availability. - Behav Ecol 10: 714–725.

Götmark, F. 1992. The Effects of Investigator Disturbance on Nesting Birds. - In: Power, D. M. (ed), Current Ornithology. Current Ornithology. Springer US, pp. 63–104.

Grüebler, M. U. and Naef-Daenzer, B. 2008. Fitness Consequences of Pre- and Post-Fledging Timing Decisions in a Double-Brooded Passerine. - Ecology 89: 2736–2745.

Gurarie, E. et al. 2016. What is the animal doing? Tools for exploring behavioural structure in animal movements. - J Anim Ecol 85: 69–84.

Gustin, M. et al. 2017. Space use and flight attributes of breeding Lesser Kestrels Falco naumanni revealed by GPS tracking. - Bird Study 64: 274–277.

Hernández-Pliego, J. et al. 2017. A few long versus many short foraging trips: different foraging strategies of lesser kestrel sexes during breeding. - Movement Ecology 5: 8.

Kacelnik, A. 1984. Central Place Foraging in Starlings (Sturnus vulgaris). I. Patch Residence Time. - Journal of Animal Ecology 53: 283–299.

Lebreton, J.-D. et al. 1992. Modeling Survival and Testing Biological Hypotheses Using Marked Animals: A Unified Approach with Case Studies. - Ecological Monographs 62: 67–118.

Lyon, B. E. and Eadie, J. M. 1991. Mode of development and interspecific avian brood parasitism. - Behav Ecol 2: 309–318.

Mayer, P. M. and Ryan, M. R. 1991. Electric Fences Reduce Mammalian Predation on Piping Plover Nests and Chicks. - Wildlife Society Bulletin (1973-2006) 19: 59–63.

Mayer-Gross, H. et al. 1997. The effect of observers visiting the nests of passerines: an experimental study. - Bird Study 44: 53–65.

Morales, J. M. et al. 2004. Extracting More Out of Relocation Data: Building Movement Models as Mixtures of Random Walks. - Ecology 85: 2436–2445.

Morales, J. M. et al. 2010. Building the bridge between animal movement and population dynamics. - Philosophical Transactions of the Royal Society B: Biological Sciences 365: 2289–2301.

Mueller, T. and Fagan, W. F. 2008. Search and navigation in dynamic environments – from individual behaviors to population distributions. - Oikos 117: 654–664.

Nathan, R. et al. 2008. A movement ecology paradigm for unifying organismal movement research. - PNAS 105: 19052–19059.

Nelson, S. K. and Hamer, T. E. 1995. Chapter 8: Nest Success and the Effects of Predation on Marbled Murrelets. - In: Ralph, C. John; Hunt, George L., Jr.; Raphael, Martin G.; Piatt, John F., Technical Editors. 1995. Ecology and conservation of the Marbled Murrelet. Gen. Tech. Rep. PSW-GTR-152. Albany, CA: Pacific Southwest Research Station, Forest Service, U.S. Department of Agriculture; p. 89-98 in press.

Nelson, T. et al. 2015. A time geographic approach for delineating areas of sustained wildlife use. - Annals of GIS 21: 81–90.

Nicholson, K. L. et al. 2019. Using fine-scale movement patterns to infer ungulate parturition. - Ecological Indicators 101: 22–30.

Olson, L. E. et al. 2011. Den use and Activity Patterns in Female Canada Lynx (Lynx canadensis) in the Northern Rocky Mountains. - nwsc 85: 455–462.

Orians, G. H. and Pearson, N. E. 1979. On the theory of central place foraging. - In: Analysis of Ecological Systems. Ohio University Press, pp. 154–177.

Pettorelli, N. et al. 2014. Satellite remote sensing for applied ecologists: opportunities and challenges. - Journal of Applied Ecology 51: 839–848.

Riotte-Lambert, L. et al. 2013. Periodicity analysis of movement recursions. - Journal of Theoretical Biology 317: 238–243.

Riotte-Lambert, L. et al. 2017. From randomness to traplining: a framework for the study of routine movement behavior. - Behav Ecol 28: 280–287.

Rodgers, J. A. et al. 1987. Population Dynamics of Wood Storks in North and Central Florida, USA. - Colonial Waterbirds 10: 151–156.

Sabine, J. B. et al. 2006. Nest Fate and Productivity of American Oystercatchers, Cumberland Island National Seashore, Georgia. - cowa 29: 308–314.

Schaub, M. and Royle, J. A. 2014. Estimating true instead of apparent survival using spatial Cormack–Jolly–Seber models. - Methods in Ecology and Evolution 5: 1316–1326.

Schick, R. S. et al. 2008. Understanding movement data and movement processes: current and emerging directions. - Ecology Letters 11: 1338–1350.

Singh, N. J. and Ericsson, G. 2014. Changing motivations during migration: linking movement speed to reproductive status in a migratory large mammal. - Biology Letters 10: 20140379.

Snow, D. W. et al. 1997. The birds of the Western Palearctic. concise edition. in press.

Stake, M. M. and Cimprich, D. A. 2003. Using Video to Monitor Predation at Black-Capped Vireo Nests. - Condor 105: 348–357.

Starck, J. M. et al. 1998. Avian Growth and Development: Evolution Within the Altricial-precocial Spectrum. - Oxford University Press.

Stenning, M. J. 1996. Hatching asynchrony, brood reduction and other rapidly reproducing hypotheses. - Trends in Ecology & Evolution 11: 243–246.

Tarwater, C. E. and Brawn, J. D. 2010. The post-fledging period in a tropical bird: patterns of parental care and survival. - Journal of Avian Biology 41: 479–487.

Van Moorter, B. et al. 2016. Movement is the glue connecting home ranges and habitat selection. - Journal of Animal Ecology 85: 21–31.

Weimerskirch, H. et al. 2001. Nest attendance and foraging movements of northern fulmars rearing chicks at Bjørnøya Barents Sea. - Polar Biol 24: 83–88.

Wiley, R. H. and Wiley, M. S. 1980. Spacing and Timing in the Nesting Ecology of a Tropical Blackbird: Comparison of Populations in Different Environments. - Ecological Monographs 50: 153–178.

Wilmers, C. C. et al. 2015. The golden age of bio-logging: how animal-borne sensors are advancing the frontiers of ecology. - Ecology 96: 1741–1753.

Zangmeister, J. L. et al. 2009. Incubation failure and nest abandonment by Leach's Storm-Petrels detected using PIT tags and temperature loggers. - Journal of Field Ornithology 80: 373–379.

585   **Figures and Tables**

587   Table 1 – **Datasets.**

			Wood stork	Lesser kestrel	Mediterranean gull
Spatial resolution (m)			18	<10	<10
Temporal resolution (min)			60	15 (summer)/30 (winter)	15
Fix failure rate			High	Low	Low
Tagged at			Fledging /non-breeding	Early nestling-rearing	Incubation
Number of tracks (individual-years)		Total	148	56	29
		Known nest location	107	50	24
		Known fate	0	21	12
		Non-breeders	41 (subadults)	16 (winter)	16 (winter)
Breeding season			Nov-	Apr-Jul	Apr-Jul

	Aug (varies with latitude)		
Breeding cycle (days)	110	60	60

588   Differences among GPS-tracking datasets for wood storks, lesser kestrels, and Mediterranean
589   gulls in terms of sample sizes, data characteristics, and species seasonality and ecology.

591   Table 2 – **Nest-detection performance metrics.**

	Wood stork	Lesser kestrel	Mediterranean gull
Positive predictive value	87.04%	100.00%	91.67%
Sensitivity	87.85%	92.00%	91.67%
False negative rate	12.15%	8.00%	8.33%
False positive rate	2.45%	50.00%	0.00%

592   Performance metrics of the nest identification algorithm (see Methods for definitions).

594   Figure 1 – **Method workflow.** Workflow of the analysis to identify nest sites (Part I) and

595 estimate reproductive fate (Part II) from telemetry data. The R package 'nestR' includes functions to

596 tackle each of the steps depicted in the boxes.

Figure 2 – **Identification of revisited locations and breeding sites among them.** Schematic diagram depicting how revisited locations (and breeding sites among them) are identified along individual tracks. A) A buffer of user-defined radius r is placed around each location of the trajectory. Buffers that contain any other locations in addition to the one they are centered on are revisited locations. A fraction of these will be breeding sites. B) To detect breeding sites specifically, revisited locations are filtered based on their temporal patterns of revisitation using parameter values that are diagnostic of breeding sites. Parameter values can be chosen based on prior knowledge or based on the comparison of known breeding and non-breeding sites with CART.

Figure 3 – **Output of Classification and Regression Trees on parameters describing nest**

revisitation patterns. Output of CART to discriminate nest and non-nest sites in A) wood stork, B)

lesser kestrel, C) Mediterranean gull. Within each node (box), the number of known non-nest and nest

sites are reported on the left and right, respectively. The root node (top) is recursively split into two

until the terminal nodes (bottom). The criterion used to split each node is shown on the corresponding

stems (bold font). The label on each node represents the class that was assigned to the content of that

node (nest site for "yes" boxes, non-nest site for "no" boxes). Thus, the number on the right in "yes"

terminal nodes and the number on the left in "no" terminal nodes correspond to correct classifications,

while the number on the left in "yes" terminal nodes and on the right in "no" terminal nodes correspond

to incorrect classifications.

Figure 4 – **Probability of survival and visit detection.** Probability of visit detection (top row) and survival (bottom row) through time (days of the attempt) estimated at the population level for A) wood stork, B) lesser kestrel, C) Mediterranean gull. Ribbons indicating 95% credible intervals are shaded.

623   Figure 5 – **Nest-survival estimates.** Estimates of probabilities of success, i.e., $P(z_T = 1)$ for

624 breeding attempts of A) wood stork, B) lesser kestrel, C) Mediterranean gull. For kestrels and gulls,

625 estimates are plotted in relation to their true fate. True fate was unknown for storks. Raw data points are

626 shown as dots (purple for failures, green for successes, gray when true fate is unknown) overlaid to

627 boxplots (black).