A data-driven method to locate nest sites and estimate reproductive outcome from avian telemetry data

Simona Picardi[1*], Brian J. Smith[2], Matthew E. Boone[1], Peter C. Frederick[3], Jacopo G. Cecere[4], Diego Rubolini[5], Lorenzo Serra[4], Simone Pirrello[4], Rena R. Borkhataria[6], Mathieu Basille[1]

[1]Department of Wildlife Ecology and Conservation, Fort Lauderdale Research and Education Center, University of Florida, 3205 College Ave, Davie, FL 33314, USA

[2]Cherokee Nation Technologies, contracted to the U.S. Geological Survey, Wetland and Aquatic Research Center, 3321 College Ave, Davie, FL 33314, USA

[3]Department of Wildlife Ecology and Conservation, University of Florida, 368 Newins-Ziegler Hall, Gainesville, FL 32611, USA

[4]Area Avifauna Migratrice, Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA), via Cà Fornacetta 9, I-40064 Ozzano Emilia (BO), Italy

[5]Dipartimento di Scienze e Politiche Ambientali, Università degli Studi di Milano, via Celoria 26, I'20133 Milan, Italy

[6]Department of Wildlife Ecology and Conservation, Everglades Research and Education Center, University of Florida, 3200 E Palm Beach Rd, Belle Glade, FL 33430, USA

[*]Corresponding author – simona.picardi@ufl.edu

Running headline: Estimation of avian reproductive outcome from telemetry data

Abstract

The use of tracking devices for collecting animal movement data has become widespread in recent decades. In parallel, this has sparked a proliferation of methods to infer individual behavior from tracking data. Being able to learn more than only the movement trajectories of animals from telemetry data is one of the major forces pushing the field of movement ecology forward. One application that has been poorly explored is the use of movement data to estimate reproductive success. We pioneered this application in birds by introducing a method to locate nesting attempts and estimate their outcome from tracking data. We implemented our method in the R package nestR and illustrate its application to three bird species: the wood stork (*Mycteria americana*), the lesser kestrel (*Falco naumanni*), and the Mediterranean gull (*Ichthyaetus melanocephalus*). We identified nest site locations based on the analysis of recursive movement patterns of breeding individuals acting as central place foragers. Using trajectories with known breeding attempts, we estimated a set of species-specific criteria for the identification of nest sites, using non-reproductive individuals as controls. We estimated nest survival as a measure of reproductive outcome from nest site re-visitation histories, using a Bayesian hierarchical modeling approach that accounted for temporally variable re-visitation patterns, probability of visit detection, and missing data. Provided the availability of adequate tracking data, our method can be broadly applied to estimate reproductive outcome in a variety of central place forager species. Inferring reproductive outcome from tracking data will allow ecologists to bridge the gap between movement and space use behavior, environmental factors, and their fitness consequences.

Key-words

movement ecology – GPS-tracking – fitness – Bayesian hierarchical models – R package nestR – wood stork – lesser kestrel – Mediterranean gull – nest survival – birds

Introduction

Technological advancements have resulted in an exponential increase in available animal tracking data in recent decades (Urbano et al., 2010; Kays, Crofoot, Jetz, & Wikelski, 2015; Wilmers et al., 2015).

The ability to obtain movement data at very fine resolutions brings us closer to understanding the

behaviors underlying movement (Cagnacci, Boitani, Powell, & Boyce, 2010; Tomkiewicz, Fuller, Kie,

& Bates, 2010). Inferring more than just the movement trajectories of animals from telemetry data is

one of the major forces pushing the field of movement ecology forward (Nathan et al., 2008; Schick et

al., 2008; Wilmers et al., 2015).

Knowing what an animal is doing when it is moving a certain way can improve our understanding of

the links between movement and resource dynamics, species interactions, distribution, and individual

fitness, which is a fundamental driver of population dynamics and evolutionary processes (Mueller &

Fagan, 2008; Schick et al., 2008; Morales et al., 2010). The importance of this pursuit justifies the

recent and ongoing proliferation of analytical techniques to infer behavior from movement (Gurarie et

al., 2016; Thiebault, Dubroca, Mullers, Tremblay, & Pistorius, 2018). Many of these approaches seek to

identify behavioral modes by splitting movement trajectories into behaviorally homogeneous bouts

based on quantitative properties of the track (Edelhoff, Signer, & Balkenhol, 2016). For instance,

topology-based segmentation techniques classify locations along a track into distinct groups based on

the distribution of path metrics, such as turning angles and step length (Van Moorter, Visscher, Jerde,

Frair, & Merrill, 2010) or speed (Garriga, Palmer, Oltra, & Bartumeus, 2016). Time-series analyses,

such as behavioral change-point analysis (Gurarie, Andrews, & Laidre, 2009) or wavelet analysis

(Soleymani, Pennekamp, Dodge, & Weibel, 2017) decompose trajectories by detecting significant

changes of a path-signal in time. State-space models use a mechanistic approach to identify latent states

in the data corresponding to different behavioral modes (Beyer, Morales, Murray, & Fortin, 2013).

These analytical tools differ from one another in their functioning, but all of them provide ways to

relate movement patterns to the underlying behavioral processes (Edelhoff et al., 2016; Gurarie et al.,

2016). For example, Polansky et al. (2015) used behavioral change point analysis to split African

elephant (*Loxodonta africana*) trajectories into phases of goal-oriented, directed movement in the

proximity of waterholes and phases of slower, more tortuous movement when resting or foraging.

Similarly, Austin et al. (2019) identified three different behavioral modes, corresponding to foraging,

directed flight, and resting/searching, in magnificent frigatebird (*Fregata magnificens*) movement

tracks using hidden Markov models.

Specific movement patterns can also diagnose events of interest: for example, Webb et al. (2008) used

spatio-temporal clustering of wolf (*Canis lupus*) GPS-tracking data to identify kill sites. Other studies

used changes in movement patterns to detect parturition events along ungulate GPS tracks, particularly

moose (*Alces alces*; Severud et al., 2015; Nicholson et al., 2019) and caribou (*Rangifer tarandus*;

DeMars, Auger   Méthé, Schlägel, & Boutin, 2013; Bonar, Ellington, Lewis, & Vander Wal, 2018).

These event-oriented applications aim to isolate focal episodes along a movement track, rather than

dividing the entire trajectory into behavioral phases. Whether the objective is segmentation or the

detection of events of interest, the signal used depends on what type of behavioral processes are being

targeted (Nams, 2014). For wolf kill sites, the signal of interest would be a phase of restricted

movement around a carcass (Webb et al., 2008). For calving moose and caribou, it would be a slow-

down in movements due to the reduced mobility of newborns (DeMars et al., 2013; Nicholson et al.,

2019).

In some contexts, recursive movement patterns can be an indicative signal of underlying processes

(Berger-Tal & Bar-David, 2015; Bracis, Bildstein, & Mueller, 2018). Many animals keep returning to

places of ecological significance, such as dens, nests, and foraging patches (Bracis et al., 2018). The

periodicity of such recursive movements can provide insight into ecological processes (Riotte-Lambert,

Benhamou, & Chamaillé-Jammes, 2013). For example, recursive movement patterns have been studied

in herbivores with the objective of understanding movement responses to spatio-temporal variability of

resources, such as cycles of forage depletion and greening (Bar-David et al., 2009). Bracis et al. (2018)

analyzed recursive movement patterns of a turkey vulture (*Cathartes aura*) to identify roost sites.

Recursive movement patterns can provide insight into behavior and life-history as well (Bracis et al.,

2018). Some animals are tied to specific locations in specific phases of their life cycle and exhibit

recursive movements to and from those locations (Bracis et al., 2018). These include altricial breeding

bird species. While breeding, altricial bird species act as central place foragers (*sensu* Orians &

Pearson, 1979), recursively departing from their nest site to embark on foraging trips and returning to

incubate their eggs or provide food for their nestlings (Andersson, 1981; Alonso, Carrascal, Muñoz-

Pulido, & Alonso, 1994; Burke & Montevecchi, 2009). Back-and-forth trips can therefore be observed

in movement data and used to identify when and where a bird is nesting. Once the location of a nest has

been identified, patterns of nest site re-visitation may reveal information about the fate of a breeding

attempt and may be used to infer reproductive outcome. Being able to estimate reproductive outcome

from tracking data connects movement to reproduction, an important component of individual fitness

(Morales et al., 2010; Singh & Ericsson, 2014). This could be especially valuable whenever gathering

data on reproductive outcome through direct observation is logistically difficult or impractical (Etterson

et al., 2011; Wilmers et al., 2015; Nicholson et al., 2019). Moreover, inferring reproductive outcome

from movement establishes a direct link between environmental dynamics and components of

individual fitness if tracking data is coupled with remote sensing data (Cagnacci et al., 2010; Pettorelli

et al., 2014). While tools to infer behavioral structure along animal trajectories have been extensively

used to evaluate responses to environmental factors or internal state, few studies so far have attempted

the use of movement data to assess reproductive success (DeMars et al., 2013). The applications of path

segmentation to detect ungulate parturition have been the first attempts at the task (DeMars et al.,

2013). Besides these examples, efforts to apply movement pattern detection to the estimation of

reproductive metrics have been limited. To our knowledge, no study has attempted to infer avian

reproductive fitness from movement data.

Here, we introduce a data-driven method to locate breeding attempts along movement trajectories of

altricial bird species and estimate their outcome based on patterns of nest re-visitation. Our approach

combines event-oriented pattern detection with the analysis of recursive movement patterns. Our

workflow is implemented in the R package nestR (https://github.com/picardis/nestR). We demonstrate

the broad applicability of our approach by illustrating its use on GPS-tracking data for three altricial

species that broadly differ in their breeding habitat and ecology: the wood stork (*Mycteria americana*),

the lesser kestrel (*Falco naumanni*), and the Mediterranean gull (*Ichthyaetus melanocephalus*).

Example datasets for all three species are available in nestR.

Materials and Methods

<u>Method description</u>

Nest site detection – Figure 1 provides a roadmap of our workflow. Nest sites are identified as

repeatedly visited locations along individual trajectories (Figure 1). Returns to a location are defined as

returns to a circular area of a user-defined radius, obtained by placing a buffer around each point of the

trajectory. Due to both behavior and GPS error, GPS points at a revisited location may be spatially

scattered around the true center (Frair et al., 2010). Defining locations as buffers around points helps

account for this scattering. The size of the buffer sets the spatial scale at which re-visitation patterns

will be calculated. Re-visitation patterns are described by the following set of parameters: the

maximum number of consecutive days a location is visited; the percentage of days visited between the

first and last visit; and the percent fixes at the nest on the day with maximum attendance. Parameter

values are then used as diagnostic features to filter nest sites among re-visited locations, based on the

rationale that re-visitation patterns of nest sites differ from those of non-nest sites. Nest sites are often

visited for longer stretches of consecutive days, on more days, and more frequently or for longer within

a day than other types of re-visited locations. Especially during incubation, many bird species attend

their nests intensively (e.g., Norton, 1972; van Vessem & Draulans, 1986; Ojowski, Eidtmann, Furness,

& Garthe, 2001). The parameters we use to describe re-visitation patterns are meant to capture these

diagnostic behaviors and are used to identify nest sites among re-visited locations. In nestR, the

identification of nests is implemented in the function find_nests(). Besides the tracking data and the set

of filtering parameters, this function also takes as input a series of species- and data-specific parameters

used to tailor the algorithm to different case studies. These include arguments that restrict the analysis

within the breeding season for a given species, and arguments that help account for data sampling rate

and fix failure rate. For a more thorough discussion of function arguments, we direct the reader to the

nestR package vignette.

Unless prior knowledge is available about re-visitation patterns to nest sites, researchers will need ways

to inform their choice of parameter values used for filtering nest sites among re-visited locations

(Figure 1). If the true location of a nest is known for a subset of the data, researchers can compare re-

visitation patterns at known nest sites to those of non-nest sites and find one or more sets of parameter

values to discriminate between them. If no on-ground data is directly available, researchers can visually

explore the data and identify trusted nest sites, where possible. For example, likely nest sites can be

recognized based on habitat features or proximity to known breeding colonies. An interactive

visualization tool is available in nestR in the form of a Shiny (https://shiny.rstudio.com) app that lets

users dynamically explore re-visited locations on satellite imagery while manipulating parameters in

real time. Once known or trusted nest sites are identified, non-nest sites can be selected based on a

criterion of temporal overlap; revisited locations that were visited simultaneously with the breeding

attempt can be assumed to not be nest sites, assuming birds cannot breed in two places at the same

time. Running the function find_nests() with non-constraining values for the filtering parameters will

return any re-visited locations, among which the user can pick known nest and non-nest sites to

compare (Figure 1).

Several approaches are possible to identify sets of parameter values to distinguish nest from non-nest

sites. In nestR, we implemented one possible approach based on classification and regression trees

(CART; De'ath & Fabricius, 2000). The function discriminate_nests() applies a CART algorithm to a

dataset of known nest and non-nest sites given as input. The tree gets pruned to the optimal number of

nodes based on a minimum relative error criterion (De'ath & Fabricius, 2000). If input data is split into

training and testing fractions, the CART also provides estimates of classification error rates through

cross-validation.

The CART identifies one or more sets of parameter values that best discriminate between nest and non-

nest sites (Figure 1). Researchers can then plug these parameter values into the filtering arguments of

find_nests() to identify nest sites only among re-visited locations (Figure 1). Researchers can decide

how to deal with candidate breeding attempts that temporally overlap with each other. Assuming birds

cannot breed in two places at the same time, the recommended option is to pick the top candidate

among any set of temporally overlapping ones and discard the rest. For each nest site, find_nests()

outputs a summary of re-visitation patterns and a history of nest re-visitation (in the form of a presence/

absence time series, with GPS fixes within the nest buffer assigned to the status "present" and GPS

fixes outside of the nest buffer to "absent"). This time series is then used to estimate the outcome of

breeding attempts.

Reproductive outcome estimation – The outcome of each identified breeding attempt is estimated using

a Bayesian hierarchical modeling approach (Figure 1). A breeding attempt is considered successful if

the nest site was visited until the end of a complete breeding cycle for the focal species. The model

estimates nest survival based on patterns of nest site re-visitation, taking into account imperfect

detection of nest visits and missing fixes. The MCMC algorithm is implemented in JAGS (Plummer,

2003) via the R package rjags.

The model specification includes two processes: the survival process, which is not directly observable,

and the observation process, which is the signal observed in the re-visitation histories. Much like a

Bayesian implementation of a Cormack-Jolly-Seber capture-mark-recapture model (Lebreton,

Burnham, Clobert, & Anderson, 1992), the latent nest survival variable is modeled at the daily scale as

a function of survival status at the previous time-step and daily survival probability:

$$z_t \sim Bern(z_{t-1} \times \phi_{t-1})$$

Observed visits on a given day are modeled as a function of current nest survival status, probability of

visit detection on that day, and number of GPS fixes available on that day:

$$Y_t \sim Bin(z_t \times p_t, N_t)$$

Where the probability of detection is:

$$p_t = Pr(visit\ detected | z_t = 1, N_t)$$

Reproductive outcome is defined as the probability that the nest was still surviving on the last day of

the theoretical duration of a complete breeding attempt:

$$Pr(z_T = 1)$$

Where T is the duration of a complete breeding attempt. Both survival and detection probability are

modeled using a binomial Generalized Linear Model as a function of the day of the attempt:

$$logit(\phi_t) = \beta_{\phi_0} + \beta_{\phi_1} \times t$$

$$logit(p_t) = \beta_{p_0} + \beta_{p_1} \times t$$

The model is fully specified by using uninformative priors on the Beta parameters, in this case a normal

distribution with a mean of 0 and precision of $1e^{-5}$. In the current implementation, daily survival and

detection are assumed to be the same for all nests in the population. The model outputs daily estimates

of survival and detection probability at the population level, as well as daily survival estimates for each

breeding attempt along with credible intervals.

<u>Applications to data</u>

We applied our workflow to GPS-tracking data for 148 individual-years for wood storks (henceforth

storks), 53 for lesser kestrels (henceforth kestrels) and 29 for Mediterranean gulls (henceforth gulls).

All tags were solar-powered and recorded fixes primarily during daytime. Details about devices,

settings, harnesses and study areas regarding storks and kestrels can be found in Borkhataria et al.

(2008) and Cecere et al. (2018), respectively. We split data into individual-years such that breeding

activities were contained within them. To find nest sites, we restricted the analysis to data within the

breeding season for each species (Table 1). While both kestrels and gulls have a well-defined breeding

season between April and August in our study areas (Snow, Perrins, Hillcoat, Gillmor, & Roselaar,

1997), storks in the southeastern U.S. can breed at slightly different times of the year depending on

latitude (Coulter, Rodgers, Ogden, & Depkin, 1999; Table 1). In this case, we only excluded the

window of time where no breeding activities were expected to occur anywhere in the range from the

analysis.

Given the spatial resolution of the GPS data (Table 1) and the expected scale of movements around the

nest site for all three species, we used a buffer of 40 m around each GPS position. We initially screened

trajectories for any re-visited locations using non-constraining values in filtering parameters (which is

equivalent to not applying any filtering). We then used on-ground data on known nest locations to

select true nests and non-nest sites among re-visited locations. Kestrels and gulls were captured at the

nest site (Table 1), so the location of the nest was known. For storks, on-ground data on nest locations

was available for 10 individual-years (Bear D., unpublished data). We explored the remaining stork

trajectories and identified those for which the top visited location was at a known breeding colony (data

from USFWS 2018). We identified 97 of these, marked them as trusted and treated them as known nest

sites for the rest of the analysis.

We used CART to compare re-visitation patterns between nest and non-nest sites. We split each of the

three datasets in two parts for training and testing (3:2 ratio). We used the resulting sets of parameter

values to filter nest sites among re-visited locations in the trajectories of breeding individuals (Table 2).

Even when the CART did not suggest that the number of consecutive days visited was an important

predictor of true nest sites, we chose a reasonable value to use as a threshold for this parameter (Table

2). We did not expect to have enough power to discern nest from non-nest sites for attempts that failed

in the first handful of days. We only retained a top candidate among any sets of breeding attempts that

were temporally overlapping. We used non-breeder trajectories (subadults in the case of storks, non-

breeding season data in the case of kestrels and gulls) to validate our results against false positives. We

calculated the positive predictive value of our algorithm as the percentage of known nest sites among

the total number of nest sites we found for each species. We calculated the sensitivity of our algorithm

as the percentage of the known nest sites that were identified. We calculated the false negative rate as

the percentage of known nest sites that we failed to identify. Finally, we calculated the false positive

rate as the percentage of non-breeding individual-years for which we erroneously identified a nest site.

We fit the nest survival model described above to estimate the outcome of the identified breeding

attempts. Since kestrels and gulls were captured after they had already started breeding, the initial part

of every breeding attempt was missing from the data. To account for this, we subtracted the average

number of days from start to hatching (for kestrels) and to late incubation (for gulls) from the full

breeding cycle when specifying the expected duration of a complete attempt (Table 1).

Results

The initial screening with no filtering identified 9954 re-visited locations (i.e., potential nest sites) for

storks, 1285 for kestrels, and 1342 for gulls. Results from the CART showed that the optimal set of

parameter values to discriminate nest from non-nest sites was 18 minimum consecutive days visited

and 90% minimum nest attendance on the top day for storks, and 52% and 45% minimum attendance

on the top day for kestrels and gulls, respectively (Figure 2). In addition to the parameters suggested by

the CART, we used a minimum of 4 and 8 consecutive days visited for kestrels and gulls, respectively

(Table 2). By filtering re-visited locations using the parameters listed in Table 2, we identified 106 nest

sites for storks, 49 for kestrels, and 28 for gulls, which closely matches the number of nest sites we

were expecting to find (Table 1). As a consequence, the positive predictive value of the algorithm

ranged between 78-94%, the sensitivity between 75-86%, and the false negative rate between 14-25%

(Table 3). The false positive rate was below 7% for storks and gulls but reached 68% for kestrels (Table

3). The probability of detecting nest visits decreased throughout the breeding attempt for all three

species (Figure 3). The distribution of survival estimates at the individual level differed between known

failed and successful attempts, although with some degree of overlap, especially for kestrels (Figure 4).

No data on true outcomes were available for storks, therefore we were unable to verify survival

estimates for this species.

Discussion

We present an original, data-driven method to identify nest site locations of altricial avian species and

estimate the outcome of breeding attempts based on GPS-tracking data. This is among the first attempts

to use telemetry data to infer a major component of fitness (DeMars et al., 2013), and the first applied

to birds. We demonstrated the broad applicability of our method by illustrating its use on GPS-tracking

data from three species representative of different ecosystems, including a subtropical wading bird, a

small steppe raptor, and a seabird. The implementation of our workflow in the R package nestR is

straightforward and requires the use of intuitive, biologically interpretable parameters.

Nest site detection – Our nest site detection method performed well on all three species, allowing us to

correctly identify most of the known nest sites from movement trajectories of breeding individuals. We

achieved high positive predictive value and sensitivity. Moreover, we may have underestimated the

positive predictive value by not being able to confirm possible second attempts following early failures.

All the species we tested our method on may attempt to breed a second time if their first clutch fails

early on, and it is possible that the nest sites we were unable to confirm included second attempts in

addition to non-nests.

Differences between our case-study datasets both in terms of species ecology and data characteristics

also presented us with different challenges. The main problem we had to deal with when applying our

method to storks was not having much on-ground data. Most of the nest locations we treated as known

were trusted nest sites for which we did not have on-ground confirmation (97 out of 107). Moreover,

we had no information regarding the outcome of the breeding attempts, and therefore we were not able

to validate results of the outcome estimation for this species. The strengths of this dataset were, first,

the large sample size; and second, the fact that birds were tagged at fledging or while non-breeding and

were tracked for several years, so we had data for the entire breeding cycle for all the attempts. For

kestrels and gulls, this was not the case. Both kestrels and gulls were tagged after the breeding attempt

had already started, so the incubation phase was partly (for gulls) or entirely (for kestrels) missing from

the data. Missing the first part of breeding attempts affected our ability to identify nest sites. Nest

attendance is high during incubation in many altricial bird species whose nestlings are not able to

thermoregulate autonomously (DuRant, Hopkins, Hepp, & Walters, 2013; see Clark, 1980 for an

example on storks). Not having tracking data for kestrels and gulls for the phase where re-visitation

patterns most obviously diagnose nesting behavior hampered our ability to identify nest sites. For

storks the lower false negative rate might have resulted from the relatively strict constraint we enforced

for the minimum number of consecutive days; it is possible that some of the stork breeding attempts we

missed failed before reaching the 18-day mark. Not identifying breeding attempts whose duration does

not exceed the minimum constraint applied is a logical implication rather than a failure. In 3 cases out

of 7, early failed attempts (within the 2 days following tagging) also explain missed detection of gull

nest sites. For kestrels, the behavior of males might also have played a role in hampering nest site

detection: 3 of the 10 nest sites that we failed to identify appear to be possible roosting sites used by

males at night in the proximity of the nesting colony. Because the use of the roosting sites is temporally

overlapping with the breeding attempt, roosting sites were likely to be selected for males instead of nest

sites whenever attendance was higher at the roost compared to the nest. When considered in the light of

these unavoidable data limitations and behavioral constraints, the performance of our nest site detection

method is well above satisfactory.

Another drawback of the kestrel and gull datasets, especially gulls, is that sample size was small. Small

sample size might have played a role in determining the high error rates we obtained as estimates from

the CART and from our post-hoc assessment. Increasing the proportion of data used for training the

CART should lead to more discerning power and more accurate results, likely at the cost of an accurate

estimation of cross-validation error rates. When constrained by small sample sizes, the choice of

whether to prioritize more accuracy in parameter selection or cross-validation is a judgement call that

depends on the study objectives and circumstances. If a post-hoc evaluation of classification

performance is deemed sufficient, researchers may choose to use the entire subset of data for which

nest locations are known to train the CART and skip the cross-validation.

Several factors may contribute to explain the discrepancies (both positive and negative) we observed

between the error rates we expected based on the CART output and the realized error rates. The error

rates estimated by the CART do not take into account an additional layer of skimming that is

introduced later in the analysis when competing attempts that temporally overlap to a top candidate are

discarded. This should reduce the realized false positive rate compared to what is expected based on

cross-validation alone. In our case, this was verified for both storks and gulls. However, spurious nest

sites were identified in some of the non-breeding kestrel tracks. This is likely explained by species-

specific behavior: non-breeding kestrels spend long stretches of time and consecutive days on a perch

while scanning for prey or resting. Distinguishing these patterns of attendance and re-visitation from

those of a nest might be challenging without applying restrictions based on seasonality and

geographical area. The underestimation of false negative rates based on the CART cross-validation

compared to our post-hoc assessment is in part due to the constraints we added in terms of minimum

consecutive days for kestrels and gulls. While decreasing our power to detect some early failed

breeding attempts, applying these additional constraints increased the positive predictive value in both

cases. Discrepancies between the expected and realized error rates also result, in part, from the

randomization selection of the sample used to train the CART.

Error rates for nest site identification vary in importance depending on the study objectives. If the

objective is to estimate reproductive outcome, ensuring that attempts are not missed should receive

priority over avoiding the selection of non-nest sites. Any re-visited location that gets erroneously

identified as a nest site would likely be classified as a failed attempt eventually anyway, and would not

affect estimates of the absolute number of successful attempts at the population level. In this case, we

suggest that researchers may want to focus on minimizing false negatives. Conversely, if the objective

of a study is, for instance, to analyze factors associated with nest site selection, minimizing false

positives should be the priority.

In an ideal situation, researchers would have access to high (or high-enough-) resolution data for the

focal species, with abundant on-ground data about nest locations and outcome to train the algorithm of

choice, estimate error rates, and validate results. Once this is done, data characteristics being equal, the

parameters found to identify nest sites can be applied to new individuals of the same species for which

on-ground information is not available. If CART is the tool of choice to inform the choice of parameter

values, we underline that classification thresholds in output should be used with caution, and

considered more as range indications than as clear-cut rules. We also recommend that parameter values

suggested by the outcome of the CART should be critically evaluated for their biological significance

before use, and that adjustments should be made as needed based on knowledge of the species biology.

Future efforts to improve our method for the identification of nest locations will include incorporating

uncertainty in our estimates of nest sites, allowing us to interpret classification results in a probabilistic

framework.

Reproductive outcome estimation – We obtained a satisfactory degree of separation in the estimated

probabilities of survival between failed and successful attempts in both gulls and kestrels, with the

estimates generally reflecting the true outcome of breeding attempts. However, having incomplete data

for each breeding attempt for kestrels and gulls affected the accuracy of our outcome estimation.

Missing the initial part of the attempts required adjustments of the parameter used to define the

duration of a breeding cycle in the nest survival models, which inevitably came with some assumptions

and generalizations. Namely, we assumed that the number of days that were missing from the data was

the same for all individuals within each species. If inaccurate, this assumption can result in wrong estimation of reproductive outcome. Assuming too short a duration of the missing initial phase can lead to underestimating reproductive outcome, by predicting true successes as failures, while assuming too long a duration can result in the opposite error, leading to the prediction of true failures as successes. Given the asynchrony of tagging dates and actual start dates of each breeding attempt, subtracting the same number of days from the breeding cycle for all individuals likely led to temporal mismatches. If the association between wrong estimation and true outcome is random, estimation errors should not affect population-level inference on the proportion of successful breeding events. In the absence of the data limitations we discussed, the performance of our method would likely improve compared to the results we presented.

An advantage of the kestrel and gull datasets, unlike the stork dataset, was the high temporal resolution. Lower sampling rates more likely result in lower detectability of nest visits, especially in some phases of the breeding attempt. For example, storks greatly reduce the frequency of their nest visits in the late nestling-rearing phase, making the detection of nest visits drop in the final part of breeding attempts with data at a 1-hour resolution. Detecting nest visits is critical for the estimation of reproductive outcome. Therefore, the higher the temporal resolution of the GPS data, the more reliable the estimates of reproductive outcome will be. Detectability of nest visits depends on the interaction of both species behavior and data resolution, as the probability of detecting a visit results from the combination of the actual frequency and duration of visits and the sampling rate. Therefore, there is no absolute rule-of-thumb as to what constitutes an adequate time resolution. Rather, sampling rate may be adjusted according to behavioral characteristic of each species when the study is designed.

The considerations we have made so far highlight how knowledge of the ecology of the focal species is necessary to successfully use the tools we presented in this paper. Researchers also need to be aware of data constraints and limitations. Overall, our application cases provide an illustration of how researchers can make the most of their imperfect data and still get useful, robust results while considering real-world shortcomings.

Synthesis and significance – The most important implication of our work is the ability to use telemetry

data to obtain estimates of bird reproductive outcome. Reproductive outcome is an important

component of fitness, and estimating it from tracking data will help bridge the long-sought connection

between movement and space use and components of demography at the individual level (Nathan et al.,

2008; Morales et al., 2010). Our method allows researchers to obtain critical information on

reproductive outcome for birds that nest in remote or inaccessible locations where it is difficult or risky

to collect on-ground data (Götmark, 1992; Mayer-Gross, Crick, & Greenwood, 1997; Etterson et al.,

2011; Wilmers et al., 2015). Getting information on reproductive outcome from tracking data also has

the advantage of allowing ready combination with environmental conditions experienced by individual

birds (Cagnacci et al., 2010; Pettorelli et al., 2014). This opens new, previously unattainable research

avenues regarding links between components of individual fitness, movement strategies, and habitat.

A limitation of our approach is that it does not provide estimation of reproductive success in terms of

number of offspring, but only in terms of overall success or failure (where success corresponds to at

least one nestling fledged and failure to none). However, estimating reproductive success at a finer

level might be possible in species whose behavioral signature in terms of nest site re-visitation patterns

differs depending on clutch size.

Besides estimating reproductive outcome, our method is useful as a tool to identify nest site locations.

Identifying nest sites is valuable in and of itself: for example, it may allow researchers to find

previously unknown nesting sites in species for which breeding locations are partly or entirely

unknown. Overall, our method can appeal to researchers with different objectives. First, it may be

useful to researchers that want to investigate reproductive outcome in relation to movement and

environmental factors. Second, it may serve researchers that want to obtain data on reproductive

outcome for species that are not easy to monitor on the ground. Third, it may be of use to researchers

that want to identify the location of nest sites for analyses of breeding site fidelity, nest-site selection

(Powell, Hodgman, Glanz, Osenton, & Fisher, 2010), or discovery of new breeding sites. Conservation

and management applications may both benefit from the availability of the tools we presented. Our

method can be applied both in situations of opportunistic use of historical tracking data or in cases

where the study is explicitly designed with these objectives in mind. Potentially, our framework can

even be adjusted for the identification of types of re-visited locations other than nest sites and taxa

other than birds.

Future research could focus on analyzing the temporal variation of the periodicity of nest visits

throughout the attempt, and possibly use that to recognize specific phases within the attempt

(incubation, early nestling-rearing, etc.). This would provide insight over the temporal component of

nesting patterns, allowing researchers to pinpoint the occurrence of events such as egg-laying or

hatching.

Conclusions

In the context of the ever-growing availability of telemetry data for more and smaller species and at

higher spatio-temporal resolution, the potential of the application we presented is vast. Our method can

be used to identify nest sites and estimate reproductive outcome of a broad variety of avian species.

When looked at from multiple angles and in different ecological contexts, the relationship between

movement and reproductive success can be investigated in a cohesive overarching framework at the

interface of avian and movement ecology.

Acknowledgments

We thank J. Hightower and C. Poli for providing helpful comments on earlier versions of this

manuscript. The wood stork tracking project was funded by the U. S. Fish and Wildlife Service, the

Army Corps of Engineers, the National Park Service, the Environmental Protection Agency (STAR

Fellowship to R.B.), the USDA National Institute of Food and Agriculture, and the Everglades

Foundation (ForEverglades Scholarship to S.P.). The Mediterranean gull tracking project was funded

by the Ministry for Environment, Land and Sea Protection of Italy and carried out with the support of

Ente Parco Delta del Po Emilia-Romagna, Reparto Carabinieri per la Biodiversità di Punta Marina, R.

Nardelli, A. Andreotti, G. Meneghini, F. Spina. Lesser kestrel tracking project activities were carried

out within the framework of the LIFE+Natura project "Un falco per amico" (LIFE11/NAT/IT000068)

with the support of S. Podofillini, E. Fulco, P. Giglio, S. C. Pellegrino, M. Lorusso (Comune di

Altamura), F. Parisi (Comune di Gravina in Puglia), and D. Ciampanella (project manager). Any use of

trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the

U. S. Government.

Authors' Contributions

S. Picardi conceived the idea. S. Picardi., B. J. S., and M. E. B. implemented the analyses. S. Picardi

wrote the paper. M. B. and P. C. F. supervised research. J. G. C., D. R., L. S., S. Pirrello, and R. R. B.

collected the data. All authors contributed to revisions.

References

Alonso, J. C., Carrascal, L. M., Muñoz-Pulido, R., & Alonso, J. A. (1994). Flock Size and Foraging
    Decisions in Central Place Foraging White Storks, *Ciconia Ciconia*. *Behaviour*, *129*(3–4), 279–
    292. doi:10.1163/156853994X00640

Andersson, M. (1981). Central Place Foraging in the Whinchat, *Saxicola Rubetra*. *Ecology*, *62*(3), 538–
    544. doi:10.2307/1937718

Austin, R., De Pascalis, F., Arnould, J., Haakonsson, J., Votier, S., Ebanks-Petrie, G., … Green, J.
    (2019). A sex-influenced flexible foraging strategy in a tropical seabird, the magnificent
    frigatebird. *Marine Ecology Progress Series*, *611*, 203–214. doi:10.3354/meps12859

Bar-David, S., Bar-David, I., Cross, P. C., Ryan, S. J., Knechtel, C. U., & Getz, W. M. (2009). Methods
    for assessing movement path recursion with application to African buffalo in South Africa.
    Ecology, *90*(9), 2467–2479. doi:10.1890/08-1532.1

Berger-Tal, O., & Bar-David, S. (2015). Recursive movement patterns: review and synthesis across
    species. *Ecosphere*, *6*(9), 1–12. doi:10.1890/ES15-00106.1

Beyer, H. L., Morales, J. M., Murray, D., & Fortin, M.-J. (2013). The effectiveness of Bayesian state-
    space models for estimating behavioural states from movement paths. *Methods in Ecology and
    Evolution*, *4*(5), 433–441. doi:10.1111/2041-210X.12026

Bonar, M., Ellington, E. H., Lewis, K. P., & Vander Wal, E. (2018). Implementing a novel movement-based approach to inferring parturition and neonate caribou calf survival. *PLOS ONE*, *13*(2), e0192204. doi:10.1371/journal.pone.0192204

Borkhataria, R. R., Frederick, P. C., Hylton, R., Bryan, A. L., & Rodgers, J. A. (2008). A Preliminary Model of Wood Stork Population Dynamics in the Southeastern United States. *Waterbirds*, *31*(sp1), 42–49. doi:10.1675/1524-4695(2008)31[42:APMOWS]2.0.CO;2

Bracis, C., Bildstein, K. L., & Mueller, T. (2018). Revisitation analysis uncovers spatio-temporal patterns in animal movement data. *Ecography*, *41*(11), 1801–1811. doi:10.1111/ecog.03618

Burke, C. M., & Montevecchi, W. A. (2009). The foraging decisions of a central place foraging seabird in response to fluctuations in local prey conditions. *Journal of Zoology*, *278*(4), 354–361. doi:10.1111/j.1469-7998.2009.00584.x

Cagnacci, F., Boitani, L., Powell, R. A., & Boyce, M. S. (2010). Animal ecology meets GPS-based radiotelemetry: a perfect storm of opportunities and challenges. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2157–2162. doi:10.1098/rstb.2010.0107

Cecere, J. G., Bondì, S., Podofillini, S., Imperio, S., Griggio, M., Fulco, E., … Rubolini, D. (2018). Spatial segregation of home ranges between neighbouring colonies in a diurnal raptor. *Scientific Reports*, *8*(1), 11762. doi:10.1038/s41598-018-29933-2

Clark, E. S. (1980). The Attentiveness and Time Budget of a Pair of Nesting Wood Storks. *Proceedings of the Colonial Waterbird Group*, *3*, 204–215.

Coulter, M. C., Rodgers, J. A., Ogden, J. C., & Depkin, F. C. (1999). Wood Stork (*Mycteria americana*). *The Birds of North America*, (409), 24.

De'ath, G., & Fabricius, K. E. (2000). Classification and Regression Trees: A Powerful yet Simple Technique for Ecological Data Analysis. *Ecology*, *81*(11), 3178–3192. doi:10.1890/0012-9658(2000)081[3178:CARTAP]2.0.CO;2

DeMars, C. A., Auger Méthé, M., Schlägel, U. E., & Boutin, S. (2013). Inferring parturition and neonate survival from movement patterns of female ungulates: a case study using woodland caribou. *Ecology and Evolution*, *3*(12), 4149–4160. doi:10.1002/ece3.785

DuRant, S. E., Hopkins, W. A., Hepp, G. R., & Walters, J. R. (2013). Ecological, evolutionary, and

    conservation implications of incubation temperature-dependent phenotypes in birds. *Biological*

    Reviews, *88*(2), 499–509. doi:10.1111/brv.12015

Edelhoff, H., Signer, J., & Balkenhol, N. (2016). Path segmentation for beginners: an overview of

    current methods for detecting changes in animal movement patterns. *Movement Ecology*, *4*, 21.

    doi:10.1186/s40462-016-0086-5

Etterson, M. A., Ellis-Felege, S. N., Evers, D., Gauthier, G., Grzybowski, J. A., Mattsson, B. J., …

    Potvien, A. (2011). Modeling fecundity in birds: Conceptual overview, current models, and

    considerations for future developments. *Ecological Modelling*, *222*(14), 2178–2190.

    doi:10.1016/j.ecolmodel.2010.10.013

Frair Jacqueline L., Fieberg John, Hebblewhite Mark, Cagnacci Francesca, DeCesare Nicholas J., &

    Pedrotti Luca. (2010). Resolving issues of imprecise and habitat-biased locations in ecological

    analyses using GPS telemetry data. *Philosophical Transactions of the Royal Society B:*

    Biological Sciences, *365*(1550), 2187–2200. doi:10.1098/rstb.2010.0084

Garriga, J., Palmer, J. R. B., Oltra, A., & Bartumeus, F. (2016). Expectation-Maximization Binary

    Clustering for Behavioural Annotation. *PLOS ONE*, *11*(3), e0151984.

    doi:10.1371/journal.pone.0151984

Götmark, F. (1992). The Effects of Investigator Disturbance on Nesting Birds. In D. M. Power (Ed.),

    Current Ornithology (pp. 63–104). Boston, MA: Springer US. doi:10.1007/978-1-4757-9921-

    7_3

Gurarie, E., Andrews, R. D., & Laidre, K. L. (2009). A novel method for identifying behavioural

    changes in animal movement data. *Ecology Letters*, *12*(5), 395–408. doi:10.1111/j.1461-

    0248.2009.01293.x

Gurarie, E., Bracis, C., Delgado, M., Meckley, T. D., Kojola, I., & Wagner, C. M. (2016). What is the

    animal doing? Tools for exploring behavioural structure in animal movements. *Journal of*

    Animal Ecology, *85*(1), 69–84. doi:10.1111/1365-2656.12379

Kays, R., Crofoot, M. C., Jetz, W., & Wikelski, M. (2015). Terrestrial animal tracking as an eye on life and planet. *Science*, *348*(6240), aaa2478. doi:10.1126/science.aaa2478

Lebreton, J.-D., Burnham, K. P., Clobert, J., & Anderson, D. R. (1992). Modeling Survival and Testing Biological Hypotheses Using Marked Animals: A Unified Approach with Case Studies. *Ecological Monographs*, *62*(1), 67–118. doi:10.2307/2937171

Mayer-Gross, H., Crick, H. Q. P., & Greenwood, J. J. D. (1997). The effect of observers visiting the nests of passerines: an experimental study. *Bird Study*, *44*(1), 53–65. doi:10.1080/00063659709461038

Morales Juan M., Moorcroft Paul R., Matthiopoulos Jason, Frair Jacqueline L., Kie John G., Powell Roger A., … Haydon Daniel T. (2010). Building the bridge between animal movement and population dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2289–2301. doi:10.1098/rstb.2010.0082

Mueller, T., & Fagan, W. F. (2008). Search and navigation in dynamic environments – from individual behaviors to population distributions. *Oikos, 117*(5), 654–664. doi:10.1111/j.0030-1299.2008.16291.x

Nams, V. O. (2014). Combining animal movements and behavioural data to detect behavioural states. *Ecology Letters*, *17*(10), 1228–1237. doi:10.1111/ele.12328

Nathan, R., Getz, W. M., Revilla, E., Holyoak, M., Kadmon, R., Saltz, D., & Smouse, P. E. (2008). A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences, 105*(49), 19052–19059. doi:10.1073/pnas.0800375105

Nicholson, K. L., Warren, M. J., Rostan, C., Månsson, J., Paragi, T. F., & Sand, H. (2019). Using fine-scale movement patterns to infer ungulate parturition. *Ecological Indicators*, *101*, 22–30. doi:10.1016/j.ecolind.2019.01.004

Norton, D. W. (1972). Incubation Schedules of Four Species of Calidridine Sandpipers at Barrow, Alaska. *The Condor*, *74*(2), 164–176. doi:10.2307/1366280

Ojowski, U., Eidtmann, C., Furness, R., & Garthe, S. (2001). Diet and nest attendance of incubating and chick-rearing northern fulmars (*Fulmarus glacialis*) in Shetland. *Marine Biology, 139*(6), 1193–1200. doi:10.1007/s002270100655

Orians, G. H., & Pearson, N. E. (1979). On the theory of central place foraging. In *Analysis of Ecological Systems* (pp. 154–177). Ohio University Press.

Pettorelli, N., Laurance, W. F., O'Brien, T. G., Wegmann, M., Nagendra, H., & Turner, W. (2014). Satellite remote sensing for applied ecologists: opportunities and challenges. *Journal of Applied Ecology, 51*(4), 839–848. doi:10.1111/1365-2664.12261

Plummer, M. (2003). JAGS: A program for analysis of Bayesian graphical models using Gibbs sampling. *Working Papers*, 8.

Polansky Leo, Kilian Werner, & Wittemyer George. (2015). Elucidating the significance of spatial memory on movement decisions by African savannah elephants using state–space models. *Proceedings of the Royal Society B: Biological Sciences, 282*(1805), 20143042. doi:10.1098/rspb.2014.3042

Powell, L. L., Hodgman, T. P., Glanz, W. E., Osenton, J. D., & Fisher, C. M. (2010). Nest-Site Selection and Nest Survival of the Rusty Blackbird: Does Timber Management Adjacent to Wetlands Create Ecological Traps? *The Condor, 112*(4), 800–809. doi:10.1525/cond.2010.100152

Riotte-Lambert, L., Benhamou, S., & Chamaillé-Jammes, S. (2013). Periodicity analysis of movement recursions. *Journal of Theoretical Biology, 317*, 238–243. doi:10.1016/j.jtbi.2012.10.026

Schaub, M., & Royle, J. A. (2014). Estimating true instead of apparent survival using spatial Cormack–Jolly–Seber models. *Methods in Ecology and Evolution, 5*(12), 1316–1326. doi:10.1111/2041-210X.12134

Schick, R. S., Loarie, S. R., Colchero, F., Best, B. D., Boustany, A., Conde, D. A., … Clark, J. S. (2008). Understanding movement data and movement processes: current and emerging directions. *Ecology Letters, 11*(12), 1338–1350. doi:10.1111/j.1461-0248.2008.01249.x

Severud, W. J., Giudice, G. D., Obermoller, T. R., Enright, T. A., Wright, R. G., & Forester, J. D. (2015). Using GPS collars to determine parturition and cause-specific mortality of moose calves. *Wildlife Society Bulletin*, *39*(3), 616–625. doi:10.1002/wsb.558

Singh Navinder J., & Ericsson Göran. (2014). Changing motivations during migration: linking movement speed to reproductive status in a migratory large mammal. *Biology Letters*, *10*(6), 20140379. doi:10.1098/rsbl.2014.0379

Snow, D. W., Perrins, C. M., Hillcoat, B., Gillmor, R., & Roselaar, C. S. (1997). The birds of the Western Palearctic. concise edition. Retrieved from https://dare.uva.nl/search?identifier=2997a701-3eac-4e5a-835f-981bfced489e

Soleymani, A., Pennekamp, F., Dodge, S., & Weibel, R. (2017). Characterizing change points and continuous transitions in movement behaviours using wavelet decomposition. *Methods in Ecology and Evolution*, *8*(9), 1113–1123. doi:10.1111/2041-210X.12755

Thiebault, A., Dubroca, L., Mullers, R. H. E., Tremblay, Y., & Pistorius, P. A. (2018). "m2b" package in r: Deriving multiple variables from movement data to predict behavioural states with random forests. *Methods in Ecology and Evolution*, *9*(6), 1548–1555. doi:10.1111/2041-210X.12989

Tomkiewicz Stanley M., Fuller Mark R., Kie John G., & Bates Kirk K. (2010). Global positioning system and associated technologies in animal behaviour and ecological research. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2163–2176. doi:10.1098/rstb.2010.0090

Urbano Ferdinando, Cagnacci Francesca, Calenge Clément, Dettki Holger, Cameron Alison, & Neteler Markus. (2010). Wildlife tracking data management: a new vision. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *365*(1550), 2177–2185. doi:10.1098/rstb.2010.0081

Van Moorter, B., Visscher, D. R., Jerde, C. L., Frair, J. L., & Merrill, E. H. (2010). Identifying Movement States From Location Data Using Cluster Analysis. *Journal of Wildlife Management*, *74*(3), 588–594. doi:10.2193/2009-155

van Vessem, J., & Draulans, D. (1986). Nest Attendance by Male and Female Gray Herons. *Journal of Field Ornithology*, *57*(1), 34–41.

Webb, N. F., Hebblewhite, M., & Merrill, E. H. (n.d.). Statistical Methods for Identifying Wolf Kill

Sites Using Global Positioning System Locations. *The Journal of Wildlife Management, 72*(3),

798–807. doi:10.2193/2006-566

Wilmers, C. C., Nickel, B., Bryce, C. M., Smith, J. A., Wheat, R. E., & Yovovich, V. (2015). The

golden age of bio-logging: how animal-borne sensors are advancing the frontiers of ecology.

Ecology, 96(7), 1741–1753. doi:10.1890/14-1401.1

Figures and Tables

Table 1 – Differences among GPS-tracking datasets for wood storks, lesser kestrels, and Mediterranean

gulls in terms of sample sizes, data characteristics, and species seasonality and ecology.

		Wood stork	Lesser kestrel	Mediterranean gull
Spatial resolution (m)		18	<10	<10
Temporal resolution (min)		60	15 summer/30 winter	15
Fix failure rate		High	Low	Low
Tagged at		Fledging/non-breeding	Early nestling-rearing	Incubation
Number of tracks (individual-years)	Total	148	53	29
	Known nest location	107	53	29
	Known outcome	0	53	29
	Non-breeders	41 (subadults)	16 (winter)	16 (winter)
Breeding season		Nov-Aug (varies with latitude)	Apr-Jul	Apr-Jul
Breeding cycle (days)		110	60	60

Table 2 – Results of Classification and Regression Trees (CART) on nest versus non-nest sites,

including parameter values used to identify nest sites among re-visited locations and cross-validated

469 error rates. Values marked with an asterisk were not taken from the CART output, but were chosen as

470 reasonable thresholds for selecting nest sites. Consecutive days: maximum number of consecutive days

471 spent at a location. Days visited: percentage of days when a location was visited between the first and

472 last visits. Attendance on top day: percentage of GPS fixes at a location on the day with maximum

473 attendance.

	Wood stork	Lesser kestrel	Mediterranean gull
Consecutive days (minimum)	18	4*	8*
Days visited (minimum)	1%	1%	1%
Attendance on top day (minimum)	90%	52%	45%
Cross-validated type I error rate (false positive)	6%	5%	30%
Cross-validated type II error rate (false negative)	8%	10%	20%

475 Table 3 – Performance metrics of the nest identification algorithm (see Methods for definitions).

	Wood stork	Lesser kestrel	Mediterranean gull
Positive predictive value	86.79%	93.48%	78.57%
Sensitivity	85.98%	81.13%	75.86%
False negative rate	14.02%	18.87%	24.14%
False positive rate	0%	68.75%	6.25%

Figure 1 – Workflow of the analysis to identify nest sites and estimate reproductive outcome from

telemetry data. The R package nestR includes functions to tackle each of the steps depicted in the

boxes.

Figure 2 – Output of CART analysis to discriminate nest and non-nest sites in A) wood stork, B) lesser

kestrel, C) Mediterranean gull.

Figure 3 – Probability of visit detection (top row) and survival (bottom row) through time estimated at

the population level for A) wood stork, B) lesser kestrel, C) Mediterranean gull.

Figure 4 – Distribution of estimated survival probabilities for breeding attempts in relation to their true

outcome for A) lesser kestrels and B) Mediterranean gull.