1 **Title:** Thyroid hormone coordinates developmental trajectories but does not underlie
2 developmental truncation in danionins
3

4 **Authors:** Yinan Hu, Angela Mauri, Joan Donahue, Rajendra Singh, Benjamin Acosta, Sarah
5 McMenamin
6
7 **Institutional Affiliation**: Boston College Biology Department
8
Corresponding Contact Information:
Sarah McMenamin
Boston College
Biology Department
140 Commonwealth Ave
Chestnut Hill MA 02467
mcmenams@bc.edu
617-552-3989

17
18 **Running Title:** Thyroid Hormone in danionins
19
20 **Key Words:** thyroid hormone, zebrafish, Danionella, danionin, paedomorphosis,
21 metamorphosis, morphometrics, lateral line, neuromasts
22
Bullet Points:
• Thyroid hormone regulates shifts in relative growth trajectories in different zebrafish tissues
• Thyroid hormone inhibits head growth in juvenile zebrafish, and regulates juvenile growth
patterns
• Thyroid hormone stimulates formation of canal neuromasts in the head and proliferation of
trunk neuromasts in zebrafish
• *Danionella*, a miniaturized danionin lineage closely related to zebrafish morphologically
resemble larval zebrafish and do not form neuromast canals
• Hypothyroidism is not the cause of paedomorphosis in *Danionella*
32

33 **Grant Sponsors:**
34 NIH R00GM105874
35 NIH R03HD091634
36 Burroughs Wellcome Collaborative Research Travel Grant 1017439
37 NSF CAREER 1845513

ABSTRACT

Background

Differences in postembryonic developmental trajectories can profoundly alter adult phenotypes and life histories. Thyroid hormone (TH) regulates metamorphosis in many vertebrate taxa with multiphasic ecologies, and alterations to TH metabolism underlie notable cases of paedomorphosis in amphibians. We tested the requirement for TH in multiple postembryonic developmental processes in zebrafish, which has a monophasic ecology, and asked if TH production was compromised in paedomorphic *Danionella*.

Results

We showed that TH regulates allometric growth in juvenile zebrafish, and inhibits relative head growth. The lateral line system showed differential requirements for TH: the hormone promotes canal neuromast formation and inhibits neuromast numbers in the head, but causes expansion of the neuromast population in the trunk. While *Danionella* morphology resembled that of larval zebrafish, the two *Danionella* species analyzed were not similar to hypothyroid zebrafish in their shape or neuromast distribution, and both possessed functional thyroid follicles.

Conclusions

Although zebrafish do not undergo a discrete ecological transformation, we found that multiple tissues undergo transitions in developmental trajectories that are dependent on TH, suggesting the TH axis and its downstream pathways as likely targets for adaptation. Nonetheless, we found no evidence that evolutionary paedomorphosis in *Danionella* is the result of compromised TH production.

INTRODUCTION

Shifts in the timing and magnitude of developmental processes during later, postembryonic stages of development can profoundly alter adult phenotypes (Gould, 1977; Raff, 1996; Smith, 2001). Revealing the pathways that regulate ontogenetic trajectories of different developmental processes will be critical in understanding how life histories and phenotypes are formed. Further, discerning the mechanisms by which different processes may be decoupled will be critical for a complete understanding of the evolution of development.

An extreme way in which postembryonic development may be changed is through paedomorphosis, wherein an organism does not complete terminal stages of development (Gould, 1977; Shaffer and Voss, 1996); organisms on the spectrum of paedomorphosis retain larval characteristics into adulthood (Johnson and Voss, 2013). Paedomorphosis is generally most obvious when it originates in clades with ecologically multiphasic life histories. Ecologically multiphasic vertebrates are those that undergo an ecological shift as they transform to juveniles, e.g. amphibians that metamorphose from aquatic larvae to terrestrial juveniles. Paedomorphosis in these clades involves loss of a markedly divergent life history stage and may be phenotypically quite obvious (McMenamin and Hadly, 2010; Johnson and Voss, 2013). Paedomorphosis in clades with ecologically monophasic life histories may be less morphologically striking, but paedomorphosis has nonetheless been posited as an important evolutionary process in many monophasic groups, including in mammals (e.g. Shea, 1983; Goodwin et al., 1997).

Since vertebrates with multiphasic life cycles often undergo a metamorphosis triggered by a
single molecular pathway (generally thyroid hormone, TH), paedomorphosis could theoretically
evolve in these groups through a single genetic change. Thus, disruptions to TH pathways cause
developmental arrest and underlie cases of paedomorphosis, as in the Mexican axolotl (see
Johnson and Voss, 2013; De Groef et al., 2018). In contrast, ecologically monophasic
vertebrates generally have more subtle relationships with the molecular signals that regulate
progression through post-embryonic development. TH functions as an important regulator of
developmental processes in ecologically monophasic vertebrates (e.g. zebrafish and mammals),
but disruptions to the thyroid axis do not result in whole-organism arrest (Brown, 1997;
McMenamin et al., 2014; Buchholz, 2015). In ecologically monophasic groups, the pathways
regulating postembryonic developmental progression are less clear, as are the mechanisms and
pathways that might underlie evolutionary paedomorphosis.
Teleosts represent the largest and most speciose group of vertebrates, exhibiting a
remarkable range of life history strategies (Finn and Kapoor, 2008; McMenamin and Parichy,
2013). Most marine fishes are ecologically multiphasic, and pelagic larvae undergo profound
physiological metamorphosis to transform into benthic juveniles (McMenamin and Parichy,
2013). Metamorphosis is stimulated by TH in flat fish (Isorna et al., 2009; Xu et al., 2016;
Campinho et al., 2018) and reef fish (Holzer et al., 2017), and likely in many other species
(Holzer and Laudet, 2015). Like in amphibians, TH stimulates the abrupt transformation of
discrete tissues across the body, causing dramatic changes in overall morphology, sensory
systems and other organs.
In contrast, most freshwater fishes, including zebrafish (*Danio rerio*), are ecologically
monophasic. These monophasic species show a more subtle and protracted transformations
from larva to juvenile. Nonetheless, specific tissues (including skeleton, pigment pattern and
the lateral line) do undergo major remodeling during postembryonic development (Parichy and
Turner, 2003; Webb and Shirey, 2003; Parichy, 2006; Parichy et al., 2009). Many of these
remodeling processes, including those in the skeleton and the pigment pattern, are dependent
on TH (Shkil et al., 2010; Shkil et al., 2012; McMenamin et al., 2014; Keer et al., 2019). Whether
in total these transformations and remodeling processes constitute a "metamorphosis"
depends on how broadly the term is defined (e.g. see Laudet, 2011; McMenamin and Parichy,
2013). A clearer picture of normal developmental trajectories in zebrafish, as well as the
relative dependencies on TH, can inform our concept of how normal postembryonic
development is regulated and whether this species exhibits a metamorphosis homologous to
that of marine species.
Many fish species have been described as paedomorphic, including ice fish (Albertson et al.,
2010), gobies (Johnson and Brothers, 1993; Giovannotti et al., 2007) and blennioid fishes
(Hastings, 2002). *Danionella* is a danionin relative of zebrafish, sister clade to the group
containing zebrafish and *Danios*, *Devarios* and *Microdeverios* (McCluskey and Postlethwait,
2014). *Danionella* are miniaturized and developmentally arrested, resembling larval zebrafish in
several ways: adult *Danionella* lack scales, barbels and many typically late-ossifying bones (Britz
et al., 2009; Britz and Conway, 2016), and retain a larva-like pigment pattern and feeding
kinematics (McMenamin et al., 2017; Galindo et al., 2018). *Danionella translucida* is emerging
as a novel model for studying behavior, neurobiology and brain function (Penalva et al., 2018;
Schulze et al., 2018), and a hybrid genome assembly was recently completed (Kadobianskyi et

al., 2019). A better understanding of the developmental relationships between *Danionella* and
zebrafish, along with information about the pathways underlying paedomorphosis in
Danionella, will offer new insights about this emerging model.
In this study we focused on two trait systems that change over the course of postembryonic
development. We first focused on body shape, which changes dramatically during
postembryonic development of many species. Vertebrates with multiphasic life cycles exhibit
considerable shape change (i.e. allometric growth) during their profound physiological
metamorphosis, followed by more proportional (i.e. isometric) growth as juveniles and adults.
We next focused on development of the lateral line system. This flow-sensing sensory
system consists of individual neuromasts distributed across the body in specific patterns (Webb
and Shirey, 2003; Coombs et al., 2014; Webb, 2014). The lateral line system is transformed
during metamorphosis in amphibians (Wahnschaffe et al., 1987) and lampreys (Gelman et al.,
2008), and in teleosts with both ecologically multiphasic (e.g. flatfishes; Harvey et al., 1992) and
ecologically monophasic lifecycles (e.g. zebrafish; Webb and Shirey, 2003).
Our goals in this study were three-fold. <u>First</u>, we aimed to characterize shape change and
the development of the lateral line system during normal zebrafish development, asking
whether these developmental trajectories show evidence of a metamorphic transformation
that corresponds with a period of increased TH production. <u>Second</u>, we tested whether TH is
required in the development of these systems, determining the extent to which the hormone is
required to coordinate shape change and/or lateral line development with overall growth.
<u>Finally</u>, we tested whether adults of two *Danionella* species recapitulate any stage of normal
zebrafish ontogeny, further testing the possibility that developmental arrest in *Danionella* is
due to a lack of TH production.
148
RESULTS AND DISCUSSION
Zebrafish show allometric shifts during postembryonic development
To test the extent to which euthyroid wild-type (WT) zebrafish undergo changes in overall
morphology during postembryonic development (see Fig 1A-B), we first assessed allometric
growth trajectories of body shape (Fig 2, green crosses). Together, the first three principal
components captured more than 88% of total shape variation; these were plotted against
individual standard length (SL; Fig 2). PC1 primarily captured body depth (Fig 2D, G). In this axis,
WT zebrafish allometrically increase body depth until the end of the larval period (10 mm SL),
then maintain a roughly isometric change in body depth (Fig 2A). PC2 captures the relative size
of the head. This component peaks at ~9 mm SL, and then decreases gradually during juvenile
and adult growth (green crosses, Fig 2B). The dorsal profile, captured by PC3 increases gradually
with growth (Fig 2 C).
To more directly test for growth in head proportion during development, we plotted the
ratio of head length/SL over the course of growth (Fig 3C, green crosses). This ratio mirrored
PC2, peaking at ~8 mm SL and then gradually declined afterwards (green crosses, Fig 3C). Eye
size had a similar developmental trajectory to head length, continued to grow proportionally
during larval phase, peaking at ~10 mm SL (green crosses, Fig 3D). Thus, WT zebrafish exhibit
shifts in allometric relationships during the transition from larva to juvenile (8-10 mm SL); these
shifts are consistent with a systemic metamorphosis. The pattern of positive allometric growth
of the head has been documented in a number of fish species (e.g. see Osse et al., 1997;

Gisbert, 1999; Gisbert and Doroshov, 2006), and likely reflects strong selection on early
development of craniofacial feeding apparatus, , respiratory and sensory systems.
171
TH inhibits head growth in the juvenile
To determine the extent to which TH is required for these allometric shifts, we measured
shape and proportion changes in transgenically thyroid ablated, hypothyroid fish (TH-; Fig 1C,
D). Changes in body depth were independent of TH, and shape changes along the PC1 axis were
indistinguishable between TH- and WT (Fig 2A). However, we found that TH influences the
relative growth of the head: in TH- fish, PC2 continues to increase along a larva-like trajectory
until ~10 mm SL (purple circles, Fig 2 B). The third PC axis reveals that TH- fish develop an
increasingly flat dorsal profile, unlike that of any stage of development in WT (Fig 2C).
In TH- fish, heads and eyes continue to grow along relatively steeper (larva-like) trajectories
(Fig 2A, B). After rapid relative growth during larval development, the relative size of the head
and eye does not decline appropriately in TH- fish (Fig 2C-D). Thus, TH is required for the
coordinated transition from a larval trajectory to the juvenile trajectory, and limits the relative
growth of the head in juveniles.
185
TH inhibits proliferation of cranial neuromasts, but promotes formation of neuromast stitches
on the body
We next focused on the development of a trait that emerges via very different processes
than allometric body growth. We tracked numbers of neuromasts of the lateral line system
across a range of body sizes (see Fig 4). In WT zebrafish, the number of trunk neuromasts
increases roughly in proportion to body size, whereas neuromasts in the head show a negative
allometric relationship to body size (green crosses, Fig 5A, B). We found that TH- adults had
~25% more total neuromasts than did WT adults (Fig 5C), and that this difference was driven
primarily by the increasing number of cranial neuromasts (Fig 5A). This increase in cranial
neuromast number is not simply due to the fact that TH- fish have larger heads, as differences
remain even when standardized by head length (Fig 5D).
A subset of cranial neuromasts acquire an elongated, rod-like shape in WT juveniles and
eventually sink into bony canals (Webb and Shirey, 2003). We found that although neuromasts
do become enclosed in canals in TH- adults, these canal neuromasts fail to acquire the mature
morphology seen in WT (Fig 4B-C versus 4F-G). The transition into mature canal neuromasts
may serve as a cue that inhibits neuromast proliferation; without this cue, neuromasts in TH-
heads may remain highly proliferative through adulthood.
In WT zebrafish, trunk neuromasts proliferate into vertical stitches starting at ~10 mm SL
(Ledent, 2002; visible in Fig 4I-J and M). In TH- fish, the trunk neuromast population grows more
slowly until stitches start forming at ~13 mm SL (see Fig 4M and Fig 5B), suggesting that TH
regulates onset of stitch formation.
207
Bioavailability of TH shifts during mid-larval development
TH production peaks during metamorphic climax in amphibians, measured as both the
prohormone thyroxine (T4, the form of TH primarily produced by the thyroid follicles, see Fig
6D) and triiodothyronine (T3, the more biologically active form that binds to nuclear TH
receptors; Regards et al., 1978; Kikuyama et al., 1993; Shi, 2000; Brown and Cai, 2007). We

wanted to test whether the physiological changes stimulated by TH in zebrafish corresponded
with periods of elevated TH production or changes in the ratio of T3/T4. We measured T4 and
T3 levels at seven stages from larval through juvenile development, staging groups of fish
according to the Standardized Standard Length staging criteria (SSL; see Parichy et al., 2009).
The relative levels of T3 and T4 remained relatively consistent from one stage to another,
except as fish transitioned from 6 to 6.4 SSL. During this period, when anal and dorsal rays first
appear (Parichy et al., 2009), we detected a drop in T4 corresponding with an increase in T3
titer (Fig 6A-C). We conclude that there is a shift in TH bioavailability during this mid-larval
period, which we have highlighted with grey bars in the graphs of Figs 2, 3, 5-7. Following this
shift, T3 availability remained elevated through juvenile development (Fig 6B).
A previous analysis identified a similar shift in T3/T4 ratio at 10-14 dpf, attributed to higher
deiodinase activity (Chang et al., 2012). Indeed, while expression of *thyroglobulin* remains
relatively stable from 10-20 dpf, *deiodinase 2* expression increases two-fold (Vergauwen et al.,
2018). However, these patterns are difficult to perfectly correlate with our measurements, as
dpf (used in these previous analyses) is an unreliable proxy for development (Parichy et al.,
2009; McMenamin et al., 2016).
Shifts in allometric trajectories of WT fish occur at 8-10 mm SL (Fig 2A, B; Fig 3C, D). In
contrast, the mid-larval shift in T3/T4 ratio occurs at a considerably smaller size (6.4 SSL) and
earlier developmental stage (Fig 6), so it seems unlikely that the change in TH ratio directly
stimulates the shifts. Nonetheless, the earliest differences between TH- and WT fish are
detectable only after ~8 mm SL in allometric relationships (Fig 3B, D) and neuromast numbers
(Fig 5B, E). Rather than stimulating a discrete period of profound physiological change (like in
metamorphic amphibians), ongoing exposure to TH during late larval periods appears to
gradually shift developmental trajectories. TH- zebrafish maintain inappropriately longer
periods of larval growth patterns; rather than maintaining larval shapes, TH- zebrafish
ultimately achieve novel adult phenotypes.
239
Adult *Danionella* resemble larval zebrafish
Danionella maintain numerous characteristics of larval zebrafish, and are described as
paedomorphic (Britz et al., 2009; Britz and Conway, 2016). We tested the prediction that
Danionella body shape should more closely resemble larval rather than adult zebrafish, by
reanalyzing our morphometric datasets to include *Danionella translucida* and *D. priapus*.
Danionella adults of both species exhibited lower PC1 scores than similarly-sized zebrafish (blue
triangles and orange squares, Fig 7), showing immature body shape relative to similarly-sized
zebrafish. Both species of *Danionella* also showed proportionally smaller heads and eyes for
their body length (blue triangles and orange squares, Fig 2). Thus, the two *Danionella* species
indeed maintain certain morphologies similar to those of larval zebrafish, and are likely to have
more isometric patterns of growth during development.
Although *D. translucida* were smaller than *D. priapus*, the two species had similar total
numbers of neuromasts (~300; Fig 5C). The size of the *D. priapus* neuromast population was
comparable to that of similarly-sized WT zebrafish, while *D. translucida* had more total
neuromasts, with a greater relative density in the trunk (Fig 5B-C, E).
255
***Danionella* do not resemble hypothyroid zebrafish, and possess functional thyroid follicles**

Since TH is involved in the most well-known examples of paedomorphosis (Gould, 1977;
Johnson and Voss, 2013; De Groef et al., 2018), we tested the possibility that TH metabolism
was disrupted in *Danionella* species. First, we asked if *Danionella* appeared more similar to TH-
than to WT zebrafish. In terms of morphometrics (Fig 7) and ratios of head and eye to SL (Fig 3),
both *Danionella* species actually more closely resembled similarly-sized WT than TH- zebrafish.
Similar to TH- zebrafish, the two *Danionella* species did not form elongated canal neuromasts
(Fig 4D, H). However, *Danionella* show fewer cranial neuromasts and more trunk neuromasts
than similarly-sized WT zebrafish, in contrast to the patterns observed in TH- zebrafish (Fig 5).
In all, the two *Danionella* species did not resemble TH- zebrafish. To further confirm that
hypothyroidism did not underlie developmental truncation in these species, we examined the
thyroid follicles, which were intact and showed T4 production comparable to that in zebrafish
(Fig 6D-F).
269
Conclusions
We identified shifts in allometric trajectories as WT zebrafish transition from larva to
juvenile, and some aspects of these shifts showed dependence on TH. However, unlike
hypothyroid amphibians, which maintain larval morphology through adulthood, TH- zebrafish
inappropriately maintain larval growth trajectories and ultimately achieve novel adult
phenotypes. We conclude that TH normally coordinates multiple aspects of the transition from
larval to juvenile developmental programs in zebrafish. We interpret these hormonally-
mediated changes to constitute a protracted, subtle metamorphosis, in contrast to the sudden
and profound metamorphosis of ecologically multiphasic species. Based on its dependence on
thyroid hormone, we posit that this subtle metamorphosis is indeed evolutionarily homologous
to the profound metamorphosis of ecologically multiphasic vertebrates.
Consistent with the idea that *Danionella* are paedomorphic, two *Danionella* species
analyzed showed several similarities to larval WT zebrafish in their body proportions and their
lateral line. However, *Danionella* bore little resemblance to TH- zebrafish, and we found that
they produced T4. We conclude that, unlike in paedomorphic amphibians, disruptions to the
thyroid axis do not underlie paedomorphosis in these species.
286
EXPERIMENTAL PROCEDURES
Fish rearing
Zebrafish were of the line *Tg(tg:nVenus-v2a-nfnB)* (McMenamin et al. 2014), which allows
conditional ablation of the thyroid. Transgenic fish were treated 4-5 days post fertilization (dpf)
with DMSO (for control euthyroid fish, WT) or DMSO + metronidazole (to ablate the thyroid
follicles and produce hypothyroid fish, TH-), following McMenamin et al (2014). All fish were
reared at 28°C with a 12:12 light:dark cycle and fed three times per day with marine rotifers,
Artemia and pure *Spirulina* flakes (Pentair). This diet was chosen to minimize introduction of
exogenous TH. *Danionella translucida* was a gift from Adam Douglass, University of Utah; *D.*
priapus was purchased through The Wet Spot (Portland OR). Both species of *Danionella* were
fed rotifers, *Artemia* and Gemma micro (Skretting).
298
Live staining of neuromasts

The fluorescent mitochondrial stain 4-(4-Diethylaminostyryl)-1-methylpyridinium iodide (DiASP,
AdipoGen) was used to visualize neuromasts (Collazo et al., 1994). Individual fish were
incubated in 0.0024% DiASP in fish system water with 0.02% Tricane (Western Chemical) for 5
minutes, then transferred to 2% methylcellulose and imaged with an Olympus DP74 camera
mounted on an Olympus SZX16 stereoscope with a FITC filter. Standard length (SL) was
measured using ImageJ (Schneider et al., 2012). Neuromasts were counted manually from
digital images; all counts were performed by the same individual to reduce variability.
307

308   **Scanning Electron Microscopy**
309   WT and TH- adult zebrafish were fixed in 4% paraformaldehyde (Sigma) overnight at 4°C, rinsed
310   several times with PBS, and dehydrated in an ascending series of ethanol. Samples were critical
311   point dried in 100% ethanol, coated with Au-Pd and imaged with a JEOL JCM-6000PLUS
312   NeoScope Benchtop SEM.
313

Fixation and skeletal staining
Specimens were euthanized by an overdose of Tricane, then fixed in 4% paraformaldehyde
(Sigma) overnight at 4°C. Specimens were rinsed several times with PBS and dehydrated in 70%
ethanol. Following (Walker and Kimmel, 2007), skeletons were stained overnight in 0.5%
Alizarin red (Sigma) and 0.02% Alcian Blue (Sigma) with 60 mM $MgCl_2$ in 70% EtOH. Stained
specimens were bleached with 3% hydrogen peroxide in 1% potassium hydroxide (KOH), then
were cleared by gradually transitioning through an ascending glycerol series from 25% to 80%.
Prior to imaging, cleared specimens were incubated in PBS overnight, and then transferred into
2% methylcellulose for positioning and imaged as above for morphometric analysis.
323

Morphometrics
Homologous anatomical landmarks and curves that outline the body were placed manually (by
a single individual) on lateral images of fish using the R package Stereomorph (Olsen and
Westneat, 2015). A list of the specific landmarks can be found in the supplementary materials
(supplementary Table S1). Landmark data was then analyzed with standard geometric
morphometrics methods using the R package Geomorph (Adams and Otárola-Castillo, 2013).
330

Head size and eye size
Head size was measured in images of cleared and stained specimens in lateral view. Head size
was measured as the distance from the tip of the snout/rostral edge of premaxilla to the
posterior tip of the occipital bone. Two eye measurements, the dorsal-ventral height, and the
anterior-posterior length were obtained, and then averaged as the eye-size measure.
Measurements were performed by a single individual to reduce variability.
337

Thyroid staining
To image thyroid follicles, adult zebrafish, *D. priapus* and *D. translucida* were fixed in 4%
paraformaldehyde (Sigma), embedded in O.C.T. Compound (Fisher Scientific), and sectioned on
to 12 μm. Slides were washed with 0.1% Tween in PBS (PBSt), blocked with 10% heat-
inactivated goat serum in PBSt, then incubated at 4°C overnight with a rabbit T4 antibody
(1:1000, Fitzgerald). Slides were then washed, incubated with a secondary antibody (goat anti-

rabbit FITC diluted 1:1000, Fitzgerald), stained with DAPI (1ug/ml) for 20 minutes, washed and
imaged on an Olympus IX83 microscope equipped with a Hamamatsu ORCA Flash 4.0LT camera.
346

Quantification of thyroxine (T4) and triiodothyronine (T3)
Live WT zebrafish were anesthetized and staged following the adult normal table (Parichy et al.
2009). Seven stages were chosen to capture major processes during the entire transition from
larva to juvenile (see Supplemental Table 2), and fish were staged by the SSL convention
(Parichy et al., 2009). Samples were prepared following McMenamin et al. (2014); briefly,
staged fish were pooled, euthanized by an overdose of tricaine, blotted dry, and weighed in
microcentrifuge tubes. Tissue was suspended in cold methanol with 1 mM 6-propyl-2-thiouracil
(PTU; Sigma) and homogenized by sonication; homogenized samples were stored at -80°C until
analysis. Samples were centrifuged, re-extracted with additional PTU-containing methanol.
Supernatants were pooled and dried with a CentriVap Concentrator (Labconco) for ~5 hours,
then resuspended in PBS. Samples were run on T4 and T3 ELISA kits (Diagnostic Automation
Inc.) with two technical replicates and with three biological replicates for each stage. Plates
were analyzed on a PerkinElmer Victor X5; mean absorbance value from three readings was
used for subsequent analysis. Standard curves were generated with 4-parameter logistic
regression via online analysis software (www.elisaanalysis.com).

Statistics
All statistical analyses were performed in in R (version 3.4.1). Differences in thyroid hormone
levels along ontogeny were first examined using ANOVA, followed by Tukey's Honest Significant
Distance Test. Developmental trajectory was estimated with the spm function in the package
SemiPar using default settings (Carroll et al., 2003). Developmental trajectories were
considered to be distinct when the ranges of standard error did not overlap between the
groups. For example, in Fig 2A, the standard error ranges always overlap (the green and purple
bands overlap) and the models are not considered different. In Fig 2B, the standard error
ranges of two models diverges at ~9 mm SL.

ACKNOWEDGEMENTS
Thank you for discussion and technical support from Catherine May, Angela Mauri Stacy
Nguyen, Pranav Parikh, Mary Xu and all the members of the McMenamin Lab. Thank you to
Jacqueline Webb and another anonymous reviewer for their insightful comments. Thank you to
Adam Douglass, David Parichy, the Boston College Statistics Department and the Boston College
Animal Care staff. We gratefully acknowledge support from NIH R00GM105874, R03HD091634,
Burroughs Wellcome Grant 1017439 and NSF CAREER 1845513.

REFERENCES

Adams DC, Otárola-Castillo E. 2013. geomorph: an R package for the collection and analysis of
geometric morphometric shape data. Methods Ecol Evol 4(4): 393-399.
Albertson RC, Yan Y-L, Titus TA, Pisano E, Vacchi M, Yelick PC, Detrich HW, Postlethwait JH.
2010. Molecular pedomorphism underlies craniofacial skeletal evolution in Antarctic
notothenioid fishes. BMC Evol Biol 10(1): 4.

Britz R, Conway KW. 2016. *Danionella dracula*, an escape from the cypriniform Bauplan via
developmental truncation? J Morphol 277(2): 147-166.
Britz R, Conway KW, Rüber L. 2009. Spectacular morphological novelty in a miniature cyprinid
fish, *Danionella dracula*. Proc R Soc Lond, Ser B: Biol Sci 276(1665): 2179-2186.
Brown DD. 1997. The role of thyroid hormone in zebrafish and axolotl development.
Proceedings of the National Academy of Sciences 94(24): 13011-13016.
Brown DD, Cai L. 2007. Amphibian metamorphosis. Dev Biol 306(1): 20.
Buchholz DR. 2015. More similar than you think: Frog metamorphosis as a model of human
perinatal endocrinology. Dev Biol 408(2): 188-195.
Campinho MA, Silva N, Martins GG, Anjos L, Florindo C, Roman-Padilla J, Garcia-Cegarra A,
Louro B, Manchado M, Power DM. 2018. A thyroid hormone regulated asymmetric responsive
centre is correlated with eye migration during flatfish metamorphosis. Scientific reports 8(1):
12267.
Carroll R, Ruppert D, Wand M (2003). Semiparametric regression, Cambridge: Cambridge
University Press.
Chang J, Wang M, Gui W, Zhao Y, Yu L, Zhu G. 2012. Changes in thyroid hormone levels during
zebrafish development. Zool Sci 29(3): 181-184.
Collazo A, Fraser SE, Mabee PM. 1994. A dual embryonic origin for vertebrate
mechanoreceptors. Science 264(5157): 426-430.
Coombs S, Bleckmann H, Fay RR, Popper AN. 2014. The lateral line system: Springer.
De Groef B, Grommen SV, Darras VM. 2018. Forever young: endocrinology of paedomorphosis
in the Mexican axolotl (*Ambystoma mexicanum*). Gen Comp Endocrinol.
Finn RN, Kapoor B. 2008. Fish larval physiology. Didcot: Taylor & Francis.
Galindo D, Sweet E, DeLeon Z, Wagner M, DeLeon A, Carter C, McMenamin S, Cooper WJ. 2018.
Thyroid hormone modulation during zebrafish development recapitulates evolved diversity in
danionin jaw protrusion mechanics. bioRxiv 418483.
Gelman S, Ayali A, Kiemel T, Sanovich E, Cohen A. 2008. Metamorphosis-related changes in the
lateral line system of lampreys, Petromyzon marinus. Journal of Comparative Physiology A
194(11): 945-956.
Giovannotti M, Cerioni PN, La Mesa M, Caputo V. 2007. Molecular phylogeny of the three
paedomorphic Mediterranean gobies (Perciformes: Gobiidae). J Exp Zool B Mol Dev Evol 308(6):
722-729.
Gisbert E. 1999. Early development and allometric growth patterns in Siberian sturgeon and
their ecological significance. J Fish Biol 54(4): 852-862.
Gisbert E, Doroshov SI. 2006. Allometric growth in green sturgeon larvae. J Appl Ichthyol 22:
202-207.
Goodwin D, Bradshaw JW, Wickens SM. 1997. Paedomorphosis affects agonistic visual signals of
domestic dogs. Anim Behav 53(2): 297-304.
Gould SJ. 1977. Ontogeny and phylogeny. Cambridge: The Belknap Press of Harvard University
Press.
Harvey R, Blaxter J, Hoyt R. 1992. Development of superficial and lateral line neuromasts in
larvae and juveniles of plaice (Pleuronectes platessa) and sole (Solea solea). J Mar Biol Assoc UK
72(3): 651-668.

Hastings PA. 2002. Evolution of morphological and behavioral ontogenies in females of a highly
dimorphic clade of blennioid fishes. Evolution 56(8): 1644-1654.
Holzer G, Besson M, Lambert A, François L, Barth P, Gillet B, Hughes S, Piganeau G, Leulier F,
Viriot L. 2017. Fish larval recruitment to reefs is a thyroid hormone-mediated metamorphosis
sensitive to the pesticide chlorpyrifos. eLife 6: e27595.
Holzer G, Laudet V. 2015. Thyroid hormones: a triple-edged sword for life history transitions.
Curr Biol 25(8): R344-R347.
Isorna E, Obregon MJ, Calvo RM, Vázquez R, Pendón C, Falcón J, Muñoz-Cueto JA. 2009.
Iodothyronine deiodinases and thyroid hormone receptors regulation during flatfish (*Solea*
senegalensis) metamorphosis. J Exp Zool B Mol Dev Evol 312(3): 231-246.
Johnson CK, Voss SR. 2013. Salamander paedomorphosis: linking thyroid hormone to life history
and life cycle evolution. Curr Top Dev Biol 103: 229-258.
Johnson DG, Brothers EB. 1993. Schindleria: A Paedomorphic Goby (Teleostei: Aobioidei). Bull
Mar Sci 52(1): 441-471.
Kadobianskyi M, Schulze L, Schuelke M, Judkewitz B. 2019. Hybrid genome assembly and
annotation of *Danionella translucida,* a transparent fish with the smallest known vertebrate
brain. bioRxiv: 539692.
Keer S, Cohen K, May C, Hu Y, McMenamin S, Hernandez LP. 2019. Anatomical assessment of
the adult skeleton of zebrafish reared under different thyroid hormone profiles. Anat Rec.
Kikuyama S, Kawamura K, Tanaka S, Yamamoto K. 1993. Aspects of amphibian metamorphosis:
hormonal control. Int Rev Cytol 145: 105-148.
Laudet V. 2011. The origins and evolution of vertebrate metamorphosis. Curr Biol 21(18): R726-
R737.
Ledent V. 2002. Postembryonic development of the posterior lateral line in zebrafish.
Development 129(3): 597-604.
McCluskey BM, Postlethwait JH. 2014. Phylogeny of zebrafish, a "model species," within Danio,
a "model genus". Mol Biol Evol 32(3): 635-652.
McMenamin S, Carter C, Cooper WJ. 2017. Thyroid Hormone Stimulates the Onset of Adult
Feeding Kinematics in Zebrafish. Zebrafish 14(6): 517-525.
McMenamin S, Chandless M, Parichy D. 2016. Working with zebrafish at postembryonic stages.
Methods Cell Biol 134: 587-607.
McMenamin SK, Bain EJ, McCann AE, Patterson LB, Eom DS, Waller ZP, Hamill JC, Kuhlman JA,
Eisen JS, Parichy DM. 2014. Thyroid hormone–dependent adult pigment cell lineage and
pattern in zebrafish. Science 345(6202): 1358-1361.
McMenamin SK, Hadly EA. 2010. Developmental dynamics of *Ambystoma tigrinum* in a
changing landscape. BMC Ecol 10(1): 10.
McMenamin SK, Parichy DM. 2013. Metamorphosis in teleosts. Curr Top Dev Biol 103: 127-165.
Olsen AM, Westneat MW. 2015. StereoMorph: an R package for the collection of 3D landmarks
and curves using a stereo camera set-up. Methods Ecol Evol 6(3): 351-356.
Osse J, Van den Boogaart J, Van Snik G, Van der Sluys L. 1997. Priorities during early growth of
fish larvae. Aquaculture 155(1-4): 249-258.
Parichy DM. 2006. Evolution of danio pigment pattern development. Heredity 97(3): 200.

Parichy DM, Elizondo MR, Mills MG, Gordon TN, Engeszer RE. 2009. Normal table of
postembryonic zebrafish development: staging by externally visible anatomy of the living fish.
Dev Dyn 238(12): 2975-3015.
Parichy DM, Turner JM. 2003. Zebrafish puma mutant decouples pigment pattern and somatic
metamorphosis. Dev Biol 256(2): 242-257.
Penalva A, Bedke J, Cook ESB, Barrios JP, Bertram EPL, Douglass AD. 2018. Establishment of the
miniature fish species *Danionella translucida* as a genetically and optically tractable
neuroscience model. bioRxiv: 444026.
Raff R. 1996. The Shape of Life: Genes, Development and the Evolution of Animal Form.
Chicago: University of Chicago Press.
Regards E, Taurog A, Nakashima T. 1978. Plasma thyroxine and triiodothyronine levels in
spontaneously metamorphosing *Rana catesbeiana* tadpoles and in adult anuran amphibia.
Endocrinology 102(3): 674-684.
Schneider CA, Rasband WS, Eliceiri KW. 2012. NIH Image to ImageJ: 25 years of image analysis.
Nat Methods 9(7): 671.
Schulze L, Henninger J, Kadobianskyi M, Chaigne T, Faustino AI, Hakiy N, Albadri S, Schuelke M,
Maler L, Del Bene F. 2018. Transparent *Danionella translucida* as a genetically tractable
vertebrate brain model. Nat Methods 15(11): 977.
Shaffer HB, Voss SR. 1996. Phylogenetic and Mechanistic Analysis of A Developmentally
Integrated Character Complex: Alternate Life History Modes in Ambystomatid Salamanders. Am
Zool 36(1): 24-35.
Shea BT. 1983. Paedomorphosis and neoteny in the pygmy chimpanzee. Science 222(4623):
521-522.
Shi Y-B. 2000. Amphibian metamorphosis: Wiley-Liss.
Shkil F, Borisov V, Abdissa B, Smirnov S. 2010. Role of thyroid hormone in the ontogeny and
morphological diversification of Barbus intermedius sensu Banister, 1973 of Lake Tana in
Ethiopia. Russian journal of developmental biology 41(6): 369-380.
Shkil F, Kapitanova D, Borisov V, Abdissa B, Smirnov S. 2012. Thyroid hormone in skeletal
development of cyprinids: effects and morphological consequences. J Appl Ichthyol 28(3): 398-
405.
Smith KK. 2001. Heterochrony revisited: the evolution of developmental sequences. Biol J Linn
Soc 73(2): 169-186.
Vergauwen L, Cavallin JE, Ankley GT, Bars C, Gabriëls IJ, Michiels ED, Fitzpatrick KR, Periz-
Stanacev J, Randolph EC, Robinson SL. 2018. Gene transcription ontogeny of hypothalamic-
pituitary-thyroid axis development in early-life stage fathead minnow and zebrafish. Gen Comp
Endocrinol.
Wahnschaffe U, Bartsch U, Fritzsch B. 1987. Metamorphic changes within the lateral-line
system of Anura. Anatomy and embryology 175(4): 431-442.
Walker M, Kimmel C. 2007. A two-color acid-free cartilage and bone stain for zebrafish larvae.
Biotechnic & Histochemistry 82(1): 23-28.
Webb JF. 2014. Lateral line morphology and development and implications for the ontogeny of
flow sensing in fishes. In: Bleckmann H, Mogdans J ,Coombs SL, editors. Flow Sensing in Air and
Water. Berlin, Springer: 247-270.

Webb JF, Shirey JE. 2003. Postembryonic development of the cranial lateral line canals and
neuromasts in zebrafish. Dev Dyn 228(3): 370-385.
Xu J, Ke Z, Xia J, He F, Bao B. 2016. Change of body height is regulated by thyroid hormone
during metamorphosis in flatfishes and zebrafish. Gen Comp Endocrinol 236: 9-16.
520

FIGURE LEGENDS

Figure 1. Cleared and stained zebrafish and *Danionella*. **A**, wild-type (WT) juvenile zebrafish. **B**,
WT adult zebrafish. **C**, TH- juvenile zebrafish. **D**, TH- adult zebrafish. **E**, adult *D. priapus*. **F**, adult
D. translucida.

Figure 2. Thyroid hormone regulates allometric trajectories in zebrafish. **A-C**, Principle
component (PC) scores of body shape plotted against standard length for both WT (green cross)
and TH- (purple open circle) zebrafish. Percent variation explained by each PC axis is shown in
the upper left corner. Developmental trajectory is shown in solid line with purple standard
error for WT and dashed line with green standard error for TH-. For comparison between
graphs, grey stripes highlight period from 6-6.4 SL, during which TH availability is elevated. **D-F**,
deformation grids showing extreme shapes along each PC. **G-I**, vector plots showing shape
change along each PC axis, with arrows pointing from negative to positive PC values.

Figure 3. Thyroid hormone regulates body proportions in zebrafish. **A**, head length as a function
of SL; **B**, eye diameter as a function of SL. **C**, Head length/SL and **D**, eye diameter/SL as functions
of SL. Showing both WT (green cross) and TH- (purple open circle) zebrafish, with comparison to
D. priapus (orange square) and *D. translucida* (blue triangle) adults. Developmental trajectory is
shown in solid line with purple standard error for WT zebrafish and dashed line with green
standard error for TH- zebrafish. For comparison between graphs, grey stripes highlight period
from 6-6.4 SL, during which TH availability is elevated.

Figure 4. The lateral line system in zebrafish and *Danionella*. **A-B**, DiASP live staining of head
neuromasts in WT juvenile (13 mm SL) and adult (18+ mm SL). Arrowhead indicates rod-shaped
canal neuromasts at adult stage. **C**, Scanning electron micrographs of a supraorbital canal
neuromast in WT adult. Dotted line outlines the neuromast; solid line outlines the bony edge of
the canal. **D**, head neuromasts in *D. priapus* adult. **E-G**, same as B-D in TH- zebrafish. **H**, facial
neuromasts in *D. translucida* adult. **I-N**, DiASP live staining of trunk neuromasts in WT juvenile
zebrafish (13 mm SL, **I**), WT adult (18+ mm SL, **J**), *D. priapus* adult (**K**), TH- juvenile (13 mm SL, **L**)
TH- adult (18+ mm SL, **M**), and *D. translucida* adult (**N**).

Figure 5. Thyroid hormone regulates lateral line development in zebrafish. **A**, number of cranial
neuromasts, as a function of SL, with comparison to *D. priapus* (orange square) and *D.*
translucida (blue triangle) adults. For zebrafish, developmental trajectory is shown in solid line
for WT and dashed line for TH-. Developmental trajectory is shown in solid line with purple
standard error for WT zebrafish and dashed line with green standard error for TH- zebrafish. For
comparison between graphs, grey stripes highlight period from 6-6.4 SL, during which TH
availability is elevated. **B-E**, same as A for number of trunk neuromasts, total number of

neuromasts (head + trunk), cranial neuromasts divided by head length, and trunk neuromasts
divided by trunk length.

Figure 6. Relative thyroid hormone content during larval development in zebrafish. **A**, T4,
thyroxine; **B**, T3, triiodothyronine; **C**, ratio of T3 to T4. All concentrations standardized to body
weight. Whiskers represents standard error at each stage. Stages that are not statistically
different (as determined by Tukey's Honest Significant Differences Test, p value cut-off was set
at 0.05) are represented with the same letter. For comparison with other figures, grey stripe
shows period from 6-6.4 SSL, during which TH availability is elevated. **D-F**: Antibody staining of
T4 in zebrafish, *D. priapus* and *D. translucida* adults showing functional thyroid follicles. Images
show 12 uM transverse sections of the thyroid region.

Figure 7. *Danionella* adults resemble larval zebrafish but not TH- zebrafish. **A-C**, Principle
component (PC) scores of body shape plotted against standard length for both WT (green cross)
and TH- (purple open circle) zebrafish, compared to *D. priapus* (orange square) and *D.*
translucida (blue triangle) adults. Percent variation explained by each PC axis is shown on the
upper left corner. Developmental trajectory is shown in solid line with purple standard error for
WT zebrafish and dashed line with green standard error for TH- zebrafish. For comparison
between graphs, grey stripes highlight period from 6-6.4 SL, during which TH availability is
elevated. **D-F**, deformation grids showing extreme shapes along each PC. **G-I**, vector plots
showing shape change along each PC axis, with arrows pointing from negative to positive PC
values.

Supplementary Table S1. Landmarks used in morphometric body shape analysis.

Landmarks	
1	Tip of snout/rostral edge of premaxilla
2	Posterior tip of supraoccipital crest
3	Dorsal insertion of the caudal fin
4	Ventral insertion of the caudal fin
5	Point of maximum curvature on the most ventral branchiolstegal ray
Semi-landmarks	
	ventral outline of the head connecting Landmark #1 and #5, n=8
	dorsal outline of the head connecting Landmark #1 and #2, n=8
	dorsal edge of the body connecting Landmark #2 and #3, n=18

589 **Supplementary Table S2.** Summary of fish samples used for ELISA.

SSL	Stage	SL (mm)	Age (± SD)	n	average weight (g) per fish
4.9	CR	4.9-5.1	12 ± 2.9	1318	0.000893649

6	aSB	5.9-6.2	13.5 ± 2.5	654	0.001709021
6.4	DR	6.4-6.6	15 ± 2.5	403	0.003138213
7.2	PB	7.2-7.5	16 ± 2.7	294	0.004842177
8.6	PR	8.5-8.7	20 ± 1.3	149	0.008157047
9.8	SP	9.5-9.6	26.5 ± 2.1	98	0.013067347
11	J	10.9-11.7	32 ± 0.5	58	0.022691379

A — WT

1mm

B — WT

C — TH-

D — TH-

E — *D. priapus*

F — *D. translucida*