Field application of an eDNA assay for the threatened white-clawed crayfish

Austropotamobius pallipes

Siobhán Atkinson[1,2]*, Jeanette E.L. Carlsson[2], Bernard Ball[2], Mary Kelly-Quinn[1], Jens Carlsson[2]

1. School of Biology and Environmental Science/Earth Institute, University College

Dublin, Dublin, Ireland

2. Area52 Research Group, School of Biology and Environmental Science/Earth Institute,

University College Dublin, Dublin, Ireland

*Corresponding author: Siobhán Atkinson, siobhan.atkinson@ucdconnect.ie

10   **Abstract**

The white-clawed crayfish *Austropotamobius pallipes* has undergone extensive declines

within its native range in the last century. Because of its threatened status, European

legislation requires the species to be regularly monitored and that Special Areas of

Conservation (SACs) be designated for it. Knowledge of the distribution of this species is vital

for addressing these needs. This study presents an environmental (e)DNA assay to detect *A.*

pallipes in water samples, based on the mitochondrial cytochrome oxidase I (COI) gene,

utilizing species-specific primers, a minor groove binding (MGB) probe and quantitative PCR.

The results of this study indicate that eDNA is an effective tool for detecting *A. pallipes* in a

lotic system, and could provide a valuable, non-invasive method for determining the

distribution of this species.

Keywords: eDNA, qPCR, conservation, non-invasive, detection, native species, river.

22   **Introduction**

The white-clawed crayfish *Austropotamobius pallipes* is a relatively large, long-lived (>10

years) crustacean that inhabits both rivers and lakes (Reynolds et al. 2010). It requires

alkaline conditions for survival and is commonly found in waterbodies overlying

carboniferous limestone bedrock (Lucey and McGarrigle 1987). *A. pallipe*s is one of the five

indigenous crayfish species in Europe (Holdich et al. 2009). This once abundant species has,

however, become greatly reduced or locally extinct across large parts of its native range

during the last century (Grandjean et al. 1997). Pollution (Demers and Reynolds 2002, Lyons

and Kelly-Quinn 2003), habitat loss and disease (Matthews and Reynolds 1992) have

contribution to this decline. Of particular concern is the crayfish plague, caused by the

fungus *Aphanomyces astaci* (Holdich et al. 2009). *A. pallipes* possesses no resistance to this

fungus, so eradication of entire populations of *A. pallipes* is possible following an outbreak

(Reynolds et al. 2010). Despite this, Ireland is considered one of the few remaining

strongholds for *A. pallipes* within Europe. One of the reasons for this is that the invasive

signal crayfish *Pacifastacus leniusculus,* which can impose wide-ranging impacts on

ecosystems and act as a vector for the crayfish plague (Vaeßen and Hollert 2015), has not

been reported in Ireland to date. However, crayfish plague has reached Ireland's rivers, and

large changes in the distribution of *A. pallipes* have been attributed, at least in part, to past

plague outbreaks (Demers et al. 2005). Additionally, there is at present an outbreak of

crayfish plague in Ireland (National Biodiversity Data Centre 2018), and hence there is a

pressing need for a non-invasive, rapid and cost-effective sampling method that can be used

to establish the distribution of the species.

Because of its vital role in freshwater ecosystem functioning and its threatened status

(Matthews and Reynolds 1992), *A. pallipes* is afforded protection under the EU Habitats

Directive and is listed under Annexes II and V. This means Ireland is required to regularly

monitor *A. pallipes*, and to designate Special Areas of Conservation (SACs) for the species

under Natura 2000 (Reynolds et al. 2010). Knowledge of the geographic distribution of *A.*

pallipes is essential for implementing these conservation measures. Traditional survey

methods include night viewing with a strong torch, modified quadrat samplers (DiStefano et

al. 2003), kick-sampling, baited traps and enclosures (Byrne and Lynch 1999), snorkeling

surveys (Reynolds et al. 2010) and SCUBA diving (Matthews and Reynolds 1992). While many

of these methods are effective, they are time-consuming, potentially costly, and may not be

effective when *A. pallipes* occur in low abundance. Furthermore, a license is required for

most traditional *A. pallipes* surveying methods. This is vital for ensuring the protection of *A.*

pallipes, however, a non-invasive methodology that does not require a license for assessing

its distribution would be beneficial. A promising new tool that has been shown to be both

effective at detecting species and to save significant time in the field (Sigsgaard et al. 2015) is

environmental DNA (eDNA) analysis. Environmental DNA is the collective term for DNA

present freely in the environment that has been shed by organisms for example in the form

of mucus, excrement, gametes, or blood (Taberlet et al. 2012, Thomsen and Willerslev

2015). The analysis has been successful in detecting species that occur in low abundance

such as rare or endangered species (Mächler et al. 2014, Laramie et al. 2015, Sigsgaard et al.

2015, Boothroyd et al. 2016, Carlsson et al. 2017), or invasive alien species (Jerde et al. 2011,

Goldberg et al. 2013, Fujiwara et al. 2016). Numerous studies have also shown eDNA to be

more sensitive than traditional methods at detecting species at low abundance (Dejean et al.

2012, Smart et al. 2015, Dougherty et al. 2016). Environmental DNA assays have been

successfully deployed for other crayfish species including *Astacus astacus*, *Astacus*

leptodactylus, *P. leniusculus* (Agersnap et al. 2017, Larson et al. 2017), *Orconectes rusticus*

(Dougherty et al. 2016, Larson et al. 2017), *Procambarus clarkii* (Cai et al. 2017) and *Faxonius*

eupunctus (Rice et al. 2018).

The aim of this study was to develop an MGB based qPCR assay to detect the presence of

the white-clawed crayfish, *A. pallipes* in water samples, and to test the reliability of the assay

by comparing the results with field observation data.

Methods

Study sites and crayfish distribution data

Eight sampling locations within seven different rivers were selected for field validation of the

assay (Table 1). The research presented here was carried out as part of a larger study

assessing the impact of river obstacles on freshwater fauna (e.g. Atlantic salmon, *Salmo*

salar, Atkinson et al. 2018). As part of this study, eDNA samples were collected above and

below a number of river obstacles (weir, ford crossing or bridge apron) in order to determine

whether they imposed migratory barriers on *S. salar*. The same samples collected for this *S.*

salar study were used for the present study. Out of the eight sites sampled, four were in

rivers with SAC status in which *A. pallipes* featured as a species of interest. Study rivers

within the River Barrow and River Nore SAC included the Delour and Dinin rivers and study

rivers within the Lower River Suir SAC included the Multeen and Duag rivers. Although the

Burren River is a tributary of the River Barrow, it does not have SAC status.

The current distribution of *A. pallipes* in these rivers was extrapolated using data provided

by the Irish Environmental Protection Agency (EPA). These data are point data based on

casual field observations made of *A. pallipes* by EPA staff whilst undertaking routine

biological monitoring between the years 2007 and 2016 (W. Trodd, Irish Environmental

Protection Agency, personal communication). In addition, it was possible at some sites to

confirm the presence of crayfish by field observations made while electrofishing for *S. salar*.

For the purposes of this study, *A. pallipes* observations were extrapolated to the wider

subcatchment level. This meant that even if only one field observation of *A. pallipes*

occurred in a subcatchment, it was assumed that there were likely to be *A. pallipes* in the

rest of that subcatchment. This was considered the least ambigious way to present the data,

because the EPA were not specifically looking for *A. pallipes* when their observations were

made, so there is no certainty that *A. pallipes* did not occur in other parts of the

subcatchment.

eDNA qPCR assay development

Species-specific primers (forward primer: 5'- GGG TTA GTG GAG AGA GGG GT -3', and

reverse primer 5'- AAT CCC CAG ATC CAC AGA CG -3') and a 5'-6-FAM labelled TaqMan®

minor groove binding probe (5'- TCA GCT ATT GCC CAC GCA -3') for *A. pallipes*, which

targeted a locus within the mitochondrial cytochrome oxidase I (COI) region, were designed

in Primer Express 3.0 (Applied Biosystems-Roche, Branchburg, NJ) using the available Irish *A.*

pallipes COI sequences in the National Centre for Biotechnology Information (NCBI -

http://www.ncbi.nlm.nih.gov/). Including primers, the total amplicon size was 96 base pairs.

To verify the species specificity for the *A. pallipes* assay in silico, probe and primer sequences

were combined and queried against the National Centre for Biotechnology Information

(NCBI - http://www.ncbi.nlm.nih.gov/) nucleotide database with BLASTn (Basic Local

Alignment Search Tool). The BLASTn analyses showed 90% sequence identity for 100% of the

assay query only for the target species. The reason for not achieving 100% sequence identity

was due to gaps between the forward and reverse primers and the probe. Non-target

species showed a maximum of 37% query cover with 100% sequence identity indicating that

only the target species *A. pallipes* would bind to all three components of the assay (forward,

reverse primers and probe). To verify the species specificity of the assay in vitro, the assay

was tested with DNA extracted from the tissue of some freshwater species co-occuring with

A. pallipes, including brown trout *S. trutta*, sea lamprey *Petromyzon marinus*, Atlantic

salmon *S. salar*, allis shad *Alsoa alosa* and twaite shad *Alosa fallax,* by using conventional

PCR followed by visualization of amplicons on agarose. The qPCR assay was optimized using

tissue extracted from *A. pallipes*.

eDNA collection, filtering and extraction

Water samples for eDNA analysis were collected in PET bottles from two locations at each

study site, above and below the river obstacle. The PET bottles were sterilized before use by

soaking them for 30 minutes in 50% sodium hypochlorite solution. Prior to collecting the

water samples, the bottles were rinsed at least five times to remove any remaining sodium

hypochlorite. Water samples were filtered through disposable filter funnels on site using a

peristaltic pump. The total volume filtered was 2 liters. To avoid contamination, water

samples were taken in an upstream direction, prior to any individuals entering the river. At

each sampling location, three replicate water samples were collected, one from each of the

left margin, right margin and the center of the river channel. In addition, one negative field

control consisting of ddH_2O was also filtered. This resulted in a total number of six water

samples for eDNA analysis and two field controls collected per study site. All water samples

were filtered through 47 mm glass microfiber filters (1.5 μm). Filters were placed in 2.0 mL

Eppendorf tubes and subsequently stored in a cooler box on ice. The filters were frozen at -

20˚C in the laboratory approximately 3-6 hours following collection. River flow conditions

were low at each study site at the time of sampling. To reduce contamination risk, all eDNA

work was performed in a dedicated Low Copy DNA laboratory. A modified CTAB

(cetyltrimethylammonium bromide) protocol (Möller et al. 1992) was employed for eDNA

extractions. Briefly, one-half of each filter was transferred to a 2.0 mL Eppendorf tube (the

other half of each filter was archived to allow for future validation or queries). A total

volume of 750 μL CTAB buffer (100 mM Tris-HCL, 20 mM EDTA, 1.4 M NaCl, 2% CTAB), and 7

μL of Proteinase K (20 mg mL^{-1}) were added to the tube. Samples were then vortexed for 10

seconds followed by incubation at 56˚C for 2 hours, after which 750 μL of

phenol/chloroform/isoamyl alcohol (25:25:1 v/v) was added. The contents of the tube were

manually mixed for 15 seconds and subsequently centrifuged (11,000 x g, 20 min). A new

tube contained 750 μL of chloroform/isoamyl alcohol (24:1 v/v) was prepared, and the

aqueous phase was transferred to it. The manual mixing followed by centrifugation steps

were repeated, and the aqueous phase was again transferred to a new tube. The DNA was

precipitated using one volume of isopropanol alcohol that was added to the aqueous phase

and incubated at -20˚C for 1 hour followed by centrifugation (11,000 x g, 20 min). The

resulting pellets were washed with 750 μL of 70% ethanol and centrifuged (11,000 x g, 5

min). The ethanol was removed, and care was taken to ensure that the pellet remained in

the tube. The tubes containing pellets were dried in a heat block (50˚C, 5 min) followed by

resuspension in 40 μL of molecular-grade ddH$_2$O. Environmental DNA concentrations were

determined with NanoDrop®-1000, Thermo Scientific, Wilmington, DE.

eDNA assay deployment

An Applied Biosystems ViiA™ 7 (Life Technologies, Inc., Applied Biosystems, Foster City,

California, U.S.A.) quantitative thermocycler was used for eDNA amplification and

quantification. Briefly, the PCR profile consisted of 50°C for 2 min and 95°C for 10 min,

followed by 40 cycles between 95°C for 15 s and 60°C for 1 min. Standard curves were

generated from DNA originating from *A. pallipes* tissue extracts (quantified with NanoDrop®-

1000, Thermo Scientific, Wilmington, DE). This served as both a positive control and as a

means of calculating the concentration of *A. pallipes* eDNA in each sample. Seven 10:1 serial

dilutions ranging from 2.9 ng μL^{-1} to 2.9 x 10^{-6} ng μL^{-1} were used for the standard curve. Each

qPCR reaction was carried out in a total volume of 30 μL, consisting of 15 μL of TaqMan®

Environmental Master Mix 2.0 (Life Technologies, Applied Biosystems, Foster City, CA), 3 μL

of each primer (final concentration of 2 μM), 3 μL probe (final concentration of 2 μM), 3 μL

eDNA/tissue extracted DNA template and 3 μL ddH$_2$O. Individual standard curves were

generated for each qPCR plate (y = -3.37x + 20.589, efficiency = 98.04 %, R^2 = 0.996 (1), y = -

3.442x + 20.392, efficiency = 95.22 %, R^2 = 0.999 (2) and y = -3.441x + 20.723, efficiency =

95.26 %, R^2 = 0.999 (3)). Each plate incorporated 2 no-template controls (NTCs) to identify

any contamination that may have occurred during preparation. Standard curve, field and

control samples were quantified in triplicate (3 technical replicates). A positive detection

was defined as being when at least 2 out of the 3 technical replicates contained amplifiable

DNA with quantitation cycle (Cq) differences not exceeding 0.5 (to allow for pipetting

errors), and when the samples were within the detection range of the standard curve. If 1

out of 3 technical replicates showed Cq differences exceeding 0.5, the replicate was

excluded from the study. In cases when the Cq value of 2 out of 3 technical replicates

differed by more than 0.5 Cq, that specific dilution series or field replicate was removed

from further study. As *S. trutta* was present in all rivers, both upstream and downstream of

the obstacles, this species was used as a positive field control to test for the presence of

amplifiable DNA in all samples. The *S. trutta* assay from previously published work

(Gustavson et al. 2015) was used for this analysis. Three replicates per location with 1

technical replicate were used.

187   **Results and Discussion**

The present study was successful in detecting *A. pallipes* in silico, in vitro and in situ. The

assay was highly sensitive and could detect *A. pallipes* eDNA concentrations as low as 0.002

ng L^{-1} at Cq 38.2 (average over 3 technical replicates, standard deviation 0.000059 ng L^{-1}). No

cross-species amplification occurred. Furthermore, zero amplification occurred in the NTCs

and field controls. All field samples analysed yielded detectable DNA with the positive field

control (*S. trutta* assay), however, this was not the case for the *A. pallipes* assay. *A. pallipes*

was only detected in 5 out of 8 sites (Table 2, Fig. 1). Within the Burren, Duag and Multeen

rivers, positive detections were observed in all field replicates above and below the river

obstacles. At the Dinin (Crettyard) site, positive detection was observed in 2 out of 3 field

replicates above the obstacle, and 3 out of 3 field replicates below the obstacle. At the Dinin

(Castlecomer) site, positive detection of *A. pallipes* occurred in only 1 field replicate, which

was located above the obstacle (Table 2). The detection of *A. pallipes* eDNA above the

obstacle but not below it was a surprising result. Intuitively, one would expect that *A.*

pallipes would have been detected below the obstacle also, as eDNA travels downstream in

a river system. It is worth noting that it was at this site that the lowest concentration of *A.*

pallipes eDNA was detected (0.002 ng L^{-1}), and amplification of eDNA occurred in only 1 out

of 6 site replicates. It is possible that eDNA concentrations in the other field samples were

too low for a positive detection, or that false negative results were obtained. The density or

biomass of *A. pallipes*, environmental conditions such as flow, temperature, pH and

sediment, or natural inhibitors such as algae or humic substances, may have contributed to

this lack of amplification (Strickler et al. 2015, Goldberg et al. 2016, Stoeckle et al. 2017).

However, as all eDNA samples yielded detectable DNA with the positive field control (e.g. *S.*

trutta), it is unlikely that in this instance the lack of amplification of *A. pallipes* DNA was a

result of inhibition of the eDNA samples.

When compared with the EPA's *A. pallipes* observation data, the eDNA results reflected

what was observed in the field. In all subcatchments where *A. pallipes* was observed by EPA

staff, the eDNA results revealed a positive detection, and in all subcatchments where *A.*

pallipes was not observed by EPA staff, the eDNA results revealed a negative detection (Fig.

1). Furthermore, it is worth noting that the eDNA results also reflected the expected

distribution of *A. pallipes* based on environmental conditions. In all sites where *A. pallipes*

was not detected by either field observation or eDNA (Delour, Brown's Beck Brook and the

Dalligan), the dominant bedrock was siliceous. *A. pallipes* are known to occur in waterbodies

with underlying calcareous bedrock rather than siliceous bedrock (Lucey and McGarrigle

1987).

It was not possible in this study to discern the impact of the river obstacles on *A. pallipes*

using eDNA. This was because in most sites, *A. pallipes* was either detected both above and

below the obstacle, or as observed at the Dinin (Castlecomer) site, above the obstacle only.

However, studies where crayfish have been physically marked or tagged have shown that

river obstacles can hinder their upstream migration (Kerby et al. 2005, Bubb et al. 2008,

Rosewarne et al. 2013). Environmental DNA assays for diadromous species such as sea

lamprey *P. marinus* (Gustavson et al. 2015), Atlantic salmon *S. salar* (Atkinson et al. 2018) or

Chinook salmon *Oncorhynchus tshawytscha* (Laramie et al. 2015) are likely to be better

indicators for assessing the migratory impact of river obstacles. Alternatively, eDNA could be

used for monitoring whether invasive species have moved above the barriers designed to

block them (Cowart et al. 2018). Nonetheless, the present research shows that eDNA is a

reliable tool for detecting *A. pallipes* in river systems. Furthermore, while the focus of this

study was in lotic systems, the assay could be readily deployed in lentic systems such as

lakes or reservoirs where field sampling may be more demanding. For example, as noted by

Reynolds et al. (2010), different survey methods for monitoring *A. pallipes* can vary in their

efficiency depending on the habitat type being surveyed. Environmental DNA monitoring

may therefore lend itself to surveying more challenging habitats such as heavily vegetated

lake margins with soft substrates, or locations where trapping or night searching is not

possible or particularly difficult due to health and safety considerations. Environmental DNA

sampling requires one site visit, which involves simply collecting water samples, and filtering

them onsite. This sampling approach is considerably less labor intensive when compared

with traditional methods. Night searches, for example, typically require two site visits (for

health and safety purposes) and are dependent on water clarity and algal growth (Reynolds

et al. 2010). Trapping is time-consuming and requires two site visits, and snorkeling and

hand searches are habitat dependent. Snorkeling is also time-consuming as it requires time

to allow for disinfection and drying out of snorkeling gear between entering different

waterbodies (Reynolds et al. 2010). Relative to traditional sampling methods, eDNA sampling

requires substantially less field equipment. Herder et al. (2014) suggested that the eDNA

method may help reduce the risk of unintentional transmission of pathogens into new areas

due to the numerous contamination prevention precautions that must be taken when

sampling. Considering this, and the continued threat of crayfish plague in Ireland and

elsewhere, eDNA sampling may be an appropriate sampling technique to employ to

minimize the risk of further spread of crayfish plague by reducing the amount of field

equipment required, and therefore potential vectors of crayfish plague. This is an important

consideration, especially if surveys are being conducted in waterbodies harboring crayfish

plague.

To conclude, this assay provides an alternative, rapid method for locating *A. pallipes* in both

lentic and lotic ecosystems, allowing for more targeted, efficient surveying strategies.

Considering the current crayfish plague outbreak in Ireland (National Biodiversity Data

Centre 2018), and indeed other countries where it is present, there is an urgent need for a

non-invasive sampling methodology for the species. Large scale deployment of this assay

would enable managers to rapidly establish, with confidence, the distribution of *A. pallipes*,

which would support the designation of new conservation areas.

Acknowledgements

This research was funded by the Irish Environmental Protection Agency (Reconnect project -

2015-W-LS-8). The authors would also like to thank the Irish Environmental Protection

Agency for providing the data on *A. pallipes* distributions.

Literature Cited

Agersnap, S., W. B. Larsen, S. W. Knudsen, D. Strand, P. F. Thomsen, M. Hesselsøe, P. B.

Mortensen, T. Vrålstad, and P. R. Møller. 2017. Monitoring of noble, signal and narrow-

clawed crayfish using environmental DNA from freshwater samples. PLoS ONE 12:1–22.

Atkinson, S., J.E.L. Carlsson, B. Ball, D. Egan, M. Kelly-Quinn, K. Whelan, and J. Carlsson. 2018.

A quantitative PCR-based environmental DNA assay for detecting Atlantic salmon (*Salmo*

salar L.). Aquatic Conservation: Marine and Freshwater Ecosystems doi: 10.1002/aqc.2931.

Boothroyd, M., N. E. Mandrak, M. Fox, and C. C. Wilson. 2016. Environmental DNA (eDNA)

detection and habitat occupancy of threatened spotted gar (*Lepisosteus oculatus*). Aquatic

Conservation: Marine and Freshwater Ecosystems 26:1107–1119.

Bubb, D. H., T. J. Thom, and M. C. Lucas. 2008. Spatial ecology of the white-clawed crayfish in

an upland stream and implications for the conservation of this endangered species. Aquatic

Conservation: Marine and Freshwater Ecosystems 18:647–657.

Byrne, C., and J. Lynch. 1999. A sampling strategy for stream populations of white-clawed

crayfish, *Austropotamobius pallipes* (Lereboullet)(Crustacea, Astacidae). Biology and

Environment: Proceedings of the Royal Irish Academy 99b:89–94.

Cai, W., Z. Ma, C. Yang, L. Wang, W. Wang, G. Zhao, Y. Geng, and D. W. Yu. 2017. Using eDNA

to detect the distribution and density of invasive crayfish in the HongheHani rice terrace

World Heritage site. PLoS ONE 12:1–13.

Carlsson, J. E. L., D. Egan, P. C. Collins, E. D. Farrell, F. Igoe, and J. Carlsson. 2017. A qPCR

MGB probe based eDNA assay for European freshwater pearl mussel (*Margaritifera*

margaritifera L.). Aquatic Conservation: Marine and Freshwater Ecosystems 27:1341-1344.

Cowart, D. A., K. G. H. Breedveld, M. J. Ellis, J. M. Hull, and E. R. Larson. 2018. Environmental

DNA (eDNA) applications for the conservation of imperiled crayfish (Decapoda: Astacidea)

through monitoring of invasive species barriers and relocated populations. Journal of

Crustacean Biology doi:10.1093/jcbiol/ruy007.

Dejean, T., A. Valentini, C. Miquel, P. Taberlet, E. Bellemain and C. Miaud. 2012. Improved

detection of an alien invasive species through environmental DNA barcoding: the example of

the American bullfrog *Lithobates catesbeianus*. Journal of Applied Ecology 49:953-959.

Demers, A., J. Lucey, M. L. McGarrigle, and J. D. Reynolds. 2005. The distribution of the

white-clawed crayfish, *Austropotamobius pallipes*, in Ireland. Biology & Environment:

Proceedings of the Royal Irish Academy 105b:65–69.

Demers, A., and J. D. Reynolds. 2002. A survey of the white-clawed crayfish,

Austropotamobius pallipes (Lereboullet), and of water quality in two catchments of eastern

Ireland. Bulletin Francais de Pêche et de Pisciculture 367:729–740.

DiStefano, R. J., C. M. Gale, B. A. Wagner, and R. D. Zweifel. 2003. A sampling method to

assess lotic crayfish communities. Journal of Crustacean Biology 23:678–690.

Dougherty, M. M., E. R. Larson, M. A. Renshaw, C. A. Gantz, S. P. Egan, D. M. Erickson, and D.

M. Lodge. 2016. Environmental DNA (eDNA) detects the invasive rusty crayfish *Orconectes*

rusticus at low abundances. Journal of Applied Ecology 53:722–732.

Fujiwara, A., S. Matsuhashi, H. Doi, S. Yamamoto, and T. Minamoto. 2016. Use of

environmental DNA to survey the distribution of an invasive submerged plant in ponds.

Freshwater Science 35:748–754.

Goldberg, C. S., A. Sepulveda, A. Ray, J. Baumgardt, and L. P. Waits. 2013. Environmental

DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus*

antipodarum). Freshwater Science 32:792–800.

Goldberg, C. S., C. R. Turner, K. Deiner, K. E. Klymus, P. F. Thomsen, M. A. Murphy, S. F.

Spear, A. McKee, S. J. Oyler-McCance, R. S. Cornman, M. B. Laramie, A. R. Mahon, R. F.

Lance, D. S. Pilliod, K. M. Strickler, L. P. Waits, A. K. Fremier, T. Takahara, J. E. Herder and P.

Taberlet. 2016. Critical considerations for the application of environmental DNA methods to

detect aquatic species. Methods in Ecology and Evolution 7: 1299-1307.

Grandjean, F., C. Souty-Grosset, R. Raimond, and D. M. Holdich. 1997. Geographical variation

of mitochondrial DNA between populations of the white-clawed crayfish *Austropotamobius*

pallipes. Freshwater Biology 37:493–501.

Gustavson, M. S., P. C. Collins, J. A. Finarelli, D. Egan, R. Ó. Conchúir, G. D. Wightman, J. J.

King, D. T. Gauthier, K. Whelan, J. E. L. Carlsson, and J. Carlsson. 2015. An eDNA assay for

Irish *Petromyzon marinus* and *Salmo trutta* and field validation in running water. Journal of

Fish Biology 87:1254–1262.

Herder, J. E., A. Valentini, E. Bellemain, T. Dejean, J. J. C. W. van Delft, P. F. Thomsen, and P.

Taberlet. 2014. Environmental DNA—a review of the possible applications for the detection

of (invasive) species. Report no. 2013–104. Stichting RAVON, Nijmegen.

Holdich, D. M., J. D. Reynolds, C. Souty-Grosset, and P. J. Sibley. 2009. A review of the ever

increasing threat to European crayfish from non-indigenous crayfish species. Knowledge and

Management of Aquatic Ecosystems 394-395:11.

Jerde, C. L., A. R. Mahon, W. L. Chadderton, and D. M. Lodge. 2011. "Sight-unseen" detection

of rare aquatic species using environmental DNA. Conservation Letters 4:150–157.

Kerby, J. L., S. P. D. Riley, L. B. Kats, and P. Wilson. 2005. Barriers and flow as limiting factors

in the spread of an invasive crayfish (*Procambarus clarkii*) in southern California streams.

Biological Conservation 126:402–409.

Laramie, M. B., D. S. Pilliod, and C. S. Goldberg. 2015. Characterizing the distribution of an

endangered salmonid using environmental DNA analysis. Biological Conservation 183:29–37.

Larson, E. R., M. A. Renshaw, C. A. Gantz, J. Umek, S. Chandra, D. M. Lodge, and S. P. Egan.

2017. Environmental DNA (eDNA) detects the invasive crayfishes *Orconectes rusticus* and

Pacifastacus leniusculus in large lakes of North America. Hydrobiologia 800:173–185.

Lucey, J., and M. L. McGarrigle. 1987. The distribution of the crayfish *Austropotamobius*

pallipes (Lereboullet) in Ireland. Irish Fisheries Investigations A29:1–13.

Lyons, R., and M. Kelly-Quinn. 2003. An investigation into the disappearance of

Austropotamobius pallipes (Lereboullet) populations in the headwaters of the Nore River,

Ireland and the correlation to water quality. Bulletin Français de la Pêche et de la Pisciculture

370-371:139–150.

Mächler, E., K. Deiner, P. Steinmann, and F. Altermatt. 2014. Utility of environmental DNA

for monitoring rare and indicator macroinvertebrate species. Freshwater Science 33:1174–

1183.

Matthews, M., and J. D. Reynolds. 1992. Ecological impact of crayfish plague in Ireland.

Hydrobiologia 234:1–6.

Möller, E. M., G. Bahnweg, H. Sandermann, and H. H. Geiger. 1992. A simple and efficient

protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and

infected plant tissues. Nucleic Acids Research 20:6115–6116.

National Biodiversity Data Centre. 2018. Crayfish plague. (Available from:

http://www.biodiversityireland.ie/projects/invasive-species/crayfish-plague/)

Reynolds, J. D., W. O'Connor, C. O'Keeffe, and D. Lynn. 2010. A technical manual for

monitoring white-clawed crayfish *Austropotamobius pallipes* in Irish lakes. (Available at:

https://www.npws.ie/sites/default/files/publications/pdf/IWM45.pdf)

Rice, C. J., E. R. Larson, and C. A. Taylor. 2018. Environmental DNA detects a rare large river

crayfish but with little relation to local abundance. Freshwater Biology 63:443–455.

Rosewarne, P. J., A. T. Piper, R. M. Wright, and A. M. Dunn. 2013. Do low-head riverine

structures hinder the spread of invasive crayfish? Case study of signal crayfish (*Pacifastacus*

leniusculus) movements at a flow gauging weir. Management of Biological Invasions 4:273–

282.

Sigsgaard, E. E., H. Carl, P. R. Møller, and P. F. Thomsen. 2015. Monitoring the near-extinct

European weather loach in Denmark based on environmental DNA from water samples.

Biological Conservation 183:46–52.

Smart, A.S., R. Tingley, A. R. Weeks, A. R. van Rooyen, and M. A. McCarthy. 2015.

Environmental DNA sampling is more sensitive than a traditional survey technique for

detecting an aquatic invader. Ecological Applications 25:1944-1952.

Stoeckle, B. C., S. Beggel, A. F. Cerwenka, E. Motivans, R. Kuehn and J. Geist. 2017. A

systematic approach to evaluate the influence of environmental conditions on eDNA

detection success in aquatic ecosystems. PloS one 12: e0189119.

Strickler, K.M., A. K. Fremier and C.S. Goldberg. 2015. Quantifying effects of UV-B,

temperature, and pH on eDNA degradation in aquatic microcosms. Biological Conservation

183: 85-92.

Taberlet, P., E. Coissac, M. Hajibabaei, and L. H. Rieseberg. 2012. Environmental DNA.

Molecular Ecology 21:1789–1793.

Thomsen, P. F., and E. Willerslev. 2015. Environmental DNA - An emerging tool in

conservation for monitoring past and present biodiversity. Biological Conservation 183:4–18.

Vaeßen, S., and H. Hollert. 2015. Impacts of the North American signal crayfish (*Pacifastacus*

leniusculus) on European ecosystems. Environmental Sciences Europe 27:1–6.

Table 1. Summary of sampling information.

River	Obstacle Type	Sampling Location	Distance from Obstacle (m)	Latitude	Longitude	Sampling Date
Burren	Weir	Upstream	1	52.83300	-6.92534	18/07/2017
		Downstream	215	52.83388	-6.92822	18/07/2017
Delour	Weir	Upstream	193	52.97709	-7.54606	19/07/2017
		Downstream	267	52.97957	-7.54100	19/07/2017
Nore (Castlecomer)	Weir	Upstream	11	52.80599	-7.20527	20/07/2017
		Downstream	57	52.80535	-7.20558	20/07/2017
Duag	Weir	Upstream	8	52.27196	-8.01319	21/07/2017
		Downstream	45	52.27174	-8.01248	21/07/2017
BBB	Ford	Upstream	1	53.01627	-6.62311	24/07/2017
		Downstream	60	53.01573	-6.62322	24/07/2017
Dalligan	Weir	Upstream	15	52.11876	-7.51938	25/07/2017

		Downstream	92	52.11862	-7.51803	25/07/2017
Muilteen	Bridge	Upstream	101	52.61304	-8.00872	27/07/2017
	Apron					
		Downstream	34	52.61182	-8.00966	27/07/2017
Nore	Bridge	Upstream	8	52.84534	-7.12758	28/07/2017
(Crettyard)	Apron					
		Downstream	68	52.84584	-7.12891	28/07/2017

388 Table 2. Environmental DNA concentrations (ng L^{-1}) from the *A. pallipes* assay at each study

389 river. Concentrations are based the average eDNA concentration in a minimum of 2 technical

390 replicates and between 1 and 3 field replicates. Confidence intervals (95%) are given, where

391 possible, for each location (upstream or downstream of the river obstacle).

River	Location	No. eDNA samples collected	No. of positive detections	eDNA concentration ng L^{-1} (95% CI)	Visual observation
Burren	Upstream	3	3	0.027 (0.01-0.02)	+
	Downstream	3	3	0.02 (0.01-0.04)	
Duag	Upstream	3	3	0.027 (0.005-0.05)	+
	Downstream	3	3	0.157 (0.09-0.23)	
Multeen	Upstream	3	3	0.034 (0.02-0.04)	+
	Downstream	3	3	0.06 (0.04-0.08)	
Dinin (Crettyard)	Upstream	3	2	0.011 (0.004-0.02)	+
	Downstream	3	3	0.006 (0.002-0.01)	
Dinin (Castlecomer)	Upstream	3	1	0.002 [a]	+
	Downstream	3	0	0	
Delour	Upstream	3	0	0	-
	Downstream	3	0	0	
Dalligan	Upstream	3	0	0	-
	Downstream	3	0	0	
Brown's Beck Brook (BBB)	Upstream	3	0	0	-
	Downstream	3	0	0	

[a] It was not possible to calculate a 95% confidence interval for this site as a positive detection was

only detected in one out of three site replicates

Figure Heading

Figure 1. Map showing the sampling locations for the eDNA analysis. Crosses indicate sites

where a positive detection of *A. pallipes* was made in at least one field replicate. Circles

indicate sites where no positive detection of *A. pallipes* was made in all field replicates. River

subcatchments in which field observations of *A. pallipes* were made between the years 2007

and 2016 are highlighted in grey. Grey circles indicate the actual locations where field

observations of *A. pallipes* were made.