Field application of an eDNA assay for the threatened white-clawed crayfish

Austropotamobius pallipes

Siobhán Atkinson[1,2,3], Jeanette E. L. Carlsson[2,4], Bernard Ball[2,5], Mary Kelly-Quinn[1,6], and

Jens Carlsson[2,7]

[1]School of Biology and Environmental Science/Earth Institute, University College Dublin,

Dublin, Ireland

[2]Area52 Research Group, School of Biology and Environmental Science/Earth Institute,

University College Dublin, Dublin, Ireland

E-mail addresses: [3]siobhan.atkinson@ucdconnect.ie, [4]jeanette.carlsson@ucd.ie,

[5]bernie.ball@ucd.ie, [6]mary.kelly-quinn@ucd.ie, [7]jens.carlsson@ucd.ie

Abstract: The white-clawed crayfish *Austropotamobius pallipes* has experienced extensive

population declines in its native range in the last century and is now a threatened species.

European legislation therefore requires *A. pallipes* populations to be monitored regularly and that

Special Areas of Conservation (SACs) be designated in part of their range. These needs cannot

be met without knowledge about the distribution of this species. However, common crayfish

sampling approaches can be time consuming, expensive, or yield false negatives at low

population densities. We therefore present an environmental DNA (eDNA) assay to detect *A.*

pallipes in water samples with species-specific primers, a minor groove binding (MGB) probe,

and quantitative PCR, based on the mitochondrial cytochrome oxidase I (COI) gene. The results

reflected the expected distribution of *A. pallipes*, based on the known locations of this species in

the catchment areas sampled. The results of this study indicate that eDNA is an effective tool for

detecting *A. pallipes* in a lotic system, and could therefore provide a valuable, non-invasive

method for determining its distribution.

Keywords: eDNA, qPCR, conservation, non-invasive, detection, native species,

Austropotamobius pallipes, Ireland, river

The white-clawed crayfish *Austropotamobius pallipes* is a relatively large, long-lived (>10 y)

crustacean that inhabits both rivers and lakes (Reynolds et al. 2010). This species requires

alkaline conditions for survival and is commonly found in waterbodies overlying carboniferous

limestone bedrock (Lucey and McGarrigle 1987). *Austropotamobius pallipe*s is 1 of the 5

indigenous crayfish species in Europe (Holdich et al. 2009) and was once abundant throughout

Europe. However, over the last century, populations of this species have been greatly reduced or

gone locally extinct across large parts of its native range (Grandjean et al. 1997). Pollution

(Demers and Reynolds 2002, Lyons and Kelly-Quinn 2003), habitat loss, and disease (Matthews

and Reynolds 1992) have all contributed to this decline. The crayfish plague, caused by the

fungus *Aphanomyces astaci* (Holdich et al. 2009) is of particular concern because *A. pallipes*

possesses no resistance to this fungus. Thus, entire populations of *A. pallipes* can be eradicated

following an outbreak (Reynolds et al. 2010).

In spite of the crayfish plague, Ireland is considered one of the few remaining strongholds

for *A. pallipes* within Europe. One reason for this may be that the invasive signal crayfish

(*Pacifastacus leniusculus*), which can cause wide-ranging impacts on freshwater ecosystems and

act as a vector for the crayfish plague (Vaeßen and Hollert 2015), has not been reported in

Ireland. However, crayfish plague has reached Irish rivers, and large changes in the distribution

of *A. pallipes* have been partially attributed to past plague outbreaks (Demers et al. 2005).

Additionally, there is currently an outbreak of crayfish plague in Ireland (National Biodiversity

Data Center 2018). Thus, there is a pressing need for a non-invasive, rapid, and cost-effective

sampling method that can be used to establish the distribution of this species.

Austropotamobius pallipes is protected under the EU Habitats Directive, and is listed

under Annexes II and V because of its vital role in freshwater ecosystem functioning and

threatened status (Matthews and Reynolds 1992). Ireland must therefore regularly monitor *A.*

pallipes populations and designate Special Areas of Conservation (SACs) for the species under

Natura 2000 (Reynolds et al. 2010). Knowledge of the geographic distribution of *A. pallipes* is

essential for implementing these conservation measures. Traditional survey methods include

night viewing with a strong light, modified quadrat sampling (DiStefano et al. 2003), kick-

sampling, baiting traps and enclosures (Byrne and Lynch 1999), snorkeling surveys (Reynolds et

al. 2010), and SCUBA diving (Matthews and Reynolds 1992). Many of these methods are

effective, but they are also time-consuming, potentially costly, and may be less effective when *A.*

pallipes occurs in low abundance. Furthermore, many of these traditional *A. pallipes* surveying

methods require a license that allows the holder to catch and disturb *A. pallipes*. Licenses for

invasive monitoring procedures are vital for ensuring the protection of *A. pallipes*, however, a

non-invasive method that does not require a license would be beneficial.

A promising new tool that can effectively detect species and save significant time in the

field (Sigsgaard et al. 2015) is environmental DNA (eDNA) analysis. eDNA is the collective

term for DNA that has been shed by organisms and is therefore present freely in the

environment. For example, eDNA can occur in the form of mucus, excrement, gametes, or blood

(Taberlet et al. 2012, Thomsen and Willerslev 2015). eDNA analysis has effectively detected

species that occur in low abundance, such as rare or endangered species (Mächler et al. 2014,

Laramie et al. 2015, Sigsgaard et al. 2015, Boothroyd et al. 2016, Carlsson et al. 2017) and

invasive alien species (Jerde et al. 2011, Goldberg et al. 2013, Fujiwara et al. 2016). In addition,

eDNA may be better able to detect species at low abundance than traditional methods (Dejean et

al. 2012, Smart et al. 2015, Dougherty et al. 2016). eDNA assays have been successfully

employed for other crayfish species, including *Astacus astacus*, *Astacus leptodactylus*, *P.*

leniusculus (Agersnap et al. 2017, Larson et al. 2017), *Orconectes rusticus* (Dougherty et al.

2016, Larson et al. 2017), *Procambarus clarkii* (Cai et al. 2017), and *Faxonius eupunctus* (Rice

et al. 2018).

The aim of this study was to develop a minor groove binding (MGB) probe-based qPCR

assay to detect the presence of *A. pallipes* in water samples and to test the reliability of this assay

by comparing the results with field distribution data.

METHODS

Study sites and crayfish distribution data

Eight sampling locations within 7 different rivers were selected for field validation of the

assay (Table 1). The research presented here was undertaken as part of a larger study assessing

the impact of river obstacles on freshwater fauna (e.g. Atlantic Salmon, *Salmo salar*, Atkinson et

al. 2018). As part of the broader study, eDNA samples were collected above and below a number

of river obstacles (weir, ford crossing, or bridge apron) to determine whether they imposed

migratory barriers on *S. salar*, and we used the same samples here. Of the 8 sites sampled, 4

were in rivers that are Special Areas of Conservation (SAC) that include *A. pallipes* as a species

of interest. We sampled the Delour and Dinin rivers, which are within the River Barrow and

River Nore Special SAC, and the Multeen and Duag rivers, which are within the Lower River

Suir SAC. The other 3 rivers that were sampled were the Burren River, Brown's Beck Brook and

the Dalligan River. River flow conditions were low at each study site at the time of sampling.

We estimated the current distribution of *A. pallipes* in these rivers with point data

provided by the Irish Environmental Protection Agency (EPA). These data were based on

opportunistic field observations (i.e. the EPA was not specifically looking for *A. papilles*) made

by EPA staff while undertaking routine biological monitoring between the years 2007 and 2016

(W. Trodd, Irish Environmental Protection Agency, personal communication). Electrofishing

surveys were carried out at each site as part of the larger study mentioned previously. At some

sites, crayfish were observed during these electrofishing surveys. These additional observations

were included in the EPA's *A. pallipes* field observation dataset. We extrapolated each *A.*

pallipes observation to the wider subcatchment level. Thus, even if *A. pallipes* was only

observed once (between the years 2007 and 2016) in a subcatchment, we assumed that they

occurred in the rest of that subcatchment.

eDNA qPCR assay development

We used *A. papilles*-specific primers (forward primer: 5'- GGG TTA GTG GAG AGA

GGG GT -3', and reverse primer 5'- AAT CCC CAG ATC CAC AGA CG -3') and a 5'-6-FAM

labeled TaqMan® MGB probe (5'- TCA GCT ATT GCC CAC GCA -3') to target a locus within

the mitochondrial cytochrome oxidase I (COI) region. We designed these primers in Primer

Express 3.0 (Applied Biosystems-Roche, Branchburg, NJ) with Irish *A. pallipes* COI sequences

from the National Center for Biotechnology Information (NCBI - http://www.ncbi.nlm.nih.gov/).

The total amplicon size was 96 base pairs, including the primers. To verify the in silico species

specificity of the *A. pallipes* assay, we combined the probe and primer sequences, and queried

them against the NCBI nucleotide database with BLASTn (Basic Local Alignment Search Tool).

The BLASTn analyses showed 90% sequence identity for 100% of the assay query only for the

target species. The sequence did not align with 100% of the sequence identity in the BLASTn

search because there were gaps between the forward and reverse primers and the probe.

However, non-target species showed a maximum of 37% query cover with 100% sequence

identity indicating that only the target species *A. pallipes* would bind to all 3 components

(forward and reverse primers, probe) of the assay. To verify the in vitro species specificity of the

assay, we tested the assay with DNA from the tissue of some freshwater species that co-occur

with *A. pallipes,* including Brown Trout (*Salmo trutta*), Sea Lamprey (*Petromyzon marinus*),

Atlantic Salmon (*S. salar*), Allis Shad (*Alsoa alosa),* and Twaite Shad (*Alosa fallax),* with

conventional PCR techniques followed by visualization of amplicons on agarose. We optimized

the qPCR assay with tissue extracted from *A. pallipes*.

eDNA collection, filtration, and extraction

We collected 2 L water samples for eDNA analysis in PET bottles from 2 locations at

each study site, above and below the river obstacle. Samples were collected between 45 and 267

m downstream of each obstacle to allow for the accumulation of eDNA from species found

potentially only below the obstacle. Upstream eDNA samples were collected between 1 and 193

m upstream of the obstacles (Table 1.). It was not possible to consistently collect samples from

the same distance at each location due to health and safety precautions. At each sampling

location, 3 replicate water samples were collected, one from each of the left margin, right

margin, and the center of the river channel. In addition, we filtered 1 negative field control of

double distilled H_2O (ddH_2O) at each location. This procedure resulted in a total of 6 water

samples for eDNA analysis and 2 field controls collected per study site. To sterilize the PET

bottles before sample collection, we soaked them for 30 min in 50% sodium hypochlorite

solution. Prior to collecting the final water samples at each location, the bottles were rinsed at

least 5 times with river water to remove any remaining sodium hypochlorite. To further avoid

sample contamination, water samples were taken in an upstream direction before any researchers

entered the river. Water samples were filtered through disposable filter funnels on site with a

peristaltic pump, through 47-mm glass microfiber filters (1.5 µm). Filters were placed in 2.0 mL

Eppendorf tubes and subsequently stored in a cooler box on ice. The filters were frozen at -20˚C

in the laboratory ~3-6 h following collection.

All eDNA work was completed in a dedicated Low Copy DNA laboratory to reduce

contamination risk. We used a modified CTAB (cetyltrimethylammonium bromide) protocol

(Möller et al. 1992) to extract the eDNA. Briefly, ½ of each filter was transferred to a 2.0 mL

Eppendorf tube, and the other ½ was archived for future validation or queries. We then added

750 µL CTAB buffer (100 mM Tris-HCL, 20 mM EDTA, 1.4 M NaCl, 2% CTAB) and 7 µL of

Proteinase K (20 mg/mL) to the tube with the sample filter. Samples were then vortexed for 10 s,

followed by incubation at 56˚C for 2 h. We then added 750 µL of a phenol, chloroform, and

isoamyl alcohol (25:25:1 v/v) mixture. We manually mixed the tube contents for 15 s and

subsequently centrifuged the tube at 11,000 x g for 20 min. We removed the aqueous solution

from the centrifuged tube, and placed it in a new tube that contained 750 µL of a chloroform and

isoamyl alcohol (24:1 v/v) mixture. We repeated the manual mixing and centrifugation steps

once, and transferred the aqueous solution to a new tube. We precipitated the DNA from the

solution by adding 1 volume of isopropanol alcohol, incubating this mixture at -20˚C for 1 h, and

centrifuging at 11,000 x g for 20 min. The resulting pellets were washed with 750 µL of 70%

ethanol and centrifuged at 11,000 x g for 5 min. We then removed the ethanol, taking care to

ensure that the pellet remained in the tube. The tubes containing pellets were dried in a heat

block (50˚C, 5 min), and the pellet was then resuspended in 40 µL of molecular-grade ddH$_2$O.

eDNA concentrations were determined with NanoDrop®-1000, Thermo Scientific, Wilmington,

DE.

eDNA assay deployment on field samples

We used an Applied Biosystems ViiA™ 7 (Life Technologies, Inc., Applied Biosystems,

Foster City, California, USA.) quantitative thermocycler to amplify and quantify eDNA. Briefly,

the PCR profile consisted of heating at 50°C for 2 min and 95°C for 10 min, followed by 40

cycles of 95°C for 15 s and 60°C for 1 min. We generated standard curves from DNA originating

from *A. pallipes* tissue extracts (quantified with NanoDrop®-1000, Thermo Scientific,

Wilmington, DE). The standard curves served as both a positive control and as a means of

calculating the concentration of *A. pallipes* eDNA in each sample. We used seven 10:1 serial

dilutions ranging from 2.9 to 2.9 x 10^{-6} ng/µL to make the standard curves. Each qPCR reaction

was carried out in a total volume of 30 µL that consisted of 15 µL of TaqMan® Environmental

Master Mix 2.0 (Life Technologies, Applied Biosystems, Foster City, CA), 3 µL of each primer

(final concentration of 2 µM), 3 µL probe (final concentration of 2 µM), 3 µL eDNA/tissue

extracted DNA template, and 3 µL ddH$_2$O. We generated individual standard curves for each

qPCR plate (y = -3.37x + 20.589, efficiency = 98.04 %, R^2 = 0.996 [1], y = -3.442x + 20.392,

efficiency = 95.22 %, R^2 = 0.999 [2] and y = -3.441x + 20.723, efficiency = 95.26 %, R^2 = 0.999

[3]). Each plate incorporated 2 no-template controls (NTCs) to identify any contamination that

occurred during preparation. Standard curve, field, and control samples were quantified in

triplicate (3 technical replicates). For a positive detection to occur at a site, at least 2 of the 3

technical replicates had to contain amplifiable DNA with quantitation cycle (Cq) differences

<0.5Cq (to allow for pipetting errors), and be within the detection range of the standard curve. If

1 out of 3 technical replicates showed Cq differences >0.5 Cq, this technical replicate was

excluded from the study. In cases when the Cq value of 2 out of 3 technical replicates differed by

>0.5 Cq, that specific dilution series or field replicate was removed from further study.

Salmo trutta were present in all rivers, both upstream and downstream of the obstacles, so

we used this species as a positive field control to test for the presence of amplifiable eDNA in all

samples. To detect the presence of *S. trutta* DNA in our samples, we used the eDNA assay

described in Gustavson et al. (2015). Standard curves were not generated, as the purpose of this

analysis was to simply establish the presence or absence of *S. trutta* DNA in our samples. Three

replicates per location with 1 technical replicate were used.

RESULTS

In this study, we successfully detected *A. pallipes* in silico, in vitro, and in situ. The *A.*

pallipes eDNA assay was highly sensitive, the dynamic range of the standard curves was

between 19.1 Cq and 39 Cq. The lowest concentration of *A. pallipes* eDNA detected within the

range of the standard curve was 0.002 ng/L at Cq 38.2 (average over 3 technical replicates, sd =

0.000059 ng/L). No cross-species amplification occurred. Furthermore, no amplification

occurred in the NTCs or field controls. *Salmo trutta* DNA (positive field control) was detected in

all the field samples, which confirmed the presence of detectable DNA in them. The eDNA

results accurately reflect the *A. pallipes* observation data the EPA collected in the field.

Austropotamobius pallipes eDNA was positively detected in all subcatchments where EPA staff

observed *A. pallipes,* and not detected in any of the subcatchments where *A. pallipes* were not

observed (Fig. 1). *Austropotamobius papilles* DNA was detected in all field replicates above and

below the river obstacles within the Burren, Duag, and Multeen rivers. At the Dinin River

(Crettyard), we detected *A. pallipes* at 2 out of 3 field replicates above the obstacle, and 3 out of

3 field replicates below the obstacle. At the Dinin River (Castlecomer), we detected *A. pallipes*

only in the field replicates above the obstacle (Table 2).

In addition, the eDNA results also reflect the expected distribution of *A. pallipes* based on

siliceous dominant bedrock. *Austropotamobius pallipes* usually occurs in waterbodies with

calcareous, rather than siliceous, bedrock (Lucey and McGarrigle 1987). The dominant bedrock

was either 100% siliceous (Brown's Beck Brook and the Dalligan River) or was 1-25%

calcareous (mixed geology, Delour River) at the sites where *A. pallipes* was not detected by

either field observation or eDNA.

DISCUSSION

This study showed that eDNA is a reliable tool for detecting *A. pallipes* in river systems.

Furthermore, this assay could also be used in lentic systems, such as lakes or reservoirs, where

field sampling can be more demanding. eDNA sampling requires only 1 site visit, which

involves collecting water samples and filtering them onsite. In contrast, survey methods for

monitoring *A. pallipes* can vary in their efficiency depending on the habitat type being surveyed,

but they all are usually more involved than what is required for collecting eDNA (Reynolds et al.

2010). Night searches, for example, typically require 2 site visits (for health and safety purposes)

and are dependent on water clarity and algal growth (Reynolds et al. 2010). Trapping is time-

consuming and requires 2 site visits, whereas snorkeling and hand searches are habitat

dependent. Thus, eDNA monitoring may lend itself to surveying more challenging habitats, such

as heavily vegetated lake margins with soft substrates, or locations where health and safety

considerations make trapping or night searching impossible or difficult.

In addition to increasing sampling efficiency, eDNA sampling requires substantially less

field equipment than traditional sampling methods. Herder et al. (2014) suggested that the eDNA

method may help reduce the risk of unintentional pathogen transmission into new areas because

of the numerous contamination-prevention precautions that must be taken when sampling.

Further, eDNA sampling, and the subsequent disinfection of field equipment is relatively quick.

For example, excluding instances where a boat is required, sampling in general only requires the

collection of a number of water samples, followed by sterilization of footwear, containers and

filter funnels. Snorkeling, SCUBA diving or trapping surveys, in contrast, require time to

disinfect and dry equipment (wetsuits, traps etc.) between entering different waterbodies

(Reynolds et al. 2010). eDNA sampling could minimize the risk of further spread of crayfish

plague by reducing the amount of field equipment required and therefore potential vectors of

crayfish plague. eDNA analysis may therefore be particularly suitable for surveys of waterbodies

that harbor crayfish plague.

It is surprising that we detected eDNA above but not below the obstacle at the Dinin

River (Castlecomer) because eDNA travels downstream in river systems (Deiner and Altermatt

2014, Deiner et al. 2016). This site had the lowest concentration of *A. pallipes* eDNA of all

locations (0.002 ng/L), and eDNA amplification occurred in only 1 out of a total of 6 site

replicates collected from above and below the obstacle. It is possible that either eDNA

concentrations in the other field samples were too low for a positive detection or we obtained

false positive results. However, we feel false positives results in this instance are unlikely, as we

detected *A. pallipes* in a total of 5 out of 6 field replicates at the Dinin River (Crettyard), which is

located approximately 9 km upstream of the Dinin River (Castlecomer) (Fig. 1). The density or

biomass of *A. pallipes*, environmental conditions such as flow, temperature, pH, and sediment, or

natural inhibitors such as algae or humic substances, could all have contributed to this lack of

amplification in the Dinin River (Castlecomer) (Strickler et al. 2015, Goldberg et al. 2016,

Stoeckle et al. 2017) if *A. pallipes* was present there. However, all eDNA samples yielded

detectable DNA with the positive field control (e.g. *S. trutta*), so it is unlikely that in this

instance the lack of amplification of *A. pallipes* DNA was a result of inhibition of the eDNA

samples.

In conclusion, this eDNA assay provides an alternative, rapid method for locating *A.*

pallipes in lotic ecosystems that allows for more targeted, efficient surveying strategies.

Considering the current crayfish plague outbreak in Ireland (National Biodiversity Data Center

2018) and other countries, there is an urgent need for a non-invasive sampling method for this

species. Large scale deployment of this assay would enable managers to rapidly establish the

distribution of *A. pallipes* with confidence, which would support the designation of new

conservation areas.

ACKNOWLEDGEMENTS

Author contributions: SA conducted some of the field and laboratory work and led the

data analyses and writing of the paper. JELC conducted some of the laboratory work. BB

designed the primers and probe and conducted the majority of the field and laboratory work.

MKQ contributed to the writing of the paper. JC conceived and designed the study and assisted

with data analyses and writing.

This research was funded by the Irish Environmental Protection Agency (Reconnect

Project - 2015-W-LS-8). The authors would also like to thank the Irish Environmental Protection

Agency for providing the data on *A. pallipes* distributions.

LITERATURE CITED

Agersnap, S., W. B. Larsen, S. W. Knudsen, D. Strand, P. F. Thomsen, M. Hesselsøe, P. B.

Mortensen, T. Vrålstad, and P. R. Møller. 2017. Monitoring of noble, signal and narrow-

clawed crayfish using environmental DNA from freshwater samples. PLoS ONE 12:1–

22.

Atkinson, S., J.E.L. Carlsson, B. Ball, D. Egan, M. Kelly-Quinn, K. Whelan, and J. Carlsson.

2018. A quantitative PCR-based environmental DNA assay for detecting Atlantic salmon

(*Salmo salar* L.). Aquatic Conservation: Marine and Freshwater Ecosystems 28:1238–

1243.

Boothroyd, M., N. E. Mandrak, M. Fox, and C. C. Wilson. 2016. Environmental DNA (eDNA)

detection and habitat occupancy of threatened spotted gar (*Lepisosteus oculatus*). Aquatic

Conservation: Marine and Freshwater Ecosystems 26:1107–1119.

Byrne, C., and J. Lynch. 1999. A sampling strategy for stream populations of white-clawed

crayfish, *Austropotamobius pallipes* (Lereboullet)(Crustacea, Astacidae). Biology and

Environment: Proceedings of the Royal Irish Academy 99b:89–94.

Cai, W., Z. Ma, C. Yang, L. Wang, W. Wang, G. Zhao, Y. Geng, and D. W. Yu. 2017. Using

eDNA to detect the distribution and density of invasive crayfish in the Honghe-Hani rice

terrace World Heritage site. PLoS ONE 12:1–13.

Carlsson, J. E. L., D. Egan, P. C. Collins, E. D. Farrell, F. Igoe, and J. Carlsson. 2017. A qPCR

MGB probe based eDNA assay for European freshwater pearl mussel (*Margaritifera*

margaritifera L.). Aquatic Conservation: Marine and Freshwater Ecosystems 27:1341-

1344.

Deiner, K., and F. Altermatt. 2014. Transport distance of invertebrate environmental DNA in a

natural river. Plos One 9:e88786.

Deiner, K., E. A. Fronhofer, E. Mächler, J-C. Walser and F. Altermatt. 2016. Environmental

DNA reveals that rivers are conveyer belts of biodiversity information. Nature

Communications 7:12544

Dejean, T., A. Valentini, C. Miquel, P. Taberlet, E. Bellemain and C. Miaud. 2012. Improved

detection of an alien invasive species through environmental DNA barcoding: the

example of the American bullfrog *Lithobates catesbeianus*. Journal of Applied Ecology

49:953-959.

Demers, A., J. Lucey, M. L. McGarrigle, and J. D. Reynolds. 2005. The distribution of the white-

clawed crayfish, *Austropotamobius pallipes*, in Ireland. Biology & Environment:

Proceedings of the Royal Irish Academy 105b:65–69.

Demers, A., and J. D. Reynolds. 2002. A survey of the white-clawed crayfish, *Austropotamobius*

pallipes (Lereboullet), and of water quality in two catchments of eastern Ireland. Bulletin

Francais de Pêche et de Pisciculture 367:729–740.

DiStefano, R. J., C. M. Gale, B. A. Wagner, and R. D. Zweifel. 2003. A sampling method to

assess lotic crayfish communities. Journal of Crustacean Biology 23:678–690.

Dougherty, M. M., E. R. Larson, M. A. Renshaw, C. A. Gantz, S. P. Egan, D. M. Erickson, and

D. M. Lodge. 2016. Environmental DNA (eDNA) detects the invasive rusty crayfish

Orconectes rusticus at low abundances. Journal of Applied Ecology 53:722–732.

Fujiwara, A., S. Matsuhashi, H. Doi, S. Yamamoto, and T. Minamoto. 2016. Use of

environmental DNA to survey the distribution of an invasive submerged plant in ponds.

Freshwater Science 35:748–754.

Goldberg, C. S., A. Sepulveda, A. Ray, J. Baumgardt, and L. P. Waits. 2013. Environmental

DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus*

antipodarum). Freshwater Science 32:792–800.

Goldberg, C. S., C. R. Turner, K. Deiner, K. E. Klymus, P. F. Thomsen, M. A. Murphy, S. F.

Spear, A. McKee, S. J. Oyler-McCance, R. S. Cornman, M. B. Laramie, A. R. Mahon, R.

F. Lance, D. S. Pilliod, K. M. Strickler, L. P. Waits, A. K. Fremier, T. Takahara, J. E.

Herder and P. Taberlet. 2016. Critical considerations for the application of environmental

DNA methods to detect aquatic species. Methods in Ecology and Evolution 7: 1299-

1307.

Grandjean, F., C. Souty-Grosset, R. Raimond, and D. M. Holdich. 1997. Geographical variation

of mitochondrial DNA between populations of the white-clawed crayfish

Austropotamobius pallipes. Freshwater Biology 37:493–501.

Gustavson, M. S., P. C. Collins, J. A. Finarelli, D. Egan, R. Ó. Conchúir, G. D. Wightman, J. J.

King, D. T. Gauthier, K. Whelan, J. E. L. Carlsson, and J. Carlsson. 2015. An eDNA

assay for Irish *Petromyzon marinus* and *Salmo trutta* and field validation in running

water. Journal of Fish Biology 87:1254–1262.

Herder, J. E., A. Valentini, E. Bellemain, T. Dejean, J. J. C. W. van Delft, P. F. Thomsen, and P.

Taberlet. 2014. Environmental DNA—a review of the possible applications for the

detection of (invasive) species. Report no. 2013–104. Stichting RAVON, Nijmegen.

Holdich, D. M., J. D. Reynolds, C. Souty-Grosset, and P. J. Sibley. 2009. A review of the ever

increasing threat to European crayfish from non-indigenous crayfish species. Knowledge

and Management of Aquatic Ecosystems 394-395:11.

Jerde, C. L., A. R. Mahon, W. L. Chadderton, and D. M. Lodge. 2011. "Sight-unseen" detection

of rare aquatic species using environmental DNA. Conservation Letters 4:150–157.

Laramie, M. B., D. S. Pilliod, and C. S. Goldberg. 2015. Characterizing the distribution of an

endangered salmonid using environmental DNA analysis. Biological Conservation

183:29–37.

Larson, E. R., M. A. Renshaw, C. A. Gantz, J. Umek, S. Chandra, D. M. Lodge, and S. P. Egan.

2017. Environmental DNA (eDNA) detects the invasive crayfishes *Orconectes rusticus*

and *Pacifastacus leniusculus* in large lakes of North America. Hydrobiologia 800:173–

185.

Lucey, J., and M. L. McGarrigle. 1987. The distribution of the crayfish *Austropotamobius*

pallipes (Lereboullet) in Ireland. Irish Fisheries Investigations A29:1–13.

Lyons, R., and M. Kelly-Quinn. 2003. An investigation into the disappearance of

Austropotamobius pallipes (Lereboullet) populations in the headwaters of the Nore River,

Ireland and the correlation to water quality. Bulletin Français de la Pêche et de la

Pisciculture 370-371:139–150.

Mächler, E., K. Deiner, P. Steinmann, and F. Altermatt. 2014. Utility of environmental DNA for

monitoring rare and indicator macroinvertebrate species. Freshwater Science 33:1174–

1183.

Matthews, M., and J. D. Reynolds. 1992. Ecological impact of crayfish plague in Ireland.

Hydrobiologia 234:1–6.

Möller, E. M., G. Bahnweg, H. Sandermann, and H. H. Geiger. 1992. A simple and efficient

protocol for isolation of high molecular weight DNA from filamentous fungi, fruit

bodies, and infected plant tissues. Nucleic Acids Research 20:6115–6116.

National Biodiversity Data Center. 2018. Crayfish plague. (Available from:

http://www.biodiversityireland.ie/projects/invasive-species/crayfish-plague/)

Reynolds, J. D., W. O'Connor, C. O'Keeffe, and D. Lynn. 2010. A technical manual for

monitoring white-clawed crayfish *Austropotamobius pallipes* in Irish lakes. (Available at:

https://www.npws.ie/sites/default/files/publications/pdf/IWM45.pdf)

Rice, C. J., E. R. Larson, and C. A. Taylor. 2018. Environmental DNA detects a rare large river

crayfish but with little relation to local abundance. Freshwater Biology 63:443–455.

Sigsgaard, E. E., H. Carl, P. R. Møller, and P. F. Thomsen. 2015. Monitoring the near-extinct

European weather loach in Denmark based on environmental DNA from water samples.

Biological Conservation 183:46–52.

Smart, A.S., R. Tingley, A. R. Weeks, A. R. van Rooyen, and M. A. McCarthy. 2015.

Environmental DNA sampling is more sensitive than a traditional survey technique for

detecting an aquatic invader. Ecological Applications 25:1944-1952.

Stoeckle, B. C., S. Beggel, A. F. Cerwenka, E. Motivans, R. Kuehn and J. Geist. 2017. A

systematic approach to evaluate the influence of environmental conditions on eDNA

detection success in aquatic ecosystems. PloS one 12: e0189119.

Strickler, K.M., A. K. Fremier and C.S. Goldberg. 2015. Quantifying effects of UV-B,

temperature, and pH on eDNA degradation in aquatic microcosms. Biological

Conservation 183: 85-92.

Taberlet, P., E. Coissac, M. Hajibabaei, and L. H. Rieseberg. 2012. Environmental DNA.

Molecular Ecology 21:1789–1793.

Thomsen, P. F., and E. Willerslev. 2015. Environmental DNA - An emerging tool in

conservation for monitoring past and present biodiversity. Biological Conservation

183:4–18.

Vaeßen, S., and H. Hollert. 2015. Impacts of the North American signal crayfish (*Pacifastacus*

leniusculus) on European ecosystems. Environmental Sciences Europe 27:1–6.

Table 1. Sampling locations from which we collected water to sample *Austropotamobius pallipes* eDNA. Each sampling site had one collection
point upstream and one downstream of an obstacle.

River	Obstacle Type	Sampling Location	Distance from Obstacle (m)	Latitude	Longitude	Sampling Date
Burren	Weir	Upstream	1	52.83300	-6.92534	18/07/2017
		Downstream	215	52.83388	-6.92822	18/07/2017
Delour	Weir	Upstream	193	52.97709	-7.54606	19/07/2017
		Downstream	267	52.97957	-7.54100	19/07/2017
Nore (Castlecomer)	Weir	Upstream	11	52.80599	-7.20527	20/07/2017
		Downstream	57	52.80535	-7.20558	20/07/2017
Duag	Weir	Upstream	8	52.27196	-8.01319	21/07/2017
		Downstream	45	52.27174	-8.01248	21/07/2017
Brown's Beck Brook (BBB)	Ford	Upstream	1	53.01627	-6.62311	24/07/2017
		Downstream	60	53.01573	-6.62322	24/07/2017
Dalligan	Weir	Upstream	15	52.11876	-7.51938	25/07/2017

		Downstream	92	52.11862	-7.51803	25/07/2017
Muilteen	Bridge	Upstream	101	52.61304	-8.00872	27/07/2017
	Apron					
		Downstream	34	52.61182	-8.00966	27/07/2017
Nore	Bridge	Upstream	8	52.84534	-7.12758	28/07/2017
(Crettyard)	Apron					
		Downstream	68	52.84584	-7.12891	28/07/2017

Table 2. Mean (± 95% CI) eDNA concentrations (ng/L) from the *A. pallipes* assay at each study river. Concentrations are based on the average eDNA concentration in a minimum of 2 technical replicates and between 1 and 3 field replicates. Confidence intervals (95%) are presented for each location where positive detections occurred in >1 of our field replicates.

River	Location	No. eDNA samples collected	No. of positive detections	eDNA concentration ng/L (95% CI)	Visual observation
Burren	Upstream	3	3	0.027 (0.01-0.02)	+
	Downstream	3	3	0.02 (0.01-0.04)	
Duag	Upstream	3	3	0.027 (0.005-0.05)	+
	Downstream	3	3	0.157 (0.09-0.23)	
Multeen	Upstream	3	3	0.034 (0.02-0.04)	+
	Downstream	3	3	0.06 (0.04-0.08)	
Dinin (Crettyard)	Upstream	3	2	0.011 (0.004-0.02)	+
	Downstream	3	3	0.006 (0.002-0.01)	
Dinin (Castlecomer)	Upstream	3	1	0.002	+
	Downstream	3	0	0	
Delour	Upstream	3	0	0	-
	Downstream	3	0	0	
Dalligan	Upstream	3	0	0	-
	Downstream	3	0	0	

Brown's Beck Brook (BBB)	Upstream	3	0	0	-
	Downstream	3	0	0	

FIGURE CAPTIONS

Fig. 1. Map showing the sampling locations for the eDNA analysis. Crosses indicate sites

where a positive detection of *A. pallipes* was made in at least 1 field replicate. Circles indicate

413   sites where *A. pallipes* was not detected in any field replicates. River subcatchments in which

414   field observations of *A. pallipes* were made between the years 2007 and 2016 are highlighted

415   in grey. Grey circles indicate the actual locations where field observations of *A. pallipes* were

416   made.