Field validation of an eDNA assay for the endangered white-clawed crayfish

Austropotamobius pallipes

Siobhán Atkinson[1,2]*, Jeanette E.L. Carlsson[2], Bernard Ball[2], Mary Kelly-Quinn[1], Jens Carlsson[2]

1. School of Biology and Environmental Science, University College Dublin, Dublin,

Ireland

2. Area52 Research Group, School of Biology and Environmental Science/Earth Institute,

University College Dublin, Dublin, Ireland

*Corresponding author: Siobhán Atkinson, siobhan.atkinson@ucdconnect.ie

Abstract

The white-clawed crayfish *Austropotamobius pallipes* has undergone extensive declines

within its native range in the last century. Because of its threatened status, European

legislation requires the species to be regularly monitored and that Special Areas of

Conservation (SACs) be designated for it. Knowledge on the distribution of this species is

vital for addressing these needs. This study presents an environmental (e)DNA assay to

detect *A. pallipes* in water samples, based on the mitochondrial cytochrome oxidase I (COI)

gene, utilizing species-specific primers, a minor groove binding (MGB) probe and

quantitative PCR. The results of this study indicate that eDNA is an effective tool for

detecting *A. pallipes* in a lotic system, and could provide a valuable, non-invasive method for

determining the distribution of this species.

Keywords: eDNA, qPCR, conservation, non-invasive, detection, native species, river.

Introduction

The white-clawed crayfish *Austropotamobius pallipes* is a relatively large, long-lived (>10

years) crustacean that inhabits both rivers and lakes (Reynolds et al. 2010). They require

alkaline conditions for survival and are commonly found in waterbodies overlying

carboniferous limestone bedrock (Lucey and McGarrigle 1987). *A. pallipe*s is one of the five

indigenous crayfish species in Europe (Holdich et al. 2009). This once abundant species has,

however, become greatly reduced or locally extinct across large parts of its native range

during the last century (Grandjean et al. 1997). Pollution (Demers and Reynolds 2002, Lyons

and Kelly-Quinn 2003), habitat loss and disease (Matthews and Reynolds 1992) have

contribution to this decline. Of particular concern is the crayfish plague, caused by the

fungus *Aphanomyces astaci* (Holdich et al. 2009). *A. pallipes* possess no resistance to this

fungus, so eradication of entire populations of *A. pallipes* is possible following an outbreak

(Reynolds et al. 2010). Despite this, Ireland is considered one of the few remaining

strongholds for *A. pallipes* within Europe. One of the reasons for this is that the invasive

signal crayfish *Pacifastacus leniusculus,* which can impose wide-ranging impacts on

ecosystems and act as a vector for the crayfish plague (Vaeßen and Hollert 2015), has not

been reported in Ireland to date. However, the crayfish plague has reached Ireland's rivers,

and large changes in the distribution of *A. pallipes* have been attributed, at least in part, to

past plague outbreaks (Demers et al. 2005). Additionally, there is at present an outbreak of

the crayfish plague in Ireland (National Biodiversity Data Centre 2018), so more than ever

there is a need for non-invasive sampling methods that do not have the potential to further

spread the crayfish plague.

Because of its vital role in freshwater ecosystem functioning and its threatened status

(Matthews and Reynolds 1992), *A. pallipes* is afforded protection under the EU Habitats

Directive and is listed under Annexes II and V. This means Ireland is required to regularly

monitor *A. pallipes*, and to designate Special Areas of Conservation (SACs) for the species

under Natura 2000 (Reynolds et al. 2010). Knowledge on the geographic distribution of *A.*

pallipes is essential for implementing these conservation measures. Traditional survey

methods include night viewing with a strong torch, modified quadrat samplers (DiStefano et

al. 2003), kick-sampling, baited traps and enclosures (Byrne and Lynch 1999) and SCUBA

diving (Matthews and Reynolds 1992). While many of these methods are effective, they are

time-consuming, potentially costly, and may not be effective when *A. pallipes* occur in low

abundance. Furthermore, as white-clawed crayfish are a protected species in Ireland, several

surveying methodologies require a licence from the National Parks and Wildlife Service.

Certainly, a low-cost, non-invasive and effective methodology for assessing the distribution

57 of *A. pallipes* is warranted. A promising new tool that has been shown to be both effective at

58 detecting species and to save significant time in the field (Sigsgaard et al. 2015) is

59 environmental DNA (eDNA) analysis. Environmental DNA is the collective term for DNA

60 present freely in the environment (Taberlet et al. 2012, Thomsen and Willerslev 2015). The

61 analysis has been successful in detecting species that occur in low abundance such as rare or

62 endangered species (Mächler et al. 2014, Laramie et al. 2015, Sigsgaard et al. 2015,

63 Boothroyd et al. 2016, Carlsson et al. 2017), or invasive alien species (Jerde et al. 2011,

64 Goldberg et al. 2013, Fujiwara et al. 2016). Futhermore, eDNA assays have been successfully

65 deployed for other crayfish species including *Astacus astacus*, *Astacus leptodactylus*, *P.*

66 *leniusculus* (Agersnap et al. 2017, Larson et al. 2017), *Orconectes rusticus* (Dougherty et al.

67 2016, Larson et al. 2017), *Procambarus clarkii* (Cai et al. 2017) and *Faxonius eupunctus* (Rice

68 et al. 2018).

69 The aim of this study was to develop an MGB based qPCR assay to detect the presence of

70 the white-clawed crayfish, *A. pallipes* in environmental samples, and to test the reliability of

71 the assay by comparing the results with field observation data.

72              **Methods**

73 *Study sites and crayfish distribution data*

74 Eight sampling locations within seven different rivers were selected for field validation of the

75 assay. This research was carried out as part of a wider study assessing the impact of river

76 obstacles on Atlantic salmon *Salmo salar*. An obstacle, (weir, ford crossing or bridge apron),

77 was therefore present in the river at each sampling location and eDNA samples were

78 collected both above and below the structure. Out of the eight sites sampled, four were in

79 rivers with SAC status in which *A. pallipes* featured as a species of interest. Study rivers

within the River Barrow and River Nore SAC included the Delour and Dinin rivers and study

rivers within the Lower River Suir SAC included the Multeen and Duag rivers. Although the

Burren River is a tributary of the River Barrow, it does not have SAC status.

The presence of *A. pallipes* in these rivers was confirmed using data provided by the Irish

Environmental Protection Agency (EPA). These data are point data based on casual field

observations made of *A. pallipes* by EPA staff whilst undertaking routine biological

monitoring between the years 2007 and 2016 (W. Trodd, Irish Environmental Protection

Agency, personal communication). In addition, it was possible at some sites to confirm the

presence of crayfish by field observations made while electrofishing for *S. salar* and *S. trutta*.

For the purposes of this study, *A. pallipes* observations were extrapolated to the wider

subcatchment level. This meant that even if only one field observation of *A. pallipes*

occurred in a subcatchment, it was assumed that there were likely to be *A. pallipes* in the

rest of that subcatchment. This was considered the most meaningful way to present the

data, because the EPA were not specifically looking for *A. pallipes* when their observations

were made, so there is no certainty that *A. pallipes* did not occur in other parts of the

subcatchment.

eDNA qPCR assay development

Species-specific primers (forward primer: 5'- GGG TTA GTG GAG AGA GGG GT -3', and

reverse primer 5'- AAT CCC CAG ATC CAC AGA CG -3') and a 5'-6-FAM labelled TaqMan®

minor groove binding probe (5'- TCA GCT ATT GCC CAC GCA -3') for *A. pallipes*, which

targeted a locus within the mitochondrial cytochrome oxidase I (COI) region, were designed

in Primer Express 3.0 (Applied Biosystems-Roche, Branchburg, NJ). Including primers, the

total amplicon size was 96 base pairs. To verify the species specificity for the *A. pallipes*

assay in silico, probe and primer sequences were queried against the National Centre for

Biotechnology Information (NCBI - http://www.ncbi.nlm.nih.gov/) nucleotide database with

BLASTn (Basic Local Alignment Search Tool). The assay was tested in vitro with freshwater

species including brown trout *S. trutta*, sea lamprey *Petromyzon marinus*, Atlantic salmon *S.*

salar, allis shad *Alsoa alosa* and twaite shad *Alosa fallax*. The qPCR assay was optimized

using tissue extracted from *A. pallipes*.

eDNA collection, filtering and extraction

Water samples for eDNA analysis were collected in sterilized 2 L PET bottles from two

locations at each study site, above and below the river obstacle, and filtered on site using a

peristaltic pump. To avoid contamination, the water samples were taken in an upstream

direction, prior to any individuals entering the river. At each sampling location, three

replicate water samples were collected, one from each of the left margin, right margin and

the center of the river channel. In addition, one negative field control consisting of ddH_20

was also filtered. This resulted in a total number of six water samples for eDNA analysis and

two field controls collected per study site. All water samples were filtered through 47 mm

glass microfiber filters (1.5 μm). Filters were stored in 2.0 mL Eppendorf tubes and

subsequently frozen at -20˚C. To reduce contamination risk, all eDNA work was performed

in a dedicated Low Copy DNA laboratory. A modified CTAB (cetyltrimethylammonium

bromide) protocol (Möller et al. 1992) was employed for eDNA extractions. Briefly, one-half

of the filter was transferred to a 2.0 mL Eppendorf tube. A total volume of 750 μL CTAB

buffer (100 mM Tris-HCL, 20 mM EDTA, 1.4 M NaCl, 2% CTAB), and 7 μL of Proteinase K (20

mg mL^{-1}) were added to the tube. Samples were then vortexed for 10 seconds followed by

incubation at 56˚C for 2 hours, after which 750 μL of phenol/chloroform/isoamyl alcohol

(25:25:1 v/v) was added. The contents of the tube were manually mixed for 15 seconds and

subsequently centrifuged (11,000 x g, 20 min). A new tube contained 750 μL of

chloroform/isoamyl alcohol (24:1 v/v) was prepared, and the aqueous phase was transferred

to it. The manual mixing followed by centrifugation steps were repeated, and the aqueous

phase was again transferred to a new tube. The DNA was precipitated using one volume of

isopropanol alcohol that was added to the aqueous phase and incubated at -20˚C for 1 hour

followed by centrifugation (11,000 x g, 20 min). The resulting pellets were washed with 750

μL of 70% ethanol and centrifuged (11,000 x g, 5 min). The ethanol was removed, and care

was taken to ensure that the pellet remained in the tube. The tubes containing pellets were

dried in a heat block (50˚C, 5 min) followed by resuspension in molecular-grade ddH$_2$O.

eDNA assay deployment

An Applied Biosystems ViiA™ 7 (Life Technologies, Inc., Applied Biosystems, Foster City,

California, U.S.A.) quantitative thermocycler was used for eDNA amplification and

quantification. Briefly, the PCR profile consisted of 50°C for 2 min and 95°C for 10 min,

followed by 40 cycles between 95°C for 15 s and 60°C for 1 min. Standard curves were

generated from DNA originating from *A. pallipes* tissue extracts (quantified with NanoDrop®-

1000, Thermo Scientific, Wilmington, DE). This served as both a positive control and as a

means of calculating the concentration of *A. pallipes* eDNA in each sample. Seven 10:1 serial

dilutions ranging from 2.9 ng μL^{-1} to 2.9 x 10^{-6} ng μL^{-1} were used for the standard curve. Each

qPCR reaction was carried out in a total volume of 30 μL, consisting of 15 μL of TaqMan®

Environmental Master Mix 2.0 (Life Technologies, Applied Biosystems, Foster City, CA), 3 μL

of each primer (final concentration of 2 μM), probe (final concentration of 2 μM), 3 μL

eDNA/tissue extracted DNA template and 3 μL ddH$_2$O. Individual standard curves were

generated for each qPCR plate (y = -3.37x + 20.589, efficiency = 98.04 %, R^2 = 0.996 (1), y = -

3.442x + 20.392, efficiency = 95.22 %, R^2 = 0.999 (2) and y = -3.441x + 20.723, efficiency =

95.26 %, R^2 = 0.999 (3)). Each plate had 2 no-template controls (NTCs) to identify any

contamination that may have occurred during preparation. Standard curve, field and control

samples were quantified in triplicate (3 technical replicates). A positive detection was

defined as being when at least 2 out of the 3 technical replicates contained amplifiable DNA

with Cq differences not exceeding 0.5, and when the samples were within the detection

range of the standard curve. If 1 out of 3 technical replicates showed Cq differences

exceeding 0.5, the replicate was excluded from the study. In cases when the Cq value of 2

out of 3 technical replicates differed by more than 0.5 Cq, that specific dilution series or field

replicate was removed from further study. As *S. trutta* was present in all rivers, both

upstream and downstream of the obstacles, this species was used as a positive field control

to test for the presence of amplifiable DNA in all samples. The *S. trutta* assay from previously

published work (Gustavson et al. 2015) was used for this analysis. Three replicates per

location with 1 technical replicate were used.

164   **Results and Discussion**

The present study was successful in detecting *A. pallipes* in silico, in vitro and in situ. The

assay was highly sensitive and could detect *A. pallipes* eDNA concentrations as low as 0.002

ng L^{-1} at Cq 38.2 (average over 3 technical replicates, standard deviation 0.000059 ng L^{-1}). No

cross-species amplification occurred when the assay was tested using conventional PCR

visualized on agarose. Furthermore, zero amplification occurred in the NTCs and field

controls. While all field samples analysed yielded detectable DNA with the positive field

control (*S. trutta* assay), this was not the case for the *A. pallipes* assay. *A. pallipes* was only

detected in 5 out of 8 sites (Table 1, Fig. 1). Within the Burren, Duag and Multeen rivers,

positive detections were observed in all field replicates above and below the river obstacles.

At the Dinin (Cretty Yard) site, positive detection was observed in 2 out of 3 field replicates

above the obstacle, and 3 out of 3 field replicates below the obstacle. At the Dinin

(Castlecomer) site, positive detection of *A. pallipes* occurred in only 1 field replicate, which

was located above the obstacle (Table 1). The detection of *A. pallipes* eDNA above the

obstacle but not below it was a surprising result. Intuitively, one would expect that *A.*

pallipes would have been detected below the obstacle also, as eDNA travels downstream in

a river system. It is worth noting that it was at this site that the lowest concentration of *A.*

pallipes eDNA was detected (0.002 ng L^{-1}), and amplification of eDNA occurred in only 1 out

of 6 site replicates. It is possible that eDNA concentrations in the other field samples were

too low for a positive detection.

When compared with the EPA's *A. pallipes* observation data, the eDNA results reflected

what was observed in the field. In all subcatchments where *A. pallipes* was observed by EPA

staff, the eDNA results revealed a positive detection, and in all subcatchments where *A.*

pallipes was not observed by EPA staff, the eDNA results revealed a negative detection (Fig.

1). Furthermore, it is worth noting that the eDNA results also reflected the expected

distribution of *A. pallipes* based on environmental conditions. In all sites where *A. pallipes*

was not detected by either field observation or eDNA (Delour, Brown's Beck Brook and the

Dalligan), the dominant bedrock was siliceous. *A. pallipes* are known to occur in waterbodies

with underlying calcareous bedrock rather than siliceous bedrock (Lucey and McGarrigle

1987).

It was not possible in this study to ascertain the impact of the river obstacles on *A. pallipes*

using eDNA. This was because in most sites, *A. pallipes* were either detected both above and

below the obstacle, or as observed at the Dinin (Castlecomer) site, above the obstacle only.

However, studies where crayfish have been physically marked or tagged have shown that

river obstacles can hinder their upstream migration (Kerby et al. 2005, Bubb et al. 2008,

Rosewarne et al. 2013). Environmental DNA assays for diadromous species such as sea

lamprey *P. marinus* (Gustavson et al. 2015) or Chinook salmon *Oncorhynchus tshawytscha*

(Laramie et al. 2015) are likely to be better indicators for assessing the migratory impact of

river obstacles. Alternatively, eDNA could be used for monitoring whether invasive species

have moved above the barriers designed to block them (Cowart et al. 2018). Nonetheless,

the present research shows that eDNA is a reliable tool for detecting *A. pallipes* in river

systems. Furthermore, while the focus of this study was in lotic systems, the assay could be

readily deployed in lentic systems such as lakes or reservoirs where field sampling may be

more challenging.

To conclude, this assay provides an alternative, rapid method for locating *A. pallipes* in both

lentic and lotic ecosystems, allowing for more targeted, efficient surveying strategies.

Considering the current crayfish plague outbreak in Ireland (National Biodiversity Data

Centre 2018), and indeed other countries where it is present, there is an urgent need for a

non-invasive sampling methodology that will minimise further spread of the pathogen. Large

scale deployment of this assay would enable managers to rapidly establish, with confidence,

the distribution of *A. pallipes*, which would support the designation of new conservation

areas for the species.

Acknowledgements

This research was funded by the Irish Environmental Protection Agency (Reconnect project -

2015-W-LS-8). The authors would also like to thank the Irish Environmental Protection

Agency for providing the data on *A. pallipes* distributions.

220   **Literature Cited**

Agersnap, S., W. B. Larsen, S. W. Knudsen, D. Strand, P. F. Thomsen, M. Hesselsøe, P. B.

Mortensen, T. Vrålstad, and P. R. Møller. 2017. Monitoring of noble, signal and narrow-

clawed crayfish using environmental DNA from freshwater samples. PLoS ONE 12:1–22.

Boothroyd, M., N. E. Mandrak, M. Fox, and C. C. Wilson. 2016. Environmental DNA (eDNA)

detection and habitat occupancy of threatened spotted gar (*Lepisosteus oculatus*). Aquatic

Conservation: Marine and Freshwater Ecosystems 26:1107–1119.

Bubb, D. H., T. J. Thom, and M. C. Lucas. 2008. Spatial ecology of the white-clawed crayfish in

an upland stream and implications for the conservation of this endangered species. Aquatic

Conservation: Marine and Freshwater Ecosystems 18:647–657.

Byrne, C., and J. Lynch. 1999. A sampling strategy for stream populations of white-clawed

crayfish, *Austropotamobius pallipes* (Lereboullet)(Crustacea, Astacidae). Biology and

Environment: Proceedings of the Royal Irish Academy 99b:89–94.

Cai, W., Z. Ma, C. Yang, L. Wang, W. Wang, G. Zhao, Y. Geng, and D. W. Yu. 2017. Using eDNA

to detect the distribution and density of invasive crayfish in the HongheHani rice terrace

World Heritage site. PLoS ONE 12:1–13.

Carlsson, J. E. L., D. Egan, P. C. Collins, E. D. Farrell, F. Igoe, and J. Carlsson. 2017. A qPCR

MGB probe based eDNA assay for European freshwater pearl mussel (*Margaritifera*

margaritifera L.). Aquatic Conservation: Marine and Freshwater Ecosystems 27:1341-1344.

Cowart, D. A., K. G. H. Breedveld, M. J. Ellis, J. M. Hull, and E. R. Larson. 2018. Environmental

DNA (eDNA) applications for the conservation of imperiled crayfish (Decapoda: Astacidea)

through monitoring of invasive species barriers and relocated populations. Journal of

Crustacean Biology doi:10.1093/jcbiol/ruy007.

Demers, A., J. Lucey, M. L. McGarrigle, and J. D. Reynolds. 2005. The distribution of the

white-clawed crayfish, *Austropotamobius pallipes*, in Ireland. Biology & Environment:

Proceedings of the Royal Irish Academy 105b:65–69.

Demers, A., and J. D. Reynolds. 2002. A survey of the white-clawed crayfish,

Austropotamobius pallipes (Lereboullet), and of water quality in two catchments of eastern

Ireland. Bulletin Francais de Pêche et de Pisciculture 367:729–740.

DiStefano, R. J., C. M. Gale, B. A. Wagner, and R. D. Zweifel. 2003. A sampling method to

assess lotic crayfish communities. Journal of Crustacean Biology 23:678–690.

Dougherty, M. M., E. R. Larson, M. A. Renshaw, C. A. Gantz, S. P. Egan, D. M. Erickson, and D.

M. Lodge. 2016. Environmental DNA (eDNA) detects the invasive rusty crayfish *Orconectes*

rusticus at low abundances. Journal of Applied Ecology 53:722–732.

Fujiwara, A., S. Matsuhashi, H. Doi, S. Yamamoto, and T. Minamoto. 2016. Use of

environmental DNA to survey the distribution of an invasive submerged plant in ponds.

Freshwater Science 35:748–754.

Goldberg, C. S., A. Sepulveda, A. Ray, J. Baumgardt, and L. P. Waits. 2013. Environmental

DNA as a new method for early detection of New Zealand mudsnails (*Potamopyrgus*

antipodarum). Freshwater Science 32:792–800.

Grandjean, F., C. Souty-Grosset, R. Raimond, and D. M. Holdich. 1997. Geographical variation

of mitochondrial DNA between populations of the white-clawed crayfish *Austropotamobius*

pallipes. Freshwater Biology 37:493–501.

Gustavson, M. S., P. C. Collins, J. A. Finarelli, D. Egan, R. Ó. Conchúir, G. D. Wightman, J. J.

King, D. T. Gauthier, K. Whelan, J. E. L. Carlsson, and J. Carlsson. 2015. An eDNA assay for

Irish *Petromyzon marinus* and *Salmo trutta* and field validation in running water. Journal of

Fish Biology 87:1254–1262.

Holdich, D. M., J. D. Reynolds, C. Souty-Grosset, and P. J. Sibley. 2009. A review of the ever

increasing threat to European crayfish from non-indigenous crayfish species. Knowledge and

Management of Aquatic Ecosystems 394-395:11.

Jerde, C. L., A. R. Mahon, W. L. Chadderton, and D. M. Lodge. 2011. "Sight-unseen" detection

of rare aquatic species using environmental DNA. Conservation Letters 4:150–157.

Kerby, J. L., S. P. D. Riley, L. B. Kats, and P. Wilson. 2005. Barriers and flow as limiting factors

in the spread of an invasive crayfish (*Procambarus clarkii*) in southern California streams.

Biological Conservation 126:402–409.

Laramie, M. B., D. S. Pilliod, and C. S. Goldberg. 2015. Characterizing the distribution of an

endangered salmonid using environmental DNA analysis. Biological Conservation 183:29–37.

Larson, E. R., M. A. Renshaw, C. A. Gantz, J. Umek, S. Chandra, D. M. Lodge, and S. P. Egan.

2017. Environmental DNA (eDNA) detects the invasive crayfishes *Orconectes rusticus* and

Pacifastacus leniusculus in large lakes of North America. Hydrobiologia 800:173–185.

Lucey, J., and M. L. McGarrigle. 1987. The distribution of the crayfish *Austropotamobius*

pallipes (Lereboullet) in Ireland. Irish Fisheries Investigations A29:1–13.

Lyons, R., and M. Kelly-Quinn. 2003. An investigation into the disappearance of

Austropotamobius pallipes (Lereboullet) populations in the headwaters of the Nore River,

Ireland and the correlation to water quality. Bulletin Français de la Pêche et de la Pisciculture

370-371:139–150.

Mächler, E., K. Deiner, P. Steinmann, and F. Altermatt. 2014. Utility of environmental DNA

for monitoring rare and indicator macroinvertebrate species. Freshwater Science 33:1174–

1183.

Matthews, M., and J. D. Reynolds. 1992. Ecological impact of crayfish plague in Ireland.

Hydrobiologia 234:1–6.

Möller, E. M., G. Bahnweg, H. Sandermann, and H. H. Geiger. 1992. A simple and efficient

protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and

infected plant tissues. Nucleic Acids Research 20:6115–6116.

National Biodiversity Data Centre. 2018. Crayfish plague. (Available from:

http://www.biodiversityireland.ie/projects/invasive-species/crayfish-plague/)

Reynolds, J. D., W. O'Connor, C. O'Keeffe, and D. Lynn. 2010. A technical manual for

monitoring white-clawed crayfish *Austropotamobius pallipes* in Irish lakes. (Available at:

https://www.npws.ie/sites/default/files/publications/pdf/IWM45.pdf)

Rice, C. J., E. R. Larson, and C. A. Taylor. 2018. Environmental DNA detects a rare large river

crayfish but with little relation to local abundance. Freshwater Biology 63:443–455.

Rosewarne, P. J., A. T. Piper, R. M. Wright, and A. M. Dunn. 2013. Do low-head riverine

structures hinder the spread of invasive crayfish? Case study of signal crayfish (*Pacifastacus*

leniusculus) movements at a flow gauging weir. Management of Biological Invasions 4:273–

282.

Sigsgaard, E. E., H. Carl, P. R. Møller, and P. F. Thomsen. 2015. Monitoring the near-extinct

European weather loach in Denmark based on environmental DNA from water samples.

Biological Conservation 183:46–52.

Taberlet, P., E. Coissac, M. Hajibabaei, and L. H. Rieseberg. 2012. Environmental DNA.

Molecular Ecology 21:1789–1793.

Thomsen, P. F., and E. Willerslev. 2015. Environmental DNA - An emerging tool in

conservation for monitoring past and present biodiversity. Biological Conservation 183:4–18.

Vaeßen, S., and H. Hollert. 2015. Impacts of the North American signal crayfish (*Pacifastacus*

leniusculus) on European ecosystems. Environmental Sciences Europe 27:1–6.

Table 1. Environmental DNA concentrations (ng L^{-1}) from the *A. pallipes* assay at each study river. Concentrations are based the average eDNA concentration in a minimum of 2 technical replicates and between 1 and 3 field replicates. Confidence intervals (95%) are given, where possible, for each location (upstream or downstream of the river obstacle).

River	Location	No. eDNA samples collected	No. of positive detections	eDNA concentration ng L^{-1} (95% CI)	Visual observation
Burren	Upstream	3	3	0.027 (0.01-0.02)	+
	Downstream	3	3	0.02 (0.01-0.04)	
Duag	Upstream	3	3	0.027 (0.005-0.05)	+
	Downstream	3	3	0.157 (0.09-0.23)	
Multeen	Upstream	3	3	0.034 (0.02-0.04)	+
	Downstream	3	3	0.06 (0.04-0.08)	
Dinin (Cretty Yard)	Upstream	3	2	0.011 (0.004-0.02)	+
	Downstream	3	3	0.006 (0.002-0.01)	
Dinin (Castlecomer)	Upstream	3	1	0.002 [a]	+
	Downstream	3	0	0	
Delour	Upstream	3	0	0	-
	Downstream	3	0	0	
Dalligan	Upstream	3	0	0	-
	Downstream	3	0	0	
Brown's Beck Brook (BBB)	Upstream	3	0	0	-

Downstream	3	0	0

[a] It was not possible to calculate a 95% confidence interval for this site as a positive detection was

only detected in one out of three site replicates

Figure Heading

Figure 1. Map showing the sampling locations for the eDNA analysis. Crosses indicate sites

where a positive detection of *A. pallipes* was made in at least one field replicate. Circles

indicate sites where no positive detection of *A. pallipes* was made in all field replicates. River

subcatchments in which field observations of *A. pallipes* were made between the years 2007

and 2016 are highlighted in grey.