SNAP23 is required for the maintenance of meiotic arrest and cortical granule exocytosis in mouse oocytes.

Running title: Exocytosis in mouse oocytes and eggs

Summary: The SNARE protein, SNAP23, is required to maintain gap junction communication between the oocyte and follicle cells that is needed to maintain oocyte meiotic arrest, as well as for cortical granule exocytosis at fertilization.

Keywords: fertilization, gamete biology, meiotic arrest, oocyte, oocyte follicle interactions, oocyte maturation, zona pellucida

Lisa M. Mehlmann,[1,2] Tracy F. Uliasz,[1] and Katie M. Lowther[1]

[1] Department of Cell Biology, UConn Health, Farmington, CT 06030. Funding for this project provided by the National Institutes of Health R56HD056366. Presented in part at the 5[th] International Conference on Oocyte Biology, 16-20 January 2019, Villefranche-sur-Mer, France.

[2] Address correspondence to Lisa Mehlmann, Department of Cell Biology, UConn Health, 263 Farmington Ave., Farmington, CT 06030. lmehlman@uchc.edu. Phone 860-679-2703; Fax 860-679-1269.

ABSTRACT

Mammalian oocytes are stored in the ovary for prolonged periods, arrested in meiotic prophase. During this period, their plasma membranes are constantly being recycled by endocytosis and exocytosis. However, the function of this membrane turnover is unknown. Here, we investigated the requirement for exocytosis in the maintenance of meiotic arrest. Using a newly developed method for rapidly and specifically depleting proteins in oocytes, we have identified the SNARE protein, SNAP23, to be required for meiotic arrest. Degradation of SNAP23 causes premature meiotic resumption in follicle-enclosed oocytes. The reduction in SNAP23 is associated with loss of gap junction communication between the oocyte and surrounding follicle cells. Reduction of SNAP23 protein also inhibits cortical granule exocytosis in response to a Ca^{2+} stimulus, as measured by lectin staining and cleavage of ZP2. Our results show an essential role for SNAP23 in two key processes that occur in mouse oocytes and eggs.

INTRODUCTION

Mammalian oocytes arrest at prophase of the first meiosis for an extended period of time, during which

they accumulate RNAs and proteins that enable them to sustain the development of an early embryo.

During this period of meiotic arrest, they constitutively secrete factors such as GDF9, BMP15, and

FGF8, which are critical for proper follicle development [1, 2]. In return, the follicle cells provide the

oocyte with substances needed for protein synthesis and metabolism, cholesterol, and importantly, the

cGMP that is necessary for the maintenance of meiotic arrest prior to the LH surge [2-5]. Prior to the LH

surge, oocytes undergo robust endocytosis that probably plays roles in the uptake of extracellular

contents and that also seems to be involved in cell signaling during meiotic arrest, as blocking receptor-

mediated endocytosis in mouse oocytes prevents oocyte maturation by elevating cAMP levels [6].

Constitutive exocytosis is probably also required to replenish lipids and membrane proteins that are

taken up during constitutive endocytosis.

During meiotic maturation, the oocyte undergoes numerous cytoplasmic changes that are required for

successful embryonic development after fertilization [7]. One change that occurs is the ability to

undergo cortical granule exocytosis in response to a large release of intracellular Ca^{2+} at fertilization [8].

Cortical granules are vesicles that reside in the cortical region of the mature egg opposite the meiotic

spindle [9, 10]. Upon exocytosis, the enzyme ovastacin, contained in the granules, cleaves the zona

pellucida protein ZP2 and renders the zona pellucida impermeable to additional sperm [11]. Thus,

cortical granule exocytosis is an important polyspermy preventing mechanism. Interestingly, immature

mouse oocytes are unable to undergo cortical granule exocytosis even when stimulated by Ca^{2+} [8, 12],

so the ability to undergo exocytosis develops during maturation.

The proteins required for exocytosis have not been well studied in mammalian oocytes or eggs. (For

simplicity, we will refer to immature, prophase-arrested oocytes herein as "oocytes," and mature,

metaphase II-stage oocytes as "eggs"). It is likely that the process is similar to exocytic events that

occur in other types of secretory cells. For example, intracellular Ca^{2+} triggers exocytosis of

neurotransmitters in neurons [13-15], and insulin receptor activation causes exocytosis of the GLUT4

transporter to the plasma membrane in adipocytes [16, 17]. These exocytic events are mediated by

SNARE (Soluble *N*-ethylmaleimide-sensitive factor attachment protein receptor) proteins in which two t-

SNARES (associated with the target membrane) form a complex with one v-SNARE (associated with

the vesicle destined for fusion; Fig. 1). The t-SNAREs are a syntaxin and either SNAP25 or SNAP23

and the v-SNARE is a vesicle-associated membrane protein (VAMP, also called synaptobrevin).

SNARE proteins contain coiled coil domains that form a very stable, parallel 4 helix bundle. One helix

is contributed by syntaxin, one by VAMP, and two by SNAP25/SNAP23.   Formation of the SNARE

complex allows the vesicle and target membranes to come into close contact and fuse, releasing the

contents of the vesicles to the extracellular space.

Cortical granule exocytosis has been studied to some extent in eggs of a variety of species. Sea urchin

eggs contain homologs of mammalian secretory proteins [18], and inhibition of the SNARE

synaptobrevin (VAMP) inhibits exocytosis [19]. In porcine eggs, cortical granules associate with

SNAP23, syntaxin 2, and VAMP1 [20] but the function of these SNARE proteins has not been

examined in detail. It was reported previously that SNAP25 is required for cortical granule exocytosis in

mouse eggs [21], based on the finding that botulinum toxin, which cleaves SNAP25 [22], inhibits

cortical granule exocytosis. Syntaxin 4 was identified in mouse eggs, but no function was investigated

[23]. Other SNARE proteins have not been identified, but proteins in the Rab family have been

implicated in cortical granule exocytosis. In particular, Rabs 3A and 27A are of interest based on their

co-localization with cortical granules [24], and on the findings that microinjection of function-blocking

antibodies [24] or protein depletion using RNAi [25] inhibits cortical granule exocytosis [24, 25]. In

addition to Rab proteins, other key players in the SNARE pathway, α-SNAP and NSF (N-

ethylmaleimide sensitive factor) [26] appear to affect cortical granule exocytosis. In this study, we

investigated the role of the SNARE protein, SNAP23, in constitutive and regulated exocytosis in mouse

oocytes and eggs, respectively. We used a newly developed method, "Trim-away," to identify SNAP23

as essential for both of these processes.

MATERIALS AND METHODS

Media and Reagents

Except where noted, all chemicals were obtained from Millipore Sigma (St. Louis, MO). The medium

used to collect oocytes was MEMα (Gibco 12000022, Thermo Fisher, Waltham, MA), supplemented

with 20 mM HEPES, 75 μg/ml penicillin G, 50 μg/ml streptomycin, 0.1% polyvinyl alcohol (PVA), and 10

μM milrinone to inhibit spontaneous meiotic resumption. For in vitro maturation, oocytes were

transferred to bicarbonate-buffered MEMα, in which the HEPES was replaced with 25 mM sodium

bicarbonate and the PVA was replaced with 5% fetal bovine serum (Invitrogen). Medium was

equilibrated with 5% CO_2 and 95% air. The medium used for follicle dissection and microinjection was

bicarbonate-buffered MEMα supplemented with 3 mg/ml bovine serum albumin, 5 μg/ml insulin, 5 μg/ml

transferrin, 5 ng/ml selenium, and 10 ng/ml follicle stimulating hormone (FSH; provided by A.F. Parlow,

National Hormone and Peptide Program, Torrance, CA).

DNA constructs and antibodies

Fluorescently tagged TRIM21 constructs (TRIM21-GFP and TRIM21-RFP) were obtained from Melina

Schuh (Max Planck Institute for Biophysical Chemistry; [27]). To make mRNA, the constructs were

linearized with XbaI and in vitro transcribed and polyadenylated using the mMessage mMachine T7

Ultra kit (Thermo Fisher). Wild type (SNAP25WT) and dominant negative (SNAP25Δ20) SNAP25

constructs were obtained from Khaled Machaca (Weill Cornell Medicine Qatar) and mRNA was

synthesized after linearizing the plasmids with NheI. SNAP23-GFP was purchased from OriGene and

subcloned into the SmaI site of the pIVT vector (provided by Carmen Williams, NIEHS)[28]. mRNA was

synthesized after linearizing the plasmid with NdeI.

The following antibodies were used: ZP2 was obtained from Jurrien Dean (NIDDK) [29]; SNAP25 was

from BD Biosciences (San Jose, CA) #610366; SNAP23 used for the blot in Fig. 2 was from Synaptic

Systems (Göttingen, Germany) #110 203, and for all other experiments was from Fisher (#PA1-738);

affinity purified antibody against the IP_3 receptor was provided by James Watras (UConn Health).

Rabbit IgG was from Sigma (#I5006). Horseradish peroxidase-conjugated secondary antibodies were

from Santa Cruz (Santa Cruz, CA) or Advansta Inc. (Menlo Park, CA).

Mouse follicle and oocyte isolation, culture, and microinjection

All experiments were done with prior approval of the Animal Care and Use Committee at UConn

Health. Preantral or antral stage follicles that were ~140-180 μm or ≥290 μm in diameter, respectively,

were manually dissected from ovaries of 23-26-day-old B6SJLF1 mice (Jackson Laboratories, Bar

Harbor, ME) as described previously [30]. Following isolation, follicles were plated on Millicell culture

plate inserts (PICMORG50, Millipore, Billerica, MA) and were cultured in a humidified atmosphere at

37° with 5% CO_2, 95% air. After a 3 hr culture period, follicle-enclosed oocytes were examined under

an upright microscope for the presence of a germinal vesicle (GV). Only follicles containing oocytes

with readily visible GVs were selected for use in these experiments. Fully grown, GV-stage oocytes

were obtained from the ovaries of 6-12-week-old NSA (CF1) mice (Envigo, South Easton, MA).

Cumulus cells were removed using a small-bore pipet.

Quantitative microinjection of follicle-enclosed oocytes was performed as described previously [30].

Four-5 follicles were loaded into a microinjection chamber between two coverslips and follicles were

removed and re-plated onto Millicells following each set of injections. In some cases, follicles were

removed from Millicells, reloaded into microinjection chambers, and the oocytes were injected with

Alexa488 (Thermo Fisher) to visualize dye transfer from the oocyte to the surrounding follicle cells.

Isolated oocytes were injected as described previously [6, 31]. Final injected concentrations were

calculated based on an oocyte volume of 200 pl.

Trim-away

Oocytes were injected with RNA encoding fluorescently tagged TRIM21 (10 pl of 0.3 μg/μl), then were

injected ~2-4 hrs later with specific antibodies (~25-50 μg/ml final injected amount, depending on the

antibody). In some cases, oocytes were co-injected with the RNA and the antibody. Purified IgG was

used for controls. Prior to injection, antibodies/IgGs were spin-dialyzed into PBS and were

concentrated using either Amicon Ultra - 0.5 ml, 10 kDa centrifugal filters (Millipore) or Vivaspin 500 50

kDa cutoff filters (GE Healthcare). Concentrated antibody/IgG solutions were diluted into PBS

containing 0.05% NP-40 to reduce stickiness [27, 32].

Western blotting

Oocytes and eggs were counted and washed into HEPES-buffered MEMα to remove serum, then were

transferred to 0.5 ml microcentrifuge tubes in a small volume of medium (1-2 μl). 15 μl of 1X Laemmli

sample buffer was added to each tube, and samples were stored at -20° until use. Western blotting

was performed as previously described [33]. Proteins were separated in 4-20% polyacrylamide gels

(Bio-Rad, Hercules, CA). Electrophoresed proteins were transferred to nitrocellulose or PVDF

membranes and were blocked either in TBST containing 5% nonfat dry milk or 5% BSA, or in Advansta

block (Advansta). Blots were incubated with primary antibodies either overnight at 4° or for 2 hrs at

room temperature, then with horseradish peroxidase-conjugated secondary antibodies. Blots were

developed using either ECL Prime (GE Healthcare, Piscataway, NJ) or WesternBright Sirius

horseradish peroxidase substrate (Advansta), using a charge-coupled device camera (G:box Chemi

XT4; Syngene (Frederick, MD).

Immunofluorescence and lectin staining

Eggs were fixed for 30 min in PBS containing 2% formaldehyde, permeabilized for 15 min in PBS

containing 0.1% PVA and 0.1% Triton X-100), and blocked for at least 15 min in PBS containing 3%

BSA and 0.01% Triton X-100. Eggs were incubated in primary antibodies, diluted in blocking buffer,

overnight at 4°C. After washing in PBS + PVA, eggs were incubated in Alexa488-conjugated

secondary antibodies for 1-2 hrs at room temperature. Eggs were washed in PBS + PVA, then were

imaged using a Zeiss Pascal confocal microscope using a 40X 1.2 NA water immersion objective (C-

Apochromat; Carl Zeiss MicroImaging, Inc., Thornwood, NY).

To label cortical granules, proteins were depleted using Trim-away as outlined above. After

RNA/antibody injections, zonae pellucidae were removed with 5 mg/ml pronase in HEPES-buffered

MEMα. Oocytes were matured overnight in bicarbonate buffered MEMα with 5% FBS. After

maturation, eggs were imaged to confirm that they were fluorescent, demonstrating that the TRIM21

protein was expressed. Eggs were then treated with 100 μM thimerosal (ICN Biochemicals, Cleveland,

OH) for 45 min in HEPES-buffered MEMα to stimulate intracellular Ca^{2+} oscillations (Mehlmann and

Kline, 1994), fixed for 30 min in PBS containing 0.1% PVA and 2% formaldehyde, blocked 30 min in

PBS containing 3 mg/ml BSA and 0.1 M glycine, permeabilized 20 min in PBS-PVA containing 0.1%

Triton X-100, washed through 3 washes PBS-PVA, incubated 30 min in 50 μg/ml TRITC-Lens culinaris

agglutinin (LCA) in PBS-PVA, washed 3 times in PBS-PVA, then imaged with the 40X 1.2 NA lens on

the confocal microscope.

Ca^{2+} imaging

SNAP23 was depleted in oocytes using Trim-away (using RFP-tagged TRIM21) and injected oocytes

were matured overnight. Uninjected eggs were used as a control. Mature eggs were loaded with

Fluo3-AM (Molecular Probes) for 30 min per the manufacturer's instructions, using PowerLoad to aid in

the uptake of the dye and probenecid to inhibit intracellular compartmentalization. After dye loading,

eggs were adhered to glass coverslips in 90 μl drops of HEPES buffered medium containing no PVA.

Five-6 eggs were imaged simultaneously using a 20X, 0.5 NA lens on the confocal microscope. After

obtaining a basal Ca^{2+} level, Ca^{2+} was recorded at one scan per second during addition of 100 μM

thimerosal (10 μl of 1 mM stock to ensure rapid mixing). Following recordings, a region of interest was

measured across a stacked image in ImageJ, values were entered into Excel, and results are

presented as the amplitude of each measurement (F) subtracted by the baseline (F_0) and divided by

the baseline signal [(F-F_0)/F_0].

RESULTS AND DISCUSSION

SNAP23, but not SNAP25, is expressed in mouse oocytes and eggs.

A previous study reported that botulinum neurotoxin A (BoNT/A) significantly inhibits cortical granule

exocytosis when injected into mouse eggs [21]. Based on this observation, and on western blots

showing expression of SNAP25 in eggs, the authors concluded that SNAP25, which is mainly

expressed throughout the nervous system [34, 35], is essential for cortical granule exocytosis.

However, BoNT/A can also cleave the closely related SNARE, SNAP23, in vitro [36] and based on the

high degree of similarity between SNAP25 and SNAP23, it is possible that antibodies might crossreact.

To investigate which SNAPs are expressed in mouse oocytes and eggs, we first tried to confirm

expression of SNAP25 using western blot. We were unable to detect SNAP25 in oocytes or eggs,

using 3 different antibodies against SNAP25 and relatively high amounts of oocyte and egg lysates

(Fig. 2A). In contrast, the closely related and ubiquitously expressed SNAP23 was abundantly

expressed (Fig. 2B). We confirmed the identity of SNAP23 by targeting the mRNA encoding it by

microinjection of siRNA. The SNAP23 siRNA reduced SNAP23 protein to 44% of the control after a 29

hr incubation following microinjection (Fig. 2B).

We localized SNAP23 in oocytes and eggs using immunofluorescence and a SNAP23-GFP fusion

protein. In oocytes, immunofluorescence showed the protein mainly concentrated at the egg cortex,

with some interior staining and staining around the nucleolus, whereas SNAP23-GFP showed staining

primarily at the cortex and around the GV (Fig. 2C). In eggs, both immunofluorescence and SNAP23-

GFP showed a thin band of SNAP23 around the entire egg cortex, with some interior staining as well.

Generally there was staining above the meiotic spindle, but in some cases there was a region near the

spindle that had less staining. This area was much smaller than the cortical granule-free area that

occurs in the vicinity of the meiotic spindle (see Fig. 5). These results demonstrate that SNAP23,

rather than SNAP25, is expressed in oocytes and eggs and show that it is located at an appropriate

region of the oocyte/egg to be involved in exocytosis.

Depleting SNAP23 stimulates meiotic resumption in follicle-enclosed oocytes.

To test a role for SNAP23 in constitutive exocytosis, we depleted it in follicle-enclosed oocytes.

Depletion of SNAP23 in oocytes using siRNA was incomplete, as levels were only reduced to ~44%

after a two day culture period (Fig. 2B). Therefore, we depleted SNAP23 using a newly developed

method called "Trim-away" [27]. This method utilizes TRIM21, an intracellular antibody receptor and E3

ubiquitin ligase. When TRIM21 is expressed in oocytes or eggs followed by injection of a specific

antibody against the protein of interest, TRIM21 binds to the antibody and recruits the protein/antibody

complex to the proteasome, causing rapid degradation [37]. Using this method, we found that SNAP23

protein was reduced to a nearly undetectable level within ~3 hrs after injecting SNAP23 antibody into

TRIM21-expressing oocytes (Fig. 3A). To test for specificity of the protein reduction, we also blotted for

the IP_3 receptor, which is highly abundant and can be detected on western blots using only a few

oocytes or eggs. There was no reduction in IP_3 receptor protein (Fig. 3A). SNAP23 depletion using

Trim-away caused germinal vesicle breakdown (GVBD) in almost 80% of follicle-enclosed oocytes after

an ~18 hr culture period, whereas control oocytes that were either uninjected or that were injected with

TRIM21/IgG did not undergo GVBD (Fig. 3B).

SNAP23 depletion disrupts communication between the oocyte and somatic cells.

Because SNAP23 depletion caused spontaneous maturation in follicle-enclosed oocytes, it is possible

that inhibiting exocytosis disrupts communication between the oocyte and surrounding granulosa cells.

To test this possibility, we first used Trim-away to deplete SNAP23 in follicle-enclosed oocytes, then

injected oocytes with a low molecular weight fluorescent dye, Alexa488, that readily passes through

gap junctions [38]. We used preantral follicles for these experiments due to the higher survival rate

than we achieve using antral follicles. The Alexa488 dye spread throughout the granulosa cells in 70%

(7/10) of control follicles (Fig. 3C,D). In contrast, the dye did not spread in any of 9 follicles in which

SNAP23 was depleted (Fig. 3C,D).

The simplest explanation for the disconnection of gap junction communication between the oocyte and

follicle cells when exocytosis is inhibited is that connexin 37 (Cx37), the protein that comprises gap

junctions between the oocyte and cumulus cells [39], cannot be replaced in the plasma membrane

following endocytosis. The half life of Cx37 is on the order of hours [40], so it needs to be continuously

exocytosed to the plasma membrane to form gap junctions. In the absence of gap junctions, the

diffusion of cGMP from the cumulus cells to the oocyte would be stopped and this would activate

PDE3A in the oocyte to lower cAMP levels to the point that the oocyte would no longer be able to

maintain meiotic arrest [3, 41].

Confirmation of the requirement of SNAP23 for maintenance of meiotic arrest, using a dominant

negative approach.

We used a dominant negative approach as a second way to inhibit SNAP23. We expressed a mutated

form of SNAP25 (SNAP25Δ20), in which 20 amino acids in the C-terminus were deleted. This mutation

disrupts the coiled-coil domain in the C-terminus which is required for the formation of the SNARE

complex [42], and inhibits exocytosis and stimulates meiotic resumption in *Xenopus* oocytes [42, 43].

Although SNAP25 was not identified in mouse oocytes, SNAP25Δ20 is likely to act as a dominant

negative SNAP in oocytes because SNAP25 and SNAP23 can complex with the same t- and v-

SNAREs [44, 45]. Both forms of SNAP25 were expressed in injected mouse oocytes (Fig. 4A).

Expressing SNAP25Δ20 in follicle-enclosed oocytes robustly stimulated meiotic resumption (Fig. 4B),

with nearly 100% of oocytes undergoing GVBD after an ~18 hr culture. Uninjected oocytes, or oocytes

injected with SNAP25WT did not resume meiosis during the culture period (Fig. 4B). These results

using both Trim-away and a dominant negative SNARE support the hypothesis that SNAP23 is

essential for constitutive exocytosis in mouse oocytes, and that constitutive exocytosis is needed to

maintain meiotic arrest.

As discussed above, gap junctions are likely to be affected by depletion of SNAP23 and this probably

inhibits the transfer of cGMP from the follicle cells to the oocyte, which causes meiotic resumption.

Another possibility to account for meiotic resumption in the oocyte when exocytosis is blocked is that

the oocyte cannot secrete factors such as GDF9 and BMP15 that act on the cumulus cells to maintain

follicle integrity. This could include the continued synthesis of connexins on the follicle side. It is also

possible that GPR3, the receptor needed for cAMP production in the oocyte [46-48], must be placed at

the plasma membrane in order to signal to make cAMP, and in the absence of GPR3 at the surface the

oocyte can no longer produce cAMP. Finally, it is possible that the oocyte secretes a ligand that

stimulates GPR3 and in the absence of this ligand the oocyte can no longer produce cAMP. GPR3 is

thought to be constitutively active and no ligand has yet been identified [49, 50], but there is still a

possibility that the oocyte produces and secretes one.

.

SNAP23 is required for cortical granule exocytosis.

During maturation, oocytes gear up for the regulated exocytic event of cortical granule exocytosis.

Because SNAP23 participates in constitutive and regulated exocytosis in other cell types [51, 52], we

investigated if it is required for cortical granule exocytosis in mouse eggs using Trim-away. For these

experiments, we co-injected immature oocytes with RNA encoding TRIM21 and an antibody against

SNAP23 and matured the oocytes for ~18 hrs. Expression of TRIM21 and the SNAP23 antibody did

not inhibit oocyte maturation and the mature eggs appeared healthy and formed first polar bodies.

After maturation, we examined cortical granule exocytosis after providing a Ca^{2+} stimulus. We initiated

Ca^{2+} release in eggs using the sulfhydryl reagent, thimerosal, which produces a rapid series of long

lasting, repetitive Ca^{2+} transients in mouse oocytes and eggs [53].

We detected cortical granule exocytosis following thimerosal treatment using two methods. In the first

method, we removed the zonae pellucidae following injection of RNA and antibody, prior to overnight

oocyte maturation. We then treated mature eggs with 100 μM thimerosal for 45 min and stained

cortical granules with a fluorescent lectin [26]. Before thimerosal treatment, we detected cortical

granules at the cortex opposite the meiotic spindle in untreated eggs in all 3 groups (Fig. 5A, left

panels). The cortical granule-free domain was sometimes quite large, comprising around 50% of the

oocyte cortex. This correlates with a recent study showing that cortical granule distribution and function

in in vitro matured eggs is reduced compared to in vivo matured eggs [54]. Following thimerosal

treatment, cortical granules were either completely absent or were greatly reduced and appeared to be

more dispersed around the surface of uninjected and control-injected eggs (Fig. 5A, right panels),

demonstrating that under our conditions, cortical granule exocytosis occurs in in vitro matured eggs. In

contrast, cortical granules were retained in the cortex of SNAP23 antibody-injected eggs, showing that

exocytosis was blocked in these eggs (Fig. 5A).

In the second method, we left the zonae intact following microinjection and overnight culture. We

treated eggs with thimerosal and examined ZP2 cleavage using western blot. As shown in Fig. 5B,

uninjected eggs that were not treated with thimerosal contained the uncleaved form of ZP2. The band

shifted following thimerosal treatment, demonstrating that ZP2 was cleaved. Likewise, ZP2 was

cleaved in TRIM21-expressing, control IgG-injected eggs that were treated with thimerosal. ZP2 was

not cleaved, however, in TRIM21-expressing, SNAP23 antibody-injected eggs treated . Probing the

ZP2 blot with SNAP23 antibody showed that SNAP23 protein was barely detectable in antibody-

injected eggs (Fig. 5B). SNAP23 depletion did not prevent Ca^{2+} release in response to thimerosal

treatment (Fig. 5C), and the peak amplitudes were similar between SNAP23-depleted eggs and

uninjected controls (data not shown). This demonstrates that SNAP23 acts downstream of Ca^{2+}

release. Taken together, these results show that SNAP23 is required for cortical granule exocytosis.

In conclusion, we show that SNAP23 is required for the maintenance of meiotic arrest and cortical

granule exocytosis in mouse oocytes and eggs, respectively. We are currently working to identify other

SNAREs that interact with SNAP23 to mediate constitutive and regulated exocytosis.

ACKNOWLEDGMENTS

We thank Melina Schuh for providing helpful advice with Trim-away, Jurrien Dean for providing ZP2

antibody, Khaled Machaca for providing the SNAP25Δ20 and SNAP25WT constructs, Carmen Williams

for providing the pIVT vector, Jim Watras for providing IP_3 receptor antibody, and Laurinda Jaffe for

helpful comments and suggestions.

FIGURE LEGENDS

Figure 1. SNARE proteins that may be involved in exocytosis in mouse oocytes and eggs. Coiled coil domains from t-SNAREs (syntaxin and SNAP23/SNAP25) zip together with a v-SNARE (synaptobrevin) to pull the vesicle membrane close to the plasma membrane and facilitate membrane fusion.

Figure 2. SNAP23, but not SNAP25, is expressed in mouse oocytes and eggs. A) Western blot showing SNAP25 expression in brain, but not in mouse oocytes or eggs; lysates from 525 oocytes or eggs (13 μg total protein) were used per lane. B) SNAP23 is expressed in mouse eggs. Blot shows lysates from 12 isolated oocytes that were not injected; injected with scrambled siRNA; or injected with siRNA targeting SNAP23 following a ~29 hr incubation and overnight maturation. C) Localization of SNAP23 in oocytes and eggs. Top panels are immunofluorescence and bottom panels are SNAP23-GFP expression. Controls = secondary antibody only (top) or uninjected egg (bottom).

Figure 3. SNAP23 is required for maintenance of meiotic arrest and gap junction communication between the oocyte and cumulus cells. A) Western blot showing SNAP23 reduction using Trim-away, while levels of IP$_3$ receptor were unaffected. B) SNAP23 depletion causes meiotic resumption in follicle-enclosed oocytes. The number of oocytes examined for each group is shown above each bar. C) Alexa488 dye spreading from oocytes to follicle cells in control or SNAP23-depleted follicle-enclosed oocytes. D) Percentages of follicles that exhibited dye spreading. The number of follicles examined is stated above each bar.

Figure 4. A dominant negative SNARE protein, SNAP25Δ20, stimulates meiotic resumption in follicle-enclosed oocytes. A) Both forms of SNAP25 were expressed in oocytes after microinjection. B) Percentage of follicle-enclosed oocytes that underwent GVBD following microinjection of SNAP25Δ20 or SNAP25WT. The number of oocytes examined is indicated above each bar.

Figure 5. SNAP23 depletion inhibits cortical granule exocytosis. SNAP23 protein was degraded using Trim-away and cortical granules were labelled with TRITC-LCA after in vitro maturation and thimerosal treatment (A); or cortical granule exocytosis was assessed by ZP2 cleavage (B). The effectiveness of SNAP23 depletion is shown in (B). C) SNAP23 depletion does not inhibit Ca^{2+} release in response to thimerosal treatment.

REFERENCES

1. Sanfins A, Rodrigues P, Albertini DF. GDF-9 and BMP-15 direct the follicle symphony. J Assist Reprod Genet 2018; 35:1741-1750.
2. Su YQ, Sugiura K, Eppig JJ. Mouse oocyte control of granulosa cell development and function: paracrine regulation of cumulus cell metabolism. Semin Reprod Med 2009; 27:32-42.
3. Norris RP, Ratzan WJ, Freudzon M, Mehlmann LM, Krall J, Movsesian MA, Wang H, Ke H, Nikolaev VO, Jaffe LA. Cyclic GMP from the surrounding somatic cells regulates cyclic AMP and meiosis in the mouse oocyte. Development 2009; 136:1869-1878.
4. Shuhaibar LC, Egbert JR, Norris RP, Lampe PD, Nikolaev VO, Thunemann M, Wen L, Feil R, Jaffe LA. Intercellular signaling via cyclic GMP diffusion through gap junctions restarts meiosis in mouse ovarian follicles. Proc Natl Acad Sci U S A 2015; 112:5527-5532.
5. Zhang M, Su YQ, Sugiura K, Xia G, Eppig JJ. Granulosa cell ligand NPPC and its receptor NPR2 maintain meiotic arrest in mouse oocytes. Science 2010; 330:366-369.
6. Lowther KM, Nikolaev VO, Mehlmann LM. Endocytosis in the mouse oocyte and its contribution to cAMP signaling during meiotic arrest. Reproduction 2011; 141:737-747.
7. Coticchio G, Dal Canto M, Mignini Renzini M, Guglielmo MC, Brambillasca F, Turchi D, Novara PV, Fadini R. Oocyte maturation: gamete-somatic cells interactions, meiotic resumption, cytoskeletal dynamics and cytoplasmic reorganization. Hum Reprod Update 2015; 21:427-454.
8. Ducibella T, Buetow J. Competence to undergo normal, fertilization-induced cortical activation develops after metaphase I of meiosis in mouse oocytes. Dev Biol 1994; 165:95-104.
9. Ducibella T, Rangarajan S, Anderson E. The development of mouse oocyte cortical reaction competence is accompanied by major changes in cortical vesicles and not cortical granule depth. Dev Biol 1988; 130:789-792.
10. Nicosia SV, Wolf DP, Inoue M. Cortical granule distribution and cell surface characteristics in mouse eggs. Dev Biol 1977; 57:56-74.
11. Burkart AD, Xiong B, Baibakov B, Jimenez-Movilla M, Dean J. Ovastacin, a cortical granule protease, cleaves ZP2 in the zona pellucida to prevent polyspermy. J Cell Biol 2012; 197:37-44.
12. Abbott AL, Fissore RA, Ducibella T. Incompetence of preovulatory mouse oocytes to undergo cortical granule exocytosis following induced calcium oscillations. Dev Biol 1999; 207:38-48.
13. Fernandez-Chacon R, Konigstorfer A, Gerber SH, Garcia J, Matos MF, Stevens CF, Brose N, Rizo J, Rosenmund C, Sudhof TC. Synaptotagmin I functions as a calcium regulator of release probability. Nature 2001; 410:41-49.
14. Geppert M, Goda Y, Hammer RE, Li C, Rosahl TW, Stevens CF, Sudhof TC. Synaptotagmin I: a major Ca^{2+} sensor for transmitter release at a central synapse. Cell 1994; 79:717-727.
15. Sudhof TC. Neurotransmitter release: the last millisecond in the life of a synaptic vesicle. Neuron 2013; 80:675-690.
16. Kioumourtzoglou D, Gould GW, Bryant NJ. Insulin stimulates syntaxin4 SNARE complex assembly via a novel regulatory mechanism. Mol Cell Biol 2014; 34:1271-1279.
17. Zhou X, Shentu P, Xu Y. Spatiotemporal Regulators for Insulin-Stimulated GLUT4 Vesicle Exocytosis. J Diabetes Res 2017; 2017:1-9.
18. Wong JL, Koppel DE, Cowan AE, Wessel GM. Membrane hemifusion is a stable intermediate of exocytosis. Dev Cell 2007; 12:653-659.
19. Avery J, Hodel A, Whitaker M. In vitro exocytosis in sea urchin eggs requires a synaptobrevin-related protein. J Cell Sci 1997; 110 (Pt 14):1555-1561.
20. Tsai PS, van Haeften T, Gadella BM. Preparation of the cortical reaction: maturation-dependent migration of SNARE proteins, clathrin, and complexin to the porcine oocyte's surface blocks membrane traffic until fertilization. Biol Reprod 2011; 84:327-335.

21.   Ikebuchi Y, Masumoto N, Matsuoka T, Yokoi T, Tahara M, Tasaka K, Miyake A, Murata Y. SNAP-25 is essential for cortical granule exocytosis in mouse eggs. Am J Physiol 1998; 274:C1496-1500.

22.   Blasi J, Chapman ER, Link E, Binz T, Yamasaki S, De Camilli P, Sudhof TC, Niemann H, Jahn R. Botulinum neurotoxin A selectively cleaves the synaptic protein SNAP-25. Nature 1993; 365:160-163.

23.   Iwahashi K, Kuji N, Fujiwara T, Tanaka H, Takahashi J, Inagaki N, Komatsu S, Yamamoto A, Yoshimura Y, Akagawa K. Expression of the exocytotic protein syntaxin in mouse oocytes. Reproduction 2003; 126:73-81.

24.   Bello OD, Cappa AI, de Paola M, Zanetti MN, Fukuda M, Fissore RA, Mayorga LS, Michaut MA. Rab3A, a possible marker of cortical granules, participates in cortical granule exocytosis in mouse eggs. Exp Cell Res 2016; 347:42-51.

25.   Wang HH, Cui Q, Zhang T, Wang ZB, Ouyang YC, Shen W, Ma JY, Schatten H, Sun QY. Rab3A, Rab27A, and Rab35 regulate different events during mouse oocyte meiotic maturation and activation. Histochem Cell Biol 2016; 145:647-657.

26.   de Paola M, Bello OD, Michaut MA. Cortical Granule Exocytosis Is Mediated by α-SNAP and N-Ethilmaleimide Sensitive Factor in Mouse Oocytes. PLoS One 2015; 10:e0135679.

27.   Clift D, McEwan WA, Labzin LI, Konieczny V, Mogessie B, James LC, Schuh M. A Method for the Acute and Rapid Degradation of Endogenous Proteins. Cell 2017; 171:1692-1706 e1618.

28.   Igarashi H, Knott JG, Schultz RM, Williams CJ. Alterations of PLCβ1 in mouse eggs change calcium oscillatory behavior following fertilization. Dev Biol 2007; 312:321-330.

29.   East IJ, Dean J. Monoclonal antibodies as probes of the distribution of ZP-2, the major sulfated glycoprotein of the murine zona pellucida. J Cell Biol 1984; 98:795-800.

30.   Jaffe LA, Norris RP, Freudzon M, Ratzan WJ, Mehlmann LM. Microinjection of follicle-enclosed mouse oocytes. Methods Mol Biol 2009; 518:157-173.

31.   Kline D. Quantitative microinjection of mouse oocytes and eggs. Methods Mol Biol 2009; 518:135-156.

32.   Clift D, So C, McEwan WA, James LC, Schuh M. Acute and rapid degradation of endogenous proteins by Trim-Away. Nat Protoc 2018; 13:2149-2175.

33.   Mehlmann LM, Carpenter G, Rhee SG, Jaffe LA. SH2 domain-mediated activation of phospholipase Cγ is not required to initiate Ca^{2+} release at fertilization of mouse eggs. Dev Biol 1998; 203:221-232.

34.   Oyler GA, Higgins GA, Hart RA, Battenberg E, Billingsley M, Bloom FE, Wilson MC. The identification of a novel synaptosomal-associated protein, SNAP-25, differentially expressed by neuronal subpopulations. J Cell Biol 1989; 109:3039-3052.

35.   Salaun C, Gould GW, Chamberlain LH. The SNARE proteins SNAP-25 and SNAP-23 display different affinities for lipid rafts in PC12 cells. Regulation by distinct cysteine-rich domains. J Biol Chem 2005; 280:1236-1240.

36.   Vaidyanathan VV, Yoshino K, Jahnz M, Dorries C, Bade S, Nauenburg S, Niemann H, Binz T. Proteolysis of SNAP-25 isoforms by botulinum neurotoxin types A, C, and E: domains and amino acid residues controlling the formation of enzyme-substrate complexes and cleavage. J Neurochem 1999; 72:327-337.

37.   Mallery DL, McEwan WA, Bidgood SR, Towers GJ, Johnson CM, James LC. Antibodies mediate intracellular immunity through tripartite motif-containing 21 (TRIM21). Proc Natl Acad Sci U S A 2010; 107:19985-19990.

38.   Norris RP, Freudzon M, Mehlmann LM, Cowan AE, Simon AM, Paul DL, Lampe PD, Jaffe LA. Luteinizing hormone causes MAP kinase-dependent phosphorylation and closure of connexin 43 gap junctions in mouse ovarian follicles: one of two paths to meiotic resumption. Development 2008; 135:3229-3238.

39.   Gershon E, Plaks V, Dekel N. Gap junctions in the ovary: expression, localization and function. Mol Cell Endocrinol 2008; 282:18-25.

40. Laird DW. Life cycle of connexins in health and disease. Biochem J 2006; 394:527-543.
41. Sela-Abramovich S, Edry I, Galiani D, Nevo N, Dekel N. Disruption of gap junctional communication within the ovarian follicle induces oocyte maturation. Endocrinology 2006; 147:2280-2286.
42. El-Jouni W, Haun S, Hodeify R, Hosein Walker A, Machaca K. Vesicular traffic at the cell membrane regulates oocyte meiotic arrest. Development 2007; 134:3307-3315.
43. Yao Y, Ferrer-Montiel AV, Montal M, Tsien RY. Activation of store-operated Ca^{2+} current in *Xenopus* oocytes requires SNAP-25 but not a diffusible messenger. Cell 1999; 98:475-485.
44. Bajohrs M, Darios F, Peak-Chew SY, Davletov B. Promiscuous interaction of SNAP-25 with all plasma membrane syntaxins in a neuroendocrine cell. Biochem J 2005; 392:283-289.
45. Ravichandran V, Chawla A, Roche PA. Identification of a novel syntaxin- and synaptobrevin/VAMP-binding protein, SNAP-23, expressed in non-neuronal tissues. J Biol Chem 1996; 271:13300-13303.
46. Ledent C, Demeestere I, Blum D, Petermans J, Hamalainen T, Smits G, Vassart G. Premature ovarian aging in mice deficient for *Gpr3*. Proc Natl Acad Sci U S A 2005; 102:8922-8926.
47. Mehlmann LM. Oocyte-specific expression of *Gpr3* is required for the maintenance of meiotic arrest in mouse oocytes. Dev Biol 2005; 288:397-404.
48. Mehlmann LM, Saeki Y, Tanaka S, Brennan TJ, Evsikov AV, Pendola FL, Knowles BB, Eppig JJ, Jaffe LA. The G_s-linked receptor GPR3 maintains meiotic arrest in mammalian oocytes. Science 2004; 306:1947-1950.
49. Eggerickx D, Denef JF, Labbe O, Hayashi Y, Refetoff S, Vassart G, Parmentier M, Libert F. Molecular cloning of an orphan G-protein-coupled receptor that constitutively activates adenylate cyclase. Biochem J 1995; 309 (Pt 3):837-843.
50. Yin H, Chu A, Li W, Wang B, Shelton F, Otero F, Nguyen DG, Caldwell JS, Chen YA. Lipid G protein-coupled receptor ligand identification using beta-arrestin PathHunter assay. J Biol Chem 2009; 284:12328-12338.
51. Kadkova A, Radecke J, Sorensen JB. The SNAP-25 Protein Family. Neuroscience 2018; In Press.
52. Reales E, Mora-Lopez F, Rivas V, Garcia-Poley A, Brieva JA, Campos-Caro A. Identification of soluble N-ethylmaleimide-sensitive factor attachment protein receptor exocytotic machinery in human plasma cells: SNAP-23 is essential for antibody secretion. J Immunol 2005; 175:6686-6693.
53. Mehlmann LM, Kline D. Regulation of intracellular calcium in the mouse egg: calcium release in response to sperm or inositol trisphosphate is enhanced after meiotic maturation. Biol Reprod 1994; 51:1088-1098.
54. Cappa AI, de Paola M, Wetten P, De Blas GA, Michaut MA. Live imaging of cortical granule exocytosis reveals that in vitro matured mouse oocytes are not fully competent to secrete their content. Biol Open 2018; In Press.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5