

1 Self-reported Health is Related to Body Height and Waist Circumference in
2 Rural Indigenous and Urbanised Latin-American Populations

3
4 Juan David Leongómez^{1*}, Oscar R. Sánchez¹, Milena Vásquez-Amézquita², Eugenio Valderrama^{1,3},
5 Andrés Castellanos-Chacón¹, Lina Morales-Sánchez^{1,4}, Javier Nieto⁵, Isaac González-Santoyo^{6*}

6
7 ¹Human Behaviour Lab, Faculty of Psychology, Universidad El Bosque. Bogota, Colombia.

8 ²Experimental Psychology Lab, Faculty of Psychology, Universidad El Bosque, Bogota, Colombia.

9 ³Current Address: LH Bailey Hortorium, Plant Biology Section, School of Integrative Plant Science,
10 Cornell University, Ithaca, NY, United States of America.

11 ⁴Current Address: Department of Psychology, Faculty of Social Sciences, Los Andes University,
12 Bogota, Colombia.

13 ⁵Laboratory of Learning and Adaptation, Faculty of Psychology, National Autonomous University of
14 Mexico, Mexico City, Mexico.

15 ⁶Neuroecology Lab, Faculty of Psychology, National Autonomous University of Mexico, Mexico City,
16 Mexico.

17

18

19 *Corresponding authors

20 E-mail address: jleongomez@unbosque.edu.co (JDL), isantoyo.unam@gmail.com (IG-S)

21

22 **Abstract**

23 Body height is a life-history component. It involves important costs for its expression and maintenance,
24 which may originate trade-offs on other costly components such as reproduction or immunity. Although
25 previous evidence has supported the idea that human height could be a sexually selected trait, the
26 explanatory mechanisms that underlie this selection are poorly understood. Despite extensive studies on
27 the association between height and attractiveness, the role of immunity in linking this relation is scarcely
28 studied, particularly in non-Western populations. Here, we tested whether human height is related to
29 health measured by self-perception, and relevant nutritional and health anthropometric indicators in
30 three Latin-American populations that widely differ in socioeconomic and ecological conditions: two
31 urbanised populations from Bogota (Colombia) and Mexico City (Mexico), and one isolated indigenous
32 population (Me'Phaa, Mexico). Results showed that self-reported health is best predicted by an
33 interaction between height and waist circumference, and the costs associated with large waist
34 circumference are height-dependent, affecting taller people more than shorter individuals. If health and
35 genetic quality cues play an important role in human mate-choice, and height and waist interact to signal
36 health, its evolutionary consequences, including cognitive and behavioural effects, should be
37 addressed in future research.

38 **Introduction**

39 In modern Western societies, it has been seen that women usually prefer men who are
40 significantly taller than average¹⁻³, while men are more tolerant in choosing women who are taller or
41 shorter than average⁴. This is consistent with the idea that male height can be adaptive⁵ and sexual
42 selection favours taller men, possibly because height may represent a honest signal of individual quality,
43 providing hereditary advantages, such as genetic quality for the offspring^{6,7}, or direct benefits,
44 provisioning resources and protection for women and their children⁸. Following these last possible
45 benefits, height has been also proposed as an indicator of resource holding potential (RHP), in terms of
46 social dominance and deference^{9,10} and socioeconomic status^{6,11}.

47 This idea is supported by evidence that the male height is directly correlated with reproductive
48 success, which is not applicable to women, suggesting an unrestricted directional selection that favours
49 very tall men but not to very tall women¹². In fact, it has been reported that taller men (but not extremely
50 tall men) are more likely to find a long-term partner and have several different long-term partners¹³,
51 while the maximum reproductive success of women is below the female average height¹⁴. Furthermore,
52 heterosexual men and women tend to adjust their preferred height of hypothetical partners according to
53 their own stature¹⁵. In general, heterosexual men and women prefer couples in which the man is taller
54 than the woman, and women show a preference for facial cues that denote a taller man¹⁶.

55 Although previous evidence has supported the idea that human height could be a sexually
56 selected trait, little is known regarding whether this role is based on the honest signalling of individual
57 quality⁶. To test this idea, it is important to consider that height needs to face a trade-off with other life
58 history components¹⁷, such as reproduction¹⁸ or immunity¹⁹, and its expression should involve certain
59 costs that not all individuals could equally afford, such that there would be an important phenotypic

60 variation on this trait. Both aspects are present in human height. First, growth in body height is a life-
61 history component^{1,20} that involves important costs for its expression and maintenance. The costs in
62 height can be measured in terms of survival and physiological expenditure¹⁹. For example, according to
63 the Hayflick limit theory of ageing²¹, our cells have a limited number of cell replications available in a
64 lifetime. A minimal increment in body height involves more cells, maybe trillions, and large numbers of
65 cell replications. These large numbers of cell replications demand a large pool of proteins to maintain
66 taller, larger bodies¹⁹, which together, with an increase of free radicals generated by normal cellular
67 metabolism, may lead to a greater likelihood of DNA damage²², thus increasing the incidence of cancer
68 and reducing longevity¹⁹.

69 Secondly, reproduction and immunity could face a trade-off with height because the effect of sex
70 hormones on these life history components; the trade-off with reproduction occurs when the notorious
71 increment of sex steroids, particularly testosterone and oestrogens, induce an accelerated growth period
72 in puberty for both sexes²³, but also a reallocation of physiological resources for reproduction posterior
73 to this period (i.e. spermatogenesis, follicular maturation, etc.), which results in a growth cessation. Both
74 steroids stimulate mineral deposition in the growth plates at the ends of the long bones, thus terminating
75 cell proliferation and resulting in the fusion of the growth plates to the shaft of the bone^{24,25}.

76 In turn, the increment of sexual steroids at sexual maturity trigger another trade-off for
77 individuals that is particularly associated with the effect of testosterone on immunity²⁶. Usually,
78 testosterone exerts suppressive effects on several, but not all, immune components²⁷. For example, it
79 may negatively affect the activity and cellular proliferation of several adaptive and innate immune
80 responses, such as Macrophages, Natural Killers, production of cytokines²⁸ and helper T cells and
81 lymphocyte activation (i.e. Th2 and Th17²⁹). In consequence, it has been documented that testosterone

82 may influence general health patterns, in relation to the severity of certain infections such as malaria,
83 leishmaniasis, amoebiasis³⁰, and tuberculosis^{31,32}

84 Therefore, as a consequence of these life-history trade-offs, height could be considered as a
85 reliable indicator of an individual's condition in terms of (1) the amount and quality of nutritional
86 resources acquired until sexual maturity, (2) the RHP to obtain resources for the somatic maintenance in
87 the adult stage, and (3) the general health condition, given the possibly costly immunosuppressive effect
88 of testosterone. Thus, height can be used for potential partners to receive information about the quality
89 of potential mates; only high-quality individuals could afford to allocate resources to this attractive
90 secondary sexual trait³³, which would result in an increased sexual preference towards taller individuals.

91 Despite extensive studies on the association between height and attractiveness, the role of height
92 as signal of biological quality has been largely studied but results are still controversial. For example,
93 one aspect that has been particularly studied is the relationship between height and various indicators of
94 health. Generally, height is positively related to measures of health in men, such as coronary heart
95 disease morbidity and mortality³⁴, limiting long-standing illness⁶ and perceived health^{6,35}. Nevertheless,
96 the implementation sex and of other indicators of health has led to conflicting results, demonstrating the
97 complexity of the question. For instance, mortality by cancer diseases has been associated more to taller
98 than shorter people of both sexes³⁶, and height somewhat predicts general health in women but
99 following a curvilinear trend⁶.

100 Moreover, most studies have been done using high-income developed populations such as
101 Western, Educated, Industrialised, Rich and Democratic (WEIRD) societies³⁷, which has led to a lack of
102 information of what is occurring in other populations with important socio-ecological differences³⁵.
103 These ecological pressures are important because although genetic allelic expression could be the main
104 factor that determines individual height differences³⁵, height is also the most sensible human anatomical

105 feature that responds to environmental and socioeconomic conditions^{18,38}. For instance, variation in
106 height across social classes is known to be greater in poorer countries³⁹ but is much reduced in countries
107 with higher standards of living⁴⁰. Economic inequality not only affects the population's nutritional
108 patterns, which are especially important during childhood to establish adult height, but also the presence
109 of infectious diseases⁴¹. Childhood disease is known to adversely affect growth. For instance, mounting
110 an immune response to fight against the infection requires concomitant increases in metabolic rate,
111 which could affect the net nutrition, and hence reduces productivity. Disease also prevents food intake,
112 impairs nutrient absorption and causes nutrient loss^{42,43}. Therefore, compared with high-income and
113 developed populations, habitants from locations with stronger ecological pressures imposed by
114 pathogens or greater nutritional deficiencies would face greater costs to robustly express this trait,
115 thereby showing stronger sexual selective pressure over height, as it signals growth rates, life-history
116 trajectories and health status more accurately. This phenotypic variation is described as developmental
117 plasticity, which is a part of the phenotypic plasticity related to growth and development, in response to
118 social, nutritional and demographic conditions, among others⁴⁴. During the last century, given a general
119 improvement in nutrition, human height has steadily increased across the globe⁴⁵, but the level of
120 dimorphism in favour of men is maintained.

121 Colombia and Mexico are two of the most socioeconomically heterogeneous countries in the
122 world with a high Human Development Index⁴⁶. Colombia and Mexico attain respective scores of 68
123 and 66 in the Healthcare Access and Quality Index⁴⁷, indicating that the populations are in relatively
124 good health compared to global standards. Yet, Colombia and Mexico have GINI coefficients of 50.8
125 and 43.4, respectively, making them the 12th and 43rd most unequal countries in the world (GINI index –
126 World Bank estimate; <https://data.worldbank.org/indicator/SI.POV.GINI>). These national-level
127 statistics, however, hide important within-country differences. In particular, Latin-American people in

128 rural areas tend to be poorer and have less access to basic services such as health and education than
129 people in urban areas.

130 According to data from the World Bank and the Colombian National Administrative Department
131 of Statistics, in 2017 Colombia was the second most unequal country in Latin-America after Brazil. In
132 rural areas, 36% of people were living in poverty and 15.4% in extreme poverty, while in urban areas,
133 these values were only 15.7% and 2.7%, respectively⁴⁸.

134 In addition to rural communities, in Latin-America indigenous people tend to have high rates of
135 poverty and extreme poverty⁴⁹, and poorer health⁵⁰, which is less susceptible to improvement by national
136 income growth⁵¹. In Mexico, there are at least 56 independent indigenous peoples whose lifestyle
137 practices differ in varying degrees from the typical ‘urbanised’ lifestyle. Among these groups, the
138 Me’Phaa people, from an isolated region known as ‘*Montaña Alta*’ of the state of Guerrero, is one of the
139 groups whose lifestyle most dramatically differs from the typical Westernised lifestyle of more
140 urbanised areas⁵². Me’Phaa communities are small groups of indigenous people, composed of 50 to 80
141 families, each with five to ten family members. Most communities are based largely on subsistence
142 farming of legumes such as beans and lentils, and the only grain cultivated is corn. Animal protein is
143 acquired by hunting and raising some fowls, and meat is only consumed during special occasions but not
144 as part of the daily diet. There is almost no access to allopathic medications, and there is no health
145 service, plumbing or water purification system. Water for washing and drinking is obtained from small
146 wells. Most of the Me’Phaa speak only their native language⁵³. In consequence, these communities have
147 the lowest income and economic development in the country, and the highest child morbidity and
148 mortality due to chronic infectious diseases⁵².

149 These three Latin-American populations can provide an interesting indication about how the
150 regional socioeconomic conditions and intensity of ecological pressures by pathogens may modulate the

151 function of height as an informative sexually selected trait of health and individual condition in each
152 sex. Therefore, the aim of the present study was to evaluate whether human height is related to health
153 measured by self-perception, and relevant nutritional and health anthropometric indicators in three
154 Latin-American populations that widely differ in socioeconomic and ecological conditions: two
155 urbanised populations from Bogota (Colombia) and Mexico City (Mexico), and one isolated indigenous
156 population (Me'Phaa, Mexico). In addition, given the possible immunological effects of testosterone,
157 and that men present higher levels than women, we predicted this relation to be different between sexes
158 in all studied populations. Therefore, we propose that height would be a stronger signal of self-reported
159 health condition in men compared to women.

160 **Methods**

161 All procedures for testing and recruitment were approved by Universidad El Bosque Institutional
162 Committee on Research Ethics (PCI.2017-9444) and National Autonomous University of Mexico
163 Committee on Research Ethics (FPSI/CE/01/2016), and run in accordance with the ethical principles and
164 guidelines of the Colombian College of Psychologists (COLPSIC) and the Official Mexican Law
165 (NOM-012-SSA3-2012). All participants read and signed a written informed consent.

166 **Participants**

167 A total of 477 adults (238 women and 239 men) participated in this study. They were from three
168 different samples: (1) Mexican indigenous population, (2) Mexican urban population and (3) Colombian
169 urban population. In Mexico, Me'Phaa indigenous participants from 'La *Montaña Alta*' were recruited
170 and participated in this study between January and March 2017, while data from participants from

171 Mexico City was collected between May and June 2017. In Colombia, data collection was carried out
172 between October 2018 and May 2019.

173 The first sample consisted of 63 subjects (mean age \pm standard deviation [SD] = 33.63 ± 9.69
174 years old) from the small Me'Phaa community – '*Plan de Gatica*' from a region known as '*Montaña*
175 *Alta*' of the state of Guerrero in Southwest Mexico. In this sample, 24 participants were women ($33.46 \pm$
176 8.61 years old) and 39 participants were men (33.74 ± 10.41 years old), who participated in a larger
177 study on immunocompetence. Both sexes were aged above 18 years old. In Mexico, people above 18
178 years old are considered adults. All measurements were collected in the participants' own community.
179 Me'Phaa communities are about 20 km apart, and it takes about three hours of travel on rural dirt roads
180 to reach the nearest large town, about 80 km away. Mexico City is about 850 km away, and the trip
181 takes about twelve hours by road. This community has the lowest income in Mexico, the highest index
182 of child morbidity and mortality by gastrointestinal and respiratory diseases (children aged 0 to 8 years
183 had the highest vulnerability and death risk⁵²), and the lowest access to health services. These conditions
184 were recorded in the National Health Information System 2016⁵².

185 The second sample consisted of 60 subjects of over 18 years old (30.27 ± 8.56 years old) from
186 the general community in Mexico City, of whom 30 were women (37.47 ± 5.61 years old) and 30 were
187 men (23.07 ± 3.22 years old). Finally, the third sample consisted of 354 undergraduate students with
188 ages ranging from 18 to 30 years old (20.39 ± 2.10 years old), 184 were women (20.16 ± 2.08 years
189 old), and 170 were men (20.64 ± 2.10 years old) from Bogota, Colombia. All urban participants were
190 recruited through public advertisements.

191 Participants from both urban population samples were taking part in two separate, larger studies
192 in each country. In Colombia, all data were collected in the morning, between 7 and 11 am, because
193 saliva samples (for hormonal analysis), as well as voice recordings, body odour samples, and facial

194 photographs were also collected as part of a separate project. Additionally, women in the Colombian and
195 Mexican samples were not hormonal contraception users, and all data were collected within the first
196 three days of their menses.

197 Participants who were under allopathic treatment and hormonal contraception users from both
198 countries were excluded from data collection. All participants completed a sociodemographic data
199 questionnaire, which included medical and psychiatric history. No women were users of hormonal
200 contraception. Although no participant reported any endocrinological or chronic disease, these health
201 issues were also considered as exclusion criteria.

202 Given that the indigenous community of '*Plan de Gatica*' consists of 60–80 families, each with
203 five to seven members, the final sample for this study could be considered as semi-representative of a
204 larger Me'Phaa population inhabiting in the same community. Nevertheless, the total population of
205 Me'Phaa people inhabiting the '*Montaña Alta*' is comprised of 20–30 communities with almost the
206 same number of families as '*Plan de Gatica*'. Therefore, it is important to mention that our sample size
207 cannot be considered representative of the total Me'Phaa people inhabiting the '*Montaña Alta*' region,
208 but from the specific '*Plan de Gatica*' community. Similar condition occurs for participants from the
209 Mexico City and Bogota samples. These participants were recruited at the National Autonomous
210 University of México and Universidad El Bosque campuses, respectively. Therefore, these samples are
211 comprised mostly of bachelor and graduate students, and cannot be considered as representative of a
212 large population of the whole city, which is comprised of about 12 million adult persons in Mexico and
213 about 5 million adults in Bogota.

214 **Procedure**

215 All participants signed the informed consent and completed the health and background
216 questionnaires. For participants from the indigenous population, the whole procedure was carried out
217 within their own communities, and participants from the Mexican and Colombian urban population
218 attended a laboratory at either the National Autonomous University of México or Universidad El Bosque
219 respectively, on individual appointments.

220 Participants from Mexico City and Bogota were recruited through public advertisements on
221 social media and poster boards located along the central campus of the National Autonomous University
222 and Universidad El Bosque. While in Mexico City, participants received either one partial course credit
223 or a payment equivalent to \$5 dollars as compensation for their participation, all participants in Bogota
224 were given academic credits for their participation.

225 For the indigenous groups, recruitment was done through the Xuajin Me'Phaa non-governmental
226 organisation, which is dedicated to the social, environmental and economic development for the
227 indigenous communities of the region (see video from this organisation, http://youtu.be/In4b9_Ek78o).
228 Xuajin Me'Phaa has extensive experience in community-based fieldwork and has built a close working
229 relationship with the community authorities. The trust and familiarity with the community customs and
230 protocols have previously led to successful academic collaborations^{53,54}. Therefore, Xuajin Me'Phaa
231 served as a liaison between the Mexican research group the and communities for the present study,
232 offering mainly two important factors in data collection: the informed consent of community members
233 and participants, and two trained interpreters of Me'Phaa and Spanish language of both sexes.

234 First, participants were asked to complete the health and sociodemographic data questionnaires.
235 Subsequently, the anthropometric measurements were taken.

236 **Self-reported health**

237 In order to obtain a standardised value of self-perception of health, we implemented in all three
238 populations the Short Form (36) health survey (SF-36; RAND Corp.; [https://www.rand.org/health-](https://www.rand.org/health-care/surveys_tools/mos/36-item-short-form/survey-instrument.html)
239 [care/surveys_tools/mos/36-item-short-form/survey-instrument.html](https://www.rand.org/health-care/surveys_tools/mos/36-item-short-form/survey-instrument.html)). The SF-36 produces eight
240 dimensions, of which we only used the last one (8) General health. Each factor is calculated by averaging
241 the recoded scores of individual items: (1) Physical functioning (items 3 to 12), (2) Role limitations due
242 to physical health (items 13 to 16), (3) Role limitations due to emotional problems (items 17 to 19), (4)
243 Energy/fatigue (items 23, 27, 29 and 31), (5) Emotional well-being (items 24, 25, 26, 28 and 30), (6)
244 Social functioning (items 20 and 32), (7) Pain (items 21 and 22) and (8) General health (items 1, 33, 34,
245 35 and 36).

246 The interpreters provided by the Xuajin Me'Phaa organisation administered the SF-36 Health
247 survey in Me'Phaa language. Interpreters used Spanish as the second language and are thoroughly
248 proficient in speaking and reading Spanish. We used the validated SF-36 survey for urban and rural
249 Mexican populations⁵⁵ for interpreters to translate Spanish to Me'Phaa language. Given the ethnical
250 customs of Me'Phaa culture, the participants were always interviewed by an interpreter of the same sex
251 to avoid bias in participant responses; for instance, men were interviewed by a male interpreter and
252 women by a female interpreter. The same interpreter interviewed all participants of his/her
253 corresponding sex.

254 For the present study, both urban and indigenous participants only answered items corresponding
255 to the dimension defined as general health (i.e. Item numbers 1, 33, 34, 35 and 36), except for item 35.
256 This item informs about the expectation for future health. Since the grammatical compositions of
257 Me'Phaa language do not consider 'infinitive' and 'future' as verbal tenses⁵⁶, an interpretation of this
258 question was not possible for the Me'Phaa people, therefore, this item was excluded.

259 In Colombia, we used a Spanish version of the SF-36 questionnaire⁵⁷, that was previously
260 validated in the same country⁵⁸.

261 To obtain the self-reported health rate, all items were recoded following the instructions on how
262 to score SF-36⁵⁷. We calculated the final factor by averaging the recoded items. To make this data
263 compatible with the Mexican database, item 35 was excluded because it cannot be answered by the
264 Mexican Indigenous population, and the general health dimension was calculated by averaging only
265 items 1, 33, 34 and 36.

266 **Anthropometric measurements**

267 All anthropometric measurements were measured thrice and subsequently averaged to obtain the
268 mean value (for agreement statistics between the three measurements of each characteristic, see section
269 1.3 in the Supplementary Material). All participants wore light clothing and had their shoes removed.
270 The same observer repeated the measurements thrice.

271 We measured the body height in cm, to the nearest mm, by using a 220 cm Zaude stadiometer,
272 with the participant's head aligned according to the Frankfurt horizontal plane, and feet together against
273 the wall.

274 Anthropomorphic measurements also included waist circumference (cm), weight (kg), fat
275 percentage, visceral fat level, muscle percentage and body mass index (BMI). The waist circumference
276 was measured midway between the lowest rib and the iliac crest in cm by using a flexible tape and was
277 recorded to the nearest mm. These anthropomorphic measurements have been used as an accurate index
278 of nutritional status and health, especially waist circumference. Metabolic syndrome is associated with
279 visceral adiposity, blood lipid disorders, inflammation, insulin resistance or full-blown diabetes and
280 increased risk of developing cardiovascular disease⁵⁹⁻⁶¹, amongst Latin-American populations⁶². Waist

281 circumference has been proposed as a crude anthropometric correlate of abdominal and visceral
282 adiposity, and it is the simplest and accurate screening variable used to identify people with the features
283 of metabolic syndrome^{63,64}. Hence, in the presence of the clinical criteria of metabolic syndrome,
284 increased waist circumference provides relevant pathophysiological information insofar as it defines the
285 prevalent form of the syndrome resulting from abdominal obesity⁶⁰.

286 Weight, fat percentage, visceral fat level, muscle percentage and BMI were obtained using an
287 Omron Healthcare HBF-510 body composition analyser, which was calibrated before each participant's
288 measurements were obtained.

289 **Statistical analysis**

290 We used linear models (LM) to test the association between height and self-reported health. The
291 dependent variable in this model was the health factor and predictor variables included participant sex,
292 age, sample (Bogota, Mexico City, Me'Phaa), height and waist and anthropometric measurements (hip,
293 weight, fat percentage, BMI and muscle percentage) as fixed, main effects, as well as all possible
294 interactions between height, waist, sample, and sex. For all models, the continuous regressors involved
295 in interactions (waist and height) were centred.

296 Although sample could be thought as a random factor (i.e. fitting linear mixed models instead),
297 we treated it as a fixed effects categorical predictor in the models because there were only three levels
298 (Bogota, Mexico City, Me'Phaa), and a minimum of five levels is recommended. To test the residual
299 distribution, generalised linear models (GLM) were fitted, but in all cases, residuals were closer to a
300 normal or gamma (inverse link) distribution, for each sample. Models here included were fitted using
301 the *lm* function in R, version 3.6.1⁶⁵.

302 The most parameterised initial model (Model 1) was then reduced, by excluding the main effects
303 of hip, weight, fat percentage, visceral fat, BMI and muscle percentage (as these are phenotypic markers
304 associated either with height or waist circumference), and keeping the main effects of age, as well as the
305 main effects and all possible interactions between any combination of height, waist, sample, and sex,
306 consistent with our predictions. This, still highly parameterised model (Model 2), was further reduced
307 using the functions
308 *dredge* (<https://www.rdocumentation.org/packages/MuMIn/versions/1.43.6/topics/dredge>) and *model.sel*
309 (<https://www.rdocumentation.org/packages/MuMIn/versions/1.43.6/topics/model.sel>) from the package
310 *MuMIn: Multi-Model Inference*⁶⁶. The *dredge* function fitted a set of 334 models with combinations
311 (subsets) of fixed effect terms from the second model, that were then compared using the function
312 *model.sel* based on the Akaike Information Criterion (AICc) and Akaike weights, allowing us to select
313 the best model (Model 3). This, best-supported model (i.e. the model with the lowest AICc with a
314 Δ AICc higher than two units from the second most adequate model), is reported⁶⁷.

315 Finally, we compared the three models selected model (Models 1, 2 and 3) using the *ICtab*
316 function from the *bbmle* package⁶⁸. Once a final model was selected, model diagnostics were performed
317 (collinearity, residual distribution and linearity of residuals in each single term effect; see section 3.3 in
318 the Supplementary Material).

319 Interactions in the final model were explored and via simple slopes analysis and Johnson-
320 Neyman intervals^{69,70}, using the R package *interactions: Comprehensive, User-Friendly Toolkit for*
321 *Probing Interactions*⁷¹. For this purpose, we implemented the functions *sim_slopes*
322 (https://www.rdocumentation.org/packages/interactions/versions/1.1.1/topics/sim_slopes), *interact_plot*
323 (https://www.rdocumentation.org/packages/interactions/versions/1.1.1/topics/interact_plot) and

324 *johnson_neyman*

325 (https://www.rdocumentation.org/packages/interactions/versions/1.1.1/topics/johnson_neyman).

326 Results

327 All data and code used to perform these analyses are openly available from the Open Science
328 Framework (OSF) project for this study (<https://osf.io/5rzfs/>).

329 Descriptives

330 Descriptive statistics of age, waist circumference, hip, height, weight, fat percentage, visceral fat,
331 BMI, muscle percentage and self-reported health and reported in Table 1.

332 **Table 1. Descriptive statistics of measured variables of all participants.**

Measured characteristic	Sample	Women						Men					
		<i>n</i>	Mean	<i>SD</i>	Median	Min	Max	<i>n</i>	Mean	<i>SD</i>	Median	Min	Max
Age	Bogota	184	20.2	2.1	20.0	18.0	30.0	170	20.6	2.1	20.0	18.0	29.0
	Me'Phaa	24	33.5	8.6	31.5	21.0	50.0	39	33.7	10.4	33.0	17.0	60.0
	Mexico City	30	37.5	5.6	38.0	25.0	46.0	30	23.1	3.2	21.5	19.0	31.0
BMI (kg/m ²)	Bogota	184	23.0	4.0	22.1	15.4	41.4	170	23.1	3.3	22.8	16.6	33.3
	Me'Phaa	24	25.4	3.1	24.9	19.7	31.7	39	25.6	4.7	24.9	19.1	40.2
	Mexico City	30	26.4	5.2	26.5	16.5	40.2	30	24.0	3.7	23.5	19.0	37.9
Fat (%)	Bogota	184	34.9	7.3	34.2	12.6	58.3	170	20.2	6.8	19.7	5.4	38.7
	Me'Phaa	24	38.8	5.3	38.0	27.4	48.4	38	24.4	8.3	23.4	9.3	44.4
	Mexico City	30	39.0	7.8	39.5	19.2	55.6	30	21.2	7.0	21.2	6.5	40.0
Height (cm)	Bogota	184	158.9	6.0	159.1	141.9	178.9	170	172.2	6.4	171.7	155.5	188.1
	Me'Phaa	24	146.2	5.5	144.0	136.0	157.0	39	159.9	6.8	161.0	143.0	173.5
	Mexico City	30	157.7	5.9	158.0	145.0	168.0	30	172.1	6.8	171.8	159.9	184.1
Hip (cm)	Bogota	184	96.6	7.7	95.7	79.8	123.0	170	98.0	7.0	97.0	83.1	122.0
	Me'Phaa	24	95.9	7.4	93.5	86.0	114.0	39	95.3	9.1	94.5	79.9	119.0
	Mexico City	30	100.1	9.7	99.6	82.2	123.6	30	96.8	8.6	96.0	78.0	126.6
Muscle (%)	Bogota	184	25.6	2.5	25.6	18.0	33.9	170	40.1	3.8	40.2	29.6	49.1
	Me'Phaa	24	25.2	2.4	25.1	20.4	29.8	36	36.9	5.2	37.4	25.6	47.9
	Mexico City	30	24.9	2.5	24.7	19.6	29.1	30	39.5	4.2	39.4	29.0	49.5

Self-reported health	Bogota	184	64.7	19.4	68.8	12.5	100.0	170	72.6	17.4	75.0	0.0	100.0
	Me'Phaa	24	50.8	10.5	50.0	31.2	68.8	39	50.3	9.2	50.0	25.0	75.0
	Mexico City	30	56.0	7.6	56.2	43.8	75.0	30	60.4	8.6	62.5	37.5	75.0
Visceral fat	Bogota	184	3.9	1.3	4.0	1.0	8.0	170	5.4	2.8	5.0	1.0	14.0
	Me'Phaa	24	6.4	1.8	6.0	3.0	11.0	35	9.4	4.7	8.0	2.0	23.0
	Mexico City	30	6.4	2.0	7.0	2.0	10.0	30	6.2	3.2	6.0	2.0	17.0
Waist circumference (cm)	Bogota	184	71.8	8.4	70.3	55.3	103.9	170	78.2	7.9	77.6	62.1	103.6
	Me'Phaa	24	87.0	8.2	86.7	73.0	106.0	39	88.6	11.9	86.4	70.5	118.0
	Mexico City	30	87.8	10.9	87.4	66.5	113.9	30	84.5	8.4	84.3	69.0	106.6
Weight (kg)	Bogota	184	57.8	10.2	55.8	39.3	93.9	170	68.2	10.5	67.4	46.5	106.6
	Me'Phaa	24	54.2	7.7	54.4	43.7	67.2	39	65.9	14.5	61.9	43.4	101.7
	Mexico City	30	65.5	12.5	65.0	41.8	100.3	30	71.0	11.5	69.0	48.7	114.1

333

334 The distribution of all measured variables is shown in Figure 1. Age, waist, height, visceral fat,
335 and self-reported health strongly varied in both women (Fig 1a) and men (Fig 1b) between samples.

336 Age, waist circumference, height, fat percentage, visceral fat, BMI and muscle percentage, were
337 significantly correlated with self-reported health ($r > 0.10$, in all cases) in both men and women
338 (bivariate Pearson correlations between all measured variables for all participants combined are shown
339 in the Supplementary Table S2, for women in Supplementary Table S3, and for men in Supplementary
340 Table S4, online).

341 Models to predict self-reported health

342 To establish the relationship between height and self-reported health, we fitted three linear
343 models (Table 2). For all models, the continuous regressors involved in interactions (waist and height)
344 were mean-centred.

345 **Table 2. Results of separate LMs testing effects of independent variables on self-reported health.**

	Model 1			Model 2			Model 3		
	<i>B</i>	<i>t</i>	<i>p</i>	<i>B</i>	<i>t</i>	<i>p</i>	<i>B</i>	<i>t</i>	<i>p</i>
(Intercept)	78.69	2.51	0.012	61.07	13.36	<0.0001	61.36	16.2	<0.0001
Age	0.19	0.86	0.393	0.17	0.85	0.393	0.17	1.0	0.299

BMI	1.18	0.63	0.53						
Fat	-0.06	-0.13	0.894						
H(c)	0.67	1.03	0.303	0.36	1.50	0.134	0.16	1.3	0.189
H(c) × S(Me'Phaa)	-1.90	-1.60	0.11	-1.78	-1.51	0.131			
H(c) × S(Mexico City)	0.17	0.24	0.813	0.14	0.19	0.846			
Hip	-0.31	-1.30	0.193						
Muscle	0.38	0.57	0.572						
S(Me'Phaa)	-46.58	-1.99	0.047	-41.54	-1.79	0.074	-16.78	-4.8	<0.0001
S(Mexico City)	-10.25	-1.45	0.148	-7.72	-1.12	0.262	-9.32	-3.2	0.001
Sex(male)	-0.15	-0.02	0.985	8.49	2.99	0.003	6.01	2.7	0.008
Sex(male) × H(c)	-0.35	-1.02	0.307	-0.41	-1.31	0.19			
Sex(male) × H(c) × S(Me'Phaa)	1.59	1.15	0.253	1.63	1.18	0.238			
Sex(male) × H(c) × S(Mexico City)	0.05	0.05	0.957	0.04	0.04	0.964			
Sex(male) × S(Me'Phaa)	19.65	0.82	0.412	17.06	0.72	0.473			
Sex(male) × S(Mexico City)	-4.68	-0.51	0.613	-4.81	-0.53	0.598			
Sex(male) × WC(c)	0.51	1.30	0.194	0.41	1.25	0.213			
Sex(male) × WC(c) × H(c)	-0.03	-0.82	0.413	-0.02	-0.68	0.496			
Sex(male) × WC(c) × H(c) × S(Me'Phaa)	-0.11	-0.90	0.368	-0.11	-0.90	0.368			
Sex(male) × WC(c) × H(c) × S(Mexico City)	0.07	0.73	0.464	0.06	0.59	0.553			
Sex(male) × WC(c) × S(Me'Phaa)	-2.71	-1.31	0.191	-2.58	-1.25	0.211			
Sex(male) × WC(c) × S(Mexico City)	-0.89	-1.14	0.256	-0.78	-1.00	0.319			
Visceral Fat	0.05	0.05	0.957						
WC(c)	-0.17	-0.58	0.561	-0.42	-2.12	0.034	-0.28	-3.2	0.001
WC(c) × H(c)	-0.01	-0.60	0.55	-0.02	-0.94	0.349	-0.02	-2.3	0.022
WC(c) × H(c) × S(Me'Phaa)	0.15	1.35	0.178	0.15	1.33	0.183			
WC(c) × H(c) × S(Mexico City)	0.02	0.28	0.778	0.02	0.27	0.788			
WC(c) × S(Me'Phaa)	2.55	1.26	0.208	2.51	1.24	0.214			
WC(c) × S(Mexico City)	0.26	0.55	0.582	0.31	0.64	0.52			
Weight	-0.35	-0.49	0.625						

346 *Note.* For Model 1, $R^2 = 0.21$; $R^2_{\text{adjusted}} = 0.156$; $F(30, 442) = 3.91$; $p < 0.001$; for Model 2, $R^2 = 0.201$; $R^2_{\text{adjusted}} =$
347 0.158 ; $F(24, 448) = 4.7$; $p < 0.001$; for Model 3, $R^2 = 0.183$; $R^2_{\text{adjusted}} = 0.171$; $F(7, 465) = 14.88$; $p < 0.001$. Female
348 participants, and Bogota were used as reference categories for Sex and Sample, respectively. For model terms: WC(c) =
349 Waist circumference (centred); H(c) = Height (centred); S = Sample. Significant effects are in bold. For a summary of each
350 model, including standard errors and 95% CIs, see Supplementary Tables S5, S6, and S7 online.
351

352 In the first model (Model 1), we included as predictors all measured variables as main effects, as
353 well as all interactions between height, waist circumference, sample, and sex. The first model was
354 initially reduced by excluding hip, weight, fat percentage, BMI and muscle percentage. We decided to
355 include waist circumference instead of visceral fat or fat percentage for two reasons: first, because these
356 three variables are strongly correlated in women and men ($r > 0.79$ in all cases; see Supplementary
357 Tables S3 and S4 online, for women and men, respectively). And second, because unlike visceral fat or
358 fat percentage, waist circumference can be directly perceived by others, and hence could have a direct

359 effect on mate-choice; fat percentage and visceral fat, on the other hand, are likely perceived and
360 assessed in social contexts through other variables, including relative waist size.

361 In the second model (Model 2), we therefore included age, height, sample, sex, waist
362 circumference, and all possible interactions between combinations of height, waist circumference,
363 sample, and sex. This second model was further reduced by the implementation of the functions *dredge*
364 and *model.sel* from the package MuMIn⁶⁶ (for details, see the Statistical analysis section in the
365 Methods). These functions fitted and compared a total of 334 models with different combinations of
366 fixed terms from Model 2; these compared models and their relative probability to be the best model are
367 shown according to their relative Akaike weights ($w_i(\text{AICc})$) in Fig. 2.

368 This analysis revealed that the best model (labelled 159 in Fig. 2), included height (centred),
369 sample, sex, waist circumference (centred) and the interaction between height (centred)
370 and waist (centred). However, to account for the age differences between samples, we selected the
371 second-best model (labelled 160 in Fig. 2), because it also included age as a regressor, and had
372 a ΔAICc of less than 2 units (≈ 0.98) compared to the best model. This model, including age, was
373 therefore selected as our final model (Model 3).

374 The three selected models were compared using the AICc, Akaike weights ($w_i(\text{AICc})$) and
375 ΔAICc (Table 3). The analyses revealed that Model 3 is not only the most parsimonious of the three
376 selected models, but has higher R^2_{adjusted} and F values (Table 2), as well as a lower AIC and higher Akaike
377 weight⁶⁷ (Table 3) than the previous two models. In fact, Model 3 is close to 464,686 times more likely
378 to be the best model compared to Model 2, and about 35,141,683 times compared to Model 1 (Model 2,
379 was around 76 times more likely compared to Model 1).

380 **Table 3. Performance criteria of the three selected models.**

Model	AICc	ΔAICc	df	$w_i(\text{AICc})$
Model 3	3999	0	9	0.99999782

Model 2	4025	26	26	0.00000215
Model 1	4034	35	32	0.00000003

381 *Note.* Models are in descending order from the best to the worst fitting. ΔAIC is the change in AIC between each model and
 382 the best model. Akaike weights ($w_i(AICc)$) are conditional probabilities for each model being the best model⁶⁷.
 383

384 Furthermore, for Model 3 (the final, minimum adequate model), Generalised Variance Inflation
 385 Factors (GVIF)⁶⁶ revealed no concerning cases of collinearity for any of the predictor terms (GVIF \leq 3,
 386 and a GVIF^{1/(2×Df)} \leq 1.6 in all cases; for details, see Supplementary Table S8 online; residual distribution
 387 by sample and linearity in each single term factor are shown in Supplementary Fig. S2 online).

388 The final model (Model 3: Table 4; Fig. 3) showed a significant, negative main effect of waist
 389 circumference ($t = -3.20$, $p = 0.001$), and significant main effects of sex (men rated their health 6.01
 390 points higher than women; $t = 2.66$, $p = 0.008$), and sample (Mexico City and Me’Phaa individuals rated
 391 their health 9.32 and 19.78 points lower than participants from Bogota; $t = -3.25$, $p = 0.001$ and ; $t = -$
 392 4.75, $p < 0.001$, respectively).

393 **Table 4. Results of the final LMM testing effects of independent variables on self-reported health**

	<i>B</i>	<i>SE(B)</i>	95% CI	<i>t</i>	<i>p</i>
(Intercept)	61.36	3.78	53.94 — 68.782	16.25	<0.0001
Age	0.17	0.16	-0.153 — 0.495	1.04	0.299
H(c)	0.16	0.12	-0.08 — 0.401	1.31	0.189
S(Mexico City)	-9.32	2.87	-14.96 — -3.681	-3.25	0.001
S(Me’Phaa)	-16.78	3.53	-23.722 — -9.83	-4.75	<0.0001
Sex(male)	6.01	2.26	1.564 — 10.458	2.66	0.008
WC(c)	-0.28	0.09	-0.459 — -0.11	-3.20	0.001
WC(c) × H(c)	-0.02	0.01	-0.036 — -0.003	-2.30	0.022
Simple slope analysis for H(c) at different values of WC(c)					
WC(c) - 1 SD = -10.53	0.37	0.15	0.075 — 0.657	2.47	0.014
WC(c) Mean = -0.12	0.16	0.12	-0.077 — 0.403	1.33	0.183
WC(c) + 1 SD = 10.28	-0.04	0.15	-0.341 — 0.261	-0.26	0.795

394 *Note.* As waist reference, the centred values used are equivalent to -1 SD (67.43 cm), mean (78.01 cm), and +1 SD (88.6 cm).
 395 Women and Bogota were used as reference categories for Sex and Sample, respectively. For model terms: WC(c) = Waist
 396 circumference (centred); H(c) = Height (centred); S = Sample. Significant predictors are in bold.
 397

398 Moreover, a significant interaction between waist and height (Table 4; $t = -2.30$, $p = 0.022$) was
399 revealed, indicating that the negative association of waist circumference with self-reported health was
400 height-dependent (Fig. 3b); the best predicted self-reported health was for tall participants with small
401 waists, and while the association between height and self-reported health is positive for people with
402 small waist circumferences, it decreases for people with increasingly large waists. Furthermore, the
403 Johnson-Neyman procedure^{69,70} (Fig. 3c), indicated that height is only a significant, always positive,
404 predictor of self-reported health for people with relatively small waist circumferences of less than 73.51
405 cm (centred: -4.51).

406 This interaction was replicated when fitting an alternative version of Model 3 (Model 3A),
407 replacing waist circumference for visceral fat, by following the same method to select it (i.e. fitting an
408 alternative Model 2, and repeating the same selection process; see section 4.3 in the Supplementary
409 Material, online). Similar to Model 3, this alternative Final Model, also included an interaction between
410 height and, in this case, visceral fat, in which height was found to be a positive significant predictor of
411 self-reported health, only for people with low levels of visceral fat (see Table S9 and Fig. S3 in the
412 Supplementary Material, online). Furthermore, in this model the interaction was more extreme than
413 when using waist circumference, and height becomes a significant, negative predictor of self-reported
414 health for people with high visceral fat (see Fig. S3c in the Supplementary Material, online).

415 **Discussion**

416 The present study provides new insights into the relationship between height and health in men
417 and women by studying three Latin-American populations, which included urban and indigenous
418 populations with marked differences in access to basic needs and services like food and health.

419 Contrary to our initial hypothesis, height was not a significant predictor of self-perceived health
420 but interacted with waist circumference. Most results in favour of a direct relationship between height
421 itself and health were carried out in small modern populations and specific Western ethnic groups more
422 than twenty years ago. New studies with non-traditional population groups have failed to verify the
423 positive relationship between height and health, especially associated with cardiovascular and
424 autoimmune diseases^{72,73}. For example, studies on Native Americans, Japanese, Indians and Pakistanis
425 showed that shorter people had a lower prevalence of cardiovascular disease than the tallest people in
426 each population⁷³. These findings were similar in Sardinian inhabitants, a European population with the
427 lowest physical stature recorded in Europe in recent years⁷².

428 Interestingly, our results suggest that there is a main negative effect of waist circumference on
429 self-perceived health. This is congruent with a broad range of studies done in different human
430 populations⁷⁴. In fact, waist circumference has been proposed as one of the most important biomarker of
431 metabolic syndrome that predicts health condition in terms of cardiovascular diseases⁶⁰. Nevertheless,
432 we found that this negative association was height-dependent in our studied samples. In other words,
433 waist circumference predicted self-reported health differently for people of different heights: while
434 being taller predicts better self-reported health for taller people with relatively small waists, being taller
435 was found to be associated with poorer perceptions of their health in people with larger waist
436 circumferences. Furthermore, while there is a cost of abdominal and visceral adiposity for tall people,
437 there is no predicted cost for shorter persons. Interestingly, epidemiological studies have widely
438 implemented an integration of both phenotypical components in the form of waist to height ratio
439 (WtHR). In general, waist circumference has stronger negative impact on health of short individuals
440 than for tall ones⁷⁵, contrasting with our results. These differences might be due to WtHR has been
441 mainly used to predict health in terms of metabolic and cardiovascular diseases (CVD), while our study

442 used a general status of health, which could include more than metabolic and CVD. In addition, we use
443 these phenotypic variables as continuous and independent predictors because the aim of our study
444 argued that human height by itself would be an honest indicator of general health, which would not be
445 able to be evaluate with WtHR as predictor. Therefore, our results argue the importance of considering a
446 phenotypic independent integration of different human features that could be involved in health or
447 physiological conditions, when a possible sexually selected trait is being evaluated as a signal of
448 individual condition.

449 On the other hand, given that height is the human anatomical feature most sensitive to
450 environmental and socioeconomic conditions^{18,38}, we expected stronger association between health and
451 height for the indigenous population where the cost to produce and maintain this costly trait is greater
452 than for inhabitants from urbanised areas. Nevertheless, we did not find inter-population differences in
453 the magnitude of this relation. Urban populations reported better health than the indigenous population,
454 and the shortest participants tended to be from the indigenous Me'Phaa sample. These results could, in
455 fact, suggest different life-history strategies. Compared with modern Western societies, different life
456 strategies could take place in harsh environments⁷⁶, for instance, investing relatively less energy in
457 growth and reallocating it towards reproduction¹⁸. In addition, a relative increase in the intensity or
458 number of infectious diseases (including paediatric diseases in Me'Phaa) and higher tendency to early
459 sexual maturity could negatively impact growth, resulting in a lower average height^{77,78}. These trends
460 could be compensations between life-history components³⁵. Finally, fast and prolonged growth imply
461 high costs for the organism¹. Rapid growth may influence mortality risk⁷⁹ and growing for a longer time
462 delays the onset of reproduction, increasing the risk of death and producing fewer offspring¹. This
463 perspective of life strategies allows us to understand the relationship between height, health and
464 reproduction. This suggests the importance of addressing factors such as ethnicity, socioeconomic

465 status, level of urbanisation in populations where there is great heterogeneity in access to food, health
466 and pressure resources for pathogens, for instance, in Latin-American populations in which this
467 relationship has barely been directly explored.

468 Although our results show that height and waist circumference are important predictors of self-
469 perception of health, we did not evaluate any immunological mechanism that may underlie the self-
470 perception responses of the participants. This limitation makes hard to directly evaluate human height as
471 an honest signal of individual condition. In the present study, the questions done to evaluate general
472 health (SF-36) is far to be a direct indicator of immune condition, since the participant's perception
473 responses could be influenced by components different than the individual's ability to deal or resist to
474 infectious pathogens, such as skeletal disorders, cancers, cardiovascular or metabolic abnormalities.
475 Nevertheless, studies that have evaluated a more direct approximation of immune condition have led to
476 controversial results. For instance, the implementation of antibody response to a hepatitis-B vaccine as a
477 marker of immune condition has been positive associated with height in men (but not in women) up to a
478 height of 185 cm, but an inverse relationship in taller men⁸⁰. Furthermore, this relation was not found
479 when different components of innate and adaptive immune system functioning were evaluated, such as
480 lysozyme activity, neutrophil function, IgA and IgG⁸¹.

481 Finally, in relation with sex differences, women reported lower average health than men in all
482 communities, which is concordant with reports and normative SF-36 data in other populations,
483 especially in younger people^{82,83}. These results could consolidate the idea that height is a reliable signal
484 of health in men³⁵, while it could reflect reproductive success in women⁸⁴ in terms of labour and birth,
485 and to a lesser extent, function as an indicator of health⁸⁵. It has been seen that taller women experience
486 fewer problems during the labour process due to a lower risk of mismatch between foetal head size and

487 size of the birth canal⁸⁵. Nevertheless, this speculative idea warrants further studies on comparing health,
488 reproductive success and female height.

489 It is important to consider that the mode of survey administration may be another limitation in
490 our study, and it could have led to confounding effects. For example, it is possible that indigenous
491 people have different understanding and thresholds about their general health perception, which we were
492 unable to evaluate without previous validation of translated items, and it could have explained the lowest
493 values of general health reported by indigenous people. Nevertheless, it could also reflect the real health
494 conditions in Me'Phaa communities and not a misunderstanding of the survey. Other national indicators
495 of health, such as morbidity and mortality by gastrointestinal and nasopharyngeal infectious diseases,
496 have reported that Me'Phaa communities also present the poorest health in Mexico⁵², which is consistent
497 with our results. In fact, items for the dimension of general health perception have the lowest standard
498 deviation and coefficient of variation in the entire SF-36 survey, in both validated Spanish^{55,58} and
499 English versions⁸⁶, which makes this dimension the most understandable one.

500 In addition, in order to consider obvious differences in language and perception of health,
501 statistical models in this study assumed these inter-population variations *a priori*. The effects of the
502 sample were considered in all performed LMs. We found that although samples differ considerably, the
503 associations between height, waist circumference and self-perceptions of health were predicted to be in
504 the same direction for all populations (i.e. not interacting with the sample).

505 Finally, we did not have any information regarding potential pregnancy history in women. This
506 is important because each pregnancy can affect waist circumference, so future studies should collect and
507 control or include this variable in all fitted models.

508 The present study contributes information which could be important in the framework of human
509 sexual selection. If health and genetic quality cues play an important role in human mate-choice⁸⁷, and

510 height and waist interact to signal health, its evolutionary consequences, including cognitive and
511 behavioural effects, should be addressed in future research. This could be done by studying the
512 interaction between waist circumference and height, in relation to reproductive and/or mating success, as
513 well as mate preferences and perceived attractiveness, in populations with both Westernised and non-
514 Westernised lifestyles.

515 **Data availability**

516 All data used for this article are openly available at the OSF⁸⁸. Code to perform all analyses, data
517 manipulation, tables and figures is available in PDF ('Supplementary_Material.pdf') and *R Markdown*
518 ('Supplementary_Material.Rmd') formats, so that it can be fully reproduced and explored in depth⁸⁹.

519 **Author contributions**

520 JDL, ORS, MV-A, EV and I.G-S. conceived and designed this study. JDL, ORS, AC-C, LM-S,
521 and IG-S collected data. JDL and IG-S analysed all data. JDL, ORS, MV-A and IG-S wrote the first
522 draft. All authors contributed to writing, approved the final version of the manuscript and gave approval
523 for publication.

524 **Acknowledgments**

525 We are grateful to Laura Rojas, Angie Ramos, Ángela Valderrama, Valentina West, Sergio
526 Camelo, Laura Quintero, Paula Garzón, María Aguirre, Andrea Pastrana, Nicola Caro, Irene Olivella,
527 Luisa Ramírez, Laura Guarín, y Henry Segura for their help in data collection in Colombia, and all our
528 participants. We also grateful to Xuajin Me'Phaa, Margarita Mucino, Julio Gatica, and Diego
529 Hernandez-Mucino for their help in the liaison with the Me'Phaa community, and for their help in data

530 collection logistics. This project was funded in Colombia by Colciencias [grant number
531 C145I004800000881-1 to JDL] and Universidad El Bosque, Vice-rectory of Research [grant number
532 PCI.2017-9444]. Logistics and data collection in Mexico were supported by UNAM-PAPIIT [grant
533 numbers IA209416, IA207019] and CONACYT Ciencia Básica [grant number 241744].

534 **Competing interests**

535 The authors declare that they have no competing interests.

536 **References**

- 537 1. Sear, R. Height and reproductive success: is bigger always better? in *Homo Novus: A Human*
538 *Without Illusions* (eds. Frey, U. J., Störmer, C. & Willführ, K. P.) vol. 44 127–143 (Springer
539 Berlin Heidelberg, 2010).
- 540 2. Pawlowski, B., Dunbar, R. I. M. & Lipowicz, A. Tall men have more reproductive success.
541 *Nature* **403**, 156 (2000).
- 542 3. Stulp, G., Buunk, A. P., Pollet, T. V., Nettle, D. & Verhulst, S. Are Human Mating Preferences
543 with Respect to Height Reflected in Actual Pairings? *PLoS One* **8**, e54186 (2013).
- 544 4. Salska, I. *et al.* Conditional mate preferences: Factors influencing preferences for height. *Pers.*
545 *Individ. Dif.* **44**, 203–215 (2008).
- 546 5. Sear, R., Allal, N. & Mace, R. Height, marriage and reproductive success in Gambian women.
547 *Res. Econ. Anthropol.* **23**, 203–224 (2004).
- 548 6. Silventoinen, K., Lahelma, E. & Rahkonen, O. Social background, adult body-height and health.
549 *Int. J. Epidemiol.* **28**, 911–918 (1999).
- 550 7. Manning, J. T. Fluctuating asymmetry and body weight in men and women: Implications for

- 551 sexual selection. *Ethol. Sociobiol.* **16**, 145–153 (1995).
- 552 8. Pawlowski, B. & Jasienska, G. Women’s preferences for sexual dimorphism in height depend on
553 menstrual cycle phase and expected duration of relationship. *Biol. Psychol.* **70**, 38–43 (2005).
- 554 9. Melamed, T. Personality correlates of physical height. *Pers. Individ. Dif.* **13**, 1349–1350 (1992).
- 555 10. Blaker, N. M. *et al.* The height leadership advantage in men and women: Testing evolutionary
556 psychology predictions about the perceptions of tall leaders. *Gr. Process. Intergr. Relations* **16**,
557 17–27 (2013).
- 558 11. Peck, M. N. & Lundberg, O. Short stature as an effect of economic and social conditions in
559 childhood. *Soc. Sci. Med.* **41**, 733–738 (1995).
- 560 12. Mueller, U. & Mazur, A. Evidence of unconstrained directional selection for male tallness.
561 *Behav. Ecol. Sociobiol.* **50**, 302–311 (2001).
- 562 13. Nettle, D. Height and reproductive success in a cohort of british men. *Hum. Nat.* **13**, 473–491
563 (2002).
- 564 14. Nettle, D. Women’s height, reproductive success and the evolution of sexual dimorphism in
565 modern humans. *Proc. R. Soc. B Biol. Sci.* **269**, 1919–1923 (2002).
- 566 15. Pawlowski, B. Variable preferences for sexual dimorphism in height as a strategy for increasing
567 the pool of potential partners in humans. *Proc. R. Soc. B Biol. Sci.* **270**, 709–712 (2003).
- 568 16. Re, D. E. & Perrett, D. I. Concordant preferences for actual height and facial cues to height. *Pers.*
569 *Individ. Dif.* **53**, 901–906 (2012).
- 570 17. Stearns, S. C. Life history evolution: successes, limitations, and prospects. *Naturwissenschaften*
571 **87**, 476–486 (2000).
- 572 18. Walker, R. *et al.* Growth rates and life histories in twenty-two small-scale societies. *Am. J. Hum.*
573 *Biol.* **18**, 295–311 (2006).

- 574 19. Samaras, T. T. How height is related to our health and longevity: A review. *Nutr. Health* **21**, 247–
575 261 (2012).
- 576 20. Wells, J. The Thrifty Phenotype Hypothesis: Thrifty Offspring or Thrifty Mother? *J. Theor. Biol.*
577 **221**, 143–161 (2003).
- 578 21. Hayflick, L. & Moorhead, P. S. The serial cultivation of human diploid cell strains. *Exp. Cell Res.*
579 **25**, 585–621 (1961).
- 580 22. Giovannelli, L. *et al.* Nutritional and lifestyle determinants of DNA oxidative damage: a study
581 in a Mediterranean population. *Carcinogenesis* **23**, 1483–1489 (2002).
- 582 23. Perry, R. J., Farquharson, C. & Ahmed, S. F. The role of sex steroids in controlling pubertal
583 growth. *Clin. Endocrinol. (Oxf)*. **68**, 4–15 (2008).
- 584 24. Ellison, P. T. *On fertile ground: A natural history of human reproduction*. (Harvard University
585 Press, 2009).
- 586 25. Iravani, M., Lagerquist, M., Ohlsson, C. & Säwendahl, L. Regulation of bone growth via ligand-
587 specific activation of estrogen receptor alpha. *J. Endocrinol.* **232**, 403–410 (2017).
- 588 26. *Eco-immunology: Evolutive Aspects and Future Perspectives*. (Springer, 2014). doi:10.1007/978-
589 94-017-8712-3.
- 590 27. Folstad, I. & Karter, A. J. Parasites, bright males, and the immunocompetence handicap. *Am. Nat.*
591 **139**, 603–622 (1992).
- 592 28. Ansar Ahmed, S., Karpuzoglu, E. & Khan, D. Effects of Sex Steroids on Innate and Adaptive
593 Immunity. in *Sex Hormones and Immunity to Infection* (eds. Klein, S. L. & Roberts, C.) 19–51
594 (Springer, 2010). doi:10.1007/978-3-642-02155-8_2.
- 595 29. Roved, J., Westerdahl, H. & Hasselquist, D. Sex differences in immune responses: Hormonal
596 effects, antagonistic selection, and evolutionary consequences. *Horm. Behav.* **88**, 95–105 (2017).

- 597 30. Bernin, H. & Lotter, H. Sex bias in the outcome of human tropical infectious diseases: Influence
598 of steroid hormones. *J. Infect. Dis.* **209**, (2014).
- 599 31. Neyrolles, O. & Quintana-Murci, L. Sexual Inequality in Tuberculosis. *PLoS Med.* **6**, e1000199
600 (2009).
- 601 32. Nhamoyebonde, S. & Leslie, A. Biological Differences Between the Sexes and Susceptibility to
602 Tuberculosis. *J. Infect. Dis.* **209**, S100–S106 (2014).
- 603 33. Sheldon, B. C. & Verhulst, S. Ecological immunology: Costly parasite defences and trade-offs in
604 evolutionary ecology. *Trends Ecol. Evol.* **11**, 317–321 (1996).
- 605 34. Paajanen, T. A., Oksala, N. K. J., Kuukasjärvi, P. & Karhunen, P. J. Short stature is associated
606 with coronary heart disease: a systematic review of the literature and a meta-analysis. *Eur. Heart*
607 *J.* **31**, 1802–1809 (2010).
- 608 35. Stulp, G. & Barrett, L. Evolutionary perspectives on human height variation. *Biol. Rev.* **91**, 206–
609 234 (2016).
- 610 36. Wormser, D. *et al.* Adult height and the risk of cause-specific death and vascular morbidity in 1
611 million people: individual participant meta-analysis. *Int. J. Epidemiol.* **41**, 1419–1433 (2012).
- 612 37. Henrich, J., Heine, S. J. & Norenzayan, A. The weirdest people in the world? *Behav. Brain Sci.*
613 **33**, 61–83 (2010).
- 614 38. Walker, R. & Hamilton, M. J. Life–History Consequences of Density Dependence and the
615 Evolution of Human Body Size. *Curr. Anthropol.* **49**, 115–122 (2008).
- 616 39. Deaton, A. Height, health, and development. *Proc. Natl. Acad. Sci.* **104**, 13232–13237 (2007).
- 617 40. Garcia, J. & Quintana-Domeque, C. The evolution of adult height in Europe: A brief note. *Econ.*
618 *Hum. Biol.* **5**, 340–349 (2007).
- 619 41. Lim, S. S. *et al.* Measuring the health-related Sustainable Development Goals in 188 countries: a

- 620 baseline analysis from the Global Burden of Disease Study 2015. *Lancet* **388**, 1813–1850 (2016).
- 621 42. Silventoinen, K. Determinants of variation in adult body height. *J. Biosoc. Sci.* **35**, 263–285
622 (2003).
- 623 43. Dowd, J. B., Zajacova, A. & Aiello, A. Early origins of health disparities: Burden of infection,
624 health, and socioeconomic status in U.S. children. *Soc. Sci. Med.* **68**, 699–707 (2009).
- 625 44. Kuzawa, C. W. & Bragg, J. M. Plasticity in Human Life History Strategy. *Curr. Anthropol.* **53**,
626 S369–S382 (2012).
- 627 45. Bentham, J. *et al.* A century of trends in adult human height. *Elife* **5**, e13410 (2016).
- 628 46. Human Development Report Office. *Human Development Indicators and Indices: 2018*
629 *Statistical Update*.
630 http://hdr.undp.org/sites/default/files/2018_human_development_statistical_update.pdf (2018).
- 631 47. Fullman, N. *et al.* Measuring performance on the Healthcare Access and Quality Index for 195
632 countries and territories and selected subnational locations: a systematic analysis from the Global
633 Burden of Disease Study 2016. *Lancet* **391**, 2236–2271 (2018).
- 634 48. Poverty and inequality. *Colombia Reports* (2018).
- 635 49. *Indigenous Peoples, Poverty and Human Development in Latin America*. (The World Bank,
636 2004). doi:10.1596/978-1-4039-9938-2.
- 637 50. Montenegro, R. A. & Stephens, C. Indigenous health in Latin America and the Caribbean. *Lancet*
638 **367**, 1859–1869 (2006).
- 639 51. Biggs, B., King, L., Basu, S. & Stuckler, D. Is wealthier always healthier? The impact of national
640 income level, inequality, and poverty on public health in Latin America. *Soc. Sci. Med.* **71**, 266–
641 273 (2010).
- 642 52. SINAIS. Sistema Nacional de Informacion en Salud. <http://www.sinais.salud.gob.mx> (2016).

- 643 53. Miramontes, O., DeSouza, O., Hernández, D. & Ceccon, E. Non-Lévy Mobility Patterns of
644 Mexican Me'Phaa Peasants Searching for Fuel Wood. *Hum. Ecol.* **40**, 167–174 (2012).
- 645 54. Hernández-Muciño, D. *et al.* La comunidad me'phaa construye su futuro: agroecología y
646 restauración como herramientas de desarrollo rural sustentable. in *Experiencias de colaboración*
647 *transdisciplinaria para la sustentabilidad* (eds. Merçon, J., Ayala-Orozco, B. & Rosell, J. A.) 66–
648 79 (CopIt ArXives, 2018).
- 649 55. Durán-Arenas, L., Gallegos-Carrillo, K., Salinas-Escudero, G. & Martínez-Salgado, H. Towards a
650 Mexican normative standard for measurement of the short format 36 health-related quality of life
651 instrument. *Salud Publica Mex.* **46**, 306–15 (2004).
- 652 56. Duncan, P. T. The Morpho-Syntax of Indefinite Pronouns in Iliatenco Me'phaa. (University of
653 Kansas, 2013).
- 654 57. Ware, J. E. & Sherbourne, C. D. The MOS 36-item short-form health survey (SF-36). I.
655 Conceptual framework and item selection. *Med. Care* **30**, 473–83 (1992).
- 656 58. Lugo A, L. H., García E, H. I. & Gómez R, C. Confiabilidad del cuestionario de calidad de vida
657 en salud SF-36 en Medellín, Colombia. *Rev. Fac. Nac. Salud Publica* **24**, 37–50 (2006).
- 658 59. Czernichow, S., Kengne, A.-P., Stamatakis, E., Hamer, M. & Batty, G. D. Body mass index,
659 waist circumference and waist-hip ratio: which is the better discriminator of cardiovascular
660 disease mortality risk? Evidence from an individual-participant meta-analysis of 82 864
661 participants from nine cohort studies. *Obes. Rev.* **12**, 680–687 (2011).
- 662 60. Després, J. P. & Lemieux, I. Abdominal obesity and metabolic syndrome. *Nature* **444**, 881–887
663 (2006).
- 664 61. Huxley, R., Mendis, S., Zheleznyakov, E., Reddy, S. & Chan, J. Body mass index, waist
665 circumference and waist:hip ratio as predictors of cardiovascular risk—a review of the literature.

- 666 *Eur. J. Clin. Nutr.* **64**, 16–22 (2010).
- 667 62. Knowles, K. M. *et al.* Waist Circumference, Body Mass Index, and Other Measures of Adiposity
668 in Predicting Cardiovascular Disease Risk Factors among Peruvian Adults. *Int. J. Hypertens.*
669 **2011**, 1–10 (2011).
- 670 63. Alberti, K. G. M., Zimmet, P. & Shaw, J. The metabolic syndrome—a new worldwide definition.
671 *Lancet* **366**, 1059–1062 (2005).
- 672 64. Expert Panel on Detection Evaluation and Treatment of High Blood Cholesterol in Adults.
673 Executive Summary of The Third Report of The National Cholesterol Education Program
674 (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In
675 Adults (Adult Treatment Panel III). *JAMA* **285**, 2486–2497 (2001).
- 676 65. R Core Team. R: A language and environment for statistical computing. (2019).
- 677 66. Fox, J. & Monette, G. Generalized Collinearity Diagnostics. *J. Am. Stat. Assoc.* **87**, 178–183
678 (1992).
- 679 67. Wagenmakers, E.-J. & Farrell, S. AIC model selection using Akaike weights. *Psychon. Bull. Rev.*
680 **11**, 192–196 (2004).
- 681 68. Bolker, B. bbmle: Tools for General Maximum Likelihood Estimation. R package version 1.0.20.
682 (2017).
- 683 69. Johnson, P. O. & Fay, L. C. The Johnson-Neyman technique, its theory and application.
684 *Psychometrika* **15**, 349–367 (1950).
- 685 70. Bauer, D. J. & Curran, P. J. Probing Interactions in Fixed and Multilevel Regression: Inferential
686 and Graphical Techniques. *Multivariate Behav. Res.* **40**, 373–400 (2005).
- 687 71. Long, J. A. interactions: Comprehensive, User-Friendly Toolkit for Probing Interactions. R
688 package version 1.1.0. (2019).

- 689 72. Pes, G. M. *et al.* The association of adult height with the risk of cardiovascular disease and cancer
690 in the population of Sardinia. *PLoS One* **13**, e0190888 (2018).
- 691 73. Samaras, T. T., Elrick, H. & Storms, L. H. Is short height really a risk factor for coronary heart
692 disease and stroke mortality? A review. *Med. Sci. Monit.* **10**, RA63-76 (2004).
- 693 74. Lean, M., Han, T. & Seidell, J. Impairment of health and quality of life in people with large waist
694 circumference. *Lancet* **351**, 853–856 (1998).
- 695 75. Schneider, H. J., Klotsche, J., Silber, S., Stalla, G. K. & Wittchen, H.-U. Measuring Abdominal
696 Obesity: Effects of Height on Distribution of Cardiometabolic Risk Factors Risk Using Waist
697 Circumference and Waist-to-Height Ratio. *Diabetes Care* **34**, e7–e7 (2011).
- 698 76. Perry, G. H. & Dominy, N. J. Evolution of the human pygmy phenotype. *Trends Ecol. Evol.* **24**,
699 218–225 (2009).
- 700 77. Harvey, P. H. & Clutton-Brock, T. H. Life History Variation in Primates. *Evolution (N. Y.)*. **39**,
701 559–581 (1985).
- 702 78. Promislow, D. E. L. & Harvey, P. H. Living fast and dying young: A comparative analysis of life-
703 history variation among mammals. *J. Zool.* **220**, 417–437 (1990).
- 704 79. Rollo, C. D. Growth negatively impacts the life span of mammals. *Evol. Dev.* **4**, 55–61 (2002).
- 705 80. Krams, I. A. *et al.* Body height affects the strength of immune response in young men, but not
706 young women. *Sci. Rep.* **4**, 1–3 (2014).
- 707 81. Pawlowski, B., Nowak, J., Borkowska, B., Augustyniak, D. & Drulis-Kawa, Z. Body height and
708 immune efficacy: testing body stature as a signal of biological quality. *Proc. R. Soc. B Biol. Sci.*
709 **284**, 20171372 (2017).
- 710 82. Hopman, W. M. *et al.* Canadian normative data for the SF-36 health survey. *CMAJ* **163**, 265–71
711 (2000).



- 712 83. Watson, E. K., Firman, D. W., Baade, P. D. & Ring, I. Telephone administration of the SF-36
713 health survey: validation studies and population norms for adults in Queensland. *Aust. N. Z. J.*
714 *Public Health* **20**, 359–363 (1996).
- 715 84. Gluckman, P. D. & Hanson, M. A. Evolution, development and timing of puberty. *Trends*
716 *Endocrinol. Metab.* **17**, 7–12 (2006).
- 717 85. Wells, J. C. K., DeSilva, J. M. & Stock, J. T. The obstetric dilemma: An ancient game of Russian
718 roulette, or a variable dilemma sensitive to ecology? *Am. J. Phys. Anthropol.* **149**, 40–71 (2012).
- 719 86. Walters, S. J. & Brazier, J. E. What is the relationship between the minimally important
720 difference and health state utility values? The case of the SF-6D. *Health Qual. Life Outcomes* **1**, 4
721 (2003).
- 722 87. Roberts, S. C. & Little, A. C. Good genes, complementary genes and human mate preferences.
723 *Genetica* **132**, 309–321 (2008).
- 724 88. Leongómez, J. D., Sánchez, O. R., Vásquez-Amézquita, M., Valderrama, E. & González-
725 Santoyo, I. Data from: Self-reported Health is Related to Body Height and Waist Circumference
726 in Rural Indigenous and Urbanised Latin-American Populations. *Open Science Framework*
727 (2019) doi:10.17605/OSF.IO/KGR5X.
- 728 89. Leongómez, J. D., Sánchez, O. R., Vásquez-Amézquita, M., Valderrama, E. & González-
729 Santoyo, I. Code and analyses for Self-reported Health is Related to Body Height and Waist
730 Circumference in Rural Indigenous and Urbanised Latin-American Populations. *Open Science*
731 *Framework* (2019) doi:10.17605/OSF.IO/9WKMT.

732 **Figure legends**

733 **Figure 1. Distribution of all measured variables by sex and sample. (a) Women. (b) Men.** For a comparison of sex
734 differences in height, waist circumference and self-reported health in the three samples, see Supplementary Figure S1 online.


735 **Figure 2. Model selection plot.** Rows represent each of the 334 compared models. Cells coloured in blue represent included
736 terms in each model, according to their Akaike weight ($w_i(AICc)$), represented as the height of each row/model. Given the
737 important age differences between samples, we selected the second-best model (labelled **160**), because it had the same
738 structure as the best model (labelled **159**), but also included Age as a regressor. Furthermore, this second-best model had a
739 $\Delta AICc$ of less than 2 units (≈ 0.98) compared to the best model. For model terms: WC(c) = Waist circumference (centred);
740 H(c) = Height (centred); S = Sample.

741 **Figure 3. Model 3 estimates and interaction between Height and Waist.** Values of Height and Waist were centred: for
742 Height, mean \pm SD = 163.83 \pm 9.85; for Waist circumference, mean \pm SD = 78.01 \pm 10.59. (a) Estimates and 95% CI for
743 each model term. For categorical predictors, women and Bogota were used as reference levels. For model terms, WC(c) =
744 Waist circumference (centred); H(c) = Height (centred); S = Sample. (b) Interaction between Height and Waist. As waist
745 reference, -1 SD (67.43 cm), mean (78.01 cm), and +1 SD (88.6 cm) values were used, showed on a blue to red colour scale.
746 (c) Johnson-Neyman plot, showing for which values of Waist (centred), the slope of Height (centred) is significant as a
747 predictor of Self-reported health; these slopes are predicted to be significant for centred Waists circumferences below -4.51
748 (73.51 cm), or above 68.73 (146.74 cm; not shown as it is a prediction for extreme values, beyond the ones found in any of
749 our samples).

a**Women****b****Men**

Sample ■ Bogota ■ Mexico City ■ Me'Phaa

Cumulative $w_i(A/Cc)$



159
160
223
1183

