Combining amoxicillin and relebactam provides new therapeutic potential

for *Mycobacterium abscessus* infection

Rose C. Lopeman[1†], James Harrison[1†], Maya Desai[2], Peter Lambert[1] and Jonathan A. G. Cox[1]*

[1]School of Life and Health Sciences, Aston University, Aston Triangle, Birmingham, UK, B4 7ET

[2]Birmingham Children's Hospital, Birmingham Women's and Children's NHS Foundation Trust, Steelhouse Lane, Birmingham, UK, B4 6NH

[†]These authors contributed equally to this work.

*Author to whom correspondence should be addressed:

Dr Jonathan A. G. Cox, School of Life and Health Sciences, Aston University, Aston Triangle,

Birmingham, UK, B4 7ET; J.a.g.cox@aston.ac.uk; 0(+44)121 204 5011

ORCID ID: 0000-0001-5208-4056

Abstract:

Infections caused by the opportunistic pathogen *Mycobacterium abscessus* are increasing in prevalence within the cystic fibrosis population. Our limited understanding of this ubiquitous environmental microorganism matched with its intrinsic resistance to most classes of antibiotic, including β-lactams, has left us with an urgent demand for new, more effective therapeutic interventions. *De novo* antimicrobial drug discovery is a lengthy process and so we have taken the approach of repurposing known antibiotics in order to provide a rapidly implementable solution to a current problem. Here we report a significant step forward in treatment potential for *M. abscessus* infection by sensitising the organism to the broad spectrum β-lactam antibiotic, amoxicillin, using the non-competitive β-lactamase inhibitor, relebactam. We demonstrate by disk diffusion and broth microdilution assay that this combination works synergistically to inhibit *M. abscessus.* We also demonstrate the direct non-competitive inhibition of the *M. abscessus* β-lactamase, Bla_{Mab} using a novel thin-layer chromatography-based assay for β-lactamase inhibition, which is subsequently kinetically validated by spectrophotometric assay using the nitrocefin reporter assay. Finally, we demonstrate the *in vitro* efficacy of this combination against a collection of *M. abscessus* clinical isolates, demonstrating the significant therapeutic potential of the amoxicillin and relebactam combination.

Introduction:

The incidence of infections with the rapidly growing non-tuberculous mycobacterial (NTM) organism *Mycobacterium abscessus* is increasing due to the increase of immunocompromised individuals (Petrini, et al. 2006) (Griffith, et al. 2014). *M. abscessus* is widely responsible for opportunistic pulmonary infections in patients with structural lung disorders such as cystic fibrosis (CF) and bronchiectasis (Griffith, et al., 2007) as well as skin and soft tissue infections (SSTIs) in humans (Moore, 1953) (Fitzgerald, et al., 1995). The ubiquitous environmental nature of *M. abscessus* may go some

way to explaining the high levels of intrinsic drug resistance to most major classes of antibiotic that is

observed clinically (Nessar, et al. 2012). Recently, macrolides were added to the ever growing list of

ineffective antimicrobial agents for *M. abscessus* infection, exemplifying the urgent need for new ways

to treat this infection (Nessar, et al. 2012). Additionally, *M. abscessus* is resistant to two of the

frontline antibiotics used for tuberculosis treatment; rifampicin and ethambutol (Nessar, et al. 2012),

and many of the new drugs being discovered for treating tuberculosis do not exhibit any antimicrobial

activity against *M. abscessus* (Soni, et al. 2016).

Pulmonary *M. abscessus* disease was historically treated with macrolide antibiotics such as

clarithromycin (Griffith, et al., 2007), however the steady emergence of antibiotic resistant strains has

prompted the implementation of combination therapy using azithromycin (macrolide) along with

amikacin (aminoglycoside) and at least one other drug of a different class (Leung, et al. 2013). The

British Thoracic Society recommends a biphasic treatment for pulmonary *M. abscessus* infection

comprising of a 1-month initiation phase including intravenously administered amikacin, tigecycline

and imipenem, supplemented with oral clarithromycin or azithromycin (depending on sensitivity

results). This is followed by a 12-month continuation phase comprising of nebulised amikacin, and a

combination of 1-4 of clofazimine, linezolid, minocycline, moxifloxacin or co-trimoxaole (Lopeman et

al. 2019) (Haworth et al 2017).

The advancements in whole genome sequencing technology have enabled the identification of three

distinct subspecies within the *M. abscessus* complex, comprising *M. abscessus* subsp. *abscessus, M.*

abscessus subsp. *bolletii,* and *M. abscessus* subsp. *massiliense,* the latter of which contains a non-

functional *erm*(41) gene, resulting in a macrolide sensitive phenotype (Bryant et al. 2013) (Tortoli et

al. 2016). Mutations in *rrs* and *rrl* 16S rRNA genes have been identified corresponding to high levels of

aminoglycoside resistance, prohibiting the effective use of amikacin in treatment for *M. abscessus*

subsp. a*bscessus* and *M. abscessus* subsp. m*assiliense* clinical isolates specifically (Kehrmann et al.

2016). All *M. abscessus* subsp. contain an endogenous class A β-lactamase (Bla$_{Mab}$) conveying intrinsic

resistance to the β-lactam antibiotics (Soroka et al 2014). The identification of Bla$_{Mab}$, a homolog of

the *Mycobacterium tuberculosis* endogenous β-lactamase, BlaC, sparked widespread investigation

into the potential of β-lactamase inhibitors to supplement the treatment of *M. abscessus* infection. A

study demonstrated the inhibitory activity of avibactam, a non-β-lactam β-lactamase inhibitor,

against Bla$_{Mab}$ significantly improving the *in vitro* and *in vivo* activity of imipenem (Lefebvre et al. 2016).

Very recently, a number of studies have been published demonstrating a number of β-lactam

combinations displaying high levels of *in vitro* synergistic activity against *M. abscessus* complex

(Pandey et al 2019) (Story-Roller et al. 2019). Furthermore, a recent study identified the *in vitro* activity

of two new non-β-lactam β-lactamase inhibitors, namely relebactam and vaborbactam,

demonstrating synergistic activity between imipenem and relebactam as well as meropenem and

vaborbactam (Kaushik et al. 2019).

In this study, we have built on previous work and identified a significant step forward in treatment

potential for *M. abscessus* infection. We have demonstrated the *in vitro* capacity for sensitising the

organism to the broad spectrum β-lactam antibiotic, amoxicillin, using the non-competitive β-

lactamase inhibitor, relebactam. We have demonstrated direct, non-competitive inhibition of

recombinant Bla$_{Mab}$ by relebactam, and demonstrated synergy between amoxicillin and relebactam as

well as activity against a range of clinical isolates. This offers a considerable therapeutic potential as

imipenem, a mainstay of current *M. abscessus* treatment, is currently undergoing phase III clinical trial

in combination with relebactam, suggesting the simple addition of amoxicillin and relebactam to the

treatment regime has the potential to significantly improve the treatment outcome.

Materials and Methods:

Bacterial isolates

A total of 16 *M. abscessus* clinical isolates from Brighton and Sussex Medical School and *M. abscessus*

NCTC 13031 were used in this study. Stock solutions of the isolates were kept in 50% glycerol (Sigma,

Dorset, UK) and Middlebrook 7H9 Broth and stored at -80 °C. Isolates were grown in Middlebrook 7H9

medium supplemented with 10% oleic acid-dextrose-catalase (OADC), 1% glycerol (50% w/v) and

0.05% Tween 80 (v/v) prior to testing. *Escherichia coli* Top 10 cells were used for propagation of

plasmid DNA. These cells were routinely grown in nutrient broth, or nutrient agar (Oxoid, UK) at 37

°C. *E. coli* BL21 (DE3) cells were used for the overproduction of recombinant protein, grown in Terrific

Broth (Melford, UK) at 37 °C.

Antimicrobials

The antimicrobial agents meropenem (MEM), amoxicillin (AMX) and phenoxymethylpenicillin

(Penicillin V/PenV) were obtained from Sigma Aldrich (Dorset, UK) and relebactam was obtained from

Carbosynth (Compton, UK). Stock solutions were prepared in sterile de-ionised water and stored at -

20 °C until use.

Disk diffusions

For the disk diffusion assays, *M. abscessus* clinical isolates were grown in Middlebrook 7H9 Broth to

logarithmic phase and 100 µL of bacterial culture was inoculated into 10 mL Middlebrook 7H9 Broth

supplemented with 0.7% bacteriological agar. This was poured as a layer in agar plates on top of 15

mL Middlebrook 7H11 Agar supplemented with 10% OADC and 1% glycerol. 6 mm sterile filter paper

diffusion disks were placed on the agar and were subsequently impregnated with antibiotic at the

following concentrations in sterile distilled water: relebactam 1 µL of 10 mg/mL, amoxicillin 3.3 µL of

3 mg/mL and meropenem 1 µL of 10 mg/mL. Plates were incubated at 30 °C for 5 days or until clear

zones of inhibition were visualised. Zones of inhibition were measured across the diameter to include

the disk itself.

Broth microdilution assay

The broth microdilution assay was performed as described previously with alterations appropriate to

this study (Institute, Clinican and Laboratory Standards 2012) (Caleffi-Ferracioli et al. 2013). Plates

were prepared by serially diluting AMX (0.128 mg/mL – 0 mg/mL) or MEM (0.128 mg/mL – 0 mg/mL)

in the *x*-axis and REL (0.128 mg/mL – 0 mg/mL) in the *y*-axis. 80 µL of *Mycobacterium abscessus* NCTC

13031 suspension adjusted to an OD_{600} of 0.1-0.2 was inoculated in each well to reach a final well

volume of 100 µL. Plates were sealed and incubated at 30 °C for 5 days or until positive control wells

reached an OD_{600} of 1.0. The MIC of amoxicillin/relebactam combination was determined by

visualising the well that had the lowest concentrations of both compounds and exhibited no bacterial

growth. The relevant well was retroactively plotted into a growth curve over time, and this growth

curve was compared to the wells containing no drug, and the wells containing relebactam only and

amoxicillin only.

Expression and purification of recombinant Bla$_{Mab}$

The *bla$_{Mab}$* gene (MAB_2875) from *Mycobacterium abscessus* NCTC 13031 was amplified by

polymerase chain reaction (PCR), with the first 90 base pairs omitted (resulting in a -30 residue N-

terminal truncated protein). Primers used were as follows (with restriction site underlined): Forward

primer   AAAAAA<u>GGATCC</u>GCGCCGGACGAACTCGCC   and   Reverse   primer

AAAAAA<u>AAGCTT</u>TCAAGCGCCGAAGGCCCG (Eurofins Genomics). Amplicons were purified and cloned

into pET28a using *Bam*HI/*Hin*DIII restriction sites and the sequence was confirmed (Eurofins

Genomics). Bla_{Mab} was expressed in *E. coli* BL21 (DE3) cells by addition of 1 mM Isopropyl β-D-1-

thiogalactopyranoside (IPTG) and incubation at 25 °C for 18 h. Bla_{Mab} was purified by Immobilised

Metal Affinity Chromatography (IMAC) and dialysed into 25 mM Tris HCl pH 7, 100 mM NaCl.

Thin Layer Chromatography (TLC) activity assay

Relebactam (2 mg/mL) was added to recombinant Bla_{Mab} (0.01 mg/mL) and incubated for 5 minutes

at room temperature, before addition of Penicillin V (4 mg/mL) for a further 10 minute incubation at

room temperature. Alongside appropriate control reactions (Figure 2), 1 μL of the reaction was

spotted onto aluminium backed silica gel plates (5735 silica gel 60 F_{254}, Merck) and dried before being

subjected to Thin Layer Chromatography (TLC) using ethyl acetate:water:acetic acid

($C_4H_8O_2$:H_2O:CH_3COOH) (3:1:1, v/v/v). Once dry, plates were visualised by being dipped into $KMnO_4$

TLC stain with light charring.

Spectrophotometric analysis of Bla_{Mab} inhibition by Relebactam

Recombinant Bla_{Mab} (0.25 nM) was mixed with an increasing concentration of relebactam (0, 0.1, 1,

10 and 100 μM) and 100 μM nitrocefin (Carbosynth, UK). The hydrolysis of nitrocefin was monitored

at 486 nm using a Multiskan Go plate reader (Thermo Scientific). Data points were subsequently

plotted as absorbance at 486 nm vs time using Graphpad Prism 7.

Results:

In this paper we have identified that *M. abscessus* can be sensitised to amoxicillin by the addition of the non-competitive β-lactamase inhibitor, relebactam. Furthermore we have shown that meropenem sensitivity can be enhanced by the addition of relebactam. Firstly, we conducted disk diffusion assays with amoxicillin and meropenem with and without relebactam. Subsequently the zones of inhibition (ZOI) were measured as a measure of sensitivity. Amoxicillin alone failed to demonstrate any kind of sensitivity, however the addition of relebactam provided detectable ZOI, demonstrating the induced sensitivity to amoxicillin by relebactam. A ZOI was visible for meropenem, however with the incorporation of relebactam, this ZOI was significantly enhanced, suggesting increased sensitivity upon combination with relebactam (1a). This led us to the hypothesis that relebactam was inhibiting the *M. abscessus* β-lactamase, Bla$_{Mab}$, preventing hydrolysis of amoxicillin to the inactive amoxicilloic acid (1b). In order to assess the clinical relevance of these observations, both amoxicillin and meropenem were tested alone and in combination with relebactam against a panel of clinical isolates of *M. abscessus* obtained from patients at Brighton and Sussex Medical School in addition to *M. abscessus* NCTC strain. In all cases, the addition of relebactam sensitised the isolate to amoxicillin (*n=3*) (1c). The same experiment was conducted with meropenem, and in most cases, an increase in sensitivity was observed (*n=3*) (1d). All necessary controls were conducted and no ZOI was observed for relebactam alone.

In order to further validate our observation, liquid cultures were exposed to increasing concentrations of amoxicillin and meropenem with and without a dose response of relebactam (1e and 1f respectively). As a control, relebactam was also tested for inhibitory activity on its own. These cultures were read every 24 hours using a spectrophotometer at OD_{600nm} up to 96 hours. Relebactam at 128 µg/mL demonstrated no inhibitory activity, giving a growth profile much the same as the compound-free control (no drug), confirming that relebactam lacks antibacterial activity against *M. abscessus*.

32 µg/mL of amoxicillin appeared to show a moderate enhancement of bacterial growth, however

when combined with 2 µg/mL of relebactam, this amoxicillin concentration was found to display

potent antibacterial activity against *M. abscessus*. The same experiment was conducted with

meropenem, wherein partial growth impairment was observed at 8 µg/mL, however when combined

with 2 µg/mL relebactam, significant antibacterial activity was observed at this concentration. End

point statistical analysis was conducted using a t-test to assess the significance of differences between

cultures with and without relebactam. In both cases, a statistically significant increase in sensitivity

was observed upon addition of 2 µg/mL of relebactam in combination with amoxicillin or meropenem.

Figure 1: Relebactam sensitises *Mycobacterium abscessus* to amoxicillin and improves sensitivity to

meropenem. A disk diffusion experiment (1a) and corresponding plate map demonstrating enhanced

sensitivity of *M. abscessus* by zone of inhibition to amoxicillin (1) with the addition of relebactam (2)

and meropenem (3) with the addition of relebactam (4). The inhibition of *M. abscessus* β-lactamase

Bla$_{Mab}$ by relebactam is summarised (1b). The disk diffusion experiments were conducted with the

NCTC *M. abscessus* strain along with a panel of clinical isolates and the diameter of the zones of

inhibition (ZOI) measured in millimetres. All strains of *M. abscessus* tested were resistant to amoxicillin

(AMX) alone, however displayed sensitivity with the addition of relebactam (REL) (1c). The same was

conducted for meropenem (MEM) and likewise, activity was clearly enhanced with the addition of

relebactam (1d). Growth curves were conducted with *M. abscessus* NCTC in medium containing

128 µg/mL relebactam and 32 µg/mL amoxicillin with and without relebactam at 2 µg/mL. Growth

inhibition was only observed with relebactam in combination with amoxicillin (1e). Likewise, the

inhibitory activity of meropenem was clearly enhanced with the addition of relebactam (1f). A t-test

was used for end point analysis between samples +/- relebactam and the results were deemed to be

significant with *p* values of <0.0001 and 0.0152 for amoxicillin and meropenem respectively.

Figure 2. Biochemical analysis of relebactam inhibition of *M. abscessus β*-lactamase, Bla_{Mab}. Our

novel Thin Layer Chromatography (TLC) assay (2a) exhibiting the activity of Bla_{Mab} in the turnover of

penicillin V (high R_f value) to penicilloic acid (lower R_f value). In the absence, or termination of activity

of Bla_{Mab} (by boiling (100 °C for 1 h) or addition of known inhibitor avibactam (Lefebvre et al., 2017)

(200 μg/mL) no lower R_f value spot corresponding to penicilloic acid is seen on the TLC plate. The

addition of relebactam to the reaction between Bla_{Mab} and penicillin V also results in the absence of

the lower Rf value spot, suggesting inhibition of Bla_{Mab} (2a). This observed inhibition is validated by a

spectrophotometric analysis, where the turnover of nitrocefin (100 μM) by Bla_{Mab} (0.25 nM) was

monitored at 486 nm with a varying concentration of relebactam (0, 0.1, 1, 10 and 100 μM). The

increase in concentration of relebactam resulted in partial inhibition of nitrocefin turnover at 1 μM

and total abrogation at 10 μM (2b). This inhibitory activity was seen within 10 seconds of pre-

incubation of relebactam with Bla_{Mab}, before addition of penicillin V, in a time course TLC assay.

However, the addition of relebactam at the same time (t=0) as penicillin V resulted in a lack of

inhibitory activity (2c). The minimum concentration of relebactam required for inhibition of Bla_{Mab} in

the TLC activity assay was assessed using a range of concentrations (200, 20, 2 and 0.2 μg/mL). Activity

of Bla_{Mab} was maintained below a relebactam concentration of 2 μg/mL in the TLC activity assay,

suggesting a minimal concentration of relebactam required to inhibit Bla_{Mab} in the assay is within the

range of 20 to 2 μg/mL (2d).

In order to validate the inhibitory activity observed phenotypically in Figure 1, we conducted

biochemical analysis of the activity of relebactam on the *M. abscessus* endogenous *β*-lactamase,

Bla_{Mab}. The gene was amplified by PCR, digested, ligated into pET28a and sequenced, before

transformation into chemically competent *E. coli* BL21. Bla_{Mab} was expressed, cells harvested and the

enzyme purified by IMAC. We subsequently devised a novel TLC-based assay for assessing *β*-

lactamase activity by separating the penicillin V substrate from the penicilloic acid product. This assay

enabled us to demonstrate the β-lactamase activity of our purified Bla$_{Mab}$, as well as assay the efficacy

of inhibitors against the enzyme (2a). We used avibactam as a positive control, as its inhibitory activity

against Bla$_{Mab}$ has previously been described by Lefebvre et al. 2017. Lane 1 contained protein

purification buffer only, and lane 2 had the addition of enzyme (0.01 mg/ml). Following

chromatography, two lower spots are observed in lanes 1 and 2, indicative of buffer. Penicillin V was

added to lane 3 and gave a characteristic spot of high R$_f$ value, demonstrating unhydrolysed penicillin

resulted in a spot just below the solvent front. Lane 4 was identical to lane 3 with the addition of Bla$_{Mab}$

protein. The hydrolysis of penicillin V to penicilloic acid by Bla$_{Mab}$ resulted in a spot with reduced R$_f$

value. The pre-incubation of enzyme with avibactam in lane 5 resulted in a loss of the lower penicilloic

acid spot, demonstrating the inhibition of Bla$_{Mab}$ activity. In lane 6, relebactam alone did not resolve

on the TLC, but its pre-incubation with Bla$_{Mab}$ before addition to the penicillin substrate in lane 7

resulted in total inhibition of hydrolysis as observed with avibactam (lane 5). Finally, the inhibition of

Bla$_{Mab}$ is further validated by the repeat of lane 4 and lane 7 conditions with heat-denatured Bla$_{Mab}$

(lanes 8 and 9 respectively). This result confirms the direct inhibition of Bla$_{Mab}$ by relebactam (*n=5*).

We further validated this result by using a commercially available β-lactamase substrate, nitrocefin,

which is selectively hydrolysed by β-lactamases, resulting in an increase in absorbance at 486 nm. By

pre-incubating Bla$_{Mab}$ enzyme with a dose response of relebactam (from 0-100 μM), before initiation

of the absorbance assay with the addition of nitrocefin substrate, we observed partial inhibition at

1 μM (0.348 μg/mL) and a complete loss of activity at 10 μM (3.48 μg/mL), confirming the direct

inhibitory activity of relebactam on Bla$_{Mab}$ (2b).

Our TLC-based β-lactamase assay enabled us to further explore the parameters of the inhibitory

activity of relebactam by varying the time of pre-incubation of relebactam with Bla$_{Mab}$ (2c) and the

minimum inhibitory concentration required to abrogate catalytic turnover of the penicillin V substrate

to the penicilloic acid substrate (2d). We found that penicillin V turnover was rapid and that only

addition of relebactam at the same time as the substrate demonstrated turnover of penicillin V,

suggesting non-competitive, irreversible inhibition of Bla_{Mab} by relebactam. The dose response of

relebactam demonstrated total inhibition down to 20 μg/mL, and activity of Bla_{Mab} was maintained

below a relebactam concentration of 2 μg/mL. This suggested a minimal concentration of relebactam

required to inhibit Bla_{Mab} in the assay is within the range of 20 to 2 μg/mL. This corresponds to less

than or equal to 100 fold stoichiometric excess of relebactam required to completely inhibit Bla_{Mab}

(0.5 μM BlaMab to 57.5 μM relebactam corresponding to 20 μg/mL).

Discussion:

In this paper we have demonstrated the inhibitory activity of the non-β-lactam based β-lactamase

inhibitor, relebactam against *M. abscessus* endogenous β-lactamase, Bla_{Mab}. Alone, this is not

sufficient to kill *M. abscessus,* however its use as part of a combination opens up considerable

therapeutic potential. We have identified that a low concentration of relebactam co-administration is

capable of sensitising *M. abscessus* NCTC and clinical isolates to amoxicillin, well within the

therapeutic range for this versatile and widely available antibiotic. Furthermore, we demonstrate

relebactam provides a significant increase in inhibitory activity of meropenem, a mainstay of *M.*

abscessus clinical intervention. Our study introduces a completely novel TLC-based β-lactamase

inhibition assay, validated with the commercially available and widely used nitrocefin assay, to

investigate the parameters of relebactam's inhibitory activity. Relebactam is currently in phase III

clinical trial for administration in combination with imipenem, another mainstay in *M. abscessus* front

line chemotherapy. Our findings therefore represent a timely and highly impactful discovery that is

likely to be translatable into the clinical setting.

Author Contributions:

R. C. L., J. H., M. D., P. L. and J. A. G. C. intellectually conceived and designed the experiments. R. C. L.,

J. H. and J. A. G. C. conducted the experiments. R. C. L., J. H. and J. A. G. C. wrote the manuscript.

Funding: This research was funded by Birmingham Women's and Children's Hospital Charity Research

Foundation (BWCHCRF) (R. C. L. 50% PhD Studentship, match funded by Aston University Prize

Scheme) and the Academy of Medical Sciences and Global Challenges Research Fund with a

Springboard Grant (SBF003\1088:).

Acknowledgements: J. A. G. C. is grateful to the Academy of Medical Sciences, Global Challenges

Research Fund and Birmingham Women's and Children's Hospital Charity Research Foundation

(BWCHCRF) for their continued support of the Mycobacterial Research Group at Aston University. The

Mycobacterium abscessus clinical isolates used in this study were provided by Dr Simon Waddell,

Brighton and Sussex Medical School.

References:

Amit Kaushik, Nicole C. Ammerman,Jin Lee,Olumide Martins, Barry N. Kreiswirth, Gyanu

Lamichhane, Nicole M. Parrish,Eric L. Nuermberger (2019). In vitro activity of the new β-

lactamase inhibitors relebactam and vaborbactam in combination with β-lactams against

Mycobacterium abscessus complex clinical isolates. Antimicrob. Agents Chemother.

doi:10.1128/AAC.02623-18

Bryant, JM, et al. 2013*Whole-genome sequencing to identify transmission of Mycobacterium*

abscessus between patients with cystic fibrosis: a retrospective cohort study. The Lancet, Vol.

381, pp. 1551-1560.

Elizabeth Story-Roller, Emily C. Maggioncalda, Gyanu Lamichhane Select β-lactam

combinations exhibit synergy against *Mycobacterium abscessus in vitro*. Antimicrobial

Agents and Chemotherapy Feb 2019, AAC.02613-18; DOI: 10.1128/AAC.02613-18

Fitzgerald, DA, et al. 1995. *Cutaneous infection with Mycobacterium abscessus.*, Journal of

Dermatology, Vol. 132, pp. 800-804.

Griffith DE. 2014. Mycobacterium abscessus subsp abscessus lung disease: 'trouble ahead,

trouble behind. . .' F1000prime Rep 6:107. https://doi.org/10.12703/P6-107.

Griffith, DE, et al. 2007. *An Official ATS/IDSA Statement: Diagnosis, Treatment, and*

Prevention of Nontuberculous Mycobacterial Diseases. 4, ATS Journals, Vol. 175.

Haworth, CS, et al. 2017. *British Thoracic Society Guideline for the management of non-*

tuberculous mycobacterial pulmonary disease (NTM-PD). BMJ Open Respiratory Research,

Vol. 4, pp. 676 1-12.

Katiany R. Caleffi-Ferracioli, Flaviane G. Maltempe, Vera Lúcia D. Siqueira, Rosilene F.

Cardoso. 2013. Fast detection of drug interaction in *Mycobacterium tuberculosis* by a

checkerboard resazurin method, Tuberculosis, Volume 93, Issue 6, Pages 660-663, ISSN

1472-9792, https://doi.org/10.1016/j.tube.2013.09.001.

Kehrmann J, Kurt N, Rueger K, Bange F-C, Buer J. 2016. GenoType NTM- DR for identifying

Mycobacterium abscessus subspecies and determining molecular resistance. J Clin Microbiol

54:1653–1655. doi:10.1128/JCM.00147-16.

Lefebvre A-L, Le Moigne V, Bernut A, Veckerlé C, Compain F, Herrmann J-L, Kremer L, Arthur

M, Mainardi J-L. 2017. Inhibition of the β-lactamase Bla by avibactam improves the *in vitro*

and *in vivo* efficacy of imipenem against *Mycobacterium abscessus*. Antimicrob Agents

Chemother 61:e02440-16. https://doi.org/ 10.1128/AAC.02440-16.

Leung, J., and Olivier, K. 2013. Nontuberculous mycobacteria in patients with cystic fibrosis.

Semin Respir Crit Care Med. 34(1):124

Lopeman, R.C.; Harrison, J.; Desai, M.; Cox, J.A.G. *Mycobacterium abscessus:* Environmental

Bacterium Turned Clinical Nightmare. *Preprints* 2019, 2019010273 (doi:

10.20944/preprints201901.0273.v1).

Moore, M and Frerichs, JB. 1953. *An Unusual Acid-Fast Infection of the Knee with*

Subcutaneous, Abscess-Like Lesions of the Gluteal Region. 2, Journal of Investigative

Dermatology, Vol. 20.

Nessar R, Cambau E, Reyrat JM, Murray A, Gicquel B. 2012. Mycobacte- rium abscessus: a

new antibiotic nightmare. J Antimicrob Chemother 67:810–818.

https://doi.org/10.1093/jac/dkr578.

Pandey R, Chen L, Manca C, Jenkins S, Glaser L, Vinnard C, Stone G, Lee J, Mathema B,

Nuermberger EL, Bonomo RA, Kreiswirth BN. 2019. Dual β-lactam combinations highly active

against *Mycobacterium abscessus* complex *in vitro*. mBio 10:e02895-18. https://

doi.org/10.1128/mBio.02895-18.

Petrini B. 2006. Mycobacterium abscessus: an emerging rapid-growing potential pathogen.

APMIS 114:319 –328. https://doi.org/10.1111/j.1600 -0463.2006.apm_390.x.

Soni, I., De Groote, M., Dasgupta, A., and Chopra, S. 2016. Challenges facing the drug

discovery pipeline for non-tuberculous mycobacteria. *J Med Microbiol.* 65(1):1-8

Soroka D, Dubee V, Soulier-Escrihuela O, Cuinet G, Hugonnet JE, Gut- mann L, Mainardi JL,

Arthur M. 2014. Characterization of broad-spectrum Mycobacterium abscessus class A beta-

lactamase. J Antimicrob Che- mother 69:691–696. https://doi.org/10.1093/jac/dkt410.

Tortoli, E, Kohl, TA and Brown-Elliot, BA. 2016. *Emended description of Mycobacterium*

abscessus, Mycobacterium abscessus subsp. abscessus and Mycobacterium abscessus subsp.

bolletii and designation of Mycobacterium abscessus subsp. massiliense comb. nov,

International Journal of Systematic and Evolutionary Microbiology, Vol. 66, pp. 4471-4479.

Wayne, PA et al. 2012. CLSI. Methods for Dilution Antimicrobial Susceptibility Tests for

Bacteria That Grow Aerobically; Approved Standard – Ninth Edition. CLSI document M07-A9:

Clinical and Laboratory Standards Institute.

2(a)

	1	2	3	4	5	6	7	8	9
Buffer	X	X	X	X	X	X	X	X	X
Penicillin V			X	X	X		X	X	X
BlaMab		X		X	X		X		
Relebactam						X	X		X
Boiled BlaMab								X	X
Avibactam					X				

2(b)

2(c)

	1	2	3	4	5	6	7	8	9	10
Buffer	X	X		X	X		X	X	X	X
Penicillin V			X	X	X		X	X	X	X
Bla$_{Mab}$		X		X	X		X	X	X	X
Avibactam					X					
Relebactam 0s						X				X
Relebactam 10s							X			
Relebactam 30s								X		
Relebactam 60s									X	

2(d)

	1	2	3	4	5	6	7	8	9	10
Buffer	X	X		X	X		X	X	X	X
Penicillin V			X	X	X		X	X	X	X
Bla$_{Mab}$		X		X	X		X	X	X	X
Avibactam					X					
Relebactam 200 µg/mL							X	X		
Relebactam 20 µg/mL									X	
Relebactam 2 µg/mL										X
Relebactam 0.2 µg/mL										X