Combining amoxicillin and relebactam provides a new therapeutic option

for *Mycobacterium abscessus* infection

Rose C. Lopeman[1†], James Harrison[1†], Savannah E. R. Gibson[1], Daniel L. Rathbone[1], Maya Desai[2],

Peter Lambert[1] and Jonathan A. G. Cox[1]*

[1]School of Life and Health Sciences, Aston University, Aston Triangle, Birmingham, UK, B4 7ET

[2]Birmingham Children's Hospital, Birmingham Women's and Children's NHS Foundation Trust,

Steelhouse Lane, Birmingham, UK, B4 6NH

[†]These authors contributed equally to this work.

*Author to whom correspondence should be addressed:

Dr Jonathan A. G. Cox, School of Life and Health Sciences, Aston University, Aston Triangle,

Birmingham, UK, B4 7ET; J.a.g.cox@aston.ac.uk; 0(+44)121 204 5011

ORCID ID: 0000-0001-5208-4056

Abstract:

Infections caused by the opportunistic pathogen *Mycobacterium abscessus* are increasing in prevalence within the cystic fibrosis population. Our limited understanding of this ubiquitous environmental microorganism matched with its intrinsic resistance to most classes of antibiotic, including β-lactams, has left us with an urgent demand for new, more effective therapeutic interventions. *De novo* antimicrobial drug discovery is a lengthy process and so we have taken the approach of repurposing known antibiotics in order to provide a rapidly implementable solution to a current problem. Here we report a significant potential step forward in the treatment of *M. abscessus* infection by sensitising the organism to the broad spectrum β-lactam antibiotic, amoxicillin, using the competitive β-lactamase inhibitor, relebactam. We demonstrate by disk diffusion and broth microdilution assay that this combination works synergistically to inhibit *M. abscessus*. We also demonstrate the direct competitive inhibition of the *M. abscessus* β-lactamase, Bla_{Mab}, using a novel thin-layer chromatography-based assay for β-lactamase inhibition, which is subsequently validated kinetically by spectrophotometric assay using the nitrocefin reporter assay and *in silico* binding studies. Furthermore, we reverse the sensitivity by overexpressing Bla_{Mab} in *M. abscessus*, demonstrating relebactam-Bla_{Mab} target engagement. Finally, we demonstrate the *in vitro* efficacy of this combination against a collection of *M. abscessus* clinical isolates, as well as the *Mycobacterium tuberculosis* model organism *Mycobacterium bovis* BCG, demonstrating the significant therapeutic potential of the amoxicillin and relebactam combination.

Introduction:

The incidence of infections with the rapidly growing non-tuberculous mycobacterial (NTM) organism *Mycobacterium abscessus* is increasing due to the increase of immunocompromised individuals (Petrini, et al. 2006) (Griffith, et al. 2014). *M. abscessus* is widely responsible for opportunistic pulmonary infections in patients with structural lung disorders such as cystic fibrosis (CF) and

bronchiectasis (Griffith, et al., 2007) as well as skin and soft tissue infections (SSTIs) in humans

(Moore, 1953) (Fitzgerald, et al., 1995). The ubiquitous environmental nature of *M. abscessus* may

go some way to explaining the high levels of intrinsic drug resistance to most major classes of

antibiotic that is observed clinically (Nessar, et al. 2012). Recently, macrolides were added to the

ever growing list of ineffective antimicrobial agents for *M. abscessus* infection, exemplifying the

urgent need for new ways to treat this infection (Nessar, et al. 2012). Additionally, *M. abscessus* is

resistant to two of the frontline antibiotics used for tuberculosis treatment; rifampicin and

ethambutol (Nessar, et al. 2012), and many of the new drugs being discovered for treating

tuberculosis do not exhibit any antimicrobial activity against *M. abscessus* (Soni, et al. 2016).

Pulmonary *M. abscessus* disease was historically treated with macrolide antibiotics such as

clarithromycin (Griffith, et al., 2007), however the steady emergence of antibiotic resistant strains

has prompted the implementation of combination therapy using azithromycin (macrolide) along

with amikacin (aminoglycoside) and at least one other drug of a different class (Leung, et al. 2013).

The British Thoracic Society recommends a biphasic treatment for pulmonary *M. abscessus* infection

comprising of a 1-month initiation phase including intravenously administered amikacin, tigecycline

and imipenem, supplemented with oral clarithromycin or azithromycin (depending on sensitivity

results). This is followed by a 12-month continuation phase comprising of nebulised amikacin, and a

combination of 1-4 of clofazimine, linezolid, minocycline, moxifloxacin or co-trimoxaole (Lopeman et

al. 2019) (Haworth et al 2017).

The advancements in whole genome sequencing technology have enabled the identification of three

distinct subspecies within the *M. abscessus* complex, comprising *M. abscessus* subsp. *abscessus, M.*

abscessus subsp. *bolletii,* and *M. abscessus* subsp. *massiliense,* the latter of which contains a non-

functional *erm*(41) gene, resulting in a macrolide sensitive phenotype (Bryant et al. 2013) (Tortoli et

al. 2016). Mutations in *rrs* and *rrl* 16S rRNA genes have been identified corresponding to high levels

of aminoglycoside resistance, prohibiting the effective use of amikacin in treatment for *M. abscessus*

subsp. a*bscessus* and *M. abscessus* subsp. m*assiliense* clinical isolates specifically (Kehrmann et al.

2016). All *M. abscessus* subsp. contain an endogenous class A β-lactamase (Bla$_{Mab}$) conveying

intrinsic resistance to the β-lactam antibiotics (Soroka et al 2014). The identification of Bla$_{Mab}$, a

homolog of the *Mycobacterium tuberculosis* endogenous β-lactamase, BlaC, sparked widespread

investigation into the potential of β-lactamase inhibitors to supplement the treatment of *M.*

abscessus infection. A study demonstrated the inhibitory activity of avibactam, a non-β-lactam β-

lactamase inhibitor, against Bla$_{Mab}$ significantly improving the *in vitro* and *in vivo* activity of

imipenem (Lefebvre et al. 2016). Recently, multiple studies have been published demonstrating a

number of β-lactam combinations displaying high levels of *in vitro* synergistic activity against *M.*

abscessus complex (Pandey et al 2019) (Story-Roller et al. 2019). Furthermore, one of these studies

identified the *in vitro* activity of two new non-β-lactam β-lactamase inhibitors, namely relebactam

and vaborbactam, demonstrating synergistic activity between imipenem and relebactam as well as

meropenem and vaborbactam (Kaushik et al. 2019).

In this study, we have built on previous work (Kaushik *et al.* 2019) and identified a significant step

forward in treatment potential for *M. abscessus* infection. We demonstrate the *in vitro* capacity for

sensitising the organism to the broad spectrum β-lactam antibiotic, amoxicillin, using the β-

lactamase inhibitor, relebactam. We establish synergy between amoxicillin and relebactam as well

as activity against a range of *M. abscessus* clinical isolates. We have identified the mechanism of

relebactam-Bla$_{Mab}$ inhibition as competitive using recombinant protein in both a novel thin-layer

chromatography (TLC)-based assay, as well as previously established spectrophotometric methods.

Furthermore, this has been delineated using *in silico* molecular docking. This offers a considerable

therapeutic potential as imipenem, a mainstay of current *M. abscessus* treatment, is currently

undergoing phase III clinical trials in combination with relebactam, suggesting the simple addition of

amoxicillin and relebactam to the treatment regime has the potential to significantly improve the

treatment outcome.

Materials and Methods:

Bacterial isolates

A total of 16 *M. abscessus* clinical isolates from Brighton and Sussex Medical School and *M.*

abscessus NCTC 13031 were used in this study, along with *Mycobacterium bovis* BCG Pasteur. Stock

solutions of the isolates were kept in 50% glycerol (Sigma, Dorset, UK) and Middlebrook 7H9 Broth

and stored at -80 °C. Isolates were grown in Middlebrook 7H9 medium supplemented with 10% oleic

acid-dextrose-catalase (OADC), 1% glycerol (50% w/v) and 0.05% Tween 80 (v/v) prior to testing.

Escherichia coli Top 10 cells were used for propagation of plasmid DNA. These cells were routinely

grown in nutrient broth, or nutrient agar (Oxoid, UK) at 37 °C. *E. coli* BL21 (DE3) cells were used for

the overproduction of recombinant protein, grown in Terrific Broth (Melford, UK) at 37 °C.

Antimicrobials

The antimicrobial agents meropenem (MEM), amoxicillin (AMX) and phenoxymethylpenicillin

(Penicillin V/PenV) were obtained from Sigma Aldrich (Dorset, UK) and relebactam (REL) was

obtained from Carbosynth (Compton, UK). Stock solutions were prepared in sterile de-ionised water

and stored at -20 °C until use.

Disk diffusions

For the disk diffusion assays, *M. abscessus* clinical isolates were grown in Middlebrook 7H9 Broth to

logarithmic phase and 100 µL of bacterial culture was inoculated into 10 mL Middlebrook 7H9 Broth

supplemented with 0.7% bacteriological agar. This was poured as a layer in agar plates on top of 15

mL Middlebrook 7H11 Agar supplemented with 10% OADC and 1% glycerol. 6 mm sterile filter paper

diffusion disks were placed on the agar and were subsequently impregnated with antibiotic at the

following concentrations in sterile distilled water: relebactam 1 µL of 10 mg/mL, amoxicillin 3.3 µL of

3 mg/mL and meropenem 1 µL of 10 mg/mL. Plates were incubated at 30 °C for 5 days or until clear

zones of inhibition were visualised. Zones of inhibition were measured across the diameter to

include the disk itself. Disk diffusions were also performed on *M. abscessus* pVV16-*bla*$_{Mab}$ and *M.*

abscessus pVV16.

Broth microdilution assay

The broth microdilution assay was performed as described previously with alterations appropriate to

this study (Institute, Clinican and Laboratory Standards 2012) (Caleffi-Ferracioli et al. 2013). Plates

were prepared by serially diluting amoxicillin (0.128 mg/mL – 0 mg/mL) or meropenem (0.128

mg/mL – 0 mg/mL) in the *x*-axis and relebactam (0.128 mg/mL – 0 mg/mL) in the *y*-axis. 80 µL of

Mycobacterium abscessus NCTC 13031 or *Mycobacterium bovis* BCG suspension adjusted to an

OD$_{600}$ of 0.1-0.2 was inoculated in each well to reach a final well volume of 100 µL. Plates were

sealed and incubated at 30 °C for 5 days for *M. abscessus* cultures or 37 °C for 10 days for *M. bovis*

BCG. The MIC of amoxicillin/relebactam combination was determined by optical density

measurement using a spectrophotometric plate reader and identified as the well that had the lowest

concentrations of both compounds and exhibited no bacterial growth. The relevant well was

retroactively plotted into a growth curve over time, and this growth curve was compared to the

wells containing no drug, and the wells containing relebactam only and amoxicillin only. The broth

microdilution assay was also performed on *M. abscessus* pVV16-*bla*$_{Mab}$ and *M. abscessus* pVV16.

Generation of *M. abscessus* pVV16-*bla~Mab~* overexpressor strain

The *bla~Mab~* gene (MAB_2875) from *Mycobacterium abscessus* NCTC 13031 was amplified by polymerase chain reaction (PCR). Primers used were as follows (with restriction site underlined): Forward primer AAAAAA<u>GGATCC</u>GCGCCGGACGAACTCGCC and Reverse primer AAAAAA<u>AAGCTT</u>AGCGCCGAAGGCCCGCAG (Eurofins Genomics). Amplicons were purified and cloned into pVV16 using *Bam*HI/*Hin*DIII restriction sites and the sequence was confirmed (Eurofins Genomics). Both the plasmid pVV16 and the construct pVV16-*bla~Mab~* were inserted into *M. abscessus* NCTC 13031 cells by electroporation (2.5kV, 2.5 µF and 1000 Ω).

Expression and purification of recombinant Bla~Mab~

The *bla~Mab~* gene, with the first 90 base pairs omitted (resulting in a -30 residue N-terminal truncated protein), was amplified by PCR using the following primers (with restriction sites underlined): Forward primer AAAAAA<u>GGATCC</u>GCGCCGGACGAACTCGCC and Reverse primer AAAAAA<u>AAGCTT</u>TCAAGCGCCGAAGGCCCG (Eurofins Genomics). Amplicons were purified and cloned into pET28a using *Bam*HI/*Hin*DIII restriction sites and the sequence was confirmed (Eurofins Genomics). Bla~Mab~ was expressed in *E. coli* BL21 (DE3) cells by addition of 1 mM Isopropyl β-D-1-thiogalactopyranoside (IPTG) and incubation at 25 °C for 18 h. Bla~Mab~ was purified by Immobilised Metal Affinity Chromatography (IMAC) and dialysed into 25 mM Tris HCl pH 7, 100 mM NaCl.

Thin Layer Chromatography (TLC) activity assay

Relebactam (2 mg/mL) was added to recombinant Bla~Mab~ (0.01 mg/mL) and incubated for 5 minutes at room temperature, before addition of Penicillin V (4 mg/mL) for a further 10 minute incubation at room temperature. Alongside appropriate control reactions (Figure 2), 1 µL of the reaction was

168 spotted onto aluminium backed silica gel plates (5735 silica gel 60 F_{254}, Merck) and dried before

169 being subjected to Thin Layer Chromatography (TLC) using ethyl acetate:water:acetic acid

170 ($C_4H_8O_2$:H_2O:CH_3COOH) (3:1:1, v/v/v). Once dry, plates were visualised by being dipped into $KMnO_4$

171 TLC stain with light charring.

173 **Biochemical analysis of Bla$_{Mab}$ inhibition by Relebactam**

174 Recombinant Bla$_{Mab}$ (0.25 nM) was mixed with an increasing concentration of relebactam (0, 0.1, 1,

175 10 and 100 µM) and 100 µM nitrocefin (Carbosynth, UK). The hydrolysis of nitrocefin was monitored

176 at 486 nm using a Multiskan Go plate reader (Thermo Scientific). This was repeated using a varying

177 concentration of nitrocefin (1-500 µM) with a shorter range of relebactam concentrations (0, 0.5,

178 0.75, 1 and 2.5 µM) and initial velocities (v_i) were plotted against substrate concentration. Data

179 analysis was performed as previously described (Dubée et al., 2014, Papp-wallace et al., 2018). Data

180 analysis was conducted using Graphpad Prism 7.

182 The decarbamylation rate was determined by incubation of Bla$_{Mab}$ (1 µM) with relebactam (20 µM)

183 for 1 hour at 25 °C. The reaction mixture was then subsequently diluted both 10,000 and 50,000 fold

184 (1.8 nM and 0.36 nM relebactam with 90 pM and 18 pM Bla$_{Mab}$ respectively), before addition of 100

185 µM nitrocefin. The reaction was monitored at 486 nm using a Multiskan Go plate reader (Thermo

186 Scientific).

188 ***In Silico* Modelling of the Possible Interaction of Relebactam with the Bla$_{Mab}$**

189 The Bla$_{Mab}$ X-ray crystal structure was obtained from the Protein Data Bank, accession code 4YFM.

190 The A chain was submitted to the GHECOM pocket-finding server (GHECOM 1.0) (Kawabata 2010;

191 Kawabata and Go 2007). The top six pockets identified were used to define the target site for

protein-ligand docking experiments between the enzyme and relebactam using CACHe Worksystem

Pro (version 7.5.0.85, Fujitsu Ltd). The amino acid residues lining theses pockets are given in Figure

3(c). Pocket 1 corresponded to the main (catalytic) site in the enzyme and for the purposes of the

docking experiment was redefined as all amino acid residues within 8 Å of serine 71. Hydrogen

atoms were added using the default settings in line with presumed protonation states for ionisable

amino acid side-chains. The positions of the added hydrogen atoms were optimised by locking the

coordinates of all the non-hydrogen atoms and subjecting the system to a molecular mechanics

(MM2) geometry optimisation. Relebactam was docked four times into each pocket, using CACHe

Worksystem Pro. The amino acid side-chains in each pocket were allowed to be flexible as were all

rotatable bonds in relebactam. The genetic algorithm settings for the docking protocol included

population size 50, maximum generations 3000, crossover rate 0.8, mutation Rate 0.2 and

convergence when the RMSD population fitness was less than 1. The best-scoring consensus

complexes from each series of dockings were taken forward for molecular dynamics simulation. The

required input files were prepared using the Antechamber module of the AMBER Tools package

(Version 14; Case et al, 2005), implementing the ff14SB force field. The system was neutralised by

addition of sodium ions and then solvated within a truncated octahedron of TIP3P water molecules

extending 8 Å from the surface of the protein. Using the Amber 14 molecular dynamics package

CUDA version (Salomon-Ferrer et al, 2013; Goetz et al, 2012; Le Grand et al, 2013), the system was

energy-minimised for 2,000 cycles using a non-bonded cut-off of 12 Å and then heated under

constant volume to 300 K over 25 ps under Langevin dynamics (time step = 1 fs). The heating was

continued at 300 K for at least 200 ns under constant pressure also using Langevin dynamics (SHAKE

on, time step = 2 fs) using the Particle–Mesh–Ewald (PME) method to treat the long range

electrostatic interactions with a 12 Å non-bonded cut-off.

Results:

In this paper we have identified that *M. abscessus* can be sensitised to amoxicillin by the addition of

the competitive β-lactamase inhibitor, relebactam. Furthermore we have shown that meropenem

sensitivity can be enhanced by the addition of relebactam. Firstly, we conducted disk diffusion

assays with amoxicillin and meropenem with and without relebactam. Subsequently the zones of

inhibition (ZOI) were measured as a marker of sensitivity. Amoxicillin alone failed to demonstrate

any kind of sensitivity, however the addition of relebactam provided a clear ZOI, demonstrating the

induced sensitivity to amoxicillin by relebactam. A ZOI was visible for meropenem, however with the

incorporation of relebactam, this ZOI was significantly enhanced, suggesting increased sensitivity

upon combination with relebactam (1a). This led us to the hypothesis that relebactam was inhibiting

the *M. abscessus* β-lactamase, Bla$_{Mab}$, preventing hydrolysis of amoxicillin to the inactive amoxicilloic

acid (1b). In order to assess the clinical relevance of these observations, both amoxicillin and

meropenem were tested alone and in combination with relebactam against a panel of clinical

isolates of *M. abscessus* obtained from patients at Brighton and Sussex Medical School in addition to

M. abscessus NCTC strain. In all cases, the addition of relebactam sensitised the isolate to amoxicillin

(*n=3*) (1c left). The same experiment was conducted with meropenem, and in most cases, an

increase in sensitivity was observed (*n=3*) (1c right). All necessary controls were conducted and no

ZOI was observed for relebactam alone.

In order to further validate our observation, liquid cultures were exposed to increasing

concentrations of amoxicillin and meropenem with and without a dose response of relebactam (1d

top and bottom respectively). As a control, relebactam was also tested for inhibitory activity on its

own. These cultures were read every 24 hours using a spectrophotometer at OD$_{600nm}$ up to 96 hours.

Relebactam at 128 µg/mL demonstrated no inhibitory activity, giving a growth profile much the

same as the compound-free control (no drug), confirming that relebactam lacks antibacterial activity

against *M. abscessus*. 32 µg/mL of amoxicillin appeared to show a moderate enhancement of

bacterial growth, however when combined with 2 µg/mL of relebactam, this amoxicillin

concentration was found to display potent antibacterial activity against *M. abscessus*. The same

experiment was conducted with meropenem, wherein partial growth impairment was observed at

8 µg/mL, however when combined with 2 µg/mL relebactam, significant antibacterial activity was

observed at this concentration. End point statistical analysis was conducted using a t-test to assess

the significance of differences between cultures with and without relebactam. In both cases, a

statistically significant increase in sensitivity was observed upon addition of 2 µg/mL of relebactam

in combination with amoxicillin or meropenem. This experiment was repeated with a Bla_{Mab}

overexpressing strain to prove direct binding of relebactam to Bla_{Mab} (1e). The *M. abscessus*

containing the Bla_{Mab} constitutive overexpression plasmid pVV16-bla_{Mab} lost sensitivity to the

amoxicillin-relebactam combination compared to the empty vector control strain (pVV16-). This

result is replicated using the ZOI method (1c), clearly demonstrated drug-target interaction between

relebactam and Bla_{Mab}. Plate images and plate map for the subsequent resistance induced by Bla_{Mab}

overexpression are shown in Supplementary Figure 1. To briefly explore the wider application for the

amoxicillin-relebactam combination against other mycobacterial infections, we tested its activity in

liquid culture against the widely used model organism for *Mycobacterium tuberculosis,*

Mycobacterium bovis BCG (Supplementary Figure 3). Our results show that *M. bovis* BCG amoxicillin

sensitivity is enhanced from 200 to 25 µg/mL upon addition of 8 µg/mL relebactam.

Figure 1: Relebactam sensitises *Mycobacterium abscessus* to amoxicillin and improves sensitivity to meropenem. A disk diffusion experiment (1a) and corresponding plate map demonstrating enhanced sensitivity of *M. abscessus* by zone of inhibition to amoxicillin (1) with the addition of relebactam (2) and meropenem (3) with the addition of relebactam (4). The inhibition of *M. abscessus* -lactamase Bla$_{Mab}$ by relebactam is summarised (1b). The disk diffusion experiments were conducted with the NCTC *M. abscessus* strain along with a panel of clinical isolates and *M. abscessus* pVV16-*bla$_{Mab}$* and *M. abscessus* pVV16. The diameter of the zones of inhibition (ZOI) measured in millimetres. All strains of *M. abscessus* tested, except *M. abscessus* pVV16-*bla$_{Mab}$*, were resistant to amoxicillin (AMX) alone, however displayed sensitivity with the addition of relebactam (REL). *M. abscessus* pVV16-*bla$_{Mab}$* was resistant to amoxicillin both with and without relebactam (1c). The same was conducted for meropenem (MEM) and likewise, activity was clearly enhanced with the addition of relebactam, even in the case of *M. abscessus* pVV16-*bla$_{Mab}$* which was resistant to meropenem alone but sensitised with the addition of relebactam (1c). Growth curves were conducted with *M. abscessus* NCTC in medium containing 128   relebactam and 32

amoxicillin with and without relebactam at 2 µg/mL. Growth inhibition was only observed with

relebactam in combination with amoxicillin (1d). Likewise, the inhibitory activity of meropenem was

clearly enhanced with the addition of relebactam (1d). A t-test was used for end point analysis

between samples +/- relebactam and the results were deemed to be significant with p values of

<0.0001 and 0.0152 for amoxicillin and meropenem respectively. The same growth curves were

performed with *M. abscessus* pVV16-*bla$_{Mab}$* and *M. abscessus* pVV16 in medium containing

32 µg/mL amoxicillin with and without relebactam at 2 µg/mL (1e). *M. abscessus* pVV16-*bla$_{Mab}$*

displayed resistance to amoxicillin and relebactam together whereas *M. abscessus* pVV16 retained

sensitivity to amoxicillin and relebactam ($p = 0.0017$).

In order to validate the inhibitory activity observed phenotypically in Figure 1, we conducted

biochemical analysis of the activity of relebactam on the *M. abscessus* endogenous β-lactamase,

Bla$_{Mab}$. The gene was amplified by PCR, digested, ligated into pET28a and sequenced, before

transformation into chemically competent *E. coli* BL21. Bla$_{Mab}$ was expressed, cells harvested and the

enzyme purified by IMAC. We subsequently devised a novel TLC-based assay for assessing β-

lactamase activity by separating the penicillin V substrate from the penicilloic acid product. This

assay enabled us to demonstrate the β-lactamase activity of our purified Bla$_{Mab}$ (2a), as well as assay

the efficacy of inhibitors against the enzyme (Supplementary 2a and 2b). We used avibactam as a

positive control, as its inhibitory activity against Bla$_{Mab}$ has previously been described by Lefebvre et

al. 2017. Lane 1 contained protein purification buffer only, and lane 2 had the addition of enzyme

(0.01 mg/mL). Following chromatography, two lower spots are observed in lanes 1 and 2, indicative

of buffer. Penicillin V was added to lane 3 and gave a characteristic spot of high R_f value,

demonstrating unhydrolysed penicillin resulted in a spot just below the solvent front. Lane 4 was

identical to lane 3 with the addition of Bla$_{Mab}$ protein. The hydrolysis of penicillin V to penicilloic acid

by Bla$_{Mab}$ resulted in a spot with reduced R_f value. The pre-incubation of enzyme with avibactam in

lane 5 resulted in a loss of the lower penicilloic acid spot, demonstrating the inhibition of Bla_{Mab}

activity. In lane 6, relebactam alone did not resolve on the TLC, but its pre-incubation with Bla_{Mab}

before addition to the penicillin substrate in lane 7 resulted in total inhibition of hydrolysis as

observed with avibactam (lane 5). Finally, the inhibition of Bla_{Mab} is further validated by the repeat of

lane 4 and lane 7 conditions with heat-denatured Bla_{Mab} (lanes 8 and 9 respectively). This result

confirms the direct inhibition of Bla_{Mab} by relebactam (*n=5*).

We investigated the kinetic parameters of the Michaelis-Menten kinetics, the apparent K_i ($K_{i\ app}$), the

second-order carbamylation rate constant (K_2/K_i) and the decarbamylation rate constant (K_{off}) of

relebactam inhibition of soluble recombinant Bla_{Mab} using a commercially available β-lactamase

substrate, nitrocefin, as a reporter substrate. Nitrocefin is selectively hydrolysed by β-lactamases,

resulting in an increase in absorbance which can be monitored at 486 nm. By pre-incubating Bla_{Mab}

enzyme with a dose response of relebactam (from 0-100 μM), before initiation of the absorbance

assay with the addition of nitrocefin substrate, we observed partial inhibition at 1 μM (0.348 μg/mL)

and a complete loss of activity at 10 μM (3.48 μg/mL), confirming the direct inhibitory activity of

relebactam on Bla_{Mab} (2b).

The initial velocities (v_i) of nitrocefin turnover were recorded for a range of nitrocefin concentrations

(1-500 μM) over a range of relebactam concentrations (0-2.5 μM). These results were plotted as

classical Michaelis-Menten curves (v_i vs [S]) (2c), resulting in Michaelis constants (K_m) between

69.43-16.96 μM, before initial velocities were abolished at 2.5 μM relebactam. However, we were

unable to derive a K_i value using this data. In response, we plotted the reciprocal initial velocities

against relebactam concentrations as a linear equation and derived $K_{i\ app}$ observed from the Y

intercept/slope, which was then normalised for the use of nitrocefin (2f) (Papp-Wallace et al., 2018).

The $K_{i\ app}$ value obtained of 183.6 nM is indicative of the high inhibitory potency of relebactam

against Bla_{Mab}. The carbamylation rate constant (K_2/K_i) of 2.32 x 10^6 $M^{-1}s^{-1}$ is similar to the observed

rate for avibactam inhibition of Bla_{Mab} (4.9 x 10^5 $M^{-1}s^{-1}$) (Dubée et al., 2014). The rate of

decarbamylation (K_{off}) of Bla_{Mab} for relebactam was 0.0146 min^{-1}, which is again similar to the rate

previously obtained for that of avibactam (0.047 min^{-1}) (Dubée et al., 2014). Our kinetics analysis

show that, like avibactam, relebactam is a potent, competitive and reversible inhibitor of Bla_{Mab},

displaying a reasonably rapid "on" rate and slow "off" rate, with only half the enzyme recovering

activity after 40 minutes in the absence of relebactam (2e) (Dubée et al., 2014).

Our TLC-based β-lactamase assay enabled us to further explore the parameters of the inhibitory

activity of relebactam by varying the time of pre-incubation of relebactam with Bla_{Mab}

(Supplementary 2a) and the minimum inhibitory concentration required to abrogate catalytic

turnover of the penicillin V substrate to the penicilloic acid substrate (Supplementary 2b). We found

that penicillin V turnover was rapid and that only addition of relebactam at the same time as the

substrate demonstrated turnover of penicillin V, suggesting competitive, reversible inhibition of

Bla_{Mab} by relebactam. The dose response of relebactam demonstrated total inhibition down to 20

µg/mL, and activity of Bla_{Mab} was maintained below a relebactam concentration of 2 µg/mL. This

suggested a minimal concentration of relebactam required to inhibit Bla_{Mab} in the assay is within the

range of 20 to 2 µg/mL. This corresponds to a less than or equal to 100 fold stoichiometric excess of

relebactam required to completely inhibit Bla_{Mab} (0.5 µM BlaMab to 57.5 µM relebactam

corresponding to 20 µg/mL).

2(a)

Penicillin V

Penicilloic acid

	1	2	3	4	5	6	7	8	9
Buffer	X	X	X	X	X	X	X	X	X
Penicillin V			X	X	X		X	X	X
BlaMab		X		X	X		X		
Relebactam						X	X		X
Boiled BlaMab								X	X
Avibactam					X				

2(b)

• 0 μM REL   ■ 0.1 μM REL   ▲ 1 μM REL

▼ 10 μM REL   ○ 100 μM REL

2(c)

—●— 0 μM REL   —▲— 0.5 μM REL
—▼— 0.75 μM REL   —◆— 1 μM REL
—●— 2.5 μM REL

2(d)

2(e)

• 0 μM REL   ■ 10,000x dilution
▲ 50,000x dilution   ▼ 0 μM BlaMab

2(f)

Kinetic parameter	Value for Bla$_{Mab}$
K_m	69.43 μM
$K_{i\,app}$	183.6 nM
K_2/K_i	2.32 x 10^6 M^{-1}s^{-1}
K_{off}	0.0146 min^{-1}

Figure 2. Biochemical analysis of relebactam inhibition of *M. abscessus* β-lactamase, Bla$_{Mab}$. Our

novel Thin Layer Chromatography (TLC) assay (2a) exhibiting the activity of Bla$_{Mab}$ in the turnover of

penicillin V (high R$_f$ value) to penicilloic acid (lower R$_f$ value). In the absence, or termination of

activity of Bla$_{Mab}$ (by boiling (100 °C) for 1 h) or addition of known inhibitor avibactam (Lefebvre et

al., 2017) (200 µg/mL) no lower R$_f$ value spot corresponding to penicilloic acid is seen on the TLC

plate. The addition of relebactam to the reaction between Bla$_{Mab}$ and penicillin V also results in the

absence of the lower Rf value spot, suggesting inhibition of Bla$_{Mab}$ (2a). This observed inhibition is

validated by a spectrophotometric analysis, where the turnover of nitrocefin (100 µM) by Bla$_{Mab}$

(0.25 nM) was monitored at 486 nm with a varying concentration of relebactam (0, 0.1, 1, 10 and

100 µM). The increase in concentration of relebactam resulted in partial inhibition of nitrocefin

turnover at 1 µM and total abrogation at 10 µM (2b). The initial velocity (v$_i$) of the reaction between

Bla$_{Mab}$ and nitrocefin was monitored for a range of substrate concentrations (1-500 µM) and

relebactam concentrations (0, 0.5, 0.75, 1 and 2.5 µM). This data was plotted v$_i$ vs [S] in order to

determine K_m values (2c/f). The values for K_{obs} were obtained as previously described (Dubée et al.,

2014) and plotted against relebactam concentrations ([I]) to deduce a carbamylation rate (K_2/K_i) for

Bla$_{Mab}$ (2d). The kinetics of Bla$_{Mab}$ decarbamylation were assessed to show the recovery of nitrocefin

hydrolysis by Bla$_{Mab}$ after inhibition by relebactam in order to derive a K_{off} value (2e). Kinetic

parameters were derived as described previously (2f) (Dubée et al., 2014).

In order to further investigate the mechanism of relebactam inhibition of Bla_{Mab}, we conducted

molecular docking simulations *in silico*. 6 potential binding sites were identified (3(c)). For pockets 2-

6 the ligand was weakly-held and generally exited the pocket after a few tens of nanoseconds. For

pocket 1 (corresponding to the main active site) the ligand reoriented itself relative to the docked

conformation and thereafter remained relatively stable within the pocket. The binding interactions

for the stable pose are shown in Figure 3(a) and 3(b) and include several polar interactions with the

sulphonamide moiety and hydrophobic interactions between the relebactam central piperidine ring

and tryptophan 106. In addition, after the initial reorientation, the relebactam carbonyl carbon

remained in the vicinity of the hydroxyl oxygen of the catalytically-active serine 71 as can be seen

after 120 ns in the distance plot given in Figure 3(d). The average distance in this period was

approximately 5.5 Å and there were many close approaches. Furthermore, the corresponding O-C=O

angle (Figure 3d) in the same period was roughly 130 ° and as such it is reasonable to assume that it

would be possible for the serine hydroxyl to attack the relebactam carbonyl and effect a ring-

opening. A further molecular dynamics simulation of the enzyme in a periodic box of explicit water

was undertaken but this time in the presence of ten copies of unbound relebactam. Over the course

of a 200 ns simulation, one ligand found its way into the main active site (pocket 1) after 130 ns and

remained stable therein. The other nine copies of the ligand found no place to reside in the enzyme.

3(a) 3(b) 3(c)

3(d)

386
Figure 3: *In Silico* **modelling of the possible interaction of relebactam with the** *M. abscessus* **β-**
lactamase, Bla$_{Mab}$. 3D and 2D protein-ligand interaction diagrams for relebactam in the main
(catalytic) active site after molecular dynamics simulation (3a and 3b respectively). Amino acid
residues featured in the top six potential binding pockets identified for Bla$_{Mab}$. Pocket 1
corresponded to the main (catalytic site) in the enzyme and for the purposes of the docking
experiment was redefined as all amino acid residues within 8 Å of serine 71 (3c). Time-courses of the
serine 71 hydroxyl oxygen – relebactam carbonyl carbon distance and the corresponding O-C=O
angle. The actual values are plotted as black points and a 50-frame moving average is over-plotted in
red (3d).

Discussion:

In this paper we have demonstrated the inhibitory activity of the non-β-lactam based β-lactamase
inhibitor, relebactam against *M. abscessus* endogenous β-lactamase, Bla$_{Mab}$. Alone, this is not
sufficient to kill *M. abscessus,* however its use as part of a combination opens up considerable

therapeutic potential. We have identified that a low concentration of relebactam co-administration

is capable of sensitising *M. abscessus* NCTC and clinical isolates to amoxicillin, well within the

therapeutic range for this versatile and widely available antibiotic. Furthermore, we demonstrate

relebactam provides a significant increase in inhibitory activity of meropenem, a mainstay of *M.*

abscessus clinical intervention. Our study introduces a completely novel TLC-based β-lactamase

inhibition assay, validated with the commercially available and widely used nitrocefin assay, to

investigate the parameters of relebactam's inhibitory activity. The results of both the kinetics

analysis and the *in silico* binding studies demonstrate the mechanism of inhibition of Bla_{Mab} by

relebactam as competitive reversible, displaying a reasonably rapid "on" rate and slow "off" rate,

indicative of the high potency of the inhibitor against Bla_{Mab}.

Relebactam is currently in phase III clinical trial for administration in combination with imipenem,

another mainstay in *M. abscessus* front line chemotherapy. Our findings therefore represent a

timely and highly impactful discovery that is easily translatable into the clinical setting, not only

providing a new therapeutic option for *M. abscessus* infections, but with the potential to treat other

mycobacterial infectious diseases such as tuberculosis.

Author Contributions:

R. C. L., J. H., M. D., D. L. R, P. L. and J. A. G. C. intellectually conceived and designed the experiments.

R. C. L., J. H., S. E. R. G, D. L. R. and J. A. G. C. conducted the experiments. R. C. L., J. H., D. L. R. and J.

A. G. C. wrote the manuscript.

Funding: This research was funded by Birmingham Women's and Children's Hospital Charity

Research Foundation (BWCHCRF) (R. C. L. 50% PhD Studentship, match funded by Aston University

Prize Scheme) and the Academy of Medical Sciences and Global Challenges Research Fund with a

Springboard Grant (SBF003\1088:).

Acknowledgements: J. A. G. C. is grateful to the Academy of Medical Sciences, Global Challenges

Research Fund and Birmingham Women's and Children's Hospital Charity Research Foundation

(BWCHCRF) for their continued support of the Mycobacterial Research Group at Aston University.

The *Mycobacterium abscessus* clinical isolates used in this study were provided by Dr Simon

Waddell, Brighton and Sussex Medical School. We are also grateful to Dr Amit Chattopadhyay, Aston

University, for technical support.

References:

Bryant, J. M., Grogono, D. M., Greaves, D., Foweraker, J., Roddick, I., Inns, T., Reacher, M., Haworth,

C. S., Curran, M. D., Harris, S. R., Peacock, S. J., Parkhill, J., Floto, R. A. (2013) Whole-genome

sequencing to identify transmission of Mycobacterium abscessus between patients with cystic

fibrosis: a retrospective cohort study. *Lancet,* 381, 1551-1560

Case, D. A., Cheatham, T. E., Darden, T., Gohlke, H., Luo, R., Merz, K. M., Onufriev, A., Simmerling,

C., Wang, B., Woods, R. J., (2005) The Amber biomolecular simulation programs. *Journal of*

Computational Chemistry, 26, 1668-1688.

Dubée, V., Bernut, A., Cortes, M., Lesne, T., Dorchene, D., Lefebvre, A., Hugonnet, J., Gutmann, L.,

Mainardi, J., Herrmann, J., Gaillard, J., Kremer, L., Arthur, M., (2014) β-Lactamase inhibition by

avibactam in Mycobacterium abscessus, Journal of Antimicrobial Chemotherapy, 70 (4), 1051–1058.

Fitzgerald, D. A., Smith, A. G., Lees, A., Yee, L., Cooper, N., Harris, S.C., Gibson, J. A. (1995) Cutaneous

infection with Mycobacterium abscessus. *British Journal of Dermatology*. 132. 800-804.

Goetz, A. W., Williamson, M. J., Xu, D., Poole, D., Le Grand, S., Walker, R. C., (2012) Routine
microsecond molecular dynamics simulations with AMBER on GPUs. 1. Generalized born. *Journal*
of Chemical Theory and Computation, 8, 1542-1555.

Griffith, D. E., Aksamit, T., Brown-Elliott, B. A., Catanzaro, A., Daley, C., Gordin, F., Holland, S. M.,
Horsburgh, R., Huitt, G., Iademarco, M. F., Iseman, M., Olivier, K., Ruoss, S., von Reyn, C. F., Wallace,
R. J., Winthrop, K. (2007) An Official ATS/IDSA Statement: Diagnosis, Treatment, and Prevention of
Nontuberculous Mycobacterial Diseases. *American Journal of Respiratory Critical Care Medicine*.
175, 367-416

Griffith, D.E. (2014) *Mycobacterium abscessus* subsp *abscessus* lung disease: 'trouble ahead, trouble
behind…'. *F1000Prime Reports*. 6, 1-6

Haworth, C. S., Banks, J., Capstick, T., Fisher, A. J., Gorsuch, T., Laurenson, I. F., Leitch, A., Loebinger,
M. R., Milburn, H. J., Nightingale, M., Ormerod, P., Singhadia, D., Smith, D., Whitehead, N., Wilson,
R., Floto, R. A. (2017) British Thoracic Society guidelines for the management of non-tuberculous
mycobacterial pulmonary disease (NTM-PD). *Thorax*, 72, 1-64

Caleffi-Ferracioli, K.R., Maltempe, F. G., Siqueira, V. L. D., Cardoso, R. F., (2013) Fast detection of
drug interaction in *Mycobacterium tuberculosis* by a checkerboard resazurin method, Tuberculosis,
Volume 93, Issue 6, Pages 660-663, ISSN 1472-9792, https://doi.org/10.1016/j.tube.2013.09.001.

Kaushik, A., Ammerman, N. C., Lee, J., Martins, O., Kreiswirth, B. N., Lamichhane, G., Parrish, N. M.,
Nuermberger, E. L. (2019) *In Vitro* Activity of the New β-Lactamase Inhibitors Relebactam and
Vaborbactam in Combination with β-Lactams against Mycobacterium abscessus Complex Clinical
Isolates. *Antimicrobial Agents and Chemotherapy,* 63, 1-11

Kawabata, T., Go, N., (2007) Detection of pockets on protein surfaces using small and large probe
spheres to find putative ligand binding sites. *Proteins*, 68,516-529.

Kawabata, T., (2010) Detection of multi-scale pockets on protein surfaces using mathematical

morphology. *Proteins*, 78, 1195-1121.

Kehrmann, J., Kurt, N., Rueger, K., Bange, F.C., Buer, J., (2016) GenoType NTM- DR for identifying

Mycobacterium abscessus subspecies and determining molecular resistance. J Clin Microbiol

54:1653–1655. doi:10.1128/JCM.00147-16.

Le Grand, S., Goetz, A. W., Walker, R. C. (2013) SPFP: Speed without compromise - a mixed

precision model for GPU accelerated molecular dynamics simulations. *Computer Physics*

Communications, 184, 374-380.

Lefebvre A-L, Le Moigne V, Bernut A, Veckerlé C, Compain F, Herrmann J-L, Kremer L, Arthur M,

Mainardi J-L. 2017. Inhibition of the β-lactamase Bla by avibactam improves the *in vitro* and *in vivo*

efficacy of imipenem against *Mycobacterium abscessus*. Antimicrob Agents Chemother 61:e02440-

16. https://doi.org/ 10.1128/AAC.02440-16.

Leung, J., and Olivier, K. 2013. Nontuberculous mycobacteria in patients with cystic fibrosis. *Semin*

Respir Crit Care Med. 34(1):124

Lopeman, R.C.; Harrison, J.; Desai, M.; Cox, J.A.G. *Mycobacterium abscessus:* Environmental

Bacterium Turned Clinical Nightmare. *Preprints* 2019, 2019010273 (doi:

10.20944/preprints201901.0273.v1).

Moore, M and Frerichs, JB. 1953. *An Unusual Acid-Fast Infection of the Knee with Subcutaneous,*

Abscess-Like Lesions of the Gluteal Region. 2, Journal of Investigative Dermatology, Vol. 20.

Nessar R, Cambau E, Reyrat JM, Murray A, Gicquel B. 2012. Mycobacterium abscessus: a new

antibiotic nightmare. J Antimicrob Chemother 67:810–818. https://doi.org/10.1093/jac/dkr578.

Pandey R, Chen L, Manca C, Jenkins S, Glaser L, Vinnard C, Stone G, Lee J, Mathema B, Nuermberger
EL, Bonomo RA, Kreiswirth BN. 2019. Dual β-lactam combinations highly active against
Mycobacterium abscessus complex *in vitro*. mBio 10:e02895-18. https://
doi.org/10.1128/mBio.02895-18.

Papp-Wallace, K. M., Barnes, M. D., Alsop, J., Taracila, M. A., Bethel, C. R., Becka, S. A., van Duin,
D., Kreiswirth, B. N., Kaye, K. S., Bonomo, R. A., (2018) Relebactam Is a Potent Inhibitor of the KPC-2
β-Lactamase and Restores Imipenem Susceptibility in KPC-Producing *Enterobacteriaceae*
Antimicrobial Agents and Chemotherapy, 62 (6) e00174-18.

Petrini B. 2006. Mycobacterium abscessus: an emerging rapid-growing potential pathogen. APMIS
114:319 –328. https://doi.org/10.1111/j.1600 -0463.2006.apm_390.x.

Salomon-Ferrer, R., Goetz, A. W., Poole, D., Le Grand, S., Walker, R. C., (2013) Routine
microsecond molecular dynamics simulations with AMBER - Part II: Particle Mesh Ewald. *Journal*
of Chemical Theory and Computation, 9, 3878-3888.

Soni, I., De Groote, M., Dasgupta, A., and Chopra, S. 2016. Challenges facing the drug discovery
pipeline for non-tuberculous mycobacteria. *J Med Microbiol.* 65(1):1-8

Soroka D, Dubee V, Soulier-Escrihuela O, Cuinet G, Hugonnet JE, Gut- mann L, Mainardi JL, Arthur M.
2014. Characterization of broad-spectrum Mycobacterium abscessus class A beta-lactamase. J
Antimicrob Che- mother 69:691–696. https://doi.org/10.1093/jac/dkt410.

Story-Roller, E., Maggioncalda, E. C., Lamichhane, G. (2019) Select β-Lactam Combinations Exhibit
Synergy against *Mycobacterium abscessus In Vitro*. *Antimicrobial Agents and Chemotherapy*. 63, 1-
11

Tortoli, E, Kohl, TA and Brown-Elliot, BA. 2016. *Emended description of Mycobacterium abscessus,*
Mycobacterium abscessus subsp. abscessus and Mycobacterium abscessus subsp. bolletii and

designation of Mycobacterium abscessus subsp. massiliense comb. nov, International Journal of

Systematic and Evolutionary Microbiology, Vol. 66, pp. 4471-4479.

Wayne, P. A. (2015) Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow

Aerobically; Approved Standard – Tenth Edition. *CLSI document M07-A10.* 32.

2(a)

Penicillin V

Penicilloic acid

	1	2	3	4	5	6	7	8	9
Buffer	X	X	X	X	X	X	X	X	X
Penicillin V			X	X	X		X	X	X
BlaMab		X		X	X		X		
Relebactam						X	X		X
Boiled BlaMab								X	X
Avibactam					X				

2(b)

2(c)

2(d)

2(e)

2(f)

Kinetic parameter	Value for Bla$_{Mab}$
K_m	69.43 μM
$K_{i\,app}$	183.6 nM
K_2/K_i	2.32 x 10^6 M^{-1}s^{-1}
K_{off}	0.0146 min^{-1}

3(a)

3(b)

Interactions
- Attractive Charge (orange)
- Conventional Hydrogen Bonding (green)
- Pi-Alkyl (pink)

3(d)

3(c)

Pocket 1	Pocket 2	Pocket 3	Pocket 4	Pocket 5	Pocket 6
LEU68	ARG45	MET69	PHE76	THR207	ARG95
MET69	GLU65	CYS70	ILE77	ASP208	TYR98
CYS70	ARG66	SER71	VAL78	TRP209	ALA107
SER71	PHE67	THR72	SER79	LEU210	PRO108
TYR72	LEU68	VAL73	ALA80	LYS211	ILE109
VAL73	MET69	LYS79	ILE81	ARG212	THR110
LYS334	CYS70	THR75	LEU82	ASN213	SER111
THR75	ARG165	PHE76	ARG83	THR214	GLN112
PHE76	LEU170	ILE77	ARG84	THR215	HIS113
ILE77	ASN171	VAL78	ARG85	GLY216	GLY117
TRP106	ILE172	ALA80	LEU86	ASP127	MET118
THR127	PRO173	THR127	SER87	GLN218	THR119
LEU128	ASP174	ALA136	GLU88	SER219	VAL120
TYR130	GLY175	ASN137	LEU91	ILE220	SER121
ASP132	ASP176	LEU139	GLN142	ARG221	GLU122
ASN133	LEU177	ILE140	LEU143	ALA222	LEU123
ALA136	ASP178	THR141	GLY145	ALA223	CYS124
GLU167	THR179	LEU143	GLY145	VAL224	ASP125
VAL168	SER180	GLY144	LYS147	PRO225	ALA126
LEU170	GLY237	GLY145	GLU148	ALA226	THR127
ASN171	PHE238	PRO146	THR149	GLY227	LEU128
ASP178	LYS239	LYS147	GLU150	TRP228	ARG129
LEU210	GLY240	GLU148	LYS151	THR229	TYR130
THR215	GLU241	THR149	PHE152	VAL230	ASP131
LYS233	THR242	GLU150	VAL153	ALA231	GLY135
THR234	ASN243	LYS151	ARG154	ASP232	TRP209
GLY235	LEU261	VAL153	SER155	LYS233	ARG212
GLY236	THR262	THR161	LEU156	THR234	
GLY237	VAL263	ARG162	GLY197	ASP244	
ASN243	PRO264	MET163	LEU198	ILE345	
ILE234		ASP164		ALA246	
		ARG165		VAL247	
		THR166		ILE348	
		GLU167		TRP249	
		GLN169		PRO250	
		LEU170		PRO251	
		ASN171		ILE256	
		LEU177		MET258	
		ASP178		ILE276	
		THR179		ALA277	
				THR380	
				ARG281	
				LEU284	
				ALA289	