Noninvasive prenatal diagnosis for Haemophilia A by a haplotype-based approach using cell-free fetal DNA in maternal plasma

Zhihui He[1], Chao Chen[2], Fengyu Guo[2], Lu Jiang[2], Yaping Zhu[2], Dan Li[1], Renhua Wu[2], Rong Lu[1], Mei Zhao[3], Fang Chen[4], Peixiang Ni[2], Weiying Jiang[5*], Jun Sun[2*], Zhiyu Peng[4*]

[1]Department of Obstetrics and Gynecology, The First Affiliated Hospital of Guangzhou Medical University, 151#, Yanjiang Road, Guangzhou, 510120, Guangdong, China;

[2]Tianjin Medical Laboratory, BGI-Tianjin, BGI-Shenzhen, Central Avenue 55, Airport Business Park East Building E3, Tianjin, 300308, China;

[3]BGI-Guangzhou Medical Laboratory, BGI-Shenzhen, Floor 4, Area B, Xinghai National Digital Family, 22#, Qinglan Street, Panyu District, Guangzhou, 510006, Guangdong, China;

[4]BGI Genomics, BGI-Shenzhen, Bei Shan Industrial Zone, Yantian District, Shenzhen, 518083, Guangdong, China;

[5]Department of Medical Genetics, Medical School, Sun Yat-Sen University, 74#, Zhongshan Rd 2, Guangzhou, 510120, Guangdong, China.

The first two authors contributed equally to this work.

Correspondence:

Zhiyu Peng (Bei Shan Industrial Zone, Yantian District, Shenzhen, China,

+8618665899278, fax number: +86(755)33023888, pengzhiyu@genomics.cn)

Jun Sun (Central Avenue 55, Airport Business Park East Building E3, Tianjin, China,

+8613622027365, fax number: none, sunjun@genomics.cn)

Weiying Jiang (74#, Zhongshan Rd 2, Guangzhou, China, +8613725336216, fax

number: +86(20)87331928, jiangwy@mail.sysu.edu.cn)

Running title: Noninvasive prenatal diagnosis for Haemophilia A

Keywords: Factor VIII; Haemophilia A; Genetic diagnosis; Haplotype; Targeted

sequencing; Noninvasive prenatal diagnosis

Abstract

Aim: This study aimed to demonstrate the feasibility of haplotype-based noninvasive prenatal diagnosis for Haemophilia A.

Methods: Two families affected by Haemophilia A participated in our study. Maternal haplotypes associated with pathogenic mutations were generated using targeted region genotypes of the mothers and the probands. Combined with the maternal pathogenic haplotypes, a hidden Markov Model was constructed to deduce fetal haplotypes using high-coverage targeted sequencing of maternal plasma DNA. The presence of a pathogenic haplotype in a male indicated an affected fetus, and that in a female indicated a carrier. Prenatal diagnoses were confirmed with amniocentesis and long-distance PCR.

Results: The haplotype-based noninvasive prenatal diagnosis of Haemophilia A was successfully performed. One fetus was identified to be Hemophilia A-negative and the other fetus was identified as a carrier; the results were confirmed by amniocentesis.

Conclusion: Our research demonstrated the feasibility of noninvasive prenatal diagnosis of Hemophilia A by a haplotype-based approach.

1 INTRODUCTION

Haemophilia A (HA) is one of the most common X-linked recessive inherited bleeding

disorders, with an incidence of approximately 1/5,000 in males[1], and is caused by a genetic

mutation of the *F8* gene that results in deficiency of coagulation factor VIII in plasma. The

clinical manifestations include spontaneous or traumatic bleeding in several organs. Prenatal

diagnosis is important for HA families because of the lack of an effective treatment. However,

an invasive prenatal diagnosis approach has a small risk of miscarriage and infection[2]. The

discovery of cell-free fetal DNA (cff-DNA) in maternal plasma makes noninvasive prenatal

diagnosis (NIPD) possible[3]. The intron 22 inversion in the *F8* gene is found in

approximately 45% of severe HA patients. Because of the background of maternal DNA[4]

and technical obstacles in determining the inversion breakpoint in plasma, the direct detection

of HA is impracticable. A haplotype-based method could potentially solve this problem by

analyzing mutation-linked polymorphism sites[5]. This study describes the few use of NIPD

for HA by a haplotype-based approach and demonstrates that our approach is an affordable

strategy for the clinically feasible NIPD of X-linked disorders.

2 MATERIALS AND METHODS

2.1 HA patient recruitment and clinical information

Two families (F01-F02) with individuals diagnosed with HA, including two pregnant women,

their brothers (who were affected by HA) and their spouses were recruited. The causative

mutation in the brothers and the pregnant women was previously identified by long-distance

PCR (LD-PCR) to be an intron 22 inversion in the *F8* gene. Genetic counseling was given to the families, and the parents decided to take our noninvasive prenatal test. All participants gave their signed informed consent. The ethics committee of The First Affiliated Hospital of Guangzhou Medical University approved this study. The 5 mL blood samples obtained from each family member, including mother, father and proband, were transported to BGI-Tianjin. Maternal plasma was separated using a double-centrifugation method and stored frozen. Amniotic fluid was obtained through amniocentesis for routine prenatal diagnosis. Invasive prenatal diagnosis and NIPD were performed in a double-blind manner by two independent investigators.

2.2 Library preparation and targeted sequencing

In this study, we used haplotype-based noninvasive prenatal testing of HA using a 110.08 kb SeqCap Kit (Roche) for the *F8* gene coding region and 445 surrounding highly heterozygous SNPs (minor allele frequency, MAF=0.3-0.5) distributed within a 1 Mb region of the X chromosome (Figure 1). Inclusion of a 10.37 kb chromosome Y-specific region facilitated the simultaneous determination of fetal sex.

Genomic DNA (gDNA) and cell-free DNA were extracted from blood or umbilical cord blood and maternal plasma using a QIAamp DNA Mini Kit (Qiagen, Germany) and a QIAamp Circulating Nucleic Acid Kit (Qiagen, Germany), respectively. The library was prepared according to our established protocol. The libraries of gDNA and cell-free DNA were pooled into one and then captured using our customized probes. Using BWA software,

the sequencing reads were aligned to the human reference genome (hg19). SNP calling was

performed using GATK software.

2.3 Determination of fetal sex and DNA fractions

The mean depth of the Y chromosome target region was utilized to determine the fetal sex. If

the fetus was female, the depth of the Y chromosome region would be extremely low,

resulting from short nonspecific region reads that could be mapped to Y chromosome-specific

region.

The fetal DNA concentration in the maternal plasma fraction was calculated by using SNP

sets with different parental homozygosity and the formula $\varepsilon = 2dF/ dF+ dM$, where ε represents

fetal DNA fraction in plasma, dF is the depth of reads supporting fetal-specific alleles and dM

is the depth of reads supporting maternal shared alleles in plasma. Assuming the maternal

genotype is AA and the paternal genotype is TT, the fetal genotype should be AT, as the

fetal-specific allele was T and the maternal shared allele was A.

2.4 Haplotype-based noninvasive prenatal diagnosis of the *F8* gene

A noninvasive haplotype-based linkage analysis method as previous described[6] was used to

determine fetal haplotype using information regarding SNPs from 1 Mb regions flanking the

coding regions of the *F8* gene from probands and parents. We defined haplotype 0 (Hap0) as

the pathogenic haplotype and haplotype 1 (Hap1) as the wildtype haplotype. Because

candidate Hap0 was inherited from maternal chromosome X, we directly phased the maternal

haplotypes of the *F8* gene. Based on maternal haplotype and SNP representation in the plasma, an hidden Markov Model was constructed. We chose the SNP sets that were heterozygous in the mother but homozygous in the father to determine the maternal inheritance. The fetal DNA fraction and sequencing error rate were utilized to determine the maternal haplotype inheritance patterns in the fetus. The probability of each state of maternal inheritance was calculated for each SNP marker in the maternal plasma. The fetal haplotype was deduced using Viterbi algorithm to identify the most hidden states.

2.5 Prenatal diagnosis of HA using amniotic fluid

LD-PCR of amniotic fluid using primers P, Q and B[7] was utilized to assess the accuracy of haplotype-based NIPD of HA. Electrophoresis showed a 12 kb band (Fragment PQ) for the wild-type, an 11 kb band (Fragment PB) for male patients and PQ and PB bands for female carriers.

3 RESULTES

3.1 Sequencing data

For maternal plasma, we obtained a mean 41.35 million data mapped to the target region with a mean depth of 375.61× and 99.48% coverage with more than 20 reads. A mean depth of 151.08× and 97.93% with 20× coverage was obtained in genomic DNA of pregnant women, their brothers and their spouses (Table 1). The data were sufficient for subsequent analysis.

3.2 Determination of fetal sex and DNA fraction

The mean depth and 20× coverage mapped to chromosome Y in F01 plasma was 26.5× and 42.59%. The fetus of F01 was inferred to be male. However, no reads were mapped to the target region of chromosome Y, so the fetus of F02 was determined to be female. The fetal DNA fractions of F01 and F02 were 11.09% and 9.37%, respectively.

3.3 Haplotype-based noninvasive prenatal diagnosis

The maternal haplotype could be constructed based on these SNPs. The number of informative phased SNPs used for NIPD for F01 and F02 was 144 and 49, respectively. In F01, the number of SNPs (Table 2, Figure 2) indicating that the fetus inherited the maternal wild-type haplotype (Hap1) was 144, while none supported the pathogenic haplotype (Hap0). The fetus could be subsequently predicted to be unaffected with HA considering the fetus's male sex. In F02, the number of SNPs (Table 2, Figure 2) indicating that the fetus inherited the maternal pathogenic haplotype (Hap0) was 49, while none supported the wild-type haplotype (Hap1). The fetus could be subsequently predicted to be a carrier of HA considering the fetus's female sex.

3.4 Invasive prenatal diagnosis

Prenatal diagnosis performed by LD-PCR of F01 was indicated to be a fetus unaffected by HA (Figure 3A). The fetus of F02 was diagnosed as a carrier of HA (Figure 3B).

4 DISCUSSION AND CONCLUSION

As a simple procedure, samples for the NIPD method are more accessible than those for invasive prenatal diagnosis. NIPD for HA would allow testing in the early weeks of gestation, giving more opportunity and time for making decisions regarding terminating the pregnancy. Moreover, it provides an option for women who refuse invasive methods to have more information about the possibility of an affected fetus. The discovery of cell-free fetal DNA (cff-DNA) in maternal plasma bring a new resource for noninvasive prenatal diagnosis. Heretofore, several studies have reported the high accuracy of noninvasive prenatal detection of fetal chromosome aneuploidy [8, 9]. In addition, whole-genome sequencing (WGS) of maternal plasma DNA can predict the fetal-specific allele with accuracy of more than 95%, thus providing the possibility of single gene detection in maternal plasma[10]. Nevertheless, the extremely high cost of whole-genome deep sequencing has limited its clinical application. Deep sequencing of target regions is a noninvasive alternative to single-gene-based disease detection, which could enable deep sequencing of interest regions at a lower cost[11]. Lam et al. developed an algorithm for noninvasive prenatal fetal diagnosis of single-gene diseases using target region capture sequencing for single-base variations and indels[12]. Using this method, two fetuses were correctly diagnosed as β-thalassemia carriers. Cell-free fetal DNA analysis in maternal plasma provides a noninvasive method for assessing fetal sex in HA-associated pregnancies, whereas without mutation-specific confirmatory analysis, the disease status of male fetuses remains unknown. Tsui et al. used microfluidics digital PCR method to perform noninvasive prenatal diagnosis of HA[13]. However, it can only detect

point mutations. The high incidence of the *F8* inversion 22 (Inv 22) mutation cannot be detected by this method. Microfluidics digital PCR is expensive as a microfluid digital PCR chip costs thousands of dollars. Our team developed a haplotype-based strategy for one-step prediction of fetal genotype and haplotype based on both parentally heterozygous sites in the maternal plasma[5].Using this method, we successfully predicted the fetal mutation status of many autosomal recessive genetic diseases using maternal plasma, including congenital adrenal hyperplasia, maple syrup urine disease, congenital deafness and Duchenne muscular dystrophy[14-16]. These are a few applications of the haplotype-based strategy for prenatal detection of HA. New technologies (for example, microfluidics digital PCR and next-generation sequencing) greatly accelerate the clinical application of NIPD, including determination of fetal sex, Rh status, and aneuploidy. Our haplotype-assisted strategy provides an opportunity for the diagnosis of single-gene diseases by NIPD.

At present, fetal gene sequencing analysis methods based on capture technologies are expensive because the capture range is large. All libraries were captured with a target region of 110.08 kb, which consisted of the *F8* gene coding area, 445 highly heterozygous SNPs distributed within 1 Mb flanking regions of *F8,* and a 10.37 kb chromosome Y-specific region. Therefore, we can reduce the cost of sequencing substantially.

Sex determination is very important for NIPD of X-linked inherited diseases. We use the specific Y chromosome probe to easily and accurately perform sex determination. The capture range and depth of the Y chromosome are markedly different between male and female

fetuses according to cff-DNA sequencing data analysis. It is approximately 0 for female fetus. NIPD may predict fetal sex accurately.

Even though NIPD of HA has many advantages compared with invasive methods, the accuracy needs to be further improved before its clinical application. As accuracy increases and costs reduce, we hypothesize that, in the future, NIPD will be used as a relatively fast and easy alternative for pregnant women who are likely to have fetuses with HA and other X-linked diseases.

ACKNOWLEDGMENTS

This work was supported by a grant from the Natural Science Foundation of Guangdong Province, China (2015A030313472). We are grateful to the participants and their families. We also thank our coinvestigators for their assistance in research design, data analysis. The authors stated that they had no interests which might be perceived as posing a conflict or bias. Zhihui He, Chao Chen, Dan Li, Renhua Wu, Rong Lu, Mei Zhao and Fang Chen performed the research. Zhihui He and Chao Chen wrote the paper. Fengyu Guo, Lu Jiang, Yaping Zhu and Peixiang Ni analysed the data. Zhihui He , Chao Chen, Weiying Jiang, Jun Sun, and Zhiyu Peng designed the research study. Zhihui He and Fang Chen received funding.

REFERENCES

1. Dai J, Lu Y, Ding Q, et al. The status of carrier and prenatal diagnosis of haemophilia in China. Haemophilia, 2012, 18 (2): 235-240.

2. Kozlowski P, Knippel A, Stressig R. Individual risk of fetal loss following routine second trimester amniocentesis: a controlled study of 20,460 cases. Ultraschall Med. 2008; 29: 165-172.

3. Lo YM, Corbetta N, Chamberlain PF, et al. Presence of fetal DNA in maternal plasma and serum. Lancet. 1997; 350: 485-487.

4. Lo YM, Tein MS, Lau TK, et al. Quantitative analysis of fetal DNA in maternal plasma and serum: implications for noninvasive prenatal diagnosis. Am J Hum Genet. 1998; 62: 768-775.

5. Chen SP, Ge HJ, Wang XB, et al. Haplotype-assisted accurate non-invasive fetal whole genome recovery through maternal plasma sequencing. Genome Med. 2013; 5: 18.

6. Xu Y, Li X, Ge HJ, et al. Haplotype-based approach for noninvasive prenatal tests of Duchenne muscular dystrophy using cell-free fetal DNA in maternal plasma. Genet Med. 2015; 17(11): 889-896.

7. He ZH, Chen SF, Chen J, Jiang WY. A Modified I-PCR to detect the factor VIII Inv22 for genetic diagnosis and prenatal diagnosis in haemophilia A. Haemophilia, 2012, 18(3): 452-456.

8. Norton ME, Brar H, Weiss J, et al. Non-Invasive Chromosomal Evaluation (NICE) Study: results of a multicenter prospective cohort study for detection of fetal trisomy 21 and trisomy 18. Am J Obstet Gynecol. 2012; 207: 137 e131-138.

9. Palomaki GE, Deciu C, Kloza EM, et al. DNA sequencing of maternal plasma reliably identifies trisomy 18 and trisomy 13 as well as Down syndrome: an international collaborative study.Genet Med. 2012; 14: 296-305.

10. Lo YM, Chan KC, Sun H, et al. Maternal plasma DNA sequencing reveals the genome-wide genetic and mutational profile of the fetus. Sci Transl Med. 2010; 2: 61ra91.

11. Liao GJ, Chan KC, Jiang P, et al. Noninvasive prenatal diagnosis of fetal trisomy 21 by allelic ratio analysis using targeted massively parallel sequencing of maternal plasma DNA. PLoS One. 2012; 7: e38154.

12. Lam KW, Jiang P, Liao GJ, et al. Noninvasive prenatal diagnosis of monogenic diseases by targeted massively parallel sequencing of maternal plasma: application to beta-thalassemia. Clin Chem. 2012; 58: 1467-1475.

13. Tsui NB, Kadir RA, Chan KC, et al. Noninvasive prenatal diagnosis of hemophilia by microfluidics digital PCR analysis of maternal plasma DNA Blood. 2011; 117(13): 3684-23691.

14. Ma DY, Ge HJ, Li XC, et al. Haplotype-based approach for noninvasive prenatal diagnosis of congenital adrenal hyperplasia by maternal plasma DNA sequencing. Gene. 2014; 2: 252-258.

15. You Y, Sun Y, Li XC, et al. Integration of targeted sequencing and NIPT into clinical practice in a Chinese family with maple syrup urine disease. Genet Med. 2014; 16: 594-600.

16. Meng M, Li XC, Chen F, et al. Noninvasive prenatal testing for autosomal recessive

conditions by maternal plasma sequencing in a case of congenital deafness. Genet Med.

2014; 16: 972-976.

TABLE 1   Sequencing data of studied families

Family	Sample	Target region data (M)	Mean depth (×)	20× Coverage fraction	Mean depth of chr Y	20× Coverage fraction of chr Y	SNPs	Fetal DNA fraction
F01	mother	21.29	193.36	98.70%	0	0		
	father	15.50	140.84	98.31%	45.28	72.2%		
	proband	15.14	137.51	98.11%	43.05	71.1%	248	11.09%
	plasma	29.93	271.87	99.37%	26.50	42.59%		
	AF	11.00	99.94	96.80%	113.67	96.74%		
F02	mother	22.73	206.49	98.79%	0	0		
	father	10.22	92.82	95.81%	102.40	95.4%		
	proband	14.91	135.47	97.83%	153.40	97.4%	106	9.37%
	plasma	52.77	479.35	99.59%	0	0		
	AF	21.06	191.30	98.63%	0	0		

Notes: AF, Amniotic fluid; the proband was the brother of each pregnant woman; "SNPs" represents the number of SNPs detected in the

studied families.

TABLE 2 **The NIPD results from *F8* gene analysis**

Family	Fetal Sex	Phased SNPs[a]	SNPs For Hap0	SNPs For Hap1	NIPD results	Predicted results
F01	male	144	0	144	Hap1	normal
F02	female	49	49	0	Hap0	carrier

Notes: "SNPs[a]" represents SNPs that were used to predict maternal haplotypes. Hap0, fetal-inherited maternal mutant haplotype; Hap1, fetal-inherited maternal wild-type haplotype.

FIGURE CAPTION

FIGURE 1 Target region of the *F8* gene and SNPs used for haplotyping. Custom NimbleGen probes were designed that targeted within the 110.08 kb region including the *F8* gene coding region and 445 surrounding highly heterozygous SNPs distributed within a 1 Mb region.

FIGURE 2 Results of haplotype-based NIPD of Haemophilia A. The X-axis represents the target region of our customized probe, and blue vertical lines represent the interval of the *F8* gene; the Y-axis represents the logarithm of the ratios of fetal-inherited maternal haplotype. Hap0 represents the pathogenic haplotype, and Hap1 represents the wild-type haplotype. The black lines above zero indicate that the fetus inherited the pathogenic haplotype (Hap0), and the black lines below zero indicate that the fetus inherited the normal haplotype (Hap1).

FIGURE 3 Results of invasive diagnosis of studied families. A) Electrophoresis results of LD-PCR of amniotic fluid (F01). *M: Marker, Lanes 1 and 2: Amniotic fluid, Lane 3: Mother, Lane 4: Carrier control from the Kit, Lane 5: Normal control from the Kit, Lane 6: Blank control. Completed by researchers of The First Affiliated Hospital of Guangzhou Medical University. B) Electrophoresis results of LD-PCR of amniotic fluid (F02). *PB was amplified by primers P and B, and PQ was amplified by primers P and Q. N: Normal control, P: Positive control (Male patient), 714R2: amniotic fluid. Completed by researchers of BGI Genomics, BGI-Shenzhen.

F01 Maternal inheritance

F02 Maternal inheritance

A

B