Title: Common gardens in teosintes reveal the establishment of a syndrome of adaptation to altitude

Short Title: Altitudinal syndrome in teosintes

M-A. Fustier[1, ☁], N.E. Martínez-Ainsworth[1, ☁], J.A. Aguirre-Liguori[2], A. Venon[1], H. Corti[1], A. Rousselet[1], F. Dumas[1], H. Dittberner[3], M.G. Camarena[4], D. Grimanelli[5], O. Ovaskainen[6,7], M. Falque[1], L. Moreau[1], J. de Meaux[3], S. Montes[4], L.E. Eguiarte[2], Y. Vigouroux[5], D. Manicacci[1,*], M.I. Tenaillon[1,*]

[1]: Génétique Quantitative et Evolution – Le Moulon, Institut National de la Recherche Agronomique, Université Paris-Sud, Centre National de la Recherche Scientifique, AgroParisTech, Université Paris-Saclay, France

[2]: Departamento de Ecología Evolutiva, Instituto de Ecología, Universidad Nacional Autónoma de México, Ciudad de México, Mexico

[3]: Institute of Botany, University of Cologne Biocenter, 47b Cologne, Germany

[4]: Campo Experimental Bajío, Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Celaya, Mexico

[5]: Université de Montpellier, Institut de Recherche pour le développement, UMR Diversité, Adaptation et Développement des plantes, Montpellier, France

[6]: Organismal and Evolutionary Biology Research Programme, PO Box, 65, FI-00014 University of Helsinki, Finland

[7]: Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

[☁]: The authors equally contributed to this work

* co-corresponding authors

E-mail: domenica.manicacci@inra.fr, maud.tenaillon@inra.fr

Abstract

In plants, local adaptation across species range is frequent. Yet, much has to be discovered on its environmental drivers, the underlying functional traits and their molecular determinants. Genome scans are popular to uncover outlier loci potentially involved in the genetic architecture of local adaptation, however links between outliers and phenotypic variation are rarely addressed. Here we focused on adaptation of teosinte populations along two elevation gradients in Mexico that display continuous environmental changes at a short geographical scale. We used two common gardens, and phenotyped 18 traits in 1664 plants from 11 populations of annual teosintes. In parallel, we genotyped these plants for 38 microsatellite markers as well as for 171 outlier single nucleotide polymorphisms (SNPs) that displayed excess of allele differentiation between pairs of lowland and highland populations and/or correlation with environmental variables. Our results revealed that phenotypic differentiation at 10 out of the 18 traits was driven by local selection. Trait covariation along the elevation gradient indicated that adaptation to altitude results from the assembly of multiple co-adapted traits into a complex syndrome: as elevation increases, plants flower earlier, produce less tillers, display lower stomata density and carry larger, longer and heavier grains. The proportion of outlier SNPs associating with phenotypic variation, however, largely depended on whether we considered a neutral structure with 5 genetic groups (73.7%) or 11 populations (13.5%), indicating that population stratification greatly affected our results. Finally, chromosomal inversions were enriched for both SNPs whose allele frequencies shifted along elevation as well as phenotypically-associated SNPs. Altogether, our results are consistent with the establishment of an altitudinal syndrome promoted by local

26   selective forces in teosinte populations in spite of detectable gene flow. Because

27   elevation mimics climate change through space, SNPs that we found underlying

28   phenotypic variation at adaptive traits may be relevant for future maize breeding.

30   **Keywords:** spatially-varying selection; F_{ST}-scan; association mapping; altitudinal

31   syndrome; pleiotropy; chromosomal inversions.

Author summary

Across their native range species encounter a diversity of habitats promoting local

adaptation of geographically distributed populations. While local adaptation is

widespread, much has yet to be discovered about the conditions of its emergence, the

targeted traits, their molecular determinants and the underlying ecological drivers.

Here we employed a reverse ecology approach, combining phenotypes and genotypes,

to mine the determinants of local adaptation of teosinte populations distributed along

two steep altitudinal gradients in Mexico. Evaluation of 11 populations in two

common gardens located at mid-elevation pointed to adaptation via an altitudinal

multivariate syndrome, in spite of gene flow. We scanned genomes to identify loci

with allele frequencies shifts along elevation, a subset of which associated to trait

variation. Because elevation mimics climate change through space, these

polymorphisms may be relevant for future maize breeding.

Introduction

Local adaptation is key for the preservation of ecologically useful genetic variation [1]. The conditions for its emergence and maintenance have been the focus of a long-standing debate nourished by ample theoretical work [2-9]. On the one hand, spatially-varying selection promotes the evolution of local adaptation, provided that there is genetic diversity underlying the variance of fitness-related traits [10]. On the other hand, opposing forces such as neutral genetic drift, temporal fluctuations of natural selection, recurrent introduction of maladaptive alleles via migration and homogenizing gene flow may hamper local adaptation (reviewed in [11]). Meta-analyzes indicate that local adaptation is pervasive in plants, with evidence of native-site fitness advantage in reciprocal transplants detected in 45% to 71% of populations [12, 13].

While local adaptation is widespread, much has yet to be discovered about the traits affected by spatially-varying selection, their molecular determinants and the underlying ecological drivers [14]. Local adaptation is predicted to increase with phenotypic, genotypic and environmental divergence among populations [6, 15, 16]. Comparisons of the quantitative genetic divergence of a trait (Q_{ST}) with the neutral genetic differentiation (F_{ST}) can provide hints on whether trait divergence is driven by spatially-divergent selection [17-20]. Striking examples of divergent selection include developmental rate in the common toad [21], drought and frost tolerance in alpine populations of the European silver fir [22], and traits related to plant phenology, size and floral display among populations of *Helianthus* species [23, 24]. These studies have reported covariation of physiological, morphological and/or life-history traits across environmental gradients which collectively define adaptive syndromes. Such

syndromes may result from several non-exclusive mechanisms: plastic responses,

pleiotropy, non-adaptive genetic correlations among traits (constraints), and joint

selection of traits encoded by different sets of genes resulting in adaptive correlations.

In some cases, the latter mechanism may involve selection and rapid spread of

chromosomal inversions that happen to capture multiple locally favored alleles [25] as

exemplified in *Mimulus guttatus* [26]. While distinction between these mechanisms is

key to decipher the evolvability of traits, empirical data on the genetic bases of

correlated traits are currently lacking [27].

The genes mediating local adaptation are usually revealed by genomic regions

harboring population-specific signatures of selection. These signatures include alleles

displaying greater-than-expected differentiation among populations [28] and can be

identified through F_{ST}-scans [29-35]. However, F_{ST}-scans and its derivative methods

[28] suffer from a number of limitations, among them a high number of false positives

(reviewed in [36, 37]) and the lack of power to detect true positives [38]. Despite

these caveats, F_{ST}-outlier approaches have helped in the discovery of emblematic

adaptive alleles such as those segregating at the *EPAS1 locus* in Tibetan human

populations adapted to high altitude [39]. An alternative to detect locally adaptive loci

is to test for genotype-environment correlations [35, 40-45]. Correlation-based

methods can be more powerful than differentiation-based methods [46], but spatial

autocorrelation of population structure and environmental variables can lead to

spurious signatures of selection [47].

Ultimately, to identify the outlier loci that have truly contributed to improve

local fitness, a link between outliers and phenotypic variation needs to be established.

The most common approach is to undertake association mapping. However, recent

literature in humans has questioned our ability to control for sample stratification in

such approach [48]. Detecting polymorphisms responsible for trait variation is

particularly challenging when trait variation and demographic history follow parallel

environmental (geographic) clines. Plants however benefit from the possibility of

conducting replicated phenotypic measurements in common gardens, where

environmental variation is controlled. Hence association mapping has been

successfully employed in the model plant species *Arabidopsis thaliana*, where

broadly distributed ecotypes evaluated in replicated common gardens have shown that

fitness-associated alleles display geographic and climatic patterns indicative of

selection [49]. Furthermore, the relative fitness of *A. thaliana* ecotypes in a given

environment could be predicted from climate-associated SNPs [50]. While climatic

selection over broad latitudinal scales produces genomic and phenotypic patterns of

local adaptation in the selfer plant *A. thaliana*, whether similar patterns exist at shorter

spatial scale in outcrossing species remains to be elucidated.

We focused here on a well-established outcrossing plant system, the teosintes,

to investigate the relationship of molecular, environmental, and phenotypic variation

in populations sampled across two elevation gradients in Mexico. The gradients

covered a relatively short yet climatically diverse, spatial scale. They encompassed

populations of two teosinte subspecies that are the closest wild relatives of maize, *Zea*

mays ssp. *parviglumis* (hereafter *parviglumis*) and *Z. mays* ssp. *mexicana* (hereafter

mexicana). The two subspecies display large effective population sizes [51], and span

a diversity of climatic conditions, from warm and mesic conditions below 1800 m for

parviglumis, to drier and cooler conditions up to 3000 m for *mexicana* [52]. Previous

studies have discovered potential determinants of local adaptation in these systems.

At a genome-wide scale, decrease in genome size correlates with increasing altitude,

which likely results from the action of natural selection on life cycle duration [53, 54].

More modest structural changes include megabase-scale inversions that harbor

clusters of SNPs whose frequencies are associated with environmental variation [55,

56]. Also, differentiation- and correlation-based genome scans in teosinte populations

succeeded in finding outlier SNPs potentially involved in local adaptation [57, 58].

But a link with phenotypic variation has yet to be established.

In this paper, we genotyped a subset of these outlier SNPs on a broad sample

of 28 teosinte populations, for which a set of neutral SNPs was also available; as well

as on an association panel encompassing 11 populations. We set up common gardens

in two locations to evaluate the association panel for 18 phenotypic traits over two

consecutive years. Individuals from this association panel were also genotyped at 38

microsatellite markers to enable associating genotypic to phenotypic variation while

controlling for sample structure and kinship among individuals. We addressed three

main questions: What is the extent of phenotypic variation within and among

populations? Can we define a set of locally-selected traits that constitute a syndrome

of adaptation to altitude? What are the genetic bases of such syndrome? We further

discuss the challenges of detecting phenotypically-associated SNPs when trait and

genetic differentiation parallel environmental clines.

Results

Trait-by-trait analysis of phenotypic variation within and among populations.

In order to investigate phenotypic variation, we set up two common garden

experiments located in Mexico to evaluate individuals from 11 teosinte populations

(Fig 1). The two experimental fields were chosen because they were located at

intermediate altitudes (S1 Fig). Although natural teosinte populations are not typically

encountered around these locations [52], we verified that environmental conditions

were compatible with both subspecies (S2 Fig). The 11 populations were sampled

among 37 populations (S1 Table) distributed along two altitudinal gradients that range

from 504 to 2176 m in altitude over ~460 kms for gradient *a*, and from 342 to 2581m

in altitude over ~350 kms for gradient *b* (S1 Fig). Lowland populations of the

subspecies *parviglumis* (n=8) and highland populations of the subspecies *mexicana*

(n=3) were climatically contrasted as can be appreciated in the Principal Component

Analysis (PCA) computed on 19 environmental variables (S2 Fig). The corresponding

set of individuals grown from seeds sampled from the 11 populations formed the

association panel.

156
Figure 1. Geographical location of sampled populations and experimental fields.

The entire set of 37 Mexican teosinte populations is shown with *parviglumis* (circles)

and *mexicana* (triangles) populations sampled along gradient *a* (white) and gradient *b*

(black). The 11 populations indicated with a purple outline constituted the association

panel. This panel was evaluated in a four-block design over two years in two

experimental fields located at mid-elevation, SENGUA and CEBAJ. Two major cities

(Mexico City and Guadalajara) are also indicated. Topographic surfaces have been

obtained from International Centre for Tropical Agriculture (Jarvis A., H.I. Reuter, A.

Nelson, E. Guevara, 2008, Hole-filled seamless SRTM data V4, International Centre

for Tropical Agriculture (CIAT), available from http://srtm.csi.cgiar.org).

167
We gathered phenotypic data during two consecutive years (2013 and 2014).

We targeted 18 phenotypic traits that included six traits related to plant architecture,

three traits related to leaves, three traits related to reproduction, five traits related to

grains, and one trait related to stomata (S2 Table). Each of the four experimental

assays (year-field combinations) encompassed four blocks. In each block, we

evaluated one offspring (half-sibs) of ~15 mother plants from each of the 11 teosinte

populations using a semi-randomized design. After filtering for missing data, the

association panel included 1664 teosinte individuals. We found significant effects of

Field, Year and/or their interaction for most traits, and a highly significant Population

effect for all of them (model M1, S3 Table).

We investigated the influence of altitude on each trait independently. All traits,

except for the number of nodes with ears (NoE), exhibited a significant effect of

altitude (S3 Table, M3 model). Note that after accounting for elevation, the

population effect remained significant for all traits, suggesting that factors other than

altitude contributed to shape phenotypic variation among populations. Traits related to

flowering time and tillering displayed a continuous decrease with elevation, and traits

related to grain size increased with elevation (Fig 2 & S3 Fig). Stomata density also

diminished with altitude (Fig 2). In contrast, plant height, height of the highest ear,

number of nodes with ear in the main tiller displayed maximum values at intermediate

altitudes (highland *parviglumis* and lowland *mexicana*) (S3 Fig).

We estimated narrow-sense heritabilites (additive genotypic effect) per

population for all traits using a mixed animal model. Average per-trait heritability

ranged from 0.150 for tassel branching to 0.664 for female flowering time, albeit with

large standard errors (S2 Table). We obtained higher heritability for grain related

traits when mother plant measurements were included in the model with 0.631 ($sd =$

0.246), 0.511 ($sd = 0.043$) and 0.274 ($sd = 0.160$) for grain length, weight and width,

respectively, suggesting that heritability was under-estimated for other traits where

mother plant values were not available.

Figure 2: Population-level box-plots of adjusted means for four traits. Traits are

female flowering time (A), male flowering time (B), grain length (C) and stomata

density (D). Populations are ranked by altitude. *Parviglumis* populations are shown in

green and *mexicana* in red, lighter colors are used for gradient 'a' and darker colors

for gradient 'b'. In the case of male and female flowering time, we report data for 9

out of 11 populations because most individuals from the two lowland populations

(P1a and P1b) did not flower in our common gardens. Covariation with elevation was

significant for the four traits. Corrections for experimental setting are detailed in the

material and methods section (Model M'1).

Multivariate analysis of phenotypic variation and correlation between traits.

Principal component analysis including all phenotypic measurements

highlighted that 21.26% of the phenotypic variation scaled along PC1 (Fig 3A), a PC

axis that is strongly collinear with altitude (Fig 3B). Although populations partly

overlapped along PC1, we observed a consistent tendency for population phenotypic

differentiation along altitude irrespective of the gradient (Fig 3C). Traits that

correlated the most to PC1 were related to grain characteristics, tillering, flowering

and to a lesser extent to stomata density (Fig 3B). PC2 correlated with traits

exhibiting a trend toward increase-with-elevation within *parviglumis*, but decrease-

with-elevation within *mexicana* (Fig 3D). Those traits were mainly related to

vegetative growth (Fig 3B). Together, both axes explained 37% of the phenotypic

variation.

Figure 3: Principal Component Analysis on phenotypic values corrected for the

experimental setting. Individuals factor map (A) and corresponding correlation

circle (B) on the first two principal components with altitude (Alt) added as a

supplementary variable (in blue). Individual phenotypic values on PC1 (C) and PC2

(D) are plotted against population ranked by altitude and color-coded following A.

For populations from the two subspecies, *parviglumis* (circles) and *mexicana*

(triangles), color intensity indicates ascending elevation in green for *parviglumis* and

red for *mexicana*. Corrections for experimental setting are detailed in the material and

methods (Model M2).

We assessed more formally pairwise-correlations between traits after

correcting for experimental design and population structure ($K=5$). We found 82

(54%) significant correlations among 153 tested pairs of traits. The following pairs of

traits had the strongest positive correlations: male and female flowering time, plant

height and height of the highest ear, height of the highest and lowest ear, grain length

with grain weight and width (S4 Fig). The correlation between flowering time (female

or male) with grain weight and length were among the strongest negative correlations

(S4 Fig).

Neutral structuring of the association panel.

We characterized the genetic structure of the association panel using SSRs.

The highest likelihood from Bayesian classification was obtained at $K=2$ and $K=5$

clusters (S5 Fig). At $K=2$, the clustering separated the lowland of gradient *a* from the

rest of the populations. From $K=3$ to $K=5$, a clear separation between the eight

parviglumis and the three *mexicana* populations emerged. Increasing K values finally

split the association panel into the 11 populations it encompassed (S6 Fig). The $K=5$

structure associated to both altitude (lowland *parviglumis* versus highland *mexicana*)

and gradients *a* and *b* (Fig 4A & B). TreeMix analysis for a subset of 10 of these

populations confirmed those results with an early split separating the lowlands from

gradient *a* (cf. *K=2*, S6 Fig) followed by the separation of the three *mexicana* from the

remaining populations (Fig 4C). TreeMix further supported three migration edges, a

model that explained 98.75% of the variance and represented a significant

improvement over a model without admixture (95.7%, Figure S7). This admixture

model was consistent with gene flow between distant lowland *parviglumis*

populations from gradient *a* and *b*, as well as between *parviglumis* and *mexicana*

populations (Fig 4C). Likewise, structure analysis also suggested admixture among

some of the lowland populations, and to a lesser extent between the two subspecies

(Fig 4B).

Figure 4: Genetic clustering, historical splits and admixture among populations

of the association panel. Genetic clustering visualization based on 38 SSRs is shown

for *K=5* (A). Colors represent the *K* clusters. Individuals (vertical lines) are

partitioned into colored segments whose length represents the membership

proportions to the *K* clusters. Populations (named after the subspecies M: *mexicana,*

P: *parviglumis* and gradient 'a' or 'b') are ranked by altitude indicated in meters

above sea level. The corresponding geographic distribution of populations along with

their average membership probabilities are plotted (B). Historical splits and

admixtures between populations were inferred from SNP data for a subset of 10

populations of the association panel (C). Admixtures are colored according to their

weight.

Identification of traits evolving under spatially-varying selection.

We estimated the posterior mean (and 95% credibility interval) of genetic differentiation (F_{ST}) among the 11 populations of the association panel using DRIFTSEL. Considering 1125 plants for which we had both individual phenotypes and individual genotypes for 38 SSRs (S4 Table), we estimated the mean F_{ST} to 0.22 (0.21-0.23). Note that we found a similar estimate on a subset of 10 of these populations with 1000 neutral SNPs (F_{ST} (CI)=0.26 (0.25-0.27)). To identify traits whose variation among populations was driven primarily by local selection, we employed the Bayesian method implemented in DRIFTSEL, that infers additive genetic values of traits from a model of population divergence under drift [59]. Selection was inferred when observed phenotypic differentiation exceeded neutral expectations for phenotypic differentiation under random genetic drift. Single-trait analyses revealed evidence for spatially-varying selection at 12 traits, with high consistency between SSRs and neutral SNPs (Table 1). Another method that contrasted genetic and phenotypic differentiation (Q_{ST}- F_{ST}) uncovered a large overlap with nine out of the 12 traits significantly deviating from the neutral model (Table 1) and one of the remaining ones displaying borderline significance (Plant height=PL, S8 Fig). Together, these two methods indicated that phenotypic divergence among populations was driven by local selective forces.

Table 1. Signals of selection (posterior probability S) for each trait considering SSR

markers (11 populations) or SNPs (10 populations).

Traits[a]	SSR[b]	SNP[b]
Plant height	0.995	0.972
Height of the lowest ear*	0.950	0.959
Height of the highest ear	0.982	0.966
Number of tillers*	1.000	1.000
Number of lateral branches*	1.000	0.990
Number of nodes with ears	0.682	0.699
Leaf length	0.888	0.875
Leaf width	0.999	0.996
Leaf color	0.633	0.583
Female flowering time*	1.000	1.000
Male flowering time*	1.000	1.000
Tassel branching*	0.925	0.908
Number of grains per ear	0.832	0.622
Grain length*	1.000	1.000
Grain width*	0.995	0.984
Grain weight*	1.000	0.999
Grain color	0.717	0.689
Stomata density*	0.999	0.999

[a]: Traits displaying signal of selection (spatially-varying traits, S > 0.95) are indicated in bold,

and marked by an asterisk when significant in Q_{ST}-$F_{ST}Comp$ analysis. We considered the

underlined traits as spatially varying. For a detailed description of traits see S2 Table.

291   Altogether, evidence of spatially varying selection at 10 traits (Table 1) as

292   well as continuous variation of a subset of traits across populations in both elevation

293   gradients (Fig 2, S3 Fig) was consistent with a syndrome where populations produced

294   less tillers, flowered earlier, displayed lower stomata density and carried larger,

295   longer and heavier grains with increasing elevation.

297   Outlier detection and correlation with environmental variables.

We successfully genotyped 218 (~81%) out of 270 outlier SNPs on a broad set

of 28 populations, of which 141 were previously detected in candidate regions for

local adaptation [58]. Candidate regions were originally identified from re-sequencing

data of only six teosinte populations (S1 Table) following an approach that included

high differentiation between highlands and lowlands, environmental correlation, and

in some cases their intersection with genomic regions involved in quantitative trait

variation in maize. The remaining outlier SNPs (77) were discovered in the present

study by performing F_{ST}-scans on the same re-sequencing data (S5 Table). We

selected outlier SNPs that were both highly differentiated between highland and

lowland populations within gradient (high/low in gradient *a* or *b* or both), and

between highland and lowland populations within subspecies in gradient *b* (high/low

within *parviglumis*, *mexicana* or both). F_{ST}-scans pinpointed three genomic regions of

particularly high differentiation (S9 Fig) that corresponded to previously described

inversions [55, 56]: one inversion on chromosome 1 (*Inv1n*), one on chromosome 4

(*Inv4m*) and one on the far end of chromosome 9 (*Inv9e*).

A substantial proportion of outlier SNPs was chosen based on their significant

correlation among six populations between variation of allele frequency and their

coordinate on the first environmental principal component [58]. We extended

environmental analyses to 171 outlier SNPs (MAF>5%) on a broader sample of 28

populations (S2 Fig) and used the two first components (PCenv1 and PCenv2) to

summarize environmental information. The first component, that explained 56% of

the variation, correlated with altitude but displayed no correlation to either latitude or

longitude. PCenv1 was defined both by temperature- and precipitation- related

variables (S2 B Fig) including Minimum Temperature of Coldest Month (T6), Mean

Temperature of Driest and Coldest Quarter (T9 and T11) and Precipitation of Driest

Month and Quarter (P14 and P17). The second PC explained 20.5% of the variation

and was mainly defined (S2 B Fig) by Isothermality (T3), Temperature Seasonality

(T4) and Temperature Annual Range (T7).

We first employed multiple regression to test for each SNP, whether the

pairwise F_{ST} matrix across 28 populations correlated to the environmental (distance

along PCenv1) and/or the geographical distance. As expected, we found a

significantly greater proportion of environmentally-correlated SNPs among outliers

compared with neutral SNPs (χ^2 =264.07, P-value=2.2 10^{-16}), a pattern not seen with

geographically-correlated SNPs. That outlier SNPs displayed a greater isolation-by-

environment than isolation-by-distance, indicated that patterns of allele frequency

differentiation among populations were primarily driven by adaptive processes. We

further tested correlations between allele frequencies and environmental variation.

Roughly 60.8% (104) of the 171 outlier SNPs associated with at least one of the two

first PCenvs, with 87 and 33 associated with PCenv1 and PCenv2, respectively, and

little overlap (S5 Table). As expected, the principal component driven by altitude

(PCenv1) correlated to allele frequency for a greater fraction of SNPs than the second

orthogonal component. Interestingly, we found enrichment of environmentally-

associated SNPs within inversions both for PCenv1 (χ^2 = 14.63, P-value=1.30 10^{-4})

and PCenv2 (χ^2 = 33.77, P-value=6.22 10^{-9}).

Associating genotypic variation to phenotypic variation.

We tested the association between phenotypes and 171 of the outlier SNPs

(MAF>5%) using the association panel. For each SNP-trait combination, the sample

size ranged from 264 to 1068, with a median of 1004 individuals (S6 Table). We used

SSRs to correct for both structure (at K=5) and kinship among individual genotypes.

This model (M5) resulted in a uniform distribution of P-values when testing the

association between genotypic variation at SSRs and phenotypic trait variation (S10

Fig). Under this model, we found that 126 outlier SNPs (73.7%) associated to at least

one trait (Fig 5 and S11 Fig) at an FDR of 10%. The number of associated SNPs per

trait varied from 0 for leaf and grain coloration, to 55 SNPs for grain length, with an

average of 22.6 SNPs per trait (S5 Table). Ninety-three (73.8%) out of the 126

associated SNPs were common to at least two traits, and the remaining 33 SNPs were

associated to a single trait (S5 Table). The ten traits displaying evidence of spatially

varying selection in the Q_{ST}-F_{ST} analyses displayed more associated SNPs per trait

(30.5 on average), than the non-spatially varying traits (12.75 on average).

A growing body of literature stresses that incomplete control of population

stratification may lead to spurious associations [61]. Hence, highly differentiated

traits along environmental gradients are expected to co-vary with any variant whose

allele frequency is differentiated along the same gradients, without underlying causal

link. We therefore expected false positives in our setting where both phenotypic traits

and outlier SNPs varied with altitude. We indeed found a slightly significant

correlation (r=0.5, P-value=0.03) between the strength of the population effect for

each trait – a measure of trait differentiation (S3 Table) – and its number of associated

SNPs (S5 Table).

To verify that additional layers of structuring among populations did not cause

an excess of associations, we repeated the association analyzes considering a

structuring with 11 populations (instead of K=5) as covariate (M5'), a proxy of the

structuring revealed at K=11 (S6 Fig). With this level of structuring, we retrieved

much less associated SNPs (S5 Table). Among the 126 SNPs associating with at least

one trait at K=5, only 22 were recovered considering 11 populations. An additional

SNP was detected with structuring at 11 populations that was absent at K=5. Eight

traits displayed no association, and the remaining traits varied from a single

associated SNP (Leaf length – LeL and the number of tillers – Til) to 8 associated

SNPs for grain weight (S5 Table). For instance, traits such as female or male

flowering time that displayed 45 and 43 associated SNPs at K=5, now displayed only

4 and 3 associated SNPs, respectively (Fig 5). Note that one trait (Leaf color)

associated with 4 SNPs considering 11 populations while displaying no association at

K=5. Significant genetic associations were therefore highly contingent on the

population structure. Noteworthy, traits under spatially varying selection still

associated with more SNPs (2.00 on average) than those with no spatially varying

selection (1.25 SNPs on average).

Figure 5: Manhattan plots of associations between 171 outlier SNPs and 6

phenotypic traits. X-axis indicates the positions of outlier SNPs on chromosomes 1

to 10, black and gray colors alternating per chromosome. Plotted on the Y-axis are the

negative Log_{10}-transformed P values obtained for the K=5 model. Significant

associations (10% FDR) are indicated considering either a structure matrix at K=5

(pink dots), 11 populations (blue dots) or both K=5 and 11 populations (purple dots)

models.

Altogether the 23 SNPs recovered considering a neutral genetic structure with

11 populations corresponded to 30 associations, 7 of the SNPs being associated to

more than one trait (S5 Table). For all these 30 associations except in two cases (FFT

with SNP_7, and MFT with SNP_28), the SNP effect did not vary among populations

(non-significant SNP-by-population interaction in model M5' when we included the

SNP interactions with year*field and population). For a subset of two SNPs, we

illustrated the regression between the trait value and the shift of allele frequencies

with altitude (Fig 6 A&B). We estimated corresponding additive and dominance

effects (S7 Table). In some cases, the intra-population effect corroborated the inter-

population variation with relatively large additive effects of the same sign (Fig 6

C&D). Note that in both examples shown in Fig 6, one or the other allele was

dominant. In other cases, the results were more difficult to interpret with negligible

additive effect but extremely strong dominance (S7 Table, SNP_210 for instance).

Figure 6: Regression of phenotypic average value on SNP allele frequency across

populations, and within-population average phenotypic value for each SNP

genotype. Per-population phenotypic average values of traits are regressed on alleles

frequencies at SNP_149 (A) and SNP_179 (B) with corresponding within-population

average phenotypic value per genotype (C & D). In A and B, the 11 populations of

the association panel are shown with *parviglumis* (green circles) and *mexicana* (red

triangles) populations sampled along gradient *a* and gradient *b*. Phenotypic average

values were corrected for the experimental design (calculated as the residues of model

M2). Pval refers to the P-value of the linear regression represented in blue. In C and

D, genotypic effects from model M5' are expressed as the average phenotypic value

of heterozygotes (1) and homozygotes for the reference (0) and the alternative allele

(2). FDR values are obtained from the association analysis on 171 SNPs with

correction for genetic structure using 11 population.

Independence of SNPs associated to phenotypes.

We computed the pairwise linkage disequilibrium (LD) as measured by r^2

between the 171 outlier SNPs using the R package LDcorSV [62]. Because we were

specifically interested by LD pattern between phenotypically-associated SNPs, as for

the association analyses we accounted for structure and kinship computed from SSRs

while estimating LD [63]. The 171 outlier SNPs were distributed along the 10

chromosomes of maize, and exhibited low level of linkage disequilibrium (LD),

except for SNPs located on chromosomes eight, nine, and a cluster of SNPs located

on chromosome 4 (S12 Fig).

Among the 171, the subset of 23 phenotypically-associated SNPs (detected

when considering the 11-population structure) displayed an excess of elevated LD

values – out of 47 pairs of SNPs phenotypically-associated to a same trait, 16 pairs

were contained in the 5% higher values of the LD distribution of all outlier SNP pairs.

Twelve out of the 16 pairs related to grain weight, the remaining four to leaf

coloration, and one pair of SNPs was associated to both traits. Noteworthy was that

inversions on chromosomes 1, 4, and 9, taken together, were enriched for

phenotypically-associated SNPs ($\chi^2 = 8.95$, P-value=0.0028). We recovered a

borderline significant enrichment with the correction K=5 ($\chi^2 = 3.82$, P-value=0.051).

Finally, we asked whether multiple SNPs contributed independently to the

phenotypic variation of a single trait. We tested a multiple SNP model where SNPs

were added incrementally when significantly associated (FDR < 0.10). We found 2, 3

and 2 SNPs for female, male flowering time and height of the highest ear,

respectively (S5 Table). Except for the latter trait, the SNPs were located on different

chromosomes.

Discussion

Plants are excellent systems to study local adaptation. First, owing to their

sessile nature, local adaptation of plant populations is pervasive [13]. Second,

environmental effects can be efficiently controlled in common garden experiments,

facilitating the identification of the physiological, morphological and phenological

traits influenced by spatially-variable selection [64]. Identification of the determinants

of complex trait variation and their covariation in natural populations is however

challenging [65]. While population genomics has brought a flurry of tools to detect

footprints of local adaptation, their reliability remains questioned [61]. In addition,

local adaptation and demographic history frequently follow the same geographic route,

making the disentangling of trait, molecular, and environmental variation, particularly

arduous. Here we investigated those links on a well-established outcrossing system,

the closest wild relatives of maize, along altitudinal gradients that display

considerable environmental shifts over short geographical scales.

The syndrome of altitudinal adaptation results from selection at multiple co-

adapted traits.

Common garden studies along elevation gradients have been conducted in

European and North American plants species [66]. Together with other studies, they

have revealed that adaptive responses to altitude are multifarious [67]. They include

physiological responses such as high photosynthetic rates [68], tolerance to frost [69],

biosynthesis of UV-induced phenolic components [70]; morphological responses with

reduced stature [71, 72], modification of leaf surface [73], increase in leaf non-

glandular trichomes [74], modification of stomata density; and phenological

responses with variation in flowering time [75], and reduced growth period [76].

Our multivariate analysis of teosinte phenotypic variation revealed a marked

differentiation between teosinte subspecies along an axis of variation (21.26% of the

total variation) that also discriminated populations by altitude (Fig 2A & B). The

combined effects of assortative mating and environmental elevation variation may

generate, in certain conditions, trait differentiation along gradients without underlying

divergent selection [77]. While we did not measure flowering time differences among

populations *in situ*, we did find evidence for long distance gene flow between

gradients and subspecies (Fig 4 A & C). In addition, several lines of arguments

suggest that the observed clinal patterns result from selection at independent traits and

is not solely driven by differences in flowering time among populations. First, two

distinct methods accounting for shared population history concur with signals of

spatially-varying selection at ten out of the 18 traits (Table 1). Nine of them exhibited

a clinal trend of increase/decrease of population phenotypic values with elevation (S3

Fig) within at least one of the two subspecies. This number is actually conservative,

because these approaches disregard the impact of selective constraints which in fact

tend to decrease inter-population differences in phenotypes. Second, while male and

female flowering times were positively correlated, they displayed only subtle

correlations ($|r|<0.16$) with other spatially-varying traits except for grain weight and

length ($|r| <0.33$). Third, we observed convergence at multiple phenotypes between

the lowland populations from the two gradients that occurred despite their

geographical and genetical distance (Fig 4) again arguing that local adaptation drives

the underlying patterns.

Spatially-varying traits that displayed altitudinal trends, collectively defined a

teosinte altitudinal syndrome of adaptation characterized by early-flowering,

production of few tillers albeit numerous lateral branches, production of heavy, long

and large grains, and decrease in stomata density. We also observed increased leaf

pigmentation with elevation, although with a less significant signal (S3 Table),

consistent with the pronounced difference in sheath color reported between

parviglumis and *mexicana* [78, 79]. Because seeds were collected from wild

populations, a potential limitation of our experimental setting is the confusion

between genetic and environmental maternal effects. Environmental maternal effects

could bias upward our heritability estimates. However, our results corroborate

previous findings of reduced number of tillers and increased grain weight in *mexicana*

compared with *parviglumis* [80]. Thus although maternal effects could not be fully

discarded, we believe they were likely to be weak.

The trend towards depleted stomata density at high altitudes (S3 Fig) could

arguably represent a physiological adaptation as stomata influence components of

plant fitness through their control of transpiration and photosynthetic rate [81]. Indeed,

in natural accessions of *A. thaliana*, stomatal traits showed signatures of local

adaptation and were associated with both climatic conditions and water-use efficiency

[82]. Furthermore, previous work has shown that in arid and hot highland

environments, densely-packed stomata may promote increased leaf cooling in

response to desiccation [83] and may also counteract limited photosynthetic rate with

decreasing pCO_2 [84]. Accordingly, increased stomata density at high elevation sites

has been reported in alpine species such as the European beech [85] as well as in

populations of *Mimulus guttatus* subjected to higher precipitations in the Sierra

Nevada [86]. In our case, higher elevations display both *arid* environment and *cooler*

temperatures during the growing season, features perhaps more comparable to other

tropical mountains for which a diversity of patterns in stomatal density variation with

altitude has been reported [87]. Further work will be needed to decipher the

mechanisms driving the pattern of declining stomata density with altitude in teosintes.

Altogether, the altitudinal syndrome was consistent with natural selection for rapid

life-cycle shift, with early-flowering in the shorter growing season of the highlands

and production of larger propagules than in the lowlands. This altitudinal syndrome

evolved in spite of detectable gene flow.

Although we did not formally measure biomass production, the lower number

of tillers and higher amount and size of grains in the highlands when compared with

the lowlands may reflect trade-offs between allocation to grain production and

vegetative growth [88]. Because grains fell at maturity and a single teosinte individual

produces hundreds of ears, we were unable to provide a proxy for total grain

production. The existence of fitness-related trade-offs therefore still needs to be

formally addressed.

Beyond trade-offs, our results more generally question the extent of

correlations between traits. In maize, for instance, we know that female and male

flowering time are positively correlated and that their genetic control is in part

determined by a common set of genes [89]. They themselves further increase with

yield-related traits [90]. Response to selection for late-flowering also led to a

correlated increase in leaf number in cultivated maize [91], and common genetic loci

have been shown to determine these traits as well [92]. Here we found strong positive

correlations between traits: male and female flowering time, grain length and width,

plant height and height of the lowest or highest ear. Strong negative correlations were

observed instead between grain weight and both male and female flowering time.

Trait correlations were therefore partly consistent with previous observations in maize,

suggesting that they were inherited from wild ancestors.

Footprints of past adaptation are relevant to detect variants involved in present

phenotypic variation.

The overall level of differentiation in our outcrossing system ($F_{ST} \approx 22\%$) fell

within the range of previous estimates (23% [93]and 33% [55] for samples

encompassing both teosinte subspecies). It is relatively low compared to other

systems such as the selfer *Arabidopsis thaliana*, where association panels typically

display maximum values of F_{ST} around 60% within 10kb-windows genome-wide [94].

Nevertheless, correction for sample structure is key for statistical associations

between genotypes and phenotypes along environmental gradients. This is because

outliers that display lowland/highland differentiation co-vary with environmental

factors, which themselves may affect traits [95]. Consistently, we found that 73.7%

SNPs associated with phenotypic variation at K=5, but only 13.5% of them did so

when considering a genetic structure with 11 populations. Except for one, the latter

set of SNPs represented a subset of the former. Because teosinte subspecies

differentiation was fully accounted for at K=5 (as shown by the clear distinction

between *mexicana* populations and the rest of the samples, Fig 4A), the inflation of

significant associations at K=5 is not due to subspecies differentiation, but rather to

residual stratification among populations within genetic groups. Likewise, recent

studies in humans, where global differentiation is comparatively low [96] have shown

that incomplete control for population structure within European samples strongly

impacts association results [61, 97]. Controlling for such structure may be even more

critical in domesticated plants, where genetic structure is inferred *a posteriori* from

genetic data (rather than *a priori* from population information) and pedigrees are

often not well described. Below, we show that considering more than one correction

using minor peaks delivered by the Evanno statistic (S5 Fig) can be informative.

Considering a structure with 5 genetic groups, the number of SNPs associated

per trait varied from 1 to 55, with no association for leaf and grain coloration (S5

Table). False positives likely represent a greater proportion of associations at K=5 as

illustrated by a slight excess of small P-values when compared with a correction with

11 populations for most traits (S10 Fig). Nevertheless, our analysis recovered credible

candidate adaptive loci that were no longer associated when a finer-grained

population structure was included in the model. For instance at K=5, we detected

Sugary1 (*Su1*), a gene encoding a starch debranching enzyme that was selected during

maize domestication and subsequent breeding [98, 99]. We found that *Su1* was

associated with variation at six traits (male and female flowering time, tassel

branching, height of the highest ear, grain weight and stomata density) pointing to

high pleiotropy. A previous study reported association of this gene to oil content in

teosintes [100]. In maize, this gene has a demonstrated role in kernel phenotypic

differences between maize genetic groups [101]. *Su1* is therefore most probably a

true-positive. That this gene was no longer recovered with the 11-population structure

correction indicated that divergent selection acted among populations. Indeed, allelic

frequency was highly contrasted among populations, with most populations fixed for

one or the other allele, and a single population with intermediate allelic frequency.

With the 11-population correction, very low power is thus left to detect the effect of

Su1 on phenotypes.

Although the confounding population structure likely influenced the genetic

associations, experimental evidence indicates that an appreciable proportion of the

variants recovered with both K=5 and 11 populations are true-positives (S5 Table).

One SNP associated with female and male flowering time, as well as with plant height

and grain length (at K=5 only for the two latter traits) maps within the *phytochrome*

B2 (SNP_210; *phyB2*) gene. Phytochromes are involved in perceiving light signals

and are essential for growth and development in plants. The maize gene *phyB2*

regulates the photoperiod-dependent floral transition, with mutants producing early

flowering phenotypes and reduced plant height [102]. Genes from the

phosphatidylethanolamine-binding proteins (PEBPs) family – *Zea mays*

CENTRORADIALIS (*ZCN*) family in maize – are also well-known to act as promotor

and repressor of the floral transition in plants [103]. Z*CN8* is the main floral activator

of maize [104], and both *ZCN8* and *ZCN5* strongly associate with flowering time

variation [101, 105]. Consistently, we found associations of male and female

flowering time with *PEBP18* (SNP_15). It is interesting to note that SNPs at two

flowering time genes, *phyB2* and *PEBP18*, influenced independently as well as in

combination both female and male flowering time variation (S5 Table).

The proportion of genic SNPs associated to phenotypic variation was not

significantly higher than that of non-genic SNPs (i.e, SNPs >1kb from a gene) ($\chi^2_{(df=1)}$

= 0.043, P-value = 0.84 at K=5 and $\chi^2_{(df=1)}$ =1.623, P-value =0.020 with 11

populations) stressing the importance of considering both types of variants [106]. For

instance, we discovered a non-genic SNP (SNP_149) that displayed a strong

association with leaf width variation as well as a pattern of allele frequency shift with

altitude among populations (Fig 6B).

Physically-linked and independent SNPs both contribute to the establishment of

adaptive genetic correlations.

We found limited LD among our outlier SNPs (S10 Fig) corroborating

previous reports (LD decay within <100bp, [58, 93]). However, the subset of

phenotypically-associated SNPs displayed greater LD, a pattern likely exacerbated by

three Mb-scale inversions located on chromosomes 1 (*Inv1n*), 4 (*Inv4m*) and 9 (*Inv9e*)

that, taken together, were enriched for SNPs associated with environmental variables

related to altitude and/or SNPs associated with phenotypic variation. Previous work

[55, 56] has shown that *Inv1n* and *Inv4m* segregate within both *parviglumis* and

mexicana, while two inversions on chromosome 9, *Inv9d* and *Inv9e*, are present only

in some of the highest *mexicana* populations; such that all four inversions also follow

an altitudinal pattern. Our findings confirmed that three of these inversions possessed

an excess of SNPs with high F_{ST} between subspecies and between low- and high-

mexicana populations for *Inv9e* [57]. Noteworthy *Inv9d* contains a large ear leaf

width quantitative trait locus in maize [106]. Corroborating these results, we found

consistent association between the only SNP located within this inversion and leaf

width variation in teosinte populations (S5 Table). Overall, our results further

strengthen the role of chromosomal inversions in teosinte altitudinal adaptation.

Because inversions suppress recombination between inverted and non-inverted

genotypes, their spread has likely contributed to the emergence and maintenance of

locally adaptive allelic combinations in the face of gene flow, as reported in a

growing number of other models (reviewed in [107]) including insects [108], fish

[109], birds [110] and plants [26, 111]. But we also found three cases of multi-SNP

determinism of traits (male and female flowering time and height of the highest ear,

Table S5) supporting selection of genetically independent loci. Consistently with

Weber et al. [100], we found that individual SNPs account for small proportions of

the phenotypic variance (S7 Table). Altogether, these observations are consistent with

joint selection of complex traits determined by several alleles of small effects, some

of which being maintained in linkage through selection of chromosomal

rearrangements.

Conclusion.

Elevation gradients provide an exceptional opportunity for investigating

variation of functional traits in response to continuous environmental factors at short

geographical scales. Here we documented patterns indicating that local adaptation,

likely facilitated by the existence of chromosomal inversions, allows teosintes to cope

with specific environmental conditions in spite of gene flow. We detected an

altitudinal syndrome in teosintes composed of sets of independent traits evolving

under spatially-varying selection. Because traits co-varied with environmental

differences along gradients, however, statistical associations between genotypes and

phenotypes largely depended on control of population stratification. Yet, several of

the variants we uncovered seem to underlie adaptive trait variation in teosintes.

Adaptive teosinte trait variation is likely relevant for maize evolution and breeding.

Whether the underlying SNPs detected in teosintes bear similar effects in maize or

whether their effects differ in domesticated backgrounds will have to be further

investigated.

Material and Methods

Description of teosinte populations and sampling.

We used 37 teosinte populations of *mexicana* (16) and *parviglumis* (21) subspecies from two previous collections [57, 58, 112] to design our sampling. These populations (S1 Table) are distributed along two altitudinal gradients (Fig 1). We plotted their altitudinal profiles using R 'raster' package [113] (S1 Fig). We further obtained 19 environmental variable layers from http://idrisi.uaemex.mx/distribucion/superficies-climaticas-para-mexico. These high-resolution layers comprised monthly values from 1910 to 2009 estimated via interpolation methods [107]. We extracted values of the 19 climatic variables for each population (S1 Table). Note that high throughput sequencing (HTS) data were obtained in a previous study for six populations out of the 37 (M6a, P1a, M7b, P2b, M1b and P8b; Fig 1, S1 Table) to detect candidate genomic regions for local adaptation [58]. The four highest and lowest of these populations were included in the association panel described below.

We defined an association panel of 11 populations on which to perform a genotype-phenotype association study (S1 Table). Our choice was guided by grain availability as well as the coverage of the whole climatic and altitudinal ranges. Hence, we computed Principal Component Analyses (PCA) for each gradient from environmental variables using the FactoMineR package in R [114] and added altitude to the PCA graphs as a supplementary variable. Our association panel comprised five populations from a first gradient (*a*) – two *mexicana* and three *parviglumis,* and six populations from a second gradient (*b*) – one *mexicana* and five *parviglumis* (Fig 1).

Finally, we extracted available SNP genotypes generated with the

MaizeSNP50 Genotyping BeadChip for 28 populations out of our 37 populations [57]

(S1 Table). From this available SNP dataset, we randomly sampled 1000 SNPs found

to display no selection footprint [57], hereafter neutral SNPs. Data for neutral SNPs

(Data S1) are available at: 10.6084/m9.figshare.9901472. We used this panel of 28

populations to investigate correlation with environmental variation. Note that 10 out

of the 28 populations were common to our association panel, and genotypes were

available for 24 to 34 individuals per population, albeit different from the ones of our

association mapping panel.

Common garden experiments

We used two common gardens for phenotypic evaluation of the association

panel (11 populations). Common gardens were located at INIFAP (Instituto Nacional

de Investigaciones Forestales, Agricolas y Pecuaria) experimental field stations in the

state of Guanajuato in Mexico, one in Celaya municipality at the Campo

Experimental Bajío (CEBAJ) (20°31'20'' N, 100°48'44''W) at 1750 meters of

elevation, and one in San Luis de la Paz municipality at the Sitio Experimental Norte

de Guanajuato (SENGUA) (21°17'55''N, 100°30'59''W) at 2017 meters of elevation.

These locations were selected because they present intermediate altitudes (S1 Fig).

The two common gardens were replicated in 2013 and 2014.

The original sampling contained 15 to 22 mother plants per population. Eight

to 12 grains per mother plant were sown each year in individual pots. After one

month, seedlings were transplanted in the field. Each of the four fields (2 locations, 2

years) was separated into four blocks encompassing 10 rows and 20 columns. We

evaluated one offspring of ~15 mother plants from each of the 11 teosinte populations

in each block, using a semi-randomized design, i.e. each row containing one or two

individuals from each population, and individuals being randomized within row,

leading to a total of 2,640 individual teosinte plants evaluated.

SSR genotyping and genetic structuring analyses on the association panel

In order to quantify the population structure and individual kinship in our

association panel, we genotyped 46 SSRs (S4 Table). Primers sequences are available

from the maize database project [115] and genotyping protocol were previously

published [116]. Genotyping was done at the GENTYANE platform (UMR INRA

1095, Clermont-Ferrand, France). Allele calling was performed on electropherograms

with the GeneMapper® Software Applied Biosystems®. Allele binning was carried

out using Autobin software [117], and further checked manually.

We employed STRUCTURE Bayesian classification software to compute a

genetic structure matrix on individual genotypes. Individuals with over 40% missing

data were excluded from analysis. For each number of clusters (K from 2 to 13), we

performed 10 independent runs of 500,000 iterations after a burn-in period of 50,000

iterations, and combined these 10 replicates using the LargeKGreedy algorithm from

the CLUMPP program [118]. We plotted the resulting clusters using DISTRUCT

software. We then used the Evanno method [119] to choose the optimal K value. We

followed the same methodology to compute a structure matrix from the outlier SNPs.

We inferred a kinship matrix **K** from the same SSRs using SPAGeDI [120].

Kinship coefficients were calculated for each pair of individuals as correlation

between allelic states [121]. Since teosintes are outcrossers and expected to exhibit an

elevated level of heterozygosity, we estimated intra-individual kinship to fill in the

diagonal. We calculated ten kinship matrices, each excluding the SSRs from one out

of the 10 chromosomes. Microsatellite data (Data S2) are available

at: 10.6084/m9.figshare.9901472

In order to gain insights into population history of divergence and admixture,

we used 1000 neutral SNPs (i.e. SNPs genotyped by Aguirre-Liguori and

collaborators [57] and that displayed patterns consistent with neutrality among 49

teosinte populations) genotyped on 10 out of the 11 populations of the association

panel to run a TreeMix analysis (TreeMix version 1.13 [122]. TreeMix models

genetic drift to infer populations splits from an outgroup as well as migration edges

along a bifurcating tree. We oriented the SNPs using the previously published

MaizeSNP50 Genotyping BeadChip data from the outgroup species *Tripsacum*

dactyloides [55]. We tested from 0 to 10 migration edges. We fitted both a simple

exponential and a non-linear least square model (threshold of 1%) to select the

optimal number of migration edges as implemented in the OptM R package [123]. We

further verified that the proportion of variance did not substantially increase beyond

the optimal selected value.

Phenotypic trait measurements

We evaluated a total of 18 phenotypic traits on the association panel (S2

Table). We measured six traits related to plant architecture (PL: Plant Height, HLE:

Height of the Lowest Ear, HHE: Height of the Highest Ear, Til: number of Tillers,

LBr: number of Lateral Branches, NoE: number of Nodes with Ears), three traits

related to leave morphologies (LeL: Leaf Length, LeW: Leaf Width, LeC: Leaf

Color), three traits related to reproduction (MFT: Male Flowering Time, FFT: Female

Flowering Time, TBr : Tassel Branching), five traits related to grains (Gr: number of

Grains per ear, GrL: Grain Length, GrWi: Grain Width, GrWe: Grain Weight, GrC:

Grain Color), and one trait related to Stomata (StD: Stomata Density). These traits

were chosen because we suspected they could contribute to differences among

teosinte populations based on a previous report of morphological characterization on

112 teosinte collections grown in five localities [124].

We measured the traits related to plant architecture and leaves after silk

emergence. Grain traits were measured at maturity. Leaf and grain coloration were

evaluated on a qualitative scale. For stomata density, we sampled three leaves per

plant and conserved them in humid paper in plastic bags. Analyses were undertaken at

the Institute for Evolution and Biodiversity (University of Münster) as followed: 5mm

blade discs were cut out from the mid length of one of the leaves and microscopic

images were taken after excitation with a 488nm laser. Nine locations ($0.15mm^2$) per

disc were captured with 10 images per location along the z-axis (vertically along the

tissue). We automatically filtered images based on quality and estimated leaf stomata

density using custom image analysis algorithms implemented in Matlab. For each

sample, we calculated the median stomata density over the (up to) nine locations. To

verify detection accuracy, manual counts were undertaken for 54 random samples.

Automatic and manual counts were highly correlated ($R^2=0.82$), indicating reliable

detection (see S1 Annex StomataDetection, Dittberner and de Meaux, for a detailed

description). The filtered data set of phenotypic measurements (Data S3) is available

at: 10.6084/m9.figshare.9901472.

Statistical analyses of phenotypic variation

In order to test for genetic effects on teosinte phenotypic variation, we

decomposed phenotypic values of each trait considering a fixed population effect plus

a random mother-plant effect (model M1):

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \delta_l + \chi_{il} + \psi_{jl} + P_{m/l} + \varepsilon_{ijklm} \qquad \text{(M1)}$$

where the response variable Y is the observed phenotypic value, μ is the total mean, α_i

is the fixed year effect (i = 2013, 2014), β_j the fixed field effect (j = field station,

SENGUA, CEBAJ), θ_{ij} is the year by field interaction, $\gamma_{k/ij}$ is the fixed block effect

(k = 1, 2, 3, 4) nested within the year-by-field combination, δ_l is the fixed effect of the

population of origin (l = 1 to 11*)*, χ_{il} is the year by population interaction, ψ_{jl} is the

field by population interaction, $P_{m/l}$ is the random effect of mother plant (m = 1 to 15)

nested within population, and ε_{ijklm} is the individual residue. Identical notations were

used in all following models. For the distribution of the effects, the same variance was

estimated within all populations. Mixed models were run using ASReml v.3.0 [125]

and MM4LMM v2.0.1 [https://rdrr.io/cran/MM4LMM/man/MM4LMM-

package.html, update by F. Laporte] R packages, which both gave very similar results,

and fixed effects were tested through Wald tests.

For each trait, we represented variation among populations using box-plots on

mean values per mother plant adjusted for the experimental design following model

M'1:

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + p_{m/l} + \varepsilon_{ijklm} \qquad \text{(M'1)}$$

where mother plant within population is considered as fixed. We used the function

predict to obtain least-square means (ls-means) of each mother plant, and looked at

the tendencies between population's values. All fixed models were computed using

lm package in R, and we visually checked the assumptions of residues independence

and normal distribution.

We performed a principal component analysis (PCA) on phenotypic values

corrected for the experimental design, using FactoMineR package in R [114] from the

residues of model M2 computed using the *lm* package in R:

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \varepsilon_{ijklm} \qquad \text{(M2)}$$

813   Finally, we tested for altitudinal effects on traits by considering the altitude of

814   the sampled population (*l*) as a covariate (ALT) and its interaction with year and field

815   in model M3:

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + c.ALT_l + a_i.ALT_l + b_j.ALT_l + P_{m/l} + \varepsilon_{ijklm}$$

(M3)

where all terms are equal to those in model M1 except that the fixed effect of the

population of origin was replaced by a regression on the population altitude (ALT_l).

Detection of selection acting on phenotypic traits

We aimed at detecting traits evolving under spatially varying selection by

comparing phenotypic to neutral genotypic differentiation. Q_{st} is a statistic analogous

to F_{ST} but for quantitative traits, which can be described as the proportion of

phenotypic variation explained by differences among populations [19, 107].

Significant differences between Q_{ST} and F_{ST} can be interpreted as evidence for

spatially-varying ($Q_{ST} > F_{ST}$) selection [126]. We used the R package *QstFstComp*

[127] that is adequate for experimental designs with randomized half-sibs in

outcrossing species. We used individuals that were both genotyped and phenotyped

on the association panel to establish the distribution of the difference between

statistics ($Q_{ST} - F_{ST}$) under the neutral hypothesis of evolution by drift - using the half-

sib dam breeding design and 1000 resamples. We next compared it to the observed

difference with 95% threshold cutoff value in order to detect traits under spatially-

varying selection.

In addition to Q_{ST}-F_{ST} analyses, we employed the DRIFTSEL R package [128]

to test for signal of selection of traits while accounting for drift-driven population

divergence and genetic relatedness among individuals (half-sib design). DRIFTSEL is a

Bayesian method that compares the probability distribution of predicted and observed

mean additive genetic values. It provides the S statistic as output, which measures the

posterior probability that the observed population divergence arose under divergent

selection (S~1), stabilizing selection (S~0) or genetic drift (intermediate S values)

[59]. It is particularly powerful for small datasets, and can distinguish between drift

and selection even when Q_{ST}-F_{ST} are equal [59]. We first applied RAFM to estimate the

F_{ST} value across populations, and the population-by-population coancestry coefficient

matrix. We next fitted both the RAFM and DRIFTSEL models with 15,000 MCMC

iterations, discarded the first 5,000 iterations as a transient, and thinned the remaining

by 10 to provide 1000 samples from the posterior distribution. Note that DRIFTSEL was

slightly modified because we had information only about the dams, but not the sires,

of the phenotyped individuals. We thus modified DRIFTSEL with the conservative

assumption of all sires being unrelated. Because DRIFTSEL does not require that the

same individuals were both genotyped and phenotyped, we used SSRs and phenotype

data of the association panel as well as the set of neutral SNPs and phenotype data on

10 out of the 11 populations. For the SNP analyses, we selected out of the 1000

neutral SNPs the 465 most informative SNPs based on the following criteria:

frequency of the less common variant at least 10%, and proportion of missing data at

most 1%. Finally, we estimated from DRIFTSEL the posterior probability of the

ancestral population mean for each trait as well as deviations of each population from

these values.

Both Q_{ST}-F_{ST} and DRIFTSEL rely on the assumption that the observed

phenotypic variation was determined by additive genotypic variation. We thus

estimated narrow-sense heritability for each trait in each population to estimate the

proportion of additive variance in performance. We calculated per population narrow-

sense heritabilites as the ratio of the estimated additive genetic variance over the total

phenotypic variance on our common garden measurements using the MCMCglmm R

package [129] where half sib family is the single random factor, and the design (block

nested within year and field) is corrected as fixed factor. For three grain-related traits,

we also ran the same model but including mother plants phenotypic values calculated

from the remaining grains not sown. We ran 100,000 iterations with 10,000 burn-in,

inverse gamma (0.001; 0.001) as priors. We then calculated the mean and standard

deviation of the 11 per population h² estimates.

Pairwise correlations between traits.

We evaluated pairwise-correlations between traits by correlating the residues

obtained from model M4, that corrects the experiment design (year, field and blocks)

as well as the underlying genetic structure estimated from SSRs:

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_j + \gamma_{k/ij} + \sum_{n=1}^{4} b_n \cdot C^n_{ijklm} + \varepsilon_{ijklm} \qquad \text{(M4)}$$

875   where b_n is the slope of the regression of Y on the nth structure covariate C^n. Structure

876   covariate values (C^n covariates, from STRUCTURE output) were calculated at the

877   individual level, i.e. for each offspring of mother plant m from population l, grown in

878   the year i field j and block k. C^n are thus declared with $ijklm$ indices, although they are

879   purely genetic covariates.

Genotyping of outlier SNPs on 28 populations

We extracted total DNA from each individual plant of the association panel as

well as 20 individuals from each of the 18 remaining populations that were not

included in the association panel (Table 1). Extractions were performed from 30 mg

of lyophilized adult leaf material following recommendations of DNeasy 96 Plant Kit

manufacturer (QIAGEN, Valencia, CA, USA). We genotyped outlier SNPs using

Kompetitive Allele Specific PCR technology (KASPar, LGC Group) [130]. Data for

outlier SNPs (Data S4 and Data S5) are available at: 10.6084/m9.figshare.9901472.

Among SNPs identified as potentially involved in local adaptation, 270 were

designed for KASPar assays, among which 218 delivered accurate quality data. Of the

218 SNPs, 141 were detected as outliers in two previous studies using a combination

of statistical methods – including F_{ST}-scans [131], Bayescan [32] and Bayenv2 [35,

132], Bayescenv [133] – applied to either six of our teosinte populations [58] or to a

broader set of 49 populations genotyped by the Illumina® MaizeSNP50 BeadChip

[57]. The remaining outlier SNPs (77) were detected by F_{ST}-scans from six

populations (S7 Fig, S5 Table), following a simplified version of the rationale in [58]

by considering only differentiation statistics: SNPs were selected if they displayed

both a high differentiation (5% highest F_{ST} values) between highland and lowland

populations in at least one of the two gradients, and a high differentiation (5% highest

F_{ST} values) between highland and lowland populations either within *parviglumis* (P2b

and P8b) or within *mexicana* (M7b and M1b) or both in gradient *b* (S1 Fig). We

thereby avoided SNPs fixed between the two subspecies.

Association mapping

We tested the association of phenotypic measurements with outlier SNPs on a

subset of individuals for which (1) phenotypic measurements were available, (2) at

least 60% of outlier SNPs were adequately genotyped, and (3) kinship and cluster

membership values were available from SSR genotyping. For association, we

removed SNPs with minor allele frequency lower than 5%.

In order to detect statistical associations between outlier SNPs and phenotypic

variation, we used the following mixed model derived from [128]:

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \sum_{n=1}^{4} b_n \cdot C_{ijklm}^n + \zeta_o + u_{ijklm} + \varepsilon_{ijklm} \qquad \text{(M5)}$$

where ζ is the fixed bi-allelic SNP factor with one level for each of the three

genotypes (o=0, 1, 2; with o=1 for heterozygous individuals), and u_{ijklm} is the random

genetic effect of the individual. We assumed that the vector of u_{ijklm} effects followed a

N(0,$\mathbf{K}$ σ^2u) distribution, where $\mathbf{K}$ is the kinship matrix computed as described above.

A variant of model M5 was employed to test for SNP association to traits,

while correcting for structure as the effect of population membership (δ_l), δ being a

factor with 11 levels (populations):

$$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \delta_l + \zeta_o + u_{ijklm} + \varepsilon_{ijklm} \qquad \text{(M5}'\text{)}$$

In order to avoid overcorrection of neutral genetic structure and improve

power, we ran the two models independently for each chromosome using a kinship

matrix $\mathbf{K}$ estimated from all SSRs except those contained in the chromosome of the

tested SNP [134]. We tested SNP effects through the Wald statistics, and applied a 10%

False Discovery Rate (FDR) threshold for each phenotype separately. In order to

validate the correction for genetic structure, the 38 multiallelic SSR genotypes were

transformed into biallelic genotypes, filtered for MAF > 5%, and used to run

associations with the complete M5 and M5' models, as well as the M5 models

excluding either kinship or both structure and kinship. For each trait, we generated

QQplots of P-values for each of these models.

Multiple SNP models were built by successively adding at each step the most

significant SNP, as long as its FDR was lower than 0.10. We controlled for population

structure considering 11 populations and used the kinship matrix that excluded the

SSR on the same chromosome as the last tested SNP.

934   Environmental correlation of outlier SNPs

We tested associations between allelic frequency at 171 outlier SNPs and

environmental variables across 28 populations, using Bayenv 2.0 [40, 111]. Because

environmental variables are highly correlated, we used the first two principal

component axes from the environmental PCA analysis (PCenv1 and PCenv2) to run

Bayenv 2.0. This software requires a neutral covariance matrix, that we computed

from the available dataset of 1000 neutral SNPs (S1 Table). We performed 100,000

iterations, saving the matrix every 500 iterations. We then tested the correlation of

these to the last matrix obtained, as well as to an F_{ST} matrix calculated with

BEDASSLE [135], as described in [57].

For each outlier SNP, we compared the posterior probability of a model that

included an environmental factor (PCenv1 or PCenv2) to a null model. We

determined a 5% threshold for significance of environmental association by running

100,000 iterations on neutral SNPs. We carried out five independent runs for each

outlier SNP and evaluated their consistency from the coefficient of variation of the

Bayes factors calculated among runs.

In order to test whether environmental distance was a better predictor of allele

frequencies at candidate SNPs than geography, we used multiple regression on

distance matrices (MRM, [136]) implemented in the ecodist R package [137] for each

outlier SNP. We used pairwise F_{ST} values as the response distance matrix and the

geographic and environmental distance matrices as explanatory matrices. We

evaluated the significance of regression coefficients by 1000 permutations and

iterations of the MRM. We determined the total number of environmentally and

geographically associated SNPs (P-value<0.05) among outliers. We employed the

same methodology for our set of 1000 neutral SNPs.

Ethics Statement

961   All the field work has been done in Mexico in collaboration with Instituto Nacional

962   de Investigaciones Forestales, Agrícolas y Pecuarias, Celaya in Celaya.

Acknowledgments

We thank Jessica Melique for her help in gathering stomata data. We are very grateful

to Angelica Cibrian at Langebio (Cinvestav, Irapuato, Mexico) who let us use her lab

to lyophilize our samples, and store them. Valeria Souza greatly helped us with

logistic support in Mexico. Insights from Delphine Legrand, Pierre de Villemereuil

and Elodie Marchadier were important for analyzing correlations between traits and to

estimate heritabilities. We would like to thank warmly five anonymous reviewers for

their insightful comments on our work.

Supporting information captions

Figure S1: Altitudinal profiles along gradients *a* and *b*. Sampled populations are plotted on parallel altitudinal profiles for gradients *a* and *b*. Darker gray lines indicate lower latitude for gradient *a* and lower longitude for gradient *b*. Sampled populations are plotted by green circles (*parviglumis*) or red triangles (*mexicana*). The altitude of the two experimental fields (CEBAJ: 1750m and SENGUA: 2017m) are marked with stars on y-axes.

Figure S2: Principal Component Analysis of 19 climate variables for 37 teosinte populations. A: Projection of *parviglumis* (in green) and *mexicana* (in red) populations on the first PCA plane with gradients *a* and *b* indicated by triangles and circles, respectively. The 11 populations evaluated in common gardens are surrounded by a purple outline. Populations that were previously sequenced to detect selection footprints are shown in bold (S1 Table). B: Correlation circle of the 19 climatic variables on the first PCA plane. Climatic variables indicated as Tn (n from 1 to 11) and Pn (n from 12 to 19) are related to temperature and precipitation, respectively. Altitude, Latitude and Longitude (in blue) were added as supplementary variables, and CEBAJ and SENGUA field locations were added as supplementary individuals.

Figure S3: Box-plots of means adjusted by field, year and block, for all traits. Populations are ranked by altitude. *parviglumis* populations are shown in green and *mexicana* in red. Lighter colors are used for gradient 'a' and darker colors for gradient 'b'. Units of measurement correspond to those defined in S2 Table. For male and

female flowering time, we report values for all 11 populations although very few individuals from the two most lowland populations (P1a and P2b) flowered. Covariation with altitude was significant for all traits except for the number of nodes with ears on the main tiller (S3 Table).

Figure S4: Pairwise correlations between phenotypic traits. Pearson coefficient sign and magnitude for significant correlations between phenotypic traits after correction for experiment design (Model M'1). X: correlations that are not significant.

Figure S5. Evanno method calculations for population number ΔK in the association panel genotyped for 38 SSRs.

Figure S6. Genetic clustering of ancestry proportions in the association panel genotyped for 38 SSRs. Genetic clustering was computed for K=2 to K=11. Vertical lines (individuals) are partitioned into coloured segments whose length represents the admixture proportions from the K clusters.

Figure S7. Determination of the migration edge number in the TreeMix model. Observed Log likelihood values are plotted against the number of migration edges tested from 0 to 10, and two models are fitted to the data (A). Both the simple exponential and the non-linear least squares delivered an optimal value of 3 for the number of migration edges (change points). The model with 3 migration edges explained 98.75% of the variance, a substantial increase from the null model with no migration edge which is 95.7% (B).

Figure S8: Significance of Q_{ST}-F_{ST} difference for each trait. The dotted blue line indicates the 95% threshold of the simulated distributions and the red line refers to the observed difference. In this analysis, we considered as spatially-varying traits those for which the observed difference fell outside the 95% threshold. Note that Plant height was borderline significant. *: Set of traits detected by DRIFTSEL.

Figure S9: Genomic F_{ST}-scans on 6 teosinte populations. We computed 4 pairwise-F_{ST} values from 6 populations previously sequenced (S1 Table). Those include F_{ST} between lowland and highland populations of each gradient (P1a-M6a, P2b-M7b) as well as within subspecies on gradient *b* (P2b-P8b, M1b-M7b). F_{ST} values are averaged across sliding windows of 20 SNPs with a step of five SNPs (from top to bottom, chromosome 1 to 10) and normalized by subtracting the F_{ST} mean and dividing by the standard deviation across pairwise comparisons. Only the top 1% values are represented. The 1‰ thresholds for each pairwise comparisons are indicated by colored horizontal lines. Horizontal black bars indicate location of inversions on chromosome 1 (*Inv1n*), chromosome 4 (*Inv4m*) and chromosome 9 (*Inv9d*). The subset of 171 outlier SNPs analyzed in the present study is indicated with black diamond marks along the X axes.

Figure S10: QQ-plots of observed P-values and expected P-values generated from 38 SSRs. We employed three versions of the model M5 with correction for neither structure nor kinship, with correction for genetic structure (at *K*=5), with correction for genetic structure (at *K*=5 and with 11 populations) and kinship.

Figure S11: Manhattan plots of associations between 171 outlier SNPs and 12 phenotypic traits. X-axis indicates the positions of outlier SNPs on chromosomes 1 to 10, black and gray colors alternating per chromosome. Plotted on the Y-axis are the negative Log_{10}-transformed P values obtained for the $K=5$ model. Significant associations (10% FDR) are indicated considering either a structure matrix at $K=5$ (pink dots), for 11 populations (blue dots), or for both $K=5$ and 11 populations models (purple dots).

Figure S12: Pairwise Linkage Disequilibrium (LD) between outlier SNPs. Pairwise LD between 171 SNPs was estimated using r^2, and corrected for structure at $K=5$ and kinship computed from 38 SSRs. Blue shaded bars show the 23 SNPs found to associate with at least one phenotype under the 11 populations structure correction.

S1 Table. Description of 37 teosinte populations and sets of populations used in the present study by data types.

S2 Table. List of the 18 phenotypic traits measured and estimates of narrow-sense heritabilities (h^2).

S3 Table. Significance of main effects for each trait as determined by models M1 and M3.

S4 Table. Description of 46 SSRs and genotyping success rate.

S5 Table. Characteristics, association with phenotypes, effects and correlation with environment of outlier SNPs.

S6 Table. Number of individuals used to test associations between 171 SNPs and 18 phenotypes.

S7 Table. **Additive and dominance effects of SNPs associated to traits after the 11-population structure correction.**

References

1.   Whitlock MC. Modern approaches to local adaptation. The American
Naturalist. 2015;186.
2.   Bradshaw AD. Ecological significance of genetic variation between
populations. Perspectives on plant population ecology. 1984:213-28.
3.   Bulmer MGG. Multiple niche polymorphism. The American Naturalist.
1972;106:254-7.
4.   Endler JA. Natural selection in the wild. 1986:354.
5.   Gay L, Crochet PA, Bell DA, Lenormand T. Comparing clines on molecular
and phenotypic traits in hybrid zones: A window on tension zone models. Evolution.
2008;62:2789-806.
6.   Lande R. Natural selection and random genetic drift in phenotypic evolution.
Evolution. 1976;30:314-34.
7.   Lenormand T. Gene flow and the limits to natural selection. Trends in
Ecology and Evolution. 2002;17:183-9.
8.   Whitlock MC, Gomulkiewicz R. Probability of fixation in a heterogeneous
environment. Genetics. 2005:1-40.
9.   Yeaman S, Otto SP. Establishment and maintenance of adaptive genetic
divergence under migration, selection and drift. Evolution. 2011;67:2123-9.
10.   Rundle HD, Nosil P. Ecological speciation. Ecology Letters. 2005;8:336-52.
11.   Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecology Letters.
2004;7:1225-41.
12.   Hereford J. A quantitative survey of local adaptation and fitness trade‐offs.
The American Naturalist. 2009;173:579-88.
13.   Leimu R, Fischer M. A meta-analysis of local adaptation in plants. PLoS
ONE. 2008;3.
14.   Tiffin P, Ross-Ibarra J. Advances and limits of using population genetics to
understand local adaptation. Trends in Ecology and Evolution. 2014;29:673-80.
15.   Garcia-Ramos G, Kirkpatrick M. Genetic models of adaptation and gene flow
in peripherial populations. Evolution. 1997;51:21-8.
16.   Slatkin M. Rare alleles as indicators of gene flow. Evolution. 1985;39(1):53-
65.
17.   Lande R. Neutral theory of quantitative genetic variance in an island model
with local extinction and colonization. Evolution. 1992;46:381-9.
18.   Whitlock MC. Neutral additive genetic variance in a metapopulation. Genetics
Research. 1999;74:215-21.
19.   Spitze K. Population structure in *Daphina obtusa*: quantitative genetic and
allozymic variation. Genetics Society of America. 1993;135:367-74.
20.   Wright S. The genetical structure of populations. Annals of Eugenics.
1951;15:323-54.
21.   Luquet E, Léna J-P, Miaud C, Plénet S. Phenotypic divergence of the common
toad (*Bufo bufo*) along an altitudinal gradient: evidence for local adaptation. Heredity.
2015;114:69-79.
22.   Roschanski AM, Csilléry K, Liepelt S, Oddou-Muratorio S, Ziegenhagen B,
Huard Fdr, et al. Evidence of divergent selection for drought and cold tolerance at
landscape and local scales in *Abies alba* Mill. in the French Mediterranean Alps.
Molecular Ecology. 2016;25:776-94.

23.   Kawakami T, Morgan TJ, Nippert JB, Ocheltree TW, Keith R, Dhakal P, et al.
Natural selection drives clinal life history patterns in the perennial sunflower species,
Helianthus maximiliani. Molecular Ecology. 2011;20:2318-28.
24.   Moyers BT, Rieseberg LH. Remarkable life history polymorphism may be
evolving under divergent selection in the silverleaf sunflower. Molecular Ecology.
2016;25:3817-30.
25.   Kirkpatrick M, Barton N. Chromosome inversions, local adaptation and
speciation. Genetics. 2006.
26.   Lowry DB, Willis JH. A widespread chromosomal inversion polymorphism
contributes to a major life-history transition, local adaptation, and reproductive
isolation. PLoS Biology. 2010;8.
27.   Legrand D, Larranaga N, Bertrand R, Ducatez S, Calvez O, Stevens VM, et al.
Evolution of a butterfly dispersal syndrome. 2016.
28.   Bierne N, Welch J, Loire E, Bonhomme F, David P. The coupling hypothesis:
Why genome scans may fail to map local adaptation genes. Molecular Ecology.
2011;20:2044-72.
29.   Lewontin RC, Krakauer J. Distribution of gene frequency as a test of the
theory of the selective neutrality of polymorphisms. Genetics. 1973;74:175-95.
30.   Beaumont MA, Nichols RA. Evaluating loci for use in the genetic analysis of
population structure. Proceedings of the Royal Society B: Biological Sciences.
1996;263:1619-26.
31.   Vitalis R, Dawson K, Boursot P. Interpretation of variation across marker loci
as evidence of selection. Genetics. 2001;158:1811-23.
32.   Foll M, Gaggiotti O. A genome scan method to identify selected loci
appropriate for both dominant and codominant markers: A Bayesian perspective.
Genetics. 2008;180:977-93.
33.   Excoffier LaH, T and Foll, Matthieu. Detecting loci under selection in a
hierarchically structured population. Heredity. 2009;103:285-98.
34.   Bonhomme M, Chevalet C, Servin B, Boitard S, Abdallah J, Blott S, et al.
Detecting selection in population trees: The Lewontin and Krakauer test extended.
Genetics. 2010;186:241-62.
35.   Günther T, Coop G. Robust identification of local adaptation from allele
frequencies. Genetics Society of America. 2013;195:205-20.
36.   Lotterhos KE, Whitlock MC. Evaluation of demographic history and neutral
parameterization on the performance of FST outlier tests. Molecular Ecology.
2014;23:2178-92.
37.   Haasl RJ, Payseur BA. Fifteen years of genomewide scans for selection:
trends, lessons and unaddressed genetic sources of complication. Molecular Ecology.
2016;25:5-23.
38.   Le Corre V, Kremer A. The genetic differentiation at quantitative trait loci
under local adaptation. Molecular Ecology. 2012;21:1548-66.
39.   Yi X, Liang Y, Huerta-Sanchez E, Jin X, Xi Ping Cuo Z, Pool JE, et al.
Sequencing of fifty human exomes reveals adaptations to high altitude. Science.
2010;329:75-8.
40.   Coop G, Witonsky D, Di Rienzo A, Pritchard JK. Using environmental
correlations to identify loci underlying local adaptation. Genetics. 2010;185:1411-23.
41.   Guillot G, Renaud S, Ledevin R, Michaux J, Claude J. A unifying model for
the analysis of phenotypic, genetic, and geographic data. 2012;61:897-911.

42.   Frichot E, Schoville SD, Bouchard G, François O. Testing for associations
between loci and environmental gradients using latent factor mixed models.
Molecular Biology and Evolution. 2013;30:1687-99.
43.   Gautier M. Genome-wide scan for adaptive divergence and association with
population-specific covariates. Genetics. 2015;201:1555-79.
44.   Joost S, Bonin A, Bruford MW, Després L, Conord C, Erhardt G, et al. A
spatial analysis method (SAM) to detect candidate loci for selection: Towards a
landscape genomics approach to adaptation. Molecular Ecology. 2007;16:3955-69.
45.   Poncet BN, Herrmann D, Gugerli F, Taberlet P, Holderegger R, Gielly L, et
al. Tracking genes of ecological relevance using a genome scan in two independent
regional population samples of *Arabis alpina*. Molecular Ecology. 2010;19:2896-907.
46.   De Mita S, Thuillet AC, Gay L, Ahmadi N, Manel S, Ronfort J, et al.
Detecting selection along environmental gradients: Analysis of eight methods and
their effectiveness for outbreeding and selfing populations. Molecular Ecology.
2013;22:1383-99.
47.   Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, et al.
Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future
directions. The American Naturalist. 2016;188:379-97.
48.   Barton N, Hermisson J, Nordborg M. Why structure matters. eLife. 2019;8.
49.   Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek
AM. A map of local adaptation in Arabidopsis thaliana. Science. 2011;334:86-9.
50.   Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG,
et al. Adaptation to climate across the *Arabidopsis thaliana* genome. Science.
2011;334:83-6.
51.   Ross-Ibarra J, Tenaillon M, Gaut BS. Historical divergence and gene flow in
the genus Zea. Genetics. 2009;181:1399-413.
52.   Hufford MB, Martínez-Meyer E, Gaut BS, Eguiarte LE, Tenaillon MI.
Inferences from the historical distribution of wild and domesticated maize provide
ecological and evolutionary insight. PLoS ONE. 2012;7.
53.   Bilinski P, Albert PS, Berg JJ, Birchler JA, Grote MN, Lorant A, et al. Parallel
altitudinal clines reveal trends in adaptive evolution of genome size in *Zea mays*.
PLoS Genetics. 2018;14.
54.   Diez CM, Gaut BS, Meca E, Scheinvar E, Montes-Hernandez S, Eguiarte LE,
et al. Genome size variation in wild and cultivated maize along altitudinal gradients.
New Phytologist. 2013;199:264-76.
55.   Pyhäjärvi T, Hufford MB, Mezmouk S, Ross-Ibarra J. Complex patterns of
local adaptation in teosinte. Genome Biology and Evolution. 2013;5:1594-609.
56.   Fang Z, Pyhäjärvi T, Weber AL, Dawe RK, Glaubitz JC, Sánchez González
JdJ, et al. Megabase-scale inversion polymorphism in the wild ancestor of maize.
Genetics. 2012;191:883-94.
57.   Aguirre-Liguori JA, Tenaillon MI, Vázquez-Lobo A, Gaut BS, Jaramillo-
Correa JP, Montes-Hernandez S, et al. Connecting genomic patterns of local
adaptation and niche suitability in teosintes. Molecular Ecology. 2017;26:4226-40.
58.   Fustier MA, Brandenburg JT, Boitard S, Lapeyronnie J, Eguiarte LE,
Vigouroux Y, et al. Signatures of local adaptation in lowland and highland teosintes
from whole-genome sequencing of pooled samples. Molecular Ecology.
2017;26:2738-56.
59.   Ovaskainen O, Karhunen M, Zheng C, Arias JMC, Merilä J. A new method to
uncover signatures of divergent and stabilizing selection in quantitative traits.
Genetics. 2011;189:621-32.

60.   McKinney GJ, Varian A, Scardina J, Nichols KM. Genetic and morphological
divergence in three strains of brook trout Salvelinus fontinalis commonly stocked in
Lake Superior. PLoS ONE. [Research Support, Non-U.S. Gov't].
2014;9(12):e113809.
61.   Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al.
Polygenic adaptation on height is overestimated due to uncorrected stratification in
genome-wide association studies. eLife. 2019;8.
62.   Desrousseaux AD, Sandron F, Siberchicot A, Cierco-Ayrolles C, Mangin B.
Package ' LDcorSV '. 2017.
63.   Mangin B, Siberchicot A, Nicolas S, Doligez A, This P, Cierco-Ayrolles C.
Novel measures of linkage disequilibrium that correct the bias due to population
structure and relatedness. Heredity. 2012;108:285-91.
64.   Savolainen O, Lascoux M, Merilä J. Ecological genomics of local adaptation.
Nature reviews Genetics. 2013;14:807-20.
65.   Anderson JT, Willis JH, Mitchell-Olds T. Evolutionary genetics of plant
adaptation. Trends in Genetics. 2011;27:258-66.
66.   Halbritter AH, Fior S, Keller I, Billeter R, Edwards PJ, Holderegger R, et al.
Trait differentiation and adaptation of plants along elevation gradients. Journal of
Evolutionary Biology. 2018;31(6):784-800.
67.   Körner C. The use of 'altitude' in ecological research. Trends in Ecology and
Evolution. 2007;22:569-74.
68.   Friend AD, Woodward FI, Switsur VR. Field measurements of
photosynthesis, stomatal conductance, leaf nitrogen and δ 13 C along altitudinal
gradients in Scotland. Functional Ecology. 1989;3:117.
69.   Neuner G. Frost resistance in alpine woody plants. Frontiers in Plant Science.
2014;5.
70.   Frohnmeyer H, Staiger D. Update on ultraviolet-B light responses ultraviolet-
B radiation-mediated responses in plants. Balancing damage and protection.
2014;133:1420-8.
71.   Byars SG, Papst W, Hoffmann AA. Local adaptation and cogradient selection
in the alpine plant, *Poa hiemata*, along a narrow altitudinal gradient. Evolution.
2007;61:2925-41.
72.   Luo Y, Widmer A, Karrenberg S. The roles of genetic drift and natural
selection in quantitative trait divergence along an altitudinal gradient in *Arabidopsis*
thaliana. Heredity. 2015;114:220-8.
73.   Guerin GR, Wen H, Lowe AJ, Guerin GR, Wen H. Leaf morphology shift
linked to climate change. Population Ecology. 2012:882-6.
74.   Kofidis G, Bosabalidis AM, Moustakas M. Contemporary seasonal and
altitudinal variations of leaf structural features in oregano (*Origanum vulgare* L.).
Annals of Botany. 2003;92(5):635-45.
75.   Mendez-Vigo B, Pico FX, Ramiro M, Martinez-Zapater JM, Alonso-Blanco
C. Altitudinal and climatic adaptation is mediated by flowering traits and FRI, FLC,
and PHYC genes in *Arabidopsis*. Plant Physiology. 2011;157:1942-55.
76.   Oleksyn J, Modrzynski J, Tjoelker MG, Zytkowaik R, Reich PB, Karolewski
P. Growth and physiology of *Picea abies* populations from elevational transects:
common garden evidence for altitudinal ecotypes and cold adaptation. 1998:573-90.
77.   Soularue JP, Kremer A. Evolutionary responses of tree phenology to the
combined effects of assortative mating, gene flow and divergent selection. Heredity.
2014;113:485-94.

78.   Doebley JF. Maize introgression into teosinte -- a reappraisal. Annals of the
Missouri Botanical Garden. 1984;71:1100-13.
79.   Lauter N, Gustus C, Westerbergh A, Doebley J. The inheritance and evolution
of leaf pigmentation and pubescence in teosinte. Genetics Society of America.
2004;167:1949-59.
80.   Smith JSC, Goodman MM, Lester RN. Variation within teosinte. I. Numerical
analysis of morphological data. Economic Botany. 1981;35:187-203.
81.   Raven J, A. Selection pressures on stomatal evolution. New Phytologist.
2002;153:371-86.
82.   Dittberner H, Korte A, Mettler-Altmann T, Weber APM, Monroe G, de
Meaux J. Natural variation in stomata size contributes to the local adaptation of
water-use efficiency in *Arabidopsis thaliana*. Molecular Ecology. 2018:4052-65.
83.   Carlson JE, Adams CA, Holsinger KE. Intraspecific variation in stomatal
traits, leaf traits and physiology reflects adaptation along aridity gradients in a South
African shrub. Annals of Botany. 2016;117:195-207.
84.   Körner C, Mayr R. Stomatal behaviour in alpine plant communities between
600 and 2600 metres above sea level. In: Grace J, Ford ED, Jarvis PG, editors. Plants
and their Atmospheric Environment: Blackwell, Oxford,; 1981. p. pp 205-18.
85.   Bresson CC, Vitasse Y, Kremer A, Delzon S. To what extent is altitudinal
variation of functional traits driven by genetic adaptation in European oak and beech?
Tree Physiology. 2011;31:1164-74.
86.   Kooyers NJ, Greenlee AB, Colicchio JM, Oh M, Blackman BK. Replicate
altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought
stress in annual *Mimulus guttatus*. New Phytologist. 2015;206:152-65.
87.   Körner C, Neumayer M, Menendez-Riedl SP, Smeets-Scheel A. Functional
morphology of mountain plants. Flora. 1989;182:353-83.
88.   Jakobsson A, Eriksson O. A comparative study of seed number, seed size,
seedling size and recruitment in grassland plants. Oikos. 2000;88:494-502.
89.   Buckler ES, Holland JB, Bradbury PJ, Acharya CB, Brown PJ, Browne C, et
al. The genetic architecture of maize flowering time. Science. 2009;325:714-8.
90.   Moreau L, Charcosset A, Gallais A. Use of trial clustering to study QTL x
environment effects for grain yield and related traits in maize. Theoretical and
Applied Genetics. 2004;110:92-105.
91.   Durand E, Bouchet S, Bertin P, Ressayre A, Jamin P, Charcosset A, et al.
Flowering time in maize: Linkage and epistasis at a major effect locus. Genetics.
2012;190:1547-62.
92.   Li D, Wang X, Zhang X, Chen Q, Xu G, Xu D, et al. The genetic architecture
of leaf number and its genetic relationship to flowering time in maize. New
Phytologist. 2015;210:256-68.
93.   Aguirre‐Liguori JA, Gaut BS, Jaramillo‐Correa JP, Tenaillon MI, Montes
‐Hernández S, García‐Oliva F, et al. Divergence with gene flow is driven by local
adaptation to temperature and soil phosphorus concentration in teosinte subspecies (
Zea mays parviglumis and *Zea mays mexicana*) Molecular Ecology. 2019:2814-30.
94.   Consortium G. 1,135 Genomes reveal the global pattern of polymorphism in
Arabidopsis thaliana. Cell. 2016;166:481-91.
95.   Novembre J, Barton NH. Tread lightly interpreting polygenic tests of
selection. Genetics. 2018; 208(4):1351-5.
96.   Guo J, Wu Y, Zhu Z, Zheng Z, Trzaskowski M, Zeng J, et al. Global genetic
differentiation of complex traits shaped by natural selection in humans. Nature
Communications. 2018; 9(1):1865.

97.   Berg JJ, Harpak A, Sinnott-Armstrong N, Joergensen AM, Mostafavi H, Field
Y, et al. Reduced signal for polygenic adaptation of height in UK Biobank. eLife.
2019;8:1-47.
98.   Whitt SR, Wilson LM, Tenaillon MI, Gaut BS, Buckler ES. Genetic diversity
and selection in the maize starch pathway. Proceedings of the National Academy of
Sciences of the United States of America. 2002;99:12959-62.
99.   Jaenicke-Despres V, Buckler ES, Smith BD, Gilbert MTP, Cooper A, Doebley
J, et al. Early allelic selection in maize as revealed by ancient DNA. 2003;302:1206-9.
100.   Weber AL, Briggs WH, Rucker J, Baltazar BM, De Jesús Sánchez-González
J, Feng P, et al. The genetic architecture of complex traits in teosinte (*Zea mays* ssp.
parviglumis): New evidence from association mapping. Genetics. 2008;180:1221-32.
101.   Bouchet S, Servin B, Bertin P, Madur D, Combes V, Dumas F, et al.
Adaptation of maize to temperate climates: Mid-density genome-wide association
genetics and diversity patterns reveal key genomic regions, with a major contribution
of the Vgt2 (ZCN8) locus. PLoS ONE. 2013;8.
102.   Sheehan MJ, Kennedy LM, Costich DE, Brutnell TP. Subfunctionalization of
PhyB1 and PhyB2 in the control of seedling and mature plant traits in maize. Plant
Journal. 2007;49:338-53.
103.   Danilevskaya ON, Meng X, Hou Z, Ananiev EV, Simmons CR. A genomic
and expression compendium of the expanded PEBP gene family from maize. Plant
Physiology. 2007;146:250-64.
104.   Meng X, Muszynski MG, Danilevskaya ON. The FT-Like ZCN8 gene
functions as a floral activator and is involved in photoperiod sensitivity in maize. The
Plant Cell. 2011;23:942-60.
105.   Li YX, Li C, Bradbury PJ, Liu X, Lu F, Romay CM, et al. Identification of
genetic variants associated with maize flowering time using an extremely large multi-
genetic background population. The Plant journal : for cell and molecular biology.
2016;86:391-402.
106.   Yu J, Li X, Zhu C, Yeh C-T, Wu W, Takacs E, et al. Genic and non-genic
contributions to natural variation of quantitative traits in maize. Genome research.
2012:2436-44.
107.   Wellenreuther M, Bernatchez L. Eco-evolutionary genomics of chromosomal
inversions. Trends in Ecology and Evolution. 2018;33:427-40.
108.   Ayala D, Ullastres A, González J. Adaptation through chromosomal
inversions in *Anopheles*. Frontiers in Genetics. 2014;5:1-10.
109.   Barth JMI, Berg PR, Jonsson PR, Bonanomi S, Corell H, Hemmer-Hansen J,
et al. Genome architecture enables local adaptation of Atlantic cod despite high
connectivity. Molecular Ecology. 2017;26:4452-66.
110.   Lundberg M, Liedvogel M, Larson K, Sigeman H, Grahn M, Wright A, et al.
Genetic differences between willow warbler migratory phenotypes are few and cluster
in large haplotype blocks. Evolution Letters. 2017:155-68.
111.   Twyford AD, Friedman J. Adaptive divergence in the monkey flower *Mimulus*
guttatus is maintained by a chromosomal inversion. Evolution. 2015;69:1476-86.
112.   Díez CM, Gaut BS, Meca E, Scheinvar E, Montes‐Hernandez S, Eguiarte
LE, et al. Genome size variation in wild and cultivated maize along altitudinal
gradients. New Phytologist. 2013;199(1):264-76.
113.   Hijmans RJ, van Etten J, Cheng J, Mattiuzzi M, Sumner M, Greenberg JA, et
al. Package 'raster': geographic data analysis and modeling. 2018.
114.   Husson F, Josse J, Le S, Mazet J. Package 'FactoMineR'. An R package.
2016:96.

115. Andorf CM, Cannon EK, Portwood JL, Gardiner JM, Harper LC, Schaeffer
ML, et al. MaizeGDB update: New tools, data and interface for the maize model
organism database. Nucleic Acids Research. 2016;44:1195-201.
116. Camus-Kulandaivelu L, Veyrieras JB, Madur D, Combes V, Fourmann M,
Barraud S, et al. Maize adaptation to temperate climate: Relationship between
population structure and polymorphism in the Dwarf8 gene. Genetics.
2006;172:2449-63.
117. Guichoux E, Lagache S, Wagner S, Chaumeil P, Léger P, Lepais O, et al.
Current trends in microsatellite genotyping. Molecular Ecology Resources.
2011;11:591-611.
118. Jakobsson M, Rosenberg NA. CLUster Matching and Permutation Program
Version 1.1.2. 2007.
119. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of
individuals using the software STRUCTURE: A simulation study. Molecular
Ecology. 2005;14:2611-20.
120. Hardy OJ, Vekemans X. spagedi: a versatile computer program to analyse
spatial genetic structure at the individual or population levels. Molecular Ecology
Notes. 2002;2:618-20.
121. Loiselle BA, Sork VL, Nason JD, Graham C. Spatial genetic structure of a
tropical understory shrub, *Psychotria officinalis* (Rubiaceae). American Journal of
Botany. 1995;82:1420-5.
122. Pickrell JK, Pritchard JK. Inference of population splits and mixtures from
genome-wide allele frequency data. PLOS Genetics. 2012;8(11):e1002967.
123. Fitak RRs. optM: an R package to optimize the number of migration edges
using threshold models. Journal of Heredity. 2019.
124. Sanchez JdJ, Kato Yamakake TA, Aguilar Sanmiguel M, Hernandez Casillas
JM, Lopez Rodriguez A, Ruiz Corral JA. Distribución y caracterización del teocintle.
1998:165.
125. Butler D, Cullis BR, Gilmour AR, Gogel BJ. ASReml-R reference manual.
Technical Report. 2007.
126. Holsinger KE, Weir BS. Genetics in geographically structured populations:
defining, estimating and interpreting F(ST). Nature reviews Genetics. 2009;10:639-
50.
127. Gilbert KJ, Whitlock MC. QST-FST comparisons with unbalanced half-sib
designs. Molecular Ecology Resources. 2015;15:262-7.
128. Karhunen M, Merilä J, Leinonen T, Cano JM, Ovaskainen O. driftsel: An R
package for detecting signals of natural selection in quantitative traits. Molecular
Ecology Resources. 2013;13:746-54.
129. Hadfield JD. MCMC methods for multi-response generalized linear mixed
models: the MCMCglmm R package. Journal of Statistical Software. 2010;33.
130. Semagn K, Babu R, Hearne S, Olsen M. Single nucleotide polymorphism
genotyping using Kompetitive Allele Specific PCR (KASP): Overview of the
technology and its application in crop improvement. Molecular Breeding. 2014;33:1-
14.
131. Weir BS, Hill WG. Estimating F-statistics. Annu Rev Genet. 2002;36:721-50.
132. Günther T, Coop G. A Short Manual for Bayenv2.0. 2016.
133. De Villemereuil P, Gaggiotti OE. A new FST-based method to uncover local
adaptation using environmental variables. Methods in Ecology and Evolution. 2015.

134.   Rincent R, Moreau L, Monod H, Kuhn E, Melchinger AE, Malvar RA, et al.
Recovering power in association mapping panels with variable levels of linkage
disequilibrium. Genetics. 2014;197:375-87.
135.   Bradburd GS, Ralph PL, Coop GM. Disentangling the effects of geographic
and ecological isolation on genetic differentiation. Evolution. 2013;67:3258-73.
136.   Lichstein JW. Multiple regression on distance matrices: A multivariate spatial
analysis tool. Plant Ecology. 2007;188:117-31.
137.   Goslee SC, Urban DL. Journal of Statistical Software The ecodist package for
dissimilarity-based analysis of ecological data. 2007.

A.

B.

C.