Title: Common gardens in teosintes reveal the establishment of a syndrome of adaptation to altitude

Short Title: Altitudinal syndrome in teosintes

M-A. Fustier[1, ☙], N.E. Martínez-Ainsworth[1, ☙], J.A. Aguirre-Liguori[2], A. Venon[1], H. Corti[1], A. Rousselet[1], F. Dumas[1], H. Dittberner[3], M.G. Camarena[4], D. Grimanelli[5], O. Ovaskainen[6,7], M. Falque[1], L. Moreau[1], J. de Meaux[2], S. Montes[4], L.E. Eguiarte[2], Y. Vigouroux[5], D. Manicacci[1,*], M.I. Tenaillon[1,*]

[1] : Génétique Quantitative et Evolution – Le Moulon, Institut National de la Recherche Agronomique, Université Paris-Sud, Centre National de la Recherche Scientifique, AgroParisTech, Université Paris-Saclay, France

[2]: Departamento de Ecología Evolutiva, Instituto de Ecología, Universidad Nacional Autónoma de México, Ciudad de México, Mexico

[3]: Institute of Botany, University of Cologne Biocenter, 47b Cologne, Germany

[4] : Campo Experimental Bajío, Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Celaya, Mexico

[5] : Université de Montpellier, Institut de Recherche pour le développement, UMR Diversité, Adaptation et Développement des plantes, Montpellier, France

[6] : Organismal and Evolutionary Biology Research Programme, PO Box, 65, FI-00014 University of Helsinki, Finland

[7]: Centre for Biodiversity Dynamics, Department of Biology, Norwegian University of Science and Technology, N-7491 Trondheim, Norway

☙: The authors equally contributed to this work

* co-corresponding authors

E-mail: domenica.manicacci@inra.fr, maud.tenaillon@inra.fr

Abstract

In plants, local adaptation across species range is frequent. Yet, much has to be discovered on its environmental drivers, the underlying functional traits and their molecular determinants. Genome scans are popular to uncover outlier loci potentially involved in the genetic architecture of local adaptation, however links between outliers and phenotypic variation are rarely addressed. Here we focused on adaptation of teosinte populations along two elevation gradients in Mexico that display continuous environmental changes at a short geographical scale. We used two common gardens, and phenotyped 18 traits in 1664 plants from 11 populations of annual teosintes. In parallel, we genotyped these plants for 38 microsatellite markers as well as for 171 outlier single nucleotide polymorphisms (SNPs) that displayed excess of allele differentiation between pairs of lowland and highland populations and/or correlation with environmental variables. Our results revealed that phenotypic differentiation at 10 out of the 18 traits was driven by local selection. Variation of these traits with elevation defines a syndrome of adaptation to altitude with plants that flower earlier, produce less tillers, display lower stomata density and carry larger, longer and heavier grains with increasing elevation. The proportion of outlier SNPs associating with phenotypic variation, however, largely depended on whether we considered a neutral structure with 5 genetic groups (73.7%) or 11 populations (13.5%), indicating that population stratification greatly affected our results. Finally, chromosomal inversions were enriched for both SNPs whose allele frequencies shifted along elevation as well as phenotypically-associated SNPs. Altogether, our results are consistent with the establishment of an altitudinal syndrome promoted by local selective forces in teosinte populations in spite of detectable gene flow. Because

elevation mimics climate change through space, SNPs that we found underlying

phenotypic variation at adaptive traits may be relevant for future maize breeding.

Keywords: spatially-varying selection; F_{st}-scan; association mapping; altitudinal

syndrome; pleiotropy; chromosomal inversions;

Author summary

Across their native range species encounter a diversity of habitats promoting local

adaptation of geographically distributed populations. While local adaptation is

widespread, much has yet to be discovered about the conditions of its emergence, the

targeted traits, their molecular determinants and the underlying ecological drivers.

Here we employed a reverse ecology approach, combining phenotypes and genotypes,

to mine the determinants of local adaptation of teosinte populations distributed along

two steep altitudinal gradients in Mexico. Evaluation of 11 populations in two

common gardens located at mid-elevation pointed to adaptation via an altitudinal

multivariate syndrome, in spite of gene flow. We scanned genomes to identify loci

with allele frequencies shifts along elevation, a subset of which associated to trait

variation. Because elevation mimics climate change through space, these

polymorphisms may be relevant for future maize breeding.

Introduction

Local adaptation is key for the preservation of ecologically useful genetic variation [1]. The conditions for its emergence and maintenance have been the focus of a long-standing debate nourished by ample theoretical work [2-9]. On the one hand, spatially-varying selection promotes the evolution of local adaptation, provided that there is genetic diversity underlying the variance of fitness-related traits [10]. On the other hand, opposing forces such as neutral genetic drift, temporal fluctuations of natural selection, recurrent introduction of maladaptive alleles via migration and homogenizing gene flow may hamper local adaptation (reviewed in [11]). Meta-analyzes indicate that local adaptation is pervasive, with evidence of native-site fitness advantage in reciprocal transplants detected in 45% to 71% of populations [12, 13].

While local adaptation is widespread, much has yet to be discovered about the traits submitted to spatially-varying selection, their molecular determinants and the underlying ecological drivers [14]. Local adaptation is predicted to increase with phenotypic, genotypic and environmental divergence among populations [6, 15, 16]. Comparisons of the quantitative genetic divergence of a trait (Q_{st}) with the neutral genetic differentiation (F_{st}) can provide hints on whether trait divergence is driven by spatially-divergent selection [17-20]. Striking examples of divergent selection include e.g. developmental rate in the common toad [21], drought and frost tolerance in alpine populations of the European silver fir [22], and traits related to plant phenology, size and floral display among populations of *Helianthus* species [23, 24].

The genes mediating local adaptation are usually revealed by genomic regions harboring population-specific signatures of selection. These signatures include alleles

displaying greater-than-expected differentiation among populations [25] and can be

identified through F_{st}-scans [26-32]. However, F_{st}-scans and its derivative methods

[25] suffer from a number of limitations, among them the high number of false

positives (reviewed in [33, 34]) and the lack of power to detect true positives [35].

Despite these caveats, F_{st}-outlier approaches have helped the discovery of emblematic

adaptive allele such as those segregating at the *EPAS1 locus* in Tibetan human

populations adapted to high altitude [36]. An alternative to detect locally adaptive loci

is to test for genotype-environment correlation [32, 37-42]. Correlation-based

methods can be more powerful than differentiation-based methods [43], but, spatial

autocorrelation of population structure and environmental variables can lead to

spurious signatures of selection [44].

Ultimately, to identify the outlier loci that have truly contributed to improve

local fitness, a link between outliers and phenotypic variation needs to be established.

The most common approach is to undertake genome wide association approach. In

plants, common garden studies, where environmental variation can be controlled,

facilitate detection of variants underlying phenotypic variation. For example,

replicated common gardens of *Arabidopsis* ecotypes have shown that fitness-

associated alleles display geographic and climatic patterns of association indicative of

selection [45]. Another study, demonstrated that the relative fitness of *Arabidopsis*

accessions in a given environment could actually be predicted from climate-

associated SNPs [46]. While it is clear that climatic selection over broad latitudinal

scales produces genomic and phenotypic patterns of local adaptation in the selfer

plant *Arabidopsis*, whether such patterns would persist in the face of gene flow, at

shorter spatial scale and/or in outcrossing species remains to be elucidated. Recent

literature in humans has questioned our ability to control for the actual variance

shown by population structure [47]. It is thus to be expected that detecting

polymorphisms directly responsible for trait variation will be challenging, when trait

variation and demographic history follow parallel environmental (geographic) clines.

We focused on a well-established outcrossing plant system, the teosintes, to

investigate the relationship of molecular, environmental, and fitness variation in

populations sampled across two elevation gradients in Mexico. Teosintes are the

closest wild relatives of maize and encompass two annual subspecies, *Zea mays* ssp.

parviglumis (hereon *parviglumis*) and *Zea mays* ssp. *mexicana* (hereon *mexicana*).

Both display large effective population sizes [48], and span a diversity of climatic

conditions, from warm and mesic conditions below 1800 m for *parviglumis*, to drier

and cooler conditions up to 3000 m for *mexicana* [49]. Previous studies have

discovered potential determinants of local adaptation in these systems. At a genome-

wide scale, decrease in genome size correlates with increasing altitude, which likely

results from the action of natural selection on life cycle duration [50, 51] . More

modest structural changes include megabase inversions that harbor clusters of SNPs

whose frequencies are associated with environmental variation [52, 53]. Also,

differentiation- and correlation-based genome scans in six teosinte populations

succeeded in finding outlier SNPs potentially involved in local adaptation [54, 55]. A

link with phenotypic variation has yet to be established.

In this paper, we used teosinte populations distributed along two altitudinal

gradients covering a relatively short yet climatically diverse, spatial scale. Both

gradients encompass populations from the two subspecies that are the closest wild

relatives of maize, *Zea mays* ssp. *parviglumis* and *Z. mays* ssp. *mexicana*. We

genotyped outlier SNPs on a sample of 28 teosinte populations, for which a set of

neutral SNPs was also available. Outlier SNPs were detected from whole genome

sequencing of six populations. In parallel, we performed common garden experiments

on a subset of 11 populations. We evaluated 18 phenotypic traits in two locations over

two consecutive years. Individuals from this association panel were also genotyped at

outlier SNPs, as well as at 38 microsatellite markers to enable a genetic association

after controlling for sample structure and kinship among individuals. We address

three main questions: What is the extent of phenotypic variation within and among

populations? Can we define a set of locally-selected traits that constitute a syndrome

of adaptation to altitude? What are the genetic bases of this syndrome? We further

discuss the challenges of detecting phenotypically-associated SNPs when trait and

genetic differentiation parallel environmental clines.

Results

Phenotypic variation within and among populations.

In order to investigate phenotypic variation, we set up two common garden

experiments located at mid-elevation in Mexico to evaluate individuals from 11

teosinte populations (Fig 1). These populations were sampled among 37 populations

(S1 Table) distributed along two altitudinal gradients from lowland populations of the

subspecies *parviglumis* to highland populations of the subspecies *mexicana* (S1 Fig).

The 11 populations encompassed 3 populations of *mexicana* and 8 of *parviglumis* and

were climatically contrasted as can be appreciated in the Principal Component

Analysis (PCA) computed on 19 environmental variables (S2 Fig). The corresponding

set of individuals from the 11 populations formed our association panel.

Figure 1. Geographical location of sampled populations and experimental fields.

The entire set of 37 Mexican populations is shown with *parviglumis* (circles) and

mexicana (triangles) populations sampled along gradient *a* (white) and gradient *b*

(black). The 11 populations indicated with a purple outline constituted our association

panel. It was evaluated in a four-block design during two years in two experimental

fields located at mid-elevation, SENGUA and CEBAJ. Two major cities (Mexico city

and Guadalajara) are also indicated.

We gathered phenotypic data during two consecutive years (2013 and 2014).

We targeted 18 phenotypic traits that included six traits related to plant architecture,

three traits related to leaves, three traits related to reproduction, five traits related to

grains, and one trait related to stomata (S2 Table). Each of the four experimental

assays (year-field combinations) encompassed four blocks. In each block, we

evaluated one offspring of ~15 mother plants (half-sibs) from each of the 11 teosinte

populations using a semi-randomized design. After filtering for missing data, our

association panel included 1664 teosinte individuals.

We found significant effects of Field, Year and/or their interaction for most

traits, and a highly significant effect of the population for all of them (model M1, S3

Table). Principal component analysis including all phenotypic measurements

highlighted that 21.26% of the phenotypic variation present both between *parviglumis*

and *mexicana* subspecies and within population scaled along PC1 (Fig 2A), a PC axis

that is strongly collinear with altitude (Fig 2B). Hence, while phenotypes of

populations largely overlapped along PC1, we observed a consistent tendency for

population phenotypic differentiation along altitude, irrespective of the gradient (Fig

2C). Traits that correlated the most to PC1 were related to grain characteristics,

tillering, flowering and to a lesser extent to stomata density (Fig 2B). PC2 correlated

with traits exhibiting a trend toward increase-with-elevation within *parviglumis*, but

decrease-with-elevation within *mexicana* (Fig 2D). Those traits were mainly related to

vegetative growth (Fig 2B). Together, both axes explained 37% of the phenotypic

variation.

Figure 2: Principal Component Analysis on phenotypic values corrected for the

experimental setting. Individuals factor map (A) and corresponding correlation

circle (B) on the first two principal components with altitude (Alt) added as a

supplementary variable (in blue). Individual phenotypic values on PC1 (C) and PC2

(D) are plotted with populations ranked by altitude and color-coded following A. For

populations from the two subspecies, *parviglumis* (circles) and *mexicana* (triangles),

color intensity indicates ascending elevation in green for *parviglumis* and red for

mexicana. Corrections for experimental setting are detailed in the material and

methods (Model M2).

We investigated in greater detail the influence of altitude on each trait

independently. All traits, except for the number of nodes with ears (NoE), exhibited a

significant effect of altitude (S3 Table, M3 model). Note that after accounting for

elevation, the population effect remained significant for all traits (data not shown),

suggesting that other factors than altitude also contributed to shape phenotypic

variation among populations. Traits related to flowering time and tillering displayed a

continuous decrease with elevation, and traits related to grain size increased with

elevation (Fig 3 & S3 Fig). Stomata density also diminished with altitude (S3 Fig). In

contrast, traits related to PC2 (e.g., plant height, height of the highest ear, number of

nodes with ear in the main tiller, Fig 2B&2D) displayed maximum values at

intermediate altitudes (highland *parviglumis* and lowland *mexicana*) (S3 Fig). We

estimated narrow-sense heritabilites (additive genotypic effect) per population for all

traits using a mixed 'animal' model. Average per-trait heritability ranged from 0.150

for tassel branching to 0.664 for female flowering time, albeit with large standard

errors (S2 Table). We obtained higher heritability for grain related traits when mother

plant measurements were included in the model with 0.631 ($sd = 0.246$), 0.511 ($sd =$

0.043) and 0.274 ($sd = 0.160$) for grain length, weight and width, respectively, as

compared to when they were not included, suggesting that heritability was under-

estimated for other traits.

Figure 3: Box-plots of adjusted means for four traits. Traits are female flowering

time (A), male flowering time (B), grain length (C) and stomata density (D).

Populations are ranked by altitude. *Parviglumis* populations are shown in green and

mexicana in red, lighter colors are used for gradient 'a' and darker colors for gradient

'b'. Units of measurement are detailed in S2 Table. In the case of flowering times, we

report data for 9 out of 11 populations because most individuals from the two lowland

populations (P1a and P1b) did not flower in our common gardens. Covariation with

elevation was significant for the four traits. Corrections for experimental setting are

detailed in the material and methods (Model M'1).

Identification of traits under spatially-varying selection.

We estimated the posterior mean (and 95% credibility interval) of genetic

differentiation (F_{st}) among the 11 populations of our association panel using

DRIFTSEL. Considering 1125 plants for which we had both individual phenotypes and

individual genotypes for 38 SSRs (S4 Table), we estimated the mean F_{st} to 0.22 (0.21-

0.23). Note that we found a similar estimate on a subset of 10 of these populations

with 1000 neutral SNPs (F_{st} (CI)=0.26 (0.25-0.27)). To identify traits whose variation

among populations was driven primarily by local selection, we employed a Bayesian

method that infers additive genetic values of traits from a model of population

divergence under drift [56]. Single-trait analyses revealed evidence for spatially-

varying selection at 12 traits, with high consistency between SSRs and neutral SNPs

(Table 1). Another method that contrasted genetic and phenotypic differentiation (Q_{st}-

F_{st}) uncovered a large overlap with nine out of the 12 traits significantly deviating

from the neutral model (Table 1) and one of the remaining ones displaying borderline

significance (Plant height=PL, S4 Fig). Together, these two methods collectively

indicated that phenotypic divergence among populations was driven by local selective

forces.

Table 1. Signals of selection (S) for each trait considering SSR markers (11 populations) or SNPs (10 populations).

Traits[a]	SSR[b]	SNP[b]
PL	0.995	0.972
HLE*	0.950	0.959
HHE	0.982	0.966
Til*	1.000	1.000
LBr*	1.000	0.990
NoE	0.682	0.699
LeL	0.888	0.875

LeW	0.999	0.996
LeC	0.633	0.583
FFT*	1.000	1.000
MFT*	1.000	1.000
TBr*	0.925	0.908
Gr	0.832	0.622
GrL*	1.000	1.000
GrWi*	0.995	0.984
GrWe*	1.000	0.999
GrC	0.717	0.689
StD*	0.999	0.999

[a]: Traits displaying signal of selection (spatially-varying traits, S>0.95) are indicated in bold, and marked by an asterisk when significant in Q_{st}-$F_{st}Comp$ analysis. We considered the underlined traits as spatially varying.

[b]: Values reported correspond to S from DRIFTSEL

Traits related to development, phenology and fitness are not independent. In

order to evaluate their correlations, we computed pairwise-correlations between traits

after correcting for experimental design and population structure (K=5) for all traits.

We found 82 (54%) significant correlations among 153 tested pairs of traits. The

following pairs of traits had the strongest positive correlations: male and female

flowering time, plant height and height of the highest ear, height of the highest and

lowest ear, grain length with grain weight and width (S11 Fig). The correlation

between flowering time (female or male) with grain weight and length were among

the strongest negative correlations (S11 Fig).

241  Altogether, evidence of spatially varying selection at 10 traits (Table 1) as

242 well as continuous variation of a subset of traits across populations in both elevation

243 gradients (Fig 3, S3 Fig) was consistent with a syndrome where populations produced

244 less tillers, flowered earlier, displayed lower stomata density and carried larger,

245 longer and heavier grains with increasing elevation.

247 **Structuring of neutral and adaptive genetic diversity.**

248  We analyzed the genetic structure of the association panel revealed by SSRs.

249 The highest likelihood was obtained at $K=2$ and $K=5$ clusters (S5 Fig). At $K=2$, the

250 clustering separated the lowland of gradient *a* from the rest of the populations. From

251 $K=3$ to $K=5$, a clear separation between the 8 *parviglumis* and the 3 *mexicana*

252 populations emerged. Increasing K values finally split the association panel into the

253 11 populations it encompassed (S6 Fig). At $K=5$, we observed a structuration

254 associated to both altitudes (lowland *parviglumis* versus highland *mexicana*) and

255 gradients *a* and *b* (Fig 4A & B). TreeMix analysis for a subset of 10 populations of

256 the association panel further confirmed those results with an early split separating the

257 lowlands from gradient *a* (cf. $K=2$, S6 Fig) as well as separation of the 3 *mexicana*

258 from the remaining populations (Fig 4C). The TreeMix analysis revealed gene flow

259 between distant lowland *parviglumis* populations from gradient *a* and *b*, as well as

260 between *parviglumis* and *mexicana* populations (Fig 4C). The structure analysis also

261 suggested admixture among some of the lowland populations, and to a lesser extent

262 between the two subspecies (Fig 4B).

Figure 4: Genetic clustering, historical splits and admixture among populations

of the association panel. Genetic clustering visualization based on 38 SSRs is shown

for *K=5* (A). Colors represent the *K* clusters. Individuals (vertical lines) are

partitioned into colored segments whose length represents the membership

proportions to the *K* clusters. Populations (named after the subspecies M: *mexicana,*

P: *parviglumis* and gradient 'a' or 'b') are ranked by altitude indicated in meters

above sea level. The corresponding geographic distribution of populations along with

their average membership probabilities are plotted (B). Historical splits and

admixtures between populations were inferred from SNP data for a subset of 10

populations of the association panel (C). We used TreeMix with 4 admixture events, a

model that explained 98.94% of the variation. Admixtures are colored according to

their weight.

We successfully genotyped 218 (~81%) out of 270 outlier SNPs on the

association panel, of which 141 were previously detected in candidate regions for

local adaptation [55]. Previously detected candidate regions were identified from re-

sequencing data of six teosinte populations (S1 Table) following an approach that

included high differentiation between highlands and lowlands, environmental

correlation, and in some cases their intersection with genomic regions involved in

quantitative trait variation in maize. The remaining ones (77) were discovered in the

present study by performing F_{st}-scans on the same re-sequencing data (S1 Table, S5

Table). We detected outlier SNPs that were both highly differentiated between

highland and lowland populations (gradients *a* or *b* or both), while displaying a

pattern of differentiation by altitude within at least one of the two subspecies. F_{st}-

scans pinpointed three genomic regions of particularly high differentiation (S7 Fig)

that corresponded to previously described inversions [52, 53]: one inversion on

chromosome 1 (*Inv1n*), one on chromosome 4 (*Inv4m*) and one on the far end of

chromosome 9 (*Inv9e*).

Associating genotypic variation at outlier SNPs to phenotypic variation.

We tested the association between phenotypes and 171 outlier SNPs

(MAF>5%) using our association panel. For each SNP-trait combination, the sample

size ranged from 264 to 1068, with a median of 1004 individuals (S6 Table). We used

SSRs to correct for both structure (at K=5) and kinship among individual genotypes.

This model (M6) resulted in a uniform distribution of P-values when testing the

association between genotypic variation at SSRs and phenotypic variation at traits (S8

Fig). Under this model, we found that 126 outlier SNPs (73.7%) associated to at least

one trait (Fig 5 and S9 Fig) at an FDR of 10%. The number of associated SNPs per

trait varied from 0 for leaf and grain coloration, to 55 SNPs for grain length, with an

average of 22.6 SNPs per trait (S5 Table). Ninety-three (73.8%) out of the 126

associated SNPs were common to at least two traits (S5 Table), and the remaining 33

SNPs were associated to a single trait (S5 Table). The ten traits displaying evidence of

spatially varying selection in the Q_{st}-F_{st} analyses displayed more associated SNPs per

trait (30.5 on average), than the non-spatially varying traits (12.75 on average).

A growing body of literature stresses that incomplete control of population

stratification may lead to spurious associations [57]. Hence, highly differentiated

traits along environmental gradients are expected to co-vary with any variant whose

allele frequency is differentiated along the same gradients, without underlying causal

link. We therefore expected false positives in our setting where both phenotypic traits

(Fig 2C) and outlier SNPs varied with altitude. We found a slightly significant

correlation (r=0.5, P-value=0.03) between the strength of the population effect for

each trait – a measure of trait differentiation (S3 Table) – and its number of associated

SNPs (S5 Table).

To verify that additional layers of structuring among populations did not cause

an excess of associations, we repeated the association analyzes including the

structuring in 11 populations as a covariate (a proxy of the structuring revealed at

K=11, S6 Fig). We retrieved much less SNPs compared to at K=5 (S5 Table). Here

only 23 SNPs associated to at least one trait, indicating that significant genetic

associations are contingent on the population structure. Eight traits showed no

association to any SNP (HLE, LBr, NoE, TBr, Gr, GrWi, GrC and StD), and the

remaining traits varied from a single associated SNP (LeL and Til) to 8 associated

SNPs for grain weight (S5 Table). For instance, traits such as female or male

flowering time that displayed 45 and 43 associated SNPs at K=5, now displayed only

4 and 3 associated SNPs, respectively (Fig 5). Traits under spatially varying selection

still associated to more SNPs (2.00 on average) than those with no spatially varying

selection (1.25 SNPs on average). Note that one trait (LeC) was associated to 4 SNPs

with structuring at 11 populations while not associated to any SNP at K=5. Among

the 126 SNPs associated to at least one trait at K=5, only 22 were recovered

considering 11 populations. And an additional SNP was detected with structuring at

11 populations while absent at K=5. One striking feature of the structuring at 11

populations was that inversions of chromosomes 1, 4, and 9, taken together, were

enriched for phenotypically-associated SNPs considering the structuring at 11

populations (χ^2 = 8.95, P-value=0.0028). We recovered a borderline significant

enrichment with the correction K=5 (χ^2 = 3.82, P-value=0.051).

Figure 5: Manhattan plots of associations between 171 outlier SNPs and 6

phenotypic traits along with their Bayenv2 outlier status for PCenv1. X-axis

indicates the positions of outlier SNPs on chromosomes 1 to 10, black and gray colors

alternating per chromosome. Significant associations (10% FDR) are indicated

considering either a structure matrix at K=5 (pink dots), for 11 populations (blue dots)

or for both K=5 and 11 populations models (purple dots). Empty dots designate SNPs

displaying significant association to PCenv1.

Altogether the 23 SNPs recovered considering a neutral genetic structure with

11 populations corresponded to 30 associations, 7 of the SNPs being associated to

more than one trait (S5 Table). For all these 30 associations but two cases (FFT with

SNP_7, and MFT with SNP_28), the SNP effect did not vary among populations

(non-significant SNP by population interaction in model M6 where we included the

SNP interactions with year*field and population). For a subset of 4 SNPs, we

illustrated the regression between the trait value and the shift of allele frequencies

with altitude (Fig 6). We estimated corresponding additive and dominance effects (S7

Table). In some cases, the intra-population effect corroborated the inter-population

variation with relatively large additive effects of the same sign (Fig 6 C & D, S7

table). Note that in both cases, one or the other allele was dominant (S7 Table). In

other cases, the results were more difficult to interpret at the inter-population level

with either no significant effect (Fig 6C) or correspondingly negligible additive effect

(Fig 6A, S7 Table) but extremely strong dominance (S7 Table).

In order to ask whether multiple SNPs contributed independently to the

phenotypic variation of a single trait, we tested a multiple SNP model where SNPs

were added incrementally when significantly associated (FDR < 0.10). We found 2, 3

and 2 SNPs for female, male flowering time and height of the highest ear,

respectively (S5 Table). Except for the latter trait, the SNPs were located on different

chromosomes.

Figure 6: Regression at the population level of phenotypic average value

corrected for the experimental design (calculated as the residual of model M2)

and allelic frequency. pval stands for the P-value of the linear regression represented

in blue.

Finally, we computed the pairwise linkage disequilibrium (LD) as measured

by r^2 between the 171 outlier SNPs using the R package LDcorSV [58] while

correcting for the underlying structure and kinship estimated from SSRs [59]. The 171

outlier SNPs were distributed along the 10 chromosomes of maize, and exhibited low

level of linkage disequilibrium (LD), except for SNPs located on chromosomes eight,

nine, and a cluster of SNPs located on chromosome 4 (S10 Fig). Among the 171, the

subset of 23 phenotypically-associated SNPs displayed an excess of elevated LD

values – out of 47 pairs of SNPs phenotypically-associated to the same trait, 16 pairs

were contained in the <5% higher values of the LD distribution of all outlier SNP

pairs. Twelve out of the 16 pairs concerned grain weight, and the remaining four

concerned leaf coloration, and one pair of SNPs was associated to both traits.

Environmental variables underlying local adaptation.

In order to analyze the environmental patterns correlating to the 218 outlier

SNPs frequencies, we genotyped an extended sample of 28 populations distributed

along the two elevation gradients. We assessed correlation between outlier SNPs and

19 climatic variables. Because climatic variables are not independent, we summarized

environmental information from the sampled populations by two principal

components (PCenv1 and PCenv2). The first component, that explained 56% of the

variation, correlated with altitude but display no correlation to either latitude or

longitude. PCenv1 was defined both by temperature- and precipitation- related

variables (S2 B Fig) including Minimum Temperature of Coldest Month (T6), Mean

Temperature of Driest and Coldest Quarter (T9 and T11), Precipitation of Driest

Month and Quarter (P14 and P17). The second PC explained 20.5% of the variation

and was mainly defined (S2 B Fig) by Isothermality (T3), Temperature Seasonality

(T4) and Temperature Annual Range (T7).

We first employed multiple regression to test whether the pairwise F_{st} matrix

across 28 populations correlated to the environmental (distance along PCenv1) and/or

the geographical distance. Overall, we found a significantly greater proportion of

environmentally-correlated SNPs among outliers compared with neutral SNPs (χ^2

=264.07, P-value=2.2 10^{-16}), a pattern not seen with geographically-correlated SNPs.

Roughly 53% (115) of the 218 outlier SNPs associated with at least one of the two

first PCenvs, with 98 and 33 associated with PCenv1 and PCenv2, respectively. As

expected, a handful of SNPs were associated to both PCenv1 and PCenv2.

Among those SNPs that were tested for phenotypic association (171) at K=5,

we found 75 (44%) that were associated to the first two environmental components,

either exclusively to PCenv1 (47) or PCenv2 (14), or both (14) (S5 Table). There was

no significant overlap between phenotypically-associated SNPs and PCenv1. Finally,

just like for phenotypically-associated SNPs, we found enrichment of

environmentally-associated SNPs within inversions both for PCenv1 (χ^2 = 16.41, P-

value=5.1 10^{-5}) and PCenv2 (χ^2 = 30.67, P-value=53.05 10^{-8}).

Discussion

Local adaptation of plant populations sampled across vast geographical ranges
has been well documented [13]. Traits underlying local adaptation, i.e. those that
evolve under spatially-varying selection, are classically identified from common
garden experiments which allow to control for environmental effects [60].
Identification of variants underlying these phenotypic changes is however
challenging. While population genomics has brought a flurry of tools to detect
footprints of local adaptation across populations, their reliability remains questioned
[57]. In addition, direct genetic links between population outliers and locally adaptive
phenotypic variation are often missing. In practice, local adaptation and demographic
history often follow the same geographic route. In such context, disentangling the
intertwining of trait, molecular, and environmental variation, which is paramount for
the study of local adaptation, is particularly arduous.

Here we investigated those links on a well-established outcrossing system, the
closest wild relatives of maize, along altitudinal gradients that display considerable
environmental shifts over short geographical scales. We used populations that range
from 504 to 2176 m in altitude over ~460 kms for gradient *a*, and from 342 to 2581m
in altitude over ~350 kms for gradient *b* (S1 Fig). We genotyped 28 populations at
218 outlier SNPs detected with population genomic tools, and determined population
values for 19 climatic variables (S1 Table). In parallel, we conducted an association
analysis using individuals derived from seeds collected from 11 of those populations.
We phenotyped individuals in two common gardens for two years, and genotyped
them at neutral markers and outlier SNPs (Fig 1).

Syndrome of adaptation to altitude.

Common garden studies along elevation gradients have been conducted in

European and North American plants species [61]. Because populations are evaluated

in the same environmental conditions, common gardens help uncover genetic clines.

Together with other studies, they have revealed that adaptive responses to altitude are

multifarious s[62]. They include physiological responses such as high photosynthetic

rates [63], tolerance to frost [64], biosynthesis of UV-induced phenolic components

[65]; morphological responses with reduced stature [66, 67], modification of leaf

surface [68], increase in leaf non-glandular trichomes [69], modification of stomata

density; and phenological responses with variation in flowering time [70], and

reduced growth period [71].

Our multivariate analysis of teosinte phenotypic variation revealed a marked

differentiation between teosinte subspecies along an axis of variation (21.26% of the

total variation) that also discriminated populations by altitude (Fig 2A & B). The

combined effects of assortative mating and environmental elevation variation may

generate, in certain conditions, trait differentiation along gradients without underlying

divergent selection [72]. While we did not measure flowering time differences among

our populations *in situ*, we did find evidence for long distance gene flow between

gradients and subspecies (Fig 4 A & C). In addition, several lines of arguments

suggest that the observed clinal patterns result from selection at independent traits and

is not solely driven by differences in flowering time among populations. First, two

distinct methods accounting for shared population history, concur with signals of

spatially-varying selection at ten out of the 18 traits (Table 1). Nine of them exhibited

a clinal trend of increase/decrease of population phenotypic values with elevation (S3

Fig) within at least one of the two subspecies. This number is actually conservative,

because these approaches disregard the impact of selective constraints which in fact

tend to decrease inter-population differences in phenotypes. Second, while male and

female flowering times were positively correlated, they displayed only subtle

correlations ($|r|<0.16$) with other spatially-varying traits except for grain weight and

length ($|r| <0.33$). Third, we observed convergence at multiple phenotypes between

the lowland populations from the two gradients that occurred despite their

geographical and genetical distance (Fig 4) again arguing that local adaptation drives

the underlying patterns.

Because seeds were directly collected from wild populations, a limitation of

our experimental setting is the confusion between genetic and environmental maternal

effects. Such control would necessitate seed production by selfing of each plant, but

was avoided since inbreeding depression is very strong in *Zea mays* [73]. However,

we are confident that our observed clinal trends in phenology and morphology reflect

adaptive genetic differentiation rather than environmental maternal effects. We

recovered indeed associations with genetic polymorphisms (Fig 5, S9 Fig). Moreover,

for two out of the three traits for which we had access to mother plant measures, grain

length and weight, we ruled out a strong mother plant effect since we estimated high

heritabilites ($h^2> 0.5$, S2 Table). In addition, a greenhouse experiment reported similar

trends for 17 accessions of annual teosintes including both *parviglumis* and *mexicana*

[74]. Consistently with our findings, the aforementioned study revealed reduced tiller

number, as well as increase in grain weight [74].

Spatially-varying traits that displayed altitudinal trends, collectively defined a

teosinte altitudinal syndrome of adaptation characterized by early-flowering,

production of few tillers albeit numerous lateral branches, production of heavy, long

and large grains, and decrease in stomata density. We also observed increased leaf

pigmentation with elevation, although with a less significant signal (S3 Table),

consistent with the pronounced difference in sheath color reported between

parviglumis and *mexicana* [75, 76]. The trend towards depleted stomata density at

high altitudes (S3 Fig) could arguably represent a physiological adaptation as stomata

influence components of plant fitness through their control of transpiration and

photosynthetic rate [77]. Indeed, in natural accessions of *A. thaliana*, stomatal traits

showed signatures of local adaptation and were associated with both climatic

conditions and water-use efficiency [78]. Furthermore, previous work has shown that

in arid and hot highland environments, densely-packed stomata may promote

increased leaf cooling in response to desiccation [79] and may also counteract limited

photosynthetic rate with decreasing pCO_2 [80]. Accordingly, increased stomata

density at high elevation sites has been reported in alpine species such as the

European beech [81] as well as in populations of *Mimulus guttatus* subjected to higher

precipitations in the Sierra Nevada [82]. In our case, higher elevations display both

arid environment and *cooler* temperatures during the growing season, features

perhaps more comparable to other tropical mountains for which a diversity of patterns

in stomatal density variation with altitude has been reported [83]. Further work will

be needed to decipher the mechanisms driving the pattern of declining stomata

density with altitude in teosintes. Altogether, the altitudinal syndrome was consistent

with natural selection for rapid life-cycle shift, with early-flowering in the shorter

growing season of the highlands and production of larger propagules than in the

lowlands. This altitudinal syndrome evolved in spite of detectable gene flow.

Although we did not formally measure biomass production, the lower number

of tillers and higher amount and size of grains in the highlands when compared with

the lowlands may reflect trade-offs between allocation to grain production and

vegetative growth [84]. Because grains fell at maturity and a single teosinte individual

produces hundreds of ears, we were unable to provide a proxy for total grain

production. The existence of fitness-related trade-offs therefore still needs to be

formally addressed.

Beyond trade-offs, our results more generally question the extent of

correlations between traits. In maize, for instance, we know that female and male

flowering time are positively correlated and that their genetic control is in part

determined by a common set of genes [85]. They themselves correlate with yield-

related traits [86]. Response to selection for late-flowering also triggers a correlated

increase in leaf number in cultivated maize [87], and common genetic loci have been

shown to determine both traits [88]. Here we found strong positive correlations

between male and female flowering time, grain length and width, plant height and

height of the lowest or highest ear. Strong negative correlations concerned grain

weight with both male and female flowering time. Together these correlations were

therefore partly consistent with previous observations in maize, suggesting that they

were inherited from wild ancestors.

531   Footprints of past adaptation are relevant to detect variants involved in present

532   phenotypic variation.

The traits measured co-varied with altitude (Fig 3, S3 Fig, Table S3) and

therefore with environmental differences along gradients. Statistical associations

between genotypes and phenotypes therefore largely depended on the correction for

underlying population stratification. We indeed found that 73.7% SNPs associated

with phenotypic variation at K=5, but only 13.5% of them did when considering a

genetic structure with 11 populations. Except for one, the latter set of SNPs

represented a subset of the former. The level of differentiation between the two

subspecies in our outcrossing system was relatively low ($F_{st} \approx 22\%$). As a comparison,

in the selfer *Arabidopsis thaliana*, differentiation among populations from the same

geographical area varies between 29% and 80% [89], and association panels typically

display maximum values of F_{st} around 60% within 10kb-windows genome-wide [90].

Subspecies differentiation was fully accounted at K=5 as shown by the clear

distinction between *mexicana* populations and the rest of the samples (Fig 4A). It is

therefore the residual stratification among populations within genetic groups that

caused the inflation of significant associations at K=5.

While false positives likely represent a substantial proportion of associations

at K=5, some hints indicate that overcorrection with 11 populations may also be at

play. For instance, at K=5 we detected *Sugary1* (*Su1*), a starch debranching enzyme

that was selected during domestication and subsequent breeding [91, 92]. We found

that *Su1* was associated with variation at six traits (male and female flowering time,

tassel branching, height of the highest ear, grain weight and stomata density) pointing

to high pleiotropy. A previous study reported association of this gene to oil content in

teosintes [93]. In maize, this gene is involved in differences of kernel phenotypes

between maize genetic groups [94] and has been a target during domestication and

breeding [91, 95]. However, *Su1* was not recovered with 11 populations indicating

that divergent selection acted among populations.

The proportion of genic SNPs associated to phenotypic variation was not

significantly higher than that of non-genic SNPs (i.e, SNPs >1kb from a gene) ($\chi^2_{(df=1)}$

= 0.043, P-value = 0.84 at K=5 and $\chi^2_{(df=1)}$ =1.623 , P-value =0.020 with 11

populations) stressing the importance of considering both types of variants [96]. For

instance, we discovered a non-genic SNP (SNP_149) that displayed a strong

association with leaf width variation as well as a pattern of allele frequency shift with

altitude among populations (Fig 6D).

Several genes that were recovered with both K=5 and 11 populations deserve

specific attention (S5 Table). One SNP within the *phytochrome B2* (SNP_210; *phyB2*)

gene was associated with female and male flowering time, as well as with plant height

and grain length (at K=5 only for the two latter traits). Phytochromes are involved in

perceiving light signals and are essential for growth and development in plants. The

maize gene *phyB2* regulates the photoperiod-dependent floral transition, with mutants

producing early flowering phenotypes and reduced plant height [97]. Genes from the

phosphatidylethanolamine-binding proteins (PEBPs) family – *Zea mays*

CENTRORADIALIS (*ZCN*) family in maize – are also well-known to act as promotor

and repressor of the floral transition in plants [98]. Z*CN8* is the main floral activator

of maize [99], and both *ZCN8* and *ZCN5* strongly associates with flowering time

variation [94, 100]. Consistently, we found associations of male and female flowering

time with *PEBP18* (SNP_210). It is interesting to note that SNPs at these two

flowering time genes, *phyB2* and *PEBP18*, influenced independently as well as in

combination both female and male flowering time variation (S5 Table).

Considering a structure with 5 genetic groups, the number of SNPs associated

per trait varied from 1 to 55, with no association for leaf and grain coloration (S5

Table). Our results therefore point to a potentially high proportion of associated SNPs.

Because we found limited LD among our SNPs (S10 Fig) corroborating previous

reports (LD decay within <100bp, [55][101]), our observations are consistent with

phenotypic variation driven by numerous independent loci with small effects (S7

Table). Consistently, Weber et al. [84] found that individual SNPs account for small

proportions of the phenotypic variance. Note that while we did not test for epistasis –

the effect of marker combinations as opposed to single marker analysis – genome-

wide association mapping studies (GWAS) in maize all concur to phenotypic

variation being dominated by additive effects [80,86]. We performed tests were

multiple associated SNPs for a given trait were added incrementally. Among all traits,

we evidenced multiple SNPs determination for three traits (male and female flowering

time and height of the highest ear, Table S5).

Environmental variables driving local adaptation.

597   A substantial proportion of our set of outlier SNPs was chosen due to their

significant correlation among six populations between variation of allele frequencies

and the projection on the first environmental principal component [55]. We showed

here that these correlations hold on our broad sample of 28 populations, with allele

frequency at 115 SNPs correlating with environmental variation. In contrast with

neutral SNPs, outlier SNPs displayed a greater isolation-by-environment than

isolation-by-distance, indicating that patterns of allele frequency differentiation

among populations was driven by adaptive processes. As expected, the principal

component driven by altitude (PCenv1) correlated to allele frequency for a greater

fraction of SNPs than the second orthogonal component. SNPs associated to PCenv1

were not enriched for SNPs associated with phenotypic variation. SNPs associated

with environmental variation but not phenotypic variation may indicate either that

they were fixed − or almost fixed − within populations of our association panel

leading to low statistical power, or that the phenotype they influence was not

measured. Conversely, SNPs associated to phenotypic but not to environmental

variation may indicate either no environmental effect or the absence of the

environmental driver from our variables. For example, it has been shown that soil is

an important driver of local adaptation in teosintes [55,101], yet we did not quantify

plant-soil interaction phenotypic traits, and neither did we evaluate soil variables.

Finally, the subset of 61 outlier SNPs (S5 Table) that were associated to both

environmental and phenotypic variation point to situations where the SNP still

segregates within populations and relates to measured phenotypes that are themselves

associated to environmental driver(s) (PCenv1 and/or PCenv2). These genomic

locations are likely of particular interest for ongoing adaptation.

Interestingly, we identified three Mb-scale inversions located on chromosomes

1 (*Inv1n*), 4 (*Inv4m*) and 9 (*Inv9e*) that, taken together, were enriched for SNPs

associated with environmental variables related to altitude and/or phenotypic variation

at multiple traits. Previous work [52, 53] has shown that *Inv1n* and *Inv4m* segregate

within both *parviglumis* and *mexicana*, while two inversions on chromosome 9, *Inv9d*

and *Inv9e*, are present only in some of the highest *mexicana* populations; such that all

four inversions also follow an altitudinal pattern. Our findings confirmed that three of

these inversions possessed an excess of SNPs with high F_{st} between subspecies and

between low- and high-*mexicana* populations for *Inv9e* [54]. Noteworthy *Inv9d*

contains a large ear leaf width quantitative trait locus in maize [96]. Corroborating

these results, we found consistent association between the only SNP located within

this inversion and leaf width variation in teosinte populations (S5 Table). Overall, our

results further strengthen the role of chromosomal inversions in teosinte altitudinal

adaptation. Because they suppress recombination between inverted and non-inverted

genotypes, their spread has likely contributed to the emergence and maintenance of

locally adaptive allelic combinations in the face of gene flow, not unlike a growing

number of examples from other models (reviewed in [102]) including insects [103],

fish [104], birds [105] and plants [106, 107].

Conclusion.

Elevation gradients provide an exceptional opportunity for investigating

variation of functional traits in response to continuous environmental factors at short

geographical scales. Here we documented patterns indicating that local adaptation,

likely facilitated by the existence of chromosomal inversions, allows teosintes to cope

with specific environmental conditions in spite of gene flow. We detected an

altitudinal syndrome in teosintes composed of sets of independent traits evolving

under spatially-varying selection. Because traits co-varied with environmental

differences along gradients, however, statistical associations between genotypes and

phenotypes largely depended on control of population stratification. Yet, several of

the variants we uncovered seem to underlie adaptive trait variation in teosintes.

Adaptive teosinte trait variation is likely relevant for maize evolution and breeding.

Whether the underlying SNPs detected in teosintes bear similar effects in maize or

whether their effects differ in domesticated backgrounds will have to be further

investigated.

Material and Methods

Description of teosinte populations and sampling.

We used 37 teosinte populations of *mexicana* (16) and *parviglumis* (21)

subspecies from two previous collections [51, 54, 55] to design our sampling. These

populations (S1 Table) are distributed along two altitudinal gradients (Fig 1). We

plotted their altitudinal profiles using R 'raster' package [108] (S1 Fig). We further

obtained 19 environmental variable layers from

http://idrisi.uaemex.mx/distribucion/superficies-climaticas-para-mexico. These high-

resolution layers comprised monthly values from 1910 to 2009 estimated via

interpolation methods [102]. We extracted values of the 19 climatic variables for each

population (S1 Table). Note that high throughput sequencing (HTS) data were

obtained in a previous study for six populations out of the 37 (M6a, P1a, M7b, P2b,

M1b and P8b; Fig 1, S1 Table) to detect candidate genomic regions for local

adaptation [55]. The four highest and lowest of these populations were included in the

association panel described below.

We defined an association panel of 11 populations on which to perform a

genotype-phenotype association study (S1 Table). Our choice was guided by grain

availability as well as the coverage of the whole climatic and altitudinal ranges.

Hence, we computed Principal Component Analyses (PCA) for each gradient from

environmental variables using the FactoMineR package in R [109] and added altitude

to the PCA graphs as a supplementary variable. Our association panel comprised five

populations from a first gradient (*a*) – two *mexicana* and three *parviglumis,* and six

populations from a second gradient (*b*) – one *mexicana* and five *parviglumis* (Fig 1).

Finally, we extracted available SNP genotypes generated with the

MaizeSNP50 Genotyping BeadChip for 28 populations out of our 37 populations [54]

(S1 Table). From this available SNP dataset, we randomly sampled 1000 SNPs found

to display no selection footprint [54], hereafter neutral SNPs. We used this panel of

28 populations to investigate correlation with environmental variation. Note that 10

out of the 28 populations were common to our association panel, and genotypes were

available for 24 to 34 individuals per population, albeit different from the ones of our

association mapping panel.

Common garden experiments

We used two common gardens for phenotypic evaluation of the association

panel (11 populations). Common gardens were located at INIFAP (Instituto Nacional

de Investigaciones Forestales, Agricolas y Pecuaria) experimental field stations in the

state of Guanajuato in Mexico, one in Celaya municipality at the Campo

Experimental Bajío (CEBAJ) (20°31'20'' N, 100°48'44''W) at 1750 meters of

elevation, and one in San Luis de la Paz municipality at the Sitio Experimental Norte

de Guanajuato (SENGUA) (21°17'55''N, 100°30'59''W) at 2017 meters of elevation.

These locations were selected because they present intermediate altitudes (S1 Fig) and

environmental conditions where both lowland and highland populations could grow.

The two common gardens were replicated in 2013 and 2014.

The original sampling contained 15 to 22 mother plants per population. Eight

to 12 grains per mother plant were sown each year in individual pots. After one

month, seedlings were transplanted in the field. Each of the four fields (2 locations, 2

years) was separated into four blocks encompassing 10 rows and 20 columns. We

evaluated one offspring of ~15 mother plants from each of the 11 teosinte populations

in each block, using a semi-randomized design, i.e. each row containing one or two

individuals from each population, and individuals being randomized within row,

leading to a total of 2,640 individual teosinte plants evaluated.

SSR genotyping and genetic structuring analyses on the association panel

In order to quantify the population structure and individual kinship in our

association panel, we genotyped 46 SSRs (S4 Table). Primers sequences are available

from the maize database project [110] and genotyping protocol were previously

published [111]. Genotyping was done at the GENTYANE platform (UMR INRA

1095, Clermont-Ferrand, France). Allele calling was performed on electropherograms

with the GeneMapper® Software Applied Biosystems®. Allele binning was carried

out using Autobin software [112], and further checked manually.

We employed STRUCTURE Bayesian classification software to compute a

genetic structure matrix on individual genotypes. Individuals with over 40% missing

data were excluded from analysis. For each number of clusters (K from 2 to 13), we

performed 10 independent runs of 500,000 iterations after a burn-in period of 50,000

iterations, and combined these 10 replicates using the LargeKGreedy algorithm from

the CLUMPP program [113]. We plotted the resulting clusters using DISTRUCT

software. We then used the Evanno method [114] to choose the optimal K value. We

followed the same methodology to compute a structure matrix from the outlier SNPs.

We inferred a kinship matrix from the same SSRs using SPAGeDI [115].

Kinship coefficients were calculated for each pair of individuals as correlation

between allelic states [116]. Since teosintes are outcrossers and expected to exhibit an

elevated level of heterozygosity, we estimated intra-individual kinship to fill in the

diagonal. We calculated ten kinship matrices, each excluding the SSRs from one out

of the 10 chromosomes. Microsatellite data are available

at: XX.XXXX/XX.figshare.XXXXXXX.

In order to gain insights into population history of divergence and admixture,

we used 1000 neutral SNPs (i.e. SNPs genotyped in [54] and that displayed patterns

consistent with neutrality among 49 teosinte populations) genotyped on 10 out of the

11 populations of the association panel to run a TreeMix analysis (TreeMix version

1.13 [117]). TreeMix models genetic drift to infer populations splits from an outgroup

as well as migration edges along a bifurcating tree. We oriented the SNPs using the

previously published MaizeSNP50 Genotyping BeadChip data from the outgroup

species *Tripsacum dactyloides* [52]. We tested 0 to 10 admixture events to build the

graph. We chose the graph and number of putative migration based on the

stabilization of the likelihood and the stabilization of the proportion of variance, i.e.

the values after which the likelihood and variance did not substantially increase.

Phenotypic trait measurements

We evaluated a total of 18 phenotypic traits on the association panel (S2

Table). We measured six traits related to plant architecture (PL: Plant Height, HLE:

Height of the Lowest Ear, HHE: Height of the Highest Ear, Til: number of Tillers,

LBr: number of Lateral Branches, NoE: number of Nodes with Ears), three traits

related to leave morphologies (LeL: Leaf Length, LeW: Leaf Width, LeC: Leaf

Color), three traits related to reproduction (MFT: Male Flowering Time, FFT: Female

Flowering Time, TBr : Tassel Branching), five traits related to grains (Gr: number of

Grains per ear, GrL: Grain Length, GrWi: Grain Width, GrWe: Grain Weight, GrC:

Grain Color), and one trait related to Stomata (StD: Stomata Density). These traits

were chosen because we suspected they could contribute to differences among

teosinte populations based on a previous report of morphological characterization on

112 teosinte collections grown in five localities [118].

We measured the traits related to plant architecture and leaves after silk

emergence. Grain traits were measured at maturity. Leaf and grain coloration were

evaluated on a qualitative scale. For stomata density, we sampled three leaves per

plant and conserved them in humid paper in plastic bags. Analyses were undertaken at

the Institute for Evolution and Biodiversity (University of Münster) as followed: 5mm

blade discs were cut out from the mid length of one of the leaves and microscopic

images were taken after excitation with a 488nm laser. Nine locations (0.15mm^2) per

disc were captured with 10 images per location along the z-axis (vertically along the

tissue). We automatically filtered images based on quality and estimated leaf stomata

density using custom image analysis algorithms implemented in Matlab. For each

sample, we calculated the median stomata density over the (up to) nine locations. To

verify detection accuracy, manual counts were undertaken for 54 random samples.

Automatic and manual counts were highly correlated ($R^2=0.82$), indicating reliable

detection (see S1 Annex StomataDetection, Dittberner and de Meaux, for a detailed

description). The filtered data set of phenotypic measurements is available

at: XX.XXXX/XX.figshare.XXXXXXX.

Statistical analyses of phenotypic variation

In order to test for genetic effects on teosinte phenotypic variation, we

decomposed phenotypic values of each trait considering a fixed population effect plus

a random mother-plant effect (model M1):

$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \delta_l + \chi_{il} + \psi_{jl} + P_{m/l} + \varepsilon_{ijklm}$ (M1)

where the response variable Y is the observed phenotypic value, μ is the total mean, α_i

is the fixed year effect (i = 2013, 2014), β_j the fixed field effect (j = field station,

SENGUA, CEBAJ), θ_{ij} is the year by field interaction, $\gamma_{k/ij}$ is the fixed block effect

(k = 1, 2, 3, 4) nested within the year-by-field combination, δ_l is the fixed effect of the

population of origin (l = 1 to 11), χ_{il} is the year by population interaction, ψ_{jl} is the

field by population interaction, $P_{m/l}$ is the random effect of mother plant (m = 1 to 15)

nested within population, and ε_{ijklm} is the individual residue. For the distribution of the

effects, the same variance was estimated within all populations. Mixed models were

run using v.3.0 [119] or MM4LMM v2.0.1

[https://rdrr.io/cran/MM4LMM/man/MM4LMM-package.html, update by F. Laporte]

R packages, which both gave very similar results, and fixed effects were tested

through Wald tests.

For each trait, we represented variation among populations using box-plots on

mean values per mother plant adjusted for the experimental design following model

M'1

$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + p_{m/l} + \varepsilon_{ijklm}$ (M'1)

where mother plant within population is considered as fixed. We used the function

predict to obtain least-square means (ls-means) of each mother plant, and looked at

the tendencies between population's values. All fixed models were computed using

lm package in R, and we visually checked the assumptions of residues independence

and normal distribution.

We performed a principal component analysis (PCA) on phenotypic values corrected

for the experimental design, using FactoMineR package in R [109] from the residues

of model M2 computed using the *lm* package in R:

$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + \varepsilon_{ijklm}$ (M2)

Finally, we tested for altitudinal effects on traits by considering the altitude of the

sampled population (l) as a covariate (ALT) and its interaction with year and field in

model M3.

$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{k/ij} + c{\cdot}ALT_l + a_i{\cdot}ALT_l + b_j{\cdot}ALT_l + P_{m/l} + \varepsilon_{ijklm}$ (M3)

where all the terms are equal to those in model M1 except that the fixed effect of the

population of origin was replaced by a regression on the population altitude (ALT$_l$).

Detection of selection acting on phenotypic traits

We aimed at detecting traits evolving under spatially varying selection by

comparing phenotypic to neutral genotypic differentiation. Q_{st} is a statistic analogous

to F_{st} but for quantitative traits, which can be described as the proportion of

phenotypic variation explained by differences among populations [19, 120].

Significant differences between Q_{st} and F_{st} can be interpreted as evidence for

spatially-varying ($Q_{st}{>}F_{st}$) selection [120]. We used the R package *QstFstComp* [121]

that is adequate for experimental designs with randomized half-sibs in outcrossing

species. We used individuals that were both genotyped and phenotyped on the

association panel to establish the distribution of the difference between statistics (Q_{st}-

F_{st}) under the neutral hypothesis of evolution by drift - using the half-sib dam

breeding design and 1000 resamples. We next compared it to the observed difference

with 95% threshold cutoff value in order to detect traits under spatially-varying

selection.

In addition to Q_{st}-F_{st} analyses, we employed the DRIFTSEL R PACKAGE [122] to

test for signal of selection of traits while accounting for drift-driven population

divergence and genetic relatedness among individuals (half-sib design). DRIFTSEL is a

Bayesian method that compares the probability distribution of predicted and observed

mean additive genetic values (S statistics, [56]). It is particularly powerful for small

datasets, and can distinguish between drift and selection even when Q_{st}-F_{st} are equal

[56]. We first applied RAFM to estimate the F_{st} value across populations, and the

population-by-population coancestry coefficient matrix. We next fitted both the RAFM

and DRIFTSEL models with 15,000 MCMC iterations, discarded the first 5,000

iterations as a transient, and thinned the remaining by 10 to provide 1000 samples

from the posterior distribution. Note that DRIFTSEL was slightly modified because we

had information only about the dams, but not the sires, of the phenotyped individuals.

We thus modified DRIFTSEL with the conservative assumption of all sires being

unrelated. Because DRIFTSEL does not require that the same individuals were both

genotyped and phenotyped, we used SSRs and phenotype data of the association

panel as well as the set of neutral SNPs and phenotype data on 10 out of the 11

populations. For the SNP analyses, we selected out of the 1000 neutral SNPs the 465

most informative SNPs based on the following criteria: frequency of the less common

variant at least 10%, and proportion of missing data at most 1%.

Finally, we estimated from DRIFTSEL the posterior probability of the ancestral

population mean for each trait as well as deviations of each population from these

values.

847   Both Q_{st}-F_{st} and DRIFTSEL rely on the assumption that the observed phenotypic

variation was determined by additive genotypic variation. We thus estimated narrow-

sense heritability for each trait in each population to estimate the proportion of

additive variance in performance. We calculated per population narrow-sense

heritabilites as the ratio of the estimated additive genetic variance over the total

phenotypic variance on our common garden measurements using the MCMCglmm R

package [123] where half sib family is the single random factor, and the design (block

nested within year and field) is corrected as fixed factor. For three grain-related traits,

we also ran the same model but including mother plants phenotypic values calculated

from the remaining grains not sown. We ran 100,000 iterations with 10,000 burn-in,

inverse gamma (0.001; 0.001) as priors. We then calculated the mean and standard

deviation of the 11 per population h² estimates.

Pairwise correlations between traits.

We evaluated pairwise-correlations between traits by correlating the residues

obtained from model M4, that corrects the experiment design (year, field and blocks)

as well as the underlying genetic structure estimated from SSRs:

$Y_{ijklm} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{ijk} + \sum_{n=1}^{4} b_{n\cdot} C^n_{ijklm} + \varepsilon_{ijklm}$ (M4)

where b_n is the slope associated to the structure covariate C^n. Membership values for

the C^n covariates were calculated at the individual level, i.e. for each offspring of

mother plant m from population l, grown in the year i field j and block k. C^n are thus

declared within $ijkl$ indices, although they are purely genetic covariates.

Genotyping of outlier SNPs on 28 populations

We extracted total DNA from each individual plant of the association panel as

well as 20 individuals from each of the 18 remaining populations that were not

included in the association panel (Table 1). Extractions were performed from 30 mg

of lyophilized adult leaf material following recommendations of DNeasy 96 Plant Kit

manufacturer (QIAGEN, Valencia, CA, USA). We genotyped outlier SNPs using

Kompetitive Allele Specific PCR technology (KASPar, LGC Group) [124]. Data for

outlier SNPs are available at: XX.XXXX/XX.figshare.XXXXXXX.

Among SNPs identified as potentially involved in local adaptation, 270 were

designed for KASPar assays, among which 218 delivered accurate quality data. Of the

218 SNPs, 141 were detected as outliers in two previous studies using a combination

of statistical methods – including F_{st}-scans [125], Bayescan [29] and Bayenv2 [32,

126], Bayescenv [127] – applied to either six of our teosinte populations [55] or to a

broader set of 49 populations genotyped by the Illumina® MaizeSNP50 BeadChip

[54]. The remaining outlier SNPs (77) were detected by F_{st}-scans from six

populations (S7 Fig, S5 Table), following a simplified version of the rationale in [55]

by considering only differentiation statistics: SNPs were selected if they displayed

both a high differentiation (5% highest F_{st} values) between highland and lowland

populations in at least one of the two gradients, and a high differentiation (5% highest

F_{st} values) between highland and lowland populations either within *parviglumis* (P2b

and P8b) or within *mexicana* (M7b and M1b) or both in gradient *b* (S1 Fig). We

thereby avoided SNPs fixed between the two subspecies.

Association mapping

We tested the association of phenotypic measurements with outlier SNPs on a

subset of individuals for which (1) phenotypic measurements were available, (2) at

least 60% of outlier SNPs were adequately genotyped, and (3) kinship and cluster

membership values were available from SSR genotyping. For association, we

removed SNPs with minor allele frequency lower than 5%.

In order to detect statistical associations between outlier SNPs and phenotypic

variation, we used the following mixed model derived from [122]:

$Y_{ijklom} = \mu + \alpha_i + \beta_j + \theta_{ij} + \gamma_{ijk} + \sum_{n=1}^{4} b_n \cdot C^n_{ijkolm} + \zeta_o + u_{ijklm} + \varepsilon_{ijklom}$ (M6)

where ζ is the fixed bi-allelic SNP factor effect with one level for each of the three

genotypes (o=0, 1, 2; with o=1 for heterozygous individuals,), u_{ijklom} being the

random genetic effect of each individual. We assumed that the u_{ijklom} vector followed

a N(0,K σ^2u) distribution, where K is the inversed kinship matrix.

In order to avoid overcorrection of neutral genetic structure and improve

power, we ran the model independently for each chromosome using a kinship matrix

estimated from all SSRs except those contained in the chromosome tested [128]. We

tested SNP effects though the Wald statistics, and applied 10% False Discovery Rate

(FDR) thresholds for each phenotype separately. In order to validate the correction for

genetic structure, the 38 multiallelic SSR genotypes were transformed to biallelic

genotypes, filtered for MAF > 5%, and used to run associations with the complete M6

model as well as the M6 model excluding either kinship or both structure and kinship.

We generated QQplots of p-values for each of these three models and the 18 traits.

Multiple SNP models were built by adding one SNP at a time as long as its

FDR was lower than 0.10. We controlled for population structure at K=11 and using

the kinship matrix that excluded the SSR on the same chromosome as the last tested

SNP.

Finally, we tested for correlation between the excess of common associated

SNPs between two traits, and the absolute value of their pairwise phenotypic

correlation. We computed both parametric (Pearson coefficient) and non-parametric

(Kendall coefficient) correlation (Fig 6B). We calculated the excess of common

associated SNPs as the difference between observation and expectation – under the

hypothesis that the proportion of associated SNPs is independent between traits.

Environmental correlation of outlier SNPs

We tested associations between allelic frequency at 218 outlier SNPs and

environmental variables across 28 populations, using Bayenv 2.0 [37,106]. Because

environmental variables are highly correlated, we used the first two principal

component axes from the environmental PCA analysis (PCenv1 and PCenv2) to run

Bayenv 2.0. This software requires a neutral covariance matrix, that we computed

from the available dataset of 1000 neutral SNPs (S1 Table). We performed 100,000

iterations, saving the matrix every 500 iterations. We then tested the correlation of

these to the last matrix obtained, as well as to an *Fst* matrix calculated with

BEDASSLE [129], as described in [54].

For each outlier SNP, we compared the posterior probability of a model that

included an environmental factor (PCenv1 or PCenv2) to a null model. We

determined a 5% threshold for significance of environmental association by running

100,000 iterations on neutral SNPs. We carried out five independent runs for each

outlier SNP and evaluated their consistency from the coefficient of variation of the

Bayes factors calculated among runs.

In order to test whether environmental distance was a better predictor of allele

frequencies at candidate SNPs than geography, we used multiple regression on

distance matrices (MRM, [130]) implemented in the ecodist R package [131] for each

outlier SNP. We used pairwise F_{st} values as the response distance matrix and the

geographic and environmental distance matrices as explanatory matrices. We

evaluated the significance of regression coefficients by 1000 permutations and

iterations of the MRM. We determined the total number of environmentally and

geographically associated SNPs (P-value<0.05) among outliers. We employed the

same methodology for our set of 1000 neutral SNPs.

Ethics Statement

All the field work has been done in Mexico in collaboration with Instituto Nacional de Investigaciones Forestales, Agrícolas y Pecuarias, Celaya in Celaya.

Acknowledgments

We thank Jessica Melique for her help in gathering stomata data. We are very grateful to Angelica Cibrian at Langebio (Cinvestav, Irapuato, Mexico) who let us use her lab to lyophilize our samples, and store them. Valeria Souza greatly helped us with logistic support in Mexico. Insights from Delphine Legrand, Pierre de Villemereuil and Elodie Marchadier were important for analyzing correlations between traits and to estimate heritabilities.

Author contributions

Conceived and designed the experiments: MIT, DM. Contributed plant material: LEE, JAA-L, SM, MIT. DNA extractions: HC, AR, M-AF, DM. Conducted field experiments and performed phenotypic measurements: M-AF, MIT, MGC, SM. Outlier SNP detection: M-AF. Microsatellite genotyping: AV, HC, FD, MF. Neutral SNP genotyping: LEE, JAA-L. Stomata density images and pipeline: JdD, HD. Statistical methodology: DM, M-AF, NEM-A, OO, LM, MIT, JAA-L. Implemented methods, wrote scripts: M-AF, NEM-A, OO, JAA-L. Analyzed the data: M-AF, NEM-A, DM. Prepared Tables and Figures: M-AF, NEM-A. Discussed interpretations: M-AF, NEM-A, DG, OO, LM, JAA-L, JdM, YV, DM, MIT. Wrote the paper: NEM-A, M-AF, MIT. Corrected and commented the manuscript: all authors.

Funding

This work was supported by two grants overseen by the French National Research

Agency (ANR) (Project ANR 12-ADAP-0002-01) to MIT and YV, and the

ECOSNord/ANUIES/CONACYT/SEP project M12A03, CONACYT-ANUIES

207571 to MIT and LEE. LEE research and JAA-L and M-AF postdoctoral salaries

were supported by CONACYT-Mexico Investigación Científica Básica

CB2011/167826 awarded to LEE. M-AF was funded by the Project ANR 12-ADAP-

0002-01 and NEM-A was funded by a CONACYT PhD fellowship grant no.

579966/410748. OO was funded by the Academy of Finland (grant 309581), the Jane

and Aatos Erkko Foundation, and the Research Council of Norway (SFF-III grant

223257). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

Supporting information captions

Figure S1: Altitudinal profiles along gradients *a* and *b*. Sampled populations are plotted on parallel altitudinal profiles for gradients *a* and *b*. Darker gray lines indicate lower latitude for gradient *a* and lower longitude for gradient *b*. Sampled populations are plotted by green circles (*parviglumis*) or red triangles (*mexicana*). The altitude of the two experimental fields (CEBAJ: 1750m and SENGUA: 2017m) are marked with stars on y-axes.

Figure S2: Principal Component Analysis of 19 climate variables for 37 teosinte populations. A: Projection of *parviglumis* (in green) and *mexicana* (in red) populations on the first PCA plane with gradients *a* and *b* indicated by triangles and circles, respectively. The 11 populations evaluated in common gardens are surrounded by a purple outline. Populations that were previously sequenced to detect selection footprints are shown in bold (S1 Table). B: Correlation circle of the 19 climatic variables on the first PCA plane. Climatic variables indicated as Tn (n from 1 to 11) and Pn (n from 12 to 19) are related to temperature and precipitation, respectively. Altitude, Latitude and Longitude (in blue) were added as supplementary variables, and SEBAJ and SENGUA field locations were added as supplementary individuals.

Figure S3: Box-plots of means adjusted by field, year and block, for all traits. Populations are ranked by altitude. *parviglumis* populations are shown in green and *mexicana* in red. Lighter colors are used for gradient 'a' and darker colors for gradient 'b'. Units of measurement correspond to those defined in S2 Table. For male and

female flowering time, we report values for all 11 populations although very few

individuals from the two most lowland populations (P1a and P2b) flowered.

Covariation with altitude was significant for all traits except for the number of nodes

with ears on the main tiller (S3 Table).

Figure S4: Significance of Q_{st}-F_{st} difference for each trait. The dotted blue line

indicates the 95% threshold of the simulated distributions and the red line refers to the

observed difference. We considered as spatially-varying traits those for which the

observed difference fell outside the 95% threshold.

Figure S5. Evanno method calculations for population number ∆K in the

association panel genotyped for 38 SSRs.

Figure S6. Genetic clustering of ancestry proportions in the association panel

genotyped for 38 SSRs. Genetic clustering was computed for K=2 to K=11. Vertical

lines (individuals) are partitioned into coloured segments whose length represents the

admixture proportions from the K clusters.

Figure S7: Genomic F_{st}-scans on 6 teosinte populations. We computed 4 pairwise-

F_{st} values from 6 populations previously sequenced (S1 Table). Those include F_{st}

between lowland and highland populations of each gradient (P1a-M6a, P2b-M7b) as

well as within subspecies on gradient *b* (P2b-P8b, M1b-M7b). F_{st} values are averaged

across sliding windows of 20 SNPs with a step of five SNPs (from top to bottom,

chromosome 1 to 10) and normalized by subtracting the *Fst* mean and dividing by the

standard deviation across pairwise comparisons. Only the top 1% values are

represented. The 1‰ thresholds for each pairwise comparisons are indicated by colored horizontal lines. Horizontal black bars indicate location of inversions on chromosome 1 (*Inv1n*), chromosome 4 (*Inv4m*) and chromosome 9 (*Inv9d*). The subset of 218 outlier SNPs analyzed in the present study is indicated with black diamond marks along the X axes.

Figure S8: QQ-plots of observed P-values and expected P-values generated from 38 SSRs. We employed three versions of the model M6 with correction for neither structure nor kinship, with correction for genetic structure (at K=5), with correction for genetic structure and kinship.

Figure S9: Manhattan plots of associations between 171 outlier SNPs and 12 phenotypic traits along with their Bayenv2 outlier status for PCenv1. X-axis indicates the positions of outlier SNPs on chromosomes 1 to 10, black and gray colors alternating per chromosome. Significant associations (10% FDR) are indicated considering either a structure matrix at K=5 (dots with a pink outline), for 11 populations (blue dots), or for both K=5 and 11 populations models (purple dots). Empty dots designate SNPs displaying significant association to PCenv1.

Figure S10: Pairwise Linkage Disequilibrium (LD) between outlier SNPs. Pairwise LD between 171 SNPs was estimated using r^2, and corrected for structure at K=5 and kinship computed from 38 SSRs. Blue shaded bars show the 23 SNPs found to associate with at least one phenotype under the 11 populations structure correction.

Figure S11: Pairwise correlations between phenotypic traits. Pearson coefficient sign and magnitude for significant correlations between phenotypic traits after correction for experiment design (Model M'1). X: correlations that are not significant.

S1 Table. Description of 37 teosinte populations and sets of populations used in the present study by data types.

S2 Table. List of the 18 phenotypic traits measured and estimates of narrow-sense heritabilities (h^2).

S3 Table. Significance of main effects for each trait as determined by models M1 and M3.

S4 Table. Description of 46 SSRs and genotyping success rate.

S5 Table. Characteristics, association with phenotypes, effects and correlation with environment of 218 outlier SNPs.

S6 Table. Number of individuals used to test associations between 171 SNPs and 18 phenotypes.

S7 Table. **Additive and dominance effects of SNPs associated to traits after the 11-population structure correction.**

References

1. Whitlock MC. Modern approaches to local adaptation. The American Naturalist. 2015;186.

2. Bradshaw AD. Ecological significance of genetic variation between populations. In: Dirzo R, Sarukhan J, editors. Perspectives on plant population ecology. Sunderland, MA.: Sinauer; 1984. p. 213-28.

3. Bulmer MGG. Multiple niche polymorphism. The American Naturalist. 1972;106:254-7.

4. Endler JA. Natural selection in the wild. 1986:354.

5. Gay L, Crochet PA, Bell DA, Lenormand T. Comparing clines on molecular and phenotypic traits in hybrid zones: A window on tension zone models. Evolution. 2008;62:2789-806.

6. Lande R. Natural selection and random genetic drift in phenotypic evolution. Evolution. 1976;30:314-34.

7. Lenormand T. Gene flow and the limits to natural selection. Trends in Ecology and Evolution. 2002;17:183-9.

8. Whitlock MC, Gomulkiewicz R. Probability of fixation in a heterogeneous environment. Genetics. 2005:1-40.

9. Yeaman S, Otto SP. Establishment and maintenance of adaptive genetic divergence under migration, selection and drift. Evolution. 2011;67:2123-9.

10. Rundle HD, Nosil P. Ecological speciation. Ecology Letters. 2005;8:336-52.

11. Kawecki TJ, Ebert D. Conceptual issues in local adaptation. Ecology Letters. 2004;7:1225-41.

12.   Hereford J. A quantitative survey of local adaptation and fitness trade⬜offs. The American Naturalist. 2009;173:579-88.

13.   Leimu R, Fischer M. A meta-analysis of local adaptation in plants. PLoS ONE. 2008;3.

14.   Tiffin P, Ross-Ibarra J. Advances and limits of using population genetics to understand local adaptation. Trends in Ecology and Evolution. 2014;29:673-80.

15.   Garcia-Ramos G, Kirkpatrick M. Genetic models of adaptation and gene flow in peripherial populations. Evolution. 1997;51:21-8.

16.   Slatkin M. Rare alleles as indicators of gene flow. Evolution. 1985;39(1):53-65.

17.   Lande R. Neutral theory of quantitative genetic variance in an island model with local extinction and colonization. Evolution. 1992;46:381-9.

18.   Whitlock MC. Neutral additive genetic variance in a metapopulation. Genetics Research. 1999;74:215-21.

19.   Spitze K. Population structure in Daphina obtusa: quantitative genetic and allozymic variation. Genetics Society of America. 1993;135:367-74.

20.   Wright S. The genetical structure of populations. Annals of Eugenics. 1951;15:323-54.

21.   Luquet E, Léna J-P, Miaud C, Plénet S. Phenotypic divergence of the common toad (Bufo bufo) along an altitudinal gradient: evidence for local adaptation. Heredity. 2015;114:69-79.

22.   Roschanski AM, Csilléry K, Liepelt S, Oddou-Muratorio S, Ziegenhagen B, Huard Fdr, et al. Evidence of divergent selection for drought and cold tolerance at landscape and local scales in Abies alba Mill. in the French Mediterranean Alps. Molecular Ecology. 2016;25:776-94.

23.   Kawakami T, Morgan TJ, Nippert JB, Ocheltree TW, Keith R, Dhakal P, et al. Natural selection drives clinal life history patterns in the perennial sunflower species, Helianthus maximiliani. Molecular Ecology. 2011;20:2318-28.

24.   Moyers BT, Rieseberg LH. Remarkable life history polymorphism may be evolving under divergent selection in the silverleaf sunflower. Molecular Ecology. 2016;25:3817-30.

25.   Bierne N, Welch J, Loire E, Bonhomme F, David P. The coupling hypothesis: Why genome scans may fail to map local adaptation genes. Molecular Ecology. 2011;20:2044-72.

26.   Lewontin RC, Krakauer J. Distribution of gene frequency as a test of the theory of the selective neutrality of polymorphisms. Genetics. 1973;74:175-95.

27.   Beaumont MA, Nichols RA. Evaluating loci for use in the genetic analysis of population structure. Proceedings of the Royal Society B: Biological Sciences. 1996;263:1619-26.

28.   Vitalis R, Dawson K, Boursot P. Interpretation of variation across marker loci as evidence of selection. Genetics. 2001;158:1811-23.

29.   Foll M, Gaggiotti O. A genome scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. Genetics. 2008;180:977-93.

30.   Excoffier L, Hofer T, Foll M. Detecting loci under selection in a hierarchically structured population. Heredity. 2009;103:285-98.

31.   Bonhomme M, Chevalet C, Servin B, Boitard S, Abdallah J, Blott S, et al. Detecting selection in population trees: The Lewontin and Krakauer test extended. Genetics. 2010;186:241-62.

32.   Günther T, Coop G. Robust identification of local adaptation from allele frequencies. Genetics Society of America. 2013;195:205-20.

33.   Lotterhos KE, Whitlock MC. Evaluation of demographic history and neutral parameterization on the performance of FST outlier tests. Molecular Ecology. 2014;23:2178-92.

34.   Haasl RJ, Payseur BA. Fifteen years of genomewide scans for selection: trends, lessons and unaddressed genetic sources of complication. Molecular Ecology. 2016;25:5-23.

35.   Le Corre V, Kremer A. The genetic differentiation at quantitative trait loci under local adaptation. Molecular Ecology. 2012;21:1548-66.

36.   Yi X, Liang Y, Huerta-Sanchez E, Jin X, Xi Ping Cuo Z, Pool JE, et al. Sequencing of fifty human exomes reveals adaptations to high altitude. Science. 2010;329:75-8.

37.   Coop G, Witonsky D, Di Rienzo A, Pritchard JK. Using environmental correlations to identify loci underlying local adaptation. Genetics. 2010;185:1411-23.

38.   Guillot G, Renaud S, Ledevin R, Michaux J, Claude J. A unifying model for the analysis of phenotypic, genetic, and geographic data. 2012;61:897-911.

39.   Frichot E, Schoville SD, Bouchard G, Franc O. Testing for Associations between Loci and Environmental Gradients Using Latent Factor Mixed Models. Molecular Biology and Evolution. 2013;30:1687-99.

40.   Gautier M. Genome-wide scan for adaptive divergence and association with population-specific covariates. Genetics. 2015;201:1555-79.

41.   Joost S, Bonin A, Bruford MW, Després L, Conord C, Erhardt G, et al. A spatial analysis method (SAM) to detect candidate loci for selection: Towards a landscape genomics approach to adaptation. Molecular Ecology. 2007;16:3955-69.

42.   Poncet BN, Herrmann D, Gugerli F, Taberlet P, Holderegger R, Gielly L, et al. Tracking genes of ecological relevance using a genome scan in two independent regional population samples of Arabis alpina. Molecular Ecology. 2010;19:2896-907.

43.   De Mita S, Thuillet AC, Gay L, Ahmadi N, Manel S, Ronfort J, et al. Detecting selection along environmental gradients: Analysis of eight methods and their effectiveness for outbreeding and selfing populations. Molecular Ecology. 2013;22:1383-99.

44.   Hoban S, Kelley JL, Lotterhos KE, Antolin MF, Bradburd G, Lowry DB, et al. Finding the genomic basis of local adaptation: pitfalls, practical solutions, and future directions. The American Naturalist. 2016;188:379-97.

45.   Fournier-Level A, Korte A, Cooper MD, Nordborg M, Schmitt J, Wilczek AM. A map of local adaptation in Arabidopsis thaliana. Science. 2011;334:86-9.

46.   Hancock AM, Brachi B, Faure N, Horton MW, Jarymowycz LB, Sperone FG, et al. Adaptation to climate across the Arabidopsis thaliana genome. Science. 2011;334:83-6.

47.   Barton N, Hermisson J, Nordborg M. Why strucutre matters. eLife. 2019;8.

48.   Ross-Ibarra J, Tenaillon M, Gaut BS. Historical divergence and gene flow in the genus Zea. Genetics. 2009;181:1399-413.

49.   Hufford MB, Martínez-Meyer E, Gaut BS, Eguiarte LE, Tenaillon MI. Inferences from the historical distribution of wild and domesticated maize provide ecological and evolutionary insight. PLoS ONE. 2012;7.

50.   Bilinski P, Albert PS, Berg JJ, Birchler JA, Grote MN, Lorant A, et al. Parallel altitudinal clines reveal trends in adaptive evolution of genome size in Zea mays. PLoS Genetics. 2018;14.

51.   Diez CM, Gaut BS, Meca E, Scheinvar E, Montes-Hernandez S, Eguiarte LE, et al. Genome size variation in wild and cultivated maize along altitudinal gradients. New Phytologist. 2013;199:264-76.

52.   Pyhäjärvi T, Hufford MB, Mezmouk S, Ross-Ibarra J. Complex patterns of local adaptation in teosinte. Genome Biology and Evolution. 2013;5:1594-609.

53.   Fang Z, Pyhäjärvi T, Weber AL, Dawe RK, Glaubitz JC, Sánchez González JdJ, et al. Megabase-scale inversion polymorphism in the wild ancestor of maize. Genetics. 2012;191:883-94.

54.   Aguirre-Liguori JA, Tenaillon MI, Vázquez-Lobo A, Gaut BS, Jaramillo-Correa JP, Montes-Hernandez S, et al. Connecting genomic patterns of local adaptation and niche suitability in teosintes. Molecular Ecology. 2017;26:4226-40.

55.   Fustier MA, Brandenburg JT, Boitard S, Lapeyronnie J, Eguiarte LE, Vigouroux Y, et al. Signatures of local adaptation in lowland and highland teosintes from whole-genome sequencing of pooled samples. Molecular Ecology. 2017;26:2738-56.

56.   Ovaskainen O, Karhunen M, Zheng C, Arias JMC, Merilä J. A new method to uncover signatures of divergent and stabilizing selection in quantitative traits. Genetics. 2011;189:621-32.

57.   Sohail M, Maier RM, Ganna A, Bloemendal A, Martin AR, Turchin MC, et al. Polygenic adaptation on height is overestimated due to uncorrected stratification in genome-wide association studies. eLife. 2019;8:1-17.

58.   Desrousseaux AD, Sandron F, Siberchicot A, Cierco-Ayrolles C, Mangin B. Package ' LDcorSV '. 2017.

59.   Mangin B, Siberchicot A, Nicolas S, Doligez A, This P, Cierco-Ayrolles C. Novel measures of linkage disequilibrium that correct the bias due to population structure and relatedness. Heredity. 2012;108:285-91.

60.   Savolainen O, Lascoux M, Merilä J. Ecological genomics of local adaptation. Nature reviews Genetics. 2013;14:807-20.

61.   Halbritter AH, Fior S, Keller I, Billeter R, Edwards PJ, Holderegger R, et al. Trait differentiation and adaptation of plants along elevation gradients. Journal of Evolutionary Biology. 2018;31(6):784-800.

62.   Körner C. The use of 'altitude' in ecological research. Trends in Ecology and Evolution. 2007;22:569-74.

63.   Friend AD, Woodward FI, Switsur VR. Field measurements of photosynthesis, stomatal conductance, leaf nitrogen and δ 13 C along altitudinal gradients in Scotland. Functional Ecology. 1989;3:117.

64.   Neuner G. Frost resistance in alpine woody plants. Frontiers in Plant Science. 2014;5.

65.   Frohnmeyer H, Staiger D. Update on ultraviolet-B light responses ultraviolet-B radiation-mediated responses in plants . Balancing damage and protection. 2014;133:1420-8.

66.   Byars SG, Papst W, Hoffmann AA. Local adaptation and cogradient selection in the alpine plant, Poa hiemata, along a narrow altitudinal gradient. Evolution. 2007;61:2925-41.

67.   Luo Y, Widmer A, Karrenberg S. The roles of genetic drift and natural selection in quantitative trait divergence along an altitudinal gradient in Arabidopsis thaliana. Heredity. 2015;114:220-8.

68.   Guerin GR, Wen H, Lowe AJ, Guerin GR, Wen H. Leaf morphology shift linked to climate change. Population Ecology. 2012:882-6.

69.   Kofidis G, Bosabalidis AM, Moustakas M. Contemporary seasonal and altitudinal variations of leaf structural features in oregano (Origanum vulgare L.). Annals of Botany. 2003;92:635-45.

70.   Mendez-Vigo B, Pico FX, Ramiro M, Martinez-Zapater JM, Alonso-Blanco C. Altitudinal and climatic adaptation is mediated by flowering traits and FRI, FLC, and PHYC genes in Arabidopsis. Plant Physiology. 2011;157:1942-55.

71.   Oleksyn J, Modrzynski J, Tjoelker MG, Zytkowaik R, Reich PB, Karolewski P. Growth and physiology of Picea abies populations from elevational transects: common garden evidence for altitudinal ecotypes and cold adaptation. 1998:573-90.

72.   Soularue JP, Kremer A. Evolutionary responses of tree phenology to the combined effects of assortative mating, gene flow and divergent selection. Heredity. 2014;113:485-94.

73.   Benson DL, Hallauer AR. Inbreeding depression rates in maize populations before and after recurrent selection. Journal of Heredity. 1994;85:122-8.

74.   Smith JSC, Goodman MM, Lester RN. Variation within teosinte. I. Numerical analysis of morphological data. Economic Botany. 1981;35:187-203.

75.   Doebley JF. Maize introgression into teosinte -- a reappraisal. Annals of the Missouri Botanical Garden. 1984;71:1100-13.

76.   Lauter N, Gustus C, Westerbergh A, Doebley J. The inheritance and evolution of leaf pigmentation and pubescence in teosinte. Genetics Society of America. 2004;167:1949-59.

77.   Raven J, A. Selection pressures on stomatal evolution. New Phytologist. 2002;153:371-86.

78.   Dittberner H, Korte A, Mettler-Altmann T, Weber APM, Monroe G, de Meaux J. Natural variation in stomata size contributes to the local adaptation of water-use efficiency in Arabidopsis thaliana. Molecular Ecology. 2018:4052-65.

79.   Carlson JE, Adams CA, Holsinger KE. Intraspecific variation in stomatal traits, leaf traits and physiology reflects adaptation along aridity gradients in a South African shrub. Annals of Botany. 2016;117:195-207.

80.   Körner C, Mayr R. Stomatal behaviour in alpine plant communities between 600 and 2600 metres above sea level. In: Grace J, Ford ED, Jarvis PG, editors. Plants and their Atmospheric Environment: Blackwell, Oxford,; 1981. p. pp 205-18.

81.   Bresson CC, Vitasse Y, Kremer A, Delzon S. To what extent is altitudinal variation of functional traits driven by genetic adaptation in European oak and beech? Tree Physiology. 2011;31:1164-74.

82.   Kooyers NJ, Greenlee AB, Colicchio JM, Oh M, Blackman BK. Replicate altitudinal clines reveal that evolutionary flexibility underlies adaptation to drought stress in annual Mimulus guttatus. New Phytologist. 2015;206:152-65.

83.   Körner C, Neumayer M, Menendez-Riedl SP, Smeets-Scheel A. Functional morphology of mountain plants. Flora. 1989;182:353-83.

84.   Jakobsson A, Eriksson O. A comparative study of seed number, seed size, seedling size and recruitment in grassland plants. Oikos. 2000;88:494-502.

85.   Buckler ES, Holland JB, Bradbury PJ, Acharya CB, Brown PJ, Browne C, et al. The genetic architecture of maize flowering time. Science. 2009;325:714-8.

86.   Moreau L, Charcosset A, Gallais A. Use of trial clustering to study QTL x environment effects for grain yield and related traits in maize. Theoretical and Applied Genetics. 2004;110:92-105.

87. Durand E, Bouchet S, Bertin P, Ressayre A, Jamin P, Charcosset A, et al. Flowering time in maize: Linkage and epistasis at a major effect locus. Genetics. 2012;190:1547-62.

88. Li D, Wang X, Zhang X, Chen Q, Xu G, Xu D, et al. The genetic architecture of leaf number and its genetic relationship to flowering time in maize. New Phytologist. 2015;210:256-68.

89. Kronholm I, Loudet O, de Meaux J. Correction: Influence of mutation rate on estimators of genetic differentiation - lessons from Arabidopsis thaliana. BMC Genetics. 2010;11:7-9.

90. Consortium G. 1,135 Genomes reveal the global pattern of polymorphism in Arabidopsis thaliana. Cell. 2016;166:481-91.

91. Whitt SR, Wilson LM, Tenaillon MI, Gaut BS, Buckler ES. Genetic diversity and selection in the maize starch pathway. Proceedings of the National Academy of Sciences of the United States of America. 2002;99:12959-62.

92. Jaenicke-Despres V, Buckler ES, Smith BD, Gilbert MTP, Cooper A, Doebley J, et al. Early allelic selection in maize as revealed by ancient DNA. 2003;302:1206-9.

93. Weber AL, Briggs WH, Rucker J, Baltazar BM, De Jesús Sánchez-González J, Feng P, et al. The genetic architecture of complex traits in teosinte (Zea mays ssp. parviglumis): New evidence from association mapping. Genetics. 2008;180:1221-32.

94. Bouchet S, Servin B, Bertin P, Madur D, Combes V, Dumas F, et al. Adaptation of maize to temperate climates: Mid-density genome-wide association genetics and diversity patterns reveal key genomic regions, with a major contribution of the Vgt2 (ZCN8) locus. PLoS ONE. 2013;8.

95.   Jaenicke-Després V, Buckler ES, Smith BD, Gilbert MTP, Cooper A, Doebley J, et al. Early allelic selection in maize as revealed by ancient DNA. Science. 2003;302:1206-8.

96.   Yu J, Li X, Zhu C, Yeh C-T, Wu W, Takacs E, et al. Genic and non-genic contributions to natural variation of quantitative traits in maize. Genome research. 2012:2436-44.

97.   Sheehan MJ, Kennedy LM, Costich DE, Brutnell TP. Subfunctionalization of PhyB1 and PhyB2 in the control of seedling and mature plant traits in maize. Plant Journal. 2007;49:338-53.

98.   Danilevskaya ON, Meng X, Hou Z, Ananiev EV, Simmons CR. A genomic and expression compendium of the expanded PEBP gene family from maize. Plant Physiology. 2007;146:250-64.

99.   Meng X, Muszynski MG, Danilevskaya ON. The FT-Like ZCN8 Gene Functions as a Floral Activator and Is Involved in Photoperiod Sensitivity in Maize. The Plant Cell. 2011;23:942-60.

100.   Li YX, Li C, Bradbury PJ, Liu X, Lu F, Romay CM, et al. Identification of genetic variants associated with maize flowering time using an extremely large multi-genetic background population. The Plant journal : for cell and molecular biology. 2016;86:391-402.

101.   Aguirre□Liguori JA, Gaut BS, Jaramillo□Correa JP, Tenaillon MI, Montes□Hernández S, García□Oliva F, et al. Divergence with gene flow is driven by local adaptation to temperature and soil phosphorus concentration in teosinte subspecies (Zea mays parviglumis and Zea mays mexicana) Molecular Ecology. 2019:2814-30.

102.   Wellenreuther M, Bernatchez L. Eco-evolutionary genomics of chromosomal inversions. Trends in Ecology and Evolution. 2018;33:427-40.

103.   Ayala D, Ullastres A, González J. Adaptation through chromosomal inversions in Anopheles. Frontiers in Genetics. 2014;5:1-10.

104.   Barth JMI, Berg PR, Jonsson PR, Bonanomi S, Corell H, Hemmer-Hansen J, et al. Genome architecture enables local adaptation of Atlantic cod despite high connectivity. Molecular Ecology. 2017;26:4452-66.

105.   Lundberg M, Liedvogel M, Larson K, Sigeman H, Grahn M, Wright A, et al. Genetic differences between willow warbler migratory phenotypes are few and cluster in large haplotype blocks. Evolution Letters. 2017:155-68.

106.   Twyford AD, Friedman J. Adaptive divergence in the monkey flower Mimulus guttatus is maintained by a chromosomal inversion. Evolution. 2015;69:1476-86.

107.   Lowry DB, Willis JH. A widespread chromosomal inversion polymorphism contributes to a major life-history transition, local adaptation, and reproductive isolation. PLoS Biology. 2010;8.

108.   Hijmans RJ, van Etten J, Cheng J, Mattiuzzi M, Sumner M, Greenberg JA, et al. Package 'raster ': geographic data analysis and modeling. 2018.

109.   Husson F, Josse J, Le S, Mazet J. Package ' FactoMineR '. An R package. 2016:96.

110.   Andorf CM, Cannon EK, Portwood JL, Gardiner JM, Harper LC, Schaeffer ML, et al. MaizeGDB update: New tools, data and interface for the maize model organism database. Nucleic Acids Research. 2016;44:1195-201.

111.   Camus-Kulandaivelu L, Veyrieras JB, Madur D, Combes V, Fourmann M, Barraud S, et al. Maize adaptation to temperate climate: Relationship between

population structure and polymorphism in the Dwarf8 gene. Genetics. 2006;172:2449-63.

112. Guichoux E, Lagache S, Wagner S, Chaumeil P, Léger P, Lepais O, et al. Current trends in microsatellite genotyping. Molecular Ecology Resources. 2011;11:591-611.

113. Jakobsson M, Rosenberg NA. CLUster Matching and Permutation Program Version 1.1.2. 2007.

114. Evanno G, Regnaut S, Goudet J. Detecting the number of clusters of individuals using the software STRUCTURE: A simulation study. Molecular Ecology. 2005;14:2611-20.

115. Hardy OJ, Vekemans X. spagedi: a versatile computer program to analyse spatial genetic structure at the individual or population levels. Molecular Ecology Notes. 2002;2:618-20.

116. Loiselle BA, Sork VL, Nason JD, Graham C. Spatial genetic structure of a tropical understory shrub, Psychotria officinalis (Rubiaceae). American Journal of Botany. 1995;82:1420-5.

117. Tian F, Bradbury PJ, Brown PJ, Hung H, Sun Q, Flint-Garcia S, et al. Genome-wide association study of leaf architecture in the maize nested association mapping population. Nature Genetics. 2011;43:159-62.

118. Sanchez JdJ, Kato Yamakake TA, Aguilar Sanmiguel M, Hernandez Casillas JM, Lopez Rodriguez A, Ruiz Corral JA. Distribucion y caracterización del teocintle. 1998:165.

119. Butler D, Cullis BR, Gilmour AR, Gogel BJ, editors. ASReml-R reference manual. Technical Report; 2007; Brisbane, Queensland, Australia.

120. Holsinger KE, Weir BS. Genetics in geographically structured populations: defining, estimating and interpreting F(ST). Nature reviews Genetics. 2009;10:639-50.

121. Gilbert KJ, Whitlock MC. QST-FST comparisons with unbalanced half-sib designs. Molecular Ecology Resources. 2015;15:262-7.

122. Karhunen M, Merilä J, Leinonen T, Cano JM, Ovaskainen O. driftsel: An R package for detecting signals of natural selection in quantitative traits. Molecular Ecology Resources. 2013;13:746-54.

123. Hadfield JD. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. Journal of Statistical Software. 2010;33.

124. Semagn K, Babu R, Hearne S, Olsen M. Single nucleotide polymorphism genotyping using Kompetitive Allele Specific PCR (KASP): Overview of the technology and its application in crop improvement. Molecular Breeding. 2014;33:1-14.

125. Weir BS, Hill WG. Estimating F-statistics. Annu Rev Genet. 2002;36:721-50.

126. Günther T, Coop G. A Short Manual for Bayenv2.0. 2016.

127. De Villemereuil P, Gaggiotti OE. A new FST-based method to uncover local adaptation using environmental variables. Methods in Ecology and Evolution. 2015.

128. Rincent R, Moreau L, Monod H, Kuhn E, Melchinger AE, Malvar RA, et al. Recovering power in association mapping panels with variable levels of linkage disequilibrium. Genetics. 2014;197:375-87.

129. Bradburd GS, Ralph PL, Coop GM. Disentangling the effects of geographic and ecological isolation on genetic differentiation. Evolution. 2013;67:3258-73.

130. Lichstein JW. Multiple regression on distance matrices: A multivariate spatial analysis tool. Plant Ecology. 2007;188:117-31.

131.   Goslee SC, Urban DL, editors. Journal of Statistical Software The ecodist

Package for Dissimilarity-based Analysis of Ecological Data2007.

A.

B.

C.