How important is it to consider lineage diversification heterogeneity in in

macroevolutionary studies: lessons from the lizard family Liolaemidae

Olave Melisa[a], Avila Luciano J. [b], Jack W. Sites, Jr.[c] and Morando Mariana[b,d]

[a]*Department of Biology, University of Konstanz, Konstanz, Germany.*

[b]*Instituto Patagónico para el Estudio de los Ecosistemas Continentales, Consejo Nacional de*

Investigaciones Científicas y Técnicas (IPEEC-CONICET), Boulevard Almirante Brown

2915, U9120ACD, Puerto Madryn, Chubut, Argentina.

[c]*Department of Biology and M.L. Bean Life Science Museum, Brigham Young University*

(BYU), Provo, UT 84602, USA; current address: Department of Biology, Austin Peay State

University, Clarksville, Tennessee, 37044.

[d]*Universidad Nacional de la Patagonia San Juan Bosco, Sede Puerto Madryn, Boulevard*

Almirante Brown 3700, U9120ACD, Puerto Madryn, Chubut, Argentina.

Abstract

Macroevolutionary studies commonly apply multiple models to test state-dependent

diversification. These models track the association between states of interest along a

phylogeny, but they do not consider whether independent shifts in character states are

associated with shifts in diversification rates. This potentially problematic issue has received

little theoretical attention, while macroevolutionary studies implementing such models in

increasing larger scale studies continue growing. A recent macroevolutionary study has found

that Andean orogeny has acted as a species pump driving diversification of the family

Liolaemidae, a highly species-rich lizard family native to temperate southern South America.

This study approaches a distribution-dependent hypothesis using the Geographic State

Speciation and Extinction model (GeoSSE). However, more recent analyses have shown that

there is a clear heterogeneous diversification pattern in the Liolaemidae, which likely biased

the GeoSSE analysis. Specifically, we show here that there are two shifts to accelered

speciation rates involving species groups that were classified as "Andean" in their

distributions. We demonstrate that this GeoSSE result is meaningless when heterogeneous

diversification rates are included. We use the lizard family Liolaemidae to demonstrate

potential risks of ignoring clade-specific differences in diversification rates in

macroevolutionary studies.

Key words: GeoSSE, diversification, speciation, extinction, macroevolution, biogeography,

Liolaemus, *Phymaturus*, *Ctenoblepharys,* Andes

Introduction

Macroevolutionary modeling of diversification plays important roles in inferring large-scale

biodiversity patterns (Schluter 2016). Several studies have focused on quantifying differences

in macroevolutionary patterns linked to geographic, ecological, life-history and other traits,

based on the variation in speciation and extinction rates (Jablonski 2008; Rabosky and

McCune 2010; Ng and Smith 2014). Given that the mechanisms underlying the correlations

between characters and diversification are generally poorly understood (Rabosky and

Goldberg 2015), models have been developed to test the role of a range of different states

promoting diversification, including binary traits (Maddison 2006), quantitative traits

(FitzJohn 2010), geographic character states (Goldberg et al. 2011), multiple characters

(FitzJohn 2012), punctuated trait changes (Goldberg and Igic 2012; Magnuson-Ford and Otto

2012), and time-dependent macroevolutionary rates (Rabosky and Glor 2010). These models

have been shown to perform very well on simulated datasets when using reasonably large

trees (FitzJohn et al. 2009; FitzJohn 2010; Rabosky and Glor 2010; Goldberg et al. 2011;

FitzJohn 2012; Goldberg and Igic 2012; Magnuson-Ford and Otto 2012; Stadler and

Bonhoeffer 2013; Davis et al. 2013), and they have been implemented in hundreds of

empirical studies (Rabosky and Goldberg 2015).

These models track associations between the states of interest and speciation and extinction

rates along a phylogenetic tree, but they do not consider whether independent shifts in trait

state are associated with shifts in diversification (Maddison and FitzJohn 2014; Rabosky and

Goldberg 2015). Therefore, even if the shift is unrelated to the state targeted, a strong

correlation with the diversification can be inferred from a rate shift (Maddison et al. 2007;

FitzJohn 2010; Maddison and FitzJohn 2014; Rabosky and Goldberg 2015). Thus, all

heterogeneity in diversification rates could potentially be linked purely to the states included

in the analysis. Consequently, while larger trees are preferred due to the presumable increase

of power, this also increases the risk of including clades with differences in states that can

affect diversification along a tree (factors such as ecological requirements, dispersal abilities

and life history [Li et al. 2018]). These potential issues have received little theoretical

attention, while macroevolutionary studies implementing such models at increasingly larger

scales continue to rise (Rabosky and Goldberg 2015).

The lizard family Liolaemidae is the most species-rich lizard clade the southern half of South

America (307 species; Reptile Database 11 February 2019). The clade includes three genera:

Ctenoblepharys, Liolaemus and *Phymaturus* (Fig. S1; Table 1)*. Ctenoblepharys* is a

monotypic genus with a distribution restricted to the coastal desert of Peru (Table 1), whereas

Liolaemus is the world's richest temperate zone genus of extant amniotes (Olave et al. 2018),

with 262 described species (Reptile Database 2 February 2019). *Liolaemus* includes a highly

diverse group of species inhabiting a wide range of different environments (Table 1). The

sister genus of *Liolaemus*, *Phymaturus* (44 species; Reptile Database 2 February 2019) is

distributed along both the eastern and western Andean slopes in Argentina and Chile

(*palluma* clade), and through Patagonia (*patagonicus* clade). *Phymaturus* are strictly

saxicolous and largely restricted to volcanic plateaus and peaks (Cei 1986).

The three genera have clear differences in species richness, ecological requirements,

behaviors, and life histories (Table 1). A recent macroevolutionary study currently has found

disparate patterns of diversification among the three genera (Olave et al. *in review*), while

another recent study has focused on the entire clade, unknowingly the shifts in the

diversification rates along the tree (Esquerré et al. 2019). This study (Esquerré et al. 2019)

represents a major contribution to evolutionary biology and herpetology in that it: (i) presents

the largest Liolaemidae time-calibrated phylogeny to date (258 taxa), (ii) the most extensive

compilation of habitats, altitudes, and temperature data for all taxa, (iii) it hypotheses

ancestral range reconstructions, and (iv) opposite to previous findings, it demonstrates that

multiple origins of viviparity are not intrinsic properties in speciation rates.

However, Esquerré et al. approach the distribution-dependent hypothesis using the

Geographic State Speciation and Extinction model (GeoSSE; Goldberg et al. 2011) to test for

differences in speciation rates in Andean vs non-Andean (low elevation) species. The

GeoSSE model detected higher speciation rates in the Andean areas, and authors infer that

the Andean orogeny has acted as a "species pump" driving diversification in the Liolaemidae.

These authors performed further analyses to support this hypothesis (e.g. ancestral

distribution reconstructions and a time variable diversification model). Nonetheless, the

GeoSSE test was clearly key to identifying the role of the Andean orogeny in driving the

diversification of this clade. However, here we show that a clearly heterogeneous

diversification history of Liolaemidae is not considered by the GeoSSE analysis. Specifically,

we detect two shifts to accelerated speciation rates involving clades that were identified as

"Andean". We show that the less diverse genus *Phymaturus* is characterized by the highest

speciation rates, and that there is a second shift within *Liolaemus*, specifically in the *L.*

elongatus clade. Consequently, the differences on speciation rates detected between Andean

vs. non-Andean species for the distribution-dependent diversification test is meaningless. We

demonstrate that the "Andean orogeny" hypothesis is not supported when the heterogeneous

diversification rates among these lizards is considered. The speciation history of the clade

Liolaemidae clearly demonstrates potential risks of the implementation of GeoSSE (and

likely other models of the family) when ignoring clade-specific differences in diversification

rates in macroevolutionary studies.

Materials and methods

Phylogenetic tree

We incorporated here the time-calibrated phylogenetic tree of Esquerré et al. (2019), which

includes the monotypic *Ctenoblepharys*, 188 described + 11 undescribed species of

Liolaemus, and 35 *Phymaturus* species (73% species coverage of all recognized

Liolaemidae). A consensus tree was obtained using TreeAnnotator 2.4 (Bouckaert et al.

2014).

Speciation and extinction rates

We estimated net diversification, speciation and extinction rates using BAMM 2.5 (Rabosky

et al. 2014). BAMM is a Bayesian approach that uses a rjMCMC to estimate lineage-specific

speciation and extinction rates, and rates of phenotypic change. Because the method

estimates rates per branch, it allows us to compare changes of these rates among clades and

species (i.e., tips) of interest. As in similar models, BAMM assumes the given topology is the

true phylogenetic tree, so to account for the topological uncertainty, we ran the analysis using

each of the last 500 trees inferred during the MCMC of BEAST. We informed the proportion

of missing taxa using *globalSamplingFraction* = 0.73, thus the program accounts for the

missing tips (i.e. 73% coverage). Priors were generated using *setBAMMpriors* in

BAMMtools (Rabosky et al. 2014), and we used all 500 obtained means for target groups

(genus, subgenus, clades and tips) to construct the final distributions used for all downstream

comparisons. All BAMM analyses were run for 5×10^6 generations, sampling every 1,000

generations, and with 25% burnin. We constructed parameter distributions per genus that

captured topological uncertainty. We calculated summary statistics using R (mean, standard

deviation and quartiles), and compared statistical differences among specific target clades

with ANOVA tests using the R function aov().

Hypothesis testing: role of the Andean orogeny in diversification of the Liolaemidae

To quantify the association between speciation and extinction rates to the Andes range, we

extracted species-specific speciation and extinction rates for different target clades, including

the whole family, genus (*Phymaturus* and *Liolaemus*), subgenus (*Eulaemus* and *Liolaemus*

sensu stricto), clades within *Phymaturus* (*P. palluma* and *P. patagonicus*) and several smaller

clades within *Liolaemus* (Table S1). We performed linear regressions using the R function

lm(), between the speciation (and extinction) rates and the maximum altitudes for all species.

The maximum altitude data were taken from the Esquerré et al. (2019) recompilation (their

Table S3). We also calculated linear models using the R function aov(), with the formula: rate

~ "target clade" * "maximum altitude".

We implemented the GeoSSE (Geographic State Speciation and Extinction) models (Goldberg

et al. 2011) to test the hypothesis of higher speciation rates associated with the Andean species,

and used the same classification and tested the same set of models as in Esquerré et al. (2019;

Table 2). However, we do not fully agree with the original classification; as an example, the

"Patagonia" group is distributed across a huge area that was assumed to be "Andean", which

we consider a poor classification for many species. For example, both the *P. patagonicus* and

the *L. lineomaculatus* clades are restricted mainly to the lowland Patagonian steppe. However,

here we respect the authors' original classification and address the issue of heterogeneous

diversification rates in our analyses and discussion. We ran all analyses for the Liolaemidae as

a single clade, and then also for different nested clades. We used ML to estimate the parameters

as a starting point for an MCMC chain of 30,000 generations with a 20% burnin. All analyses

were performed in the R package *diversitree* (FitzJohn et al. 2009).

Results

Heterogeneous diversification within the family Liolaemidae

BAMM estimation of speciation and extinction rates on the Liolaemidae phylogeny (Figure

1A-B), displays two prominent shifts (PP = 0.4; Table S3), including the origin of the genus

Phymaturus (red), and the *Liolaemus elongatus* clade (light blue). There are significant

differences in speciation and extinction rates among genera ($p < 0.001$), as clearly shown by

the distributions of parameter estimations (Fig. 2). Specifically, the genus *Phymaturus* has the

highest speciation rate, is also associated with a high extinction rate. This result is concordant

with another study currently under review, using an different phylogenetic tree (Olave et al.

in review).

Hypothesis test: the role of the Andes mountains in diversification of the Liolaemidae

We constructed linear models between the maximum altitude (MA) of species occurrence

records, and the species-specific speciation and extinction rates. When considering all species

of Liolaemidae (258 tips), we find highly significant differences among genera ($p < 2^{-16}$), and

no significant effect of MA ($p = 0.808$), or their interaction ($p = 0.207$; Table S3A; Fig. S2).

Analyses of *Liolaemus* alone (194 tips) show a highly significant subgenus effect (p = 3.10^{-8}),

but non-significant MA effect (p = 0.365) or interaction effects (p = 0.57; Table S3E).

Equivalent results (i.e. no effect of the MA, but significant clade effect) were found for the

subgenus *Liolaemus* sensu stricto (s.s.) when including (97 tips), or excluding the *L.*

elongatus clade (71 tips), as well for the *Eulaemus* subgenus (97 tips; see Table S3F-G).

We also found a significant negative linear correlation between MA and speciation (p = 1×10^{-4}

) and extinction (p-value = 0.0019) rates in the subgenus *Eulaemus*, but a poor fit of the

model (R-squared < 0.2; Fig. 3). Analyses of *Phymaturus* alone (58 tips) show a positive

linear correlation between speciation rate and MA (Fig. 3), but there is also a clear clustering

of the *P. patagonicus* and *P. palluma* clades, both detected by the linear model (clades p < 2^{-16}

) with a non-significant contribution of the MA (p = 0.158), or their interaction (p = 0.769;

Table S3B). Finally, we found a significant correlation between speciation and extinction

rates for the *Phymaturus palluma* clade alone (28 tips; Fig S3 and Table S3C).

We performed distribution-dependent diversification tests using the GeoSSE program, first

testing the entire clade Liolaemidae, and found highly significant results (p-value =

0.0001735; Fig. 4) for the constrained model of equal speciation in Andean and sub-Andean

regions (Table S4). Thus, the GeoSSE model returns significantly higher speciation rates in

the Andean clade (= 0.27) relative to "lowland" species (= 0.11). This result is consistent

with previous findings by Esquerré et al. (2019); i.e., high-elevation Andean environments

are significantly associated with high speciation rates in the Liolaemidae. This analysis

included all *Phymaturus* species as Andean (Table 2), which also displayed a speciation rate

three times higher than *Liolaemus* (Fig. 2). We re-ran the analyses for the *Liolaemus* species

only, which returned only a slightly significant p-value = 0.04114 (Fig. 4) for a higher

speciation rate in Andean species (0.1883 vs. 0.1225; Table S4). This signal disappears

completely with the removal of the *L. elongatus* clade (which was classified entirely as

Andean; p-value = 0.16863), or when running the test with the *Eulaemus* subgenus alone (p-

value = 0.182782; Fig. 4).

Discussion

Incorporating large trees for macroevolutionary studies has the advantage of providing larger

datasets, and presumably more power. However, it is important to keep in mind the

assumptions that go into such analysis: it treats all clades as evolving according to the same

model, with the same values for the rate parameters. Here, we used the lizard family

Liolaemidae to test for errors associated the use of large trees where clade-specific

differences could bias results and lead to wrong conclusions. Our results clearly indicate that

the signals of accelerated speciation rates associated with the Andean uplift in the

distribution-dependent diversification test implemented in GeoSEE are biased (Fig. 4), due to

the two diversification rate shifts along the tree (Fig. 1). Specifically, we demonstrated that

the genera *Phymaturus* and *Liolaemus* display clear disparate patterns of diversification and

that, when incorporated into this study, show that there is no apparent signal of Andean

orogeny increasing speciation rates in the Liolaemidae. Earlier studies have confounded

clade-specific rate accelerations with the distribution-dependent diversification results.

We do not argue against the implementation of GeoSSE (or any other model) in

macroevolutionary studies, and do not doubt about the utility of state-dependent

diversification models in general. However, our study calls attention to identify

diversification rate heterogeneity for subsequent partitioning for the GeoSSE model (or other

models within the family; see also Rabosky and Goldberg 2015), and we show that the

BAMM program is a good option to identify such changes.

Acknowledgments

We thank D. Esquerré for providing clarification of how they performed their analyses. We

thank all members of the Grupo de Herpetología Patagónica (IPEEC-CONICET) for

continuing support. Financial support was provided by ANPCYT-FONCYT 1252/2015

(MM), and a postdoctoral fellowship (MO) from the Alexander von Humboldt Foundation at

Meyer Lab, Konstanz, Germany.

Author contributions

MO and MM designed the study. MO carried out the analyses. MO, MM and JS wrote and

edited the manuscript. LJA and MM provided recommendations based on the biology of the

focal organism. All authors read and approved the final manuscript.

References

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C. H., Xie, D., ... & Drummond, A.

J. (2014). BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS*

computational biology, *10*(4), e1003537.

Cei, J. M. (1986). Reptiles del centro, centro-oeste y sur de la Argentina: Herpetofauna de las

zonas áridas y semiáridas.

Davis, M. P., Midford, P. E., & Maddison, W. (2013). Exploring power and parameter

estimation of the BiSSE method for analyzing species diversification. *BMC Evolutionary*

Biology, *13*(1), 38.

Esquerré, D., Brennan, I. G., Catullo, R. A., Torres-Pérez, F., & Keogh, J. S. (2019). How

mountains shape biodiversity: The role of the Andes in biogeography, diversification, and

reproductive biology in South America's most species-rich lizard radiation (Squamata:

Liolaemidae). *Evolution* (in press).

FitzJohn, R. G. (2010). Quantitative traits and diversification. *Systematic biology*, *59*(6), 619-

633.

FitzJohn, R. G. (2012). Diversitree: comparative phylogenetic analyses of diversification in

R. *Methods in Ecology and Evolution*, *3*(6), 1084-1092.

FitzJohn, R. G., Maddison, W. P., & Otto, S. P. (2009). Estimating trait-dependent speciation

and extinction rates from incompletely resolved phylogenies. *Systematic biology*, *58*(6),

595-611.

Goldberg, E. E., & Igić, B. (2012). Tempo and mode in plant breeding system

evolution. *Evolution: International Journal of Organic Evolution*, *66*(12), 3701-3709.

Goldberg, E. E., Lancaster, L. T., & Ree, R. H. (2011). Phylogenetic inference of reciprocal

effects between geographic range evolution and diversification. *Systematic Biology*, *60*(4),

451-465.

Jablonski, D. (2008). Species selection: theory and data. *Annual Review of Ecology,*

Evolution, and Systematics, *39*, 501-524.

Li, J., Huang, J. P., Sukumaran, J., & Knowles, L. L. (2018). Microevolutionary processes

impact macroevolutionary patterns. *BMC evolutionary biology*, *18*(1), 123.

Maddison, W. P. (2006). Confounding asymmetries in evolutionary diversification and

character change. *Evolution*, *60*(8), 1743-1746.

Maddison, W. P., & FitzJohn, R. G. (2014). The unsolved challenge to phylogenetic

correlation tests for categorical characters. *Systematic biology*, *64*(1), 127-136.

Maddison, W. P., Midford, P. E., & Otto, S. P. (2007). Estimating a binary character's effect

on speciation and extinction. *Systematic biology*, *56*(5), 701-710.

Magnuson-Ford, K., & Otto, S. P. (2012). Linking the investigations of character evolution

and species diversification. *The American Naturalist*, *180*(2), 225-245.

Ng, J., & Smith, S. D. (2014). How traits shape trees: new approaches for detecting character

state-dependent lineage diversification. *Journal of evolutionary biology*, *27*(10), 2035-

2045.

Olave, M., Avila, L. J., Sites Jr, J. W., & Morando, M. (2018). Hybridization could be a

common phenomenon within the highly diverse lizard genus Liolaemus. *Journal of*

evolutionary biology, *31*(6), 893-903.

Olave, M., Gonzalez Marin, A., Avila, L.J., Sites, J.W. Jr. & Morando, M. Disparate patterns

of diversification within Liolaemini lizards. In *Neotropical diversification*, edited by Rull

& Carnaval. Springer Nature (*in review*).

Rabosky, D. L., & Glor, R. E. (2010). Equilibrium speciation dynamics in a model adaptive

radiation of island lizards. *Proceedings of the National Academy of Sciences*, *107*(51),

22178-22183.

Rabosky, D. L., & Goldberg, E. E. (2015). Model inadequacy and mistaken inferences of

trait-dependent speciation. *Systematic biology*, *64*(2), 340-355.

Rabosky, D. L., Grundler, M., Anderson, C., Title, P., Shi, J. J., Brown, J. W., ... & Larson, J.

G. (2014). BAMM tools: an R package for the analysis of evolutionary dynamics on

phylogenetic trees. *Methods in Ecology and Evolution*, *5*(7), 701-707.

Schluter, D. (2016). Speciation, Ecological Opportunity, and Latitude: (American Society of

Naturalists Address). *The American Naturalist*, *187*(1), 1-18.

Stadler, T., & Bonhoeffer, S. (2013). Uncovering epidemiological dynamics in heterogeneous

host populations using phylogenetic methods. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *368*(1614), 20120198.

Tables and Figures

	Ctenoblepharys	*Phymaturus*	*Liolaemus*
Described species	1	44	262
Distribution	Perú	Argentina Chile	Argentina Chile Perú Bolivia Southern Brazil Uruguay
Habitat	coastal desert	saxicolous	terrestrial arboreous arenicolous saxicolous
Diet	insectivores	herbivores	herbivores omnivores insectivores
Time for sexual maturity	unknown	7-8 years	2 years
Reproductive mode	oviparous	viviparous	viviparous oviparous parthenogenesis

Table 1: Summary of distribution, habitat use, diet and reproductive mode among the three Liolaemidae genera.

	Considered "Andean species"			Considered "Non-Andean species"		
	Patagonia	Central Andes	Altiplanic Andes	Central Chile	Atacama Desert	Eastern lowlands
Liolaemus	63	48	56	17	14	32
Phymaturus	33	23	5	0	0	1
Ctenoblepharys	0	0	0	0	1	0

Table 2: Species count for the geographic classification from Esquerré et al. (2019). Taken from their supplementary material.

307 Figure 1: Color-coded phylogenetic trees for the speciation (A) and extinction (B) rates

308 through time for the Liolaemidae.

Figure 2: Speciation and extinction rates obtained for *Ctenoblepharys* (green), *Liolaemus*

genus (blue) and *Phymaturus* genus (red). The density plots are constructed considering the

mean obtained from each of the last 500 trees of the MCMC run for phylogenetic estimation.

The p-value corresponds to an ANOVA test comparing distributions.

318   Figure 3: Linear regressions of the speciation/extinction rates as a function of the maximum

319   altitude (meters) of the species occurrence for different target clades: *Phymaturus* genus,

320   *Eulaemus* subgenus, *Liolaemus* sensu strict (s.s.) subgenus when excluding the *L. elongatus*

321   clade and with the *L. elongatus* clade. See also the Figure S2-4 for more regressions, and

322   Table S2 for full results of the linear model.

Figure 4: GeoSSE results for the different target clades: Liolaemidae, *Liolaemus*, *Liolaemus* (excluding *L. elongatus* clade) and *Eulaemus*. See also Table S3 for more details.