Article

Evaluating the within-host dynamics of *Ranavirus* infection with mechanistic disease models and experimental data

Joseph R. Mihaljevic[1,*], Amy L. Greer[2] and Jesse L. Brunner[3]

[1] Northern Arizona University, School of Informatics, Computing, and Cyber Systems, Flagstaff, AZ, USA; joseph.mihaljevic@nau.edu

[2] University of Guelph, Department of Population Medicine, Guelph, ON, Canada; agreer@uoguelph.ca

[3] Washington State University, School of Biological Sciences, Pullman, WA, USA; jesse.brunner@wsu.edu

* Correspondence: joseph.mihaljevic@nau.edu; Tel.: 1-928-523-5125

Abstract: Mechanistic models are critical for our understanding of both within-host dynamics (i.e., pathogen population growth and immune system processes) and among-host dynamics (i.e., transmission). Rarely, however, have within-host models been synthesized with data to infer processes, validate hypotheses, or generate new theories. In this study we use mechanistic models and empirical, time-series data of viral titer to better understand the growth of ranaviruses within their amphibian hosts and the immune dynamics that limit viral replication. Specifically, we fit a suite of potential models to our data, where each model represents a hypothesis about the interactions between viral growth and immune defense. Through formal model comparison, we find a parsimonious model that captures key features of our time-series data: the viral titer rises and falls through time, likely due to an immune system response, and that the initial viral dosage affects both the peak viral titer and the timing of the peak. Importantly, our model makes several predictions, including the existence of long-term viral infections, that can be validated in future studies.

Keywords: Amphibian; *Ranavirus*; frog virus 3, Mathematical models, Bayesian inference

1. Introduction

Understanding the specific interactions between a host's immune system and an infectious agent can be critical for understanding population-level patterns of infection prevalence and the dynamics of prevalence through time [1,2]. Mechanistic models of within-host dynamics use mathematical expressions to represent the processes of pathogen population growth and immune defenses that fight against pathogen growth. These types of models have revealed key insights, especially regarding the effects of various immune strategies on the evolution of pathogen virulence [3,4]. Within-host models also help us to understand the effects of chronic and acute infections on disease progression, pathogen evolution, and the links between within-host and among-host processes [5]. For example, Kennedy et al. [6] showed that models of within-host viral growth and immune processes accurately recapitulate the effects of viral exposure dose on the probability of death in an insect host. Here we develop within-host models to understand the growth of ranaviruses in their amphibian hosts, and we compare our models' expectations to laboratory data.

Viruses in the genus *Ranavirus* can be impressively lethal, causing rapid host mortality and mass mortality events, especially in larval amphibians. However, the outcome of infection varies tremendously across host taxa [7,8] and life stage [9–11], as well as dose of exposure and numerous environmental factors (reviewed in [12]). Much of this variation is presumably due to variation in the host-virus interaction, where viral replication is countered by immune responses. Indeed, some qualitatively distinct outcomes appear to have their basis in different immune responses. Adult

Xenopus laevis, for instance, are refractory to ranavirus infections while larvae often succumb, and
this is due to their distinct immune responses [13,14]. The much more effective adult response to
infection involves rapid, substantial CD8+ cellular responses, whereas tadpoles' responses are
dramatically less effective and slower. Larval *X. laevis* produce type I and type III interferon
responses, which are effective at reducing viral replication, but are eventually overwhelmed by the
virus [15,16]. However, while there has been great progress in elucidating the components of the
immune system that interact with and help control ranavirus, many questions remain.
In particular, the dynamics of viral population growth are largely unknown outside of cell
culture experiments. What are the dynamics of the virus population within the host? Does the host's
immune system effectively control the virus population, outside of the *Xenopus* model system? Does
the immune response lead to viral clearance or can persistent infections remain? Does this outcome
depend on the initial exposure dose? We propose that mathematical models of the within-host
dynamics of ranaviruses, which provide a quantitative understanding of the host-virus interaction,
will help answer these questions and will provide a means for integrating the numerous factors that
appear to be important for the outcome of ranavirus infections. There are several additional
advantages of using a mechanistic, model-based approach. First, parameterized disease models yield
quantitative insights that can be validated by collecting new data. Second, we can simulate
parameterized models under novel scenarios to predict how the system might behave. Third, we can
use models to identify which novel sources of data would best inform our understanding under
different scenarios. Thus, our approach offers a rigorous method to generate new hypotheses and
guide future studies to understand the interactions between ranaviruses and host immune defenses.
To this end we set out to develop several versions of a mechanistic, within-host model that
embodied hypotheses about the growth of frog virus 3 (FV3) and the host's immune response to the
infection. We then fit these models to a time series of FV3 titers within bullfrog (*Lithobates catesbeianus*)
tadpoles that were experimentally infected with FV3. We used a model-comparison approach to
determine the most parsimonious model(s) that could describe the observed viral titer through time
and the effects of viral exposure dose. The salient features of these dynamics are: 1) a clear rise and
then slower decline in viral titers over time, suggesting viral clearance by the host's immune response
and motivating our model-fitting exercise, 2) an earlier and higher peak in viral titers with higher
exposure dose, and 3) low-level infections remaining detectable for seven weeks. We then use these
models to generate testable hypotheses and guide future studies to understand the interactions
between ranaviruses and host immune defenses.

2. Materials and Methods

Our goal is to test a set of hypotheses about how ranaviral populations grow within their larval
amphibian hosts and how the virus interacts with the larval host's immune system. Our strategy is
to develop mechanistic within-host growth models with various assumptions that correspond to
our hypotheses. We then attempt to fit these models to serially-collected data on within-host titers,
and we use a model-comparison approach to determine the most parsimonious model structure,
and therefore the most parsimonious support for our hypotheses.

2.1 Experimental assessment of within-host ranaviral dynamics

Details of the experiment can be found in Brunner et al. [17], but the essential details are that
tadpoles (Gosner [18] stages 25–28) were exposed in 400 mL water baths for 24 h to one of three
doses of the ranavirus FV3: a high dose (10^5 plaque-forming units [pfu] mL^{-1}; n=150), medium-
dose (10^3 pfu mL^{-1}; n=150), or a low, but unknown dose (n=90 in 'mock' exposure to inadvertently
contaminated cell culture media). The tadpoles were assigned to be euthanized and sampled on
days 2, 4, 6, 8, 14, 21, 28, 35, 42, and 49 post exposure (n=15 high-dose, n=15 low-dose, and n=9
'mock' tadpoles per time point), although some died and were sampled before their assigned date.
Viral titers in liver and kidney samples were measured with a quantitative real-time PCR assay [19].

We fit models to these time-series data that embodied two assumptions about viral growth and two
about the host's immune response. First, we fit models with exponential viral growth ($V' = \phi V$) or
with logistic growth [$V' = \phi V \left(1 - \frac{V}{K}\right)$]. In these differential equations, V' represents $\frac{dV(t)}{dt}$, where V
is the density of virus within the host, ϕ is the per capita growth rate of the virus, and K is the
viral carrying capacity. In some models, K can be viewed as the viral titer above which the virus
kills the host [5]; however, because we are modeling mean-field dynamics, we do not explicitly
account for virus-induced host mortality.

We then layer on possible immune system dynamics. From previous experimental work, we know
that larval amphibians (at least *Xenopus*) show interferon type I and type III responses to FV3,
although the latter appears more effective [15,16]. While we lack detailed knowledge of the immune
response to FV3, especially in non-model amphibian species, we begin with the reasonable
assumption that the growth of the immune components, Z, to viral infection follows the Michaelis-
Menten form of enzymatic activity [20,21]:

$$Z' = \psi Z \left(\frac{V}{V + \gamma}\right). \tag{1}$$

109   In this case, the growth rate of the immune component ramps up to a maximum, ψZ, as the virus
110   population increases. The rate of immune growth is mediated by γ, which is the half-saturation
111   constant. In this formulation the density of immune components never returns to pre-infection
112   levels. We therefore also considered a second formulation

$$Z' = (N_Z - \delta Z) + \psi Z \left(\frac{V}{V + \gamma}\right), \tag{2}$$

114   where we define $N_Z = \delta Z(0)$, such that, when no virus is present, the immune system stabilizes to
115   its homeostatic level of immune component density equal to $Z(0)$.

116   Finally, we assume a mass-action attack rate of the immune system against the virus, which is a
117   Type I Hollings' functional response, such that our most complex model becomes:

$$V' = \phi V \left(1 - \frac{V}{K}\right) - \beta V Z \tag{3.1}$$

$$Z' = (N_Z - \delta Z) + \psi Z \left(\frac{V}{V + \gamma}\right). \tag{3.2}$$

Note that all of our models are therefore nested, and Table 1 shows all model formulations that we
fit to our data set. We divide the models into two classes, A and B, which are differentiated by the
assumptions about the immune system's growth dynamics. Class A models are mostly
distinguished by the fact that, at equilibrium, if the immune system response is strong enough, the
virus is driven extinct (i.e., full viral clearance). For Class B models, the virus is not driven extinct,
such that the virus persists at low levels within the hosts (Appendix A).

On a less technical note, in some ways our models are not specific about how the immune system of
the larval amphibian functions. For example, our model could equally represent a population of
immune cells or a population of molecules generated by the immune system (e.g., interferon
molecules) that act to control viral growth more or less indirectly. In other words, although we
model a mass-action attack rate of the immune system against the virus, we do not necessarily
mean that the virus and immune component must come into direct contact. This immune action
could be indirect, for instance, by controlling apoptosis of virus-infected cells.

Table 1. Model structures and model comparisons. The bolded model (B2) is the most parsimonious based on LOO-IC selection.

Class	ID	Structure	Notes	Penalty (pLOO)	LOO-IC	ΔLOO-IC
A	A1	$V' = \phi V - \beta VZ$ $Z' = \psi Z \frac{V}{(V+\gamma)}$	Drives virus extinct. Z goes to equilibrium ($Z_{(\infty)}$), which is above $Z_{(0)}$.	8.6	1121.6	29.0
	A2	$V' = \phi V \left(1-\frac{V}{K}\right) - \beta VZ$ $Z' = \psi Z \frac{V}{(V+\gamma)}$	Conditions under which virus goes to carrying capacity. Or virus goes extinct.	7.6	1151.9	59.3
B	B1	$V' = \phi V - \beta VZ$ $Z' = (N_Z - \delta Z) + \psi ZV$	Damped oscillations to a stable point equilibrium, where virus is persistent in host. The model fit shows several oscillations before equilibrium.	10.8	1144.5	51.9
	B2	$V' = \phi V - \beta VZ$ $Z' = (N_Z - \delta Z) + \psi Z \frac{V}{(V+\gamma)}$	Spike in viral load, then decline to stable point equilibrium, where virus is persistent in host.	**10.0**	**1092.6**	**0**
	B3	$V' = \phi V \left(1-\frac{V}{K}\right) - \beta VZ$ $Z' = (N_Z - \delta Z) + \psi Z \frac{V}{(V+\gamma)}$	Over-fitting. Extra parameter (carrying capacity, K) unnecessary.	10.0	1092.9	0.3

We employ a Bayesian approach to inference to fit the suite of dynamical models to the time-series
of within-host viral titers. Then we use Bayesian information criteria to compare the within-sample
predictive accuracy of our models, while penalizing model complexity. In this way, we seek to
understand the most parsimonious model structure that captures the main features of the time
series. All model-fitting code is available on our open-source Bitbucket repository
(https://bitbucket.org/jrmihalj/ranavirus-within-host-dynamics/src/master/).

We use the open-source statistical programming language, *Stan* [22], to fit our differential equation
models to the time-series data. *Stan* employs a Hamiltonian Monte Carlo (HMC) algorithm to
sample from the model's posterior. For each model, we sampled the posterior using three Markov
chains, with a 2000 iteration warm-up, followed by 3000 iterations, thinned by 3, for a total of 1000
samples recorded from each chain. We conducted various graphical and quantitative diagnoses of
the chain behavior, including inspections for temporal auto-correlation. We assessed convergence
with the Gelman-Rubin statistic, $\widehat{R}$ [23]. All models converged after 5000 iterations.

Model dynamics are sensitive to parameter magnitudes. Therefore, in constructing our prior
probability distributions for the parameters, we often restricted parameters to vague, but realistic
ranges (see open-source model statements). All parameters were restricted to positive values, and
for parameters that could take very large values (e.g., K) we set the parameter on a natural-
logarithmic scale.

Besides estimating the model parameters, we also had to estimate the initial conditions of the
system: the initial viral titer, $V(0)$, and the initial immune component density, $Z(0)$. Because we
have three viral dosage treatments, we estimated three independent $V(0)$. We used a hierarchical
prior structure, such that each treatment's initial titer was drawn from a distribution with an
estimated mean and variance (see model code). We further assumed that the initial immune
component density, $Z(0)$, did not vary among treatments. We therefore assumed that all larval
amphibians start with the same average immune system component density that then responds to
the invasion of the virus. Because the magnitude of $V(0)$ and $Z(0)$ could be large, we again set
these parameters on a natural-logarithmic scale.

The viral titer values spanned approximately 6 orders of magnitude. To improve numerical
integration performance, by reducing the error in the integration algorithms due to such large viral
titer values, we fit the differential equation models using natural log-transformed titer. Assuming
that $v = log(V)$ and $z = log(Z)$, we can reformulate Equation 3 as follows:

168
$$v' = \phi\left(1 - \frac{e^v}{K}\right) - \beta e^z \qquad (4.1)$$

169
$$z' = \left(\frac{N_z}{e^z} - \delta\right) + \psi\left(\frac{e^v}{e^v + \gamma}\right). \qquad (4.2)$$

Thus, the scaling of the parameters does not change, just the scaling of the state variables. This also
allowed us to more reliably assume a Gaussian likelihood for log-transformed viral titer. We further
allowed the residual variance to be different for each dosage treatment.

We also note that we only included non-zero titers in our data set for the model-fitting routine. Our
reasoning is that a zero titer could represent an individual that never became infected, even upon
exposure, or an individual that cleared infection later on. Because we could not distinguish between
these two scenarios, we excluded these values. We did re-fit our models to the full data set,
including zeros, to understand the effects on our inference. Importantly, this did not change the

rank-order of our models using model comparison, but it did generally make the models fit more
poorly to the data, bringing down the average predicted titer and altering the marginal posterior
estimates of the model parameters. We therefore do not consider these model fits further.

2.4 Model comparisons

Our goal for model comparison was to determine the most parsimonious model(s) with the best
goodness-of-fit. In practice this means assessing the within-sample predictive performance of each
model and correcting for model complexity. Given that we used a Bayesian approach to fitting our
models to the data, we chose to use a Bayesian information theoretic approach to model
comparison. Therefore, we calculated the leave-one-out information criterion (LOO-IC) for each
model. Analogous to the Akaike information criterion (AIC), we can compare models based on the
LOO-IC value, following the rule-of-thumb that ΔLOO-IC> **3** designates models with significantly
different model performance, and where lower LOO-IC values indicate better-fitting, more
parsimonious models.

3. Results

Figure 1. The fit of model B2 to the experimental data. Circles are data points representing the viral
DNA copies from individual bullfrog tadpoles that were sampled on a given day (closed circles) or
that died prior to their assigned sampling day (open circles). Also shown are the median model fit
(solid red line) and 95% Bayesian credibe interval of the fit (dashed red lines).

A full qualitative and statistical analysis of the experimental data can be found in [17]. For our
purposes, it is important to note the clear rise and fall of viral titers over time in the medium and
high dosage treatments, which suggested a possible immune response that was leading towards
viral clearance, and which motivated our modeling-fitting and model comparison routine.

Model B2, our best-fitting and most parsimonious model, fits to the experimental data very well
(Figure 1). This model includes exponential growth of the virus, and an assumption of homeostatic
levels of immune components within the host, and the model's parameter estimates are shown in
Table 2. Importantly, model B2 captures key features of the data that the other models failed to
capture. First, the model correctly predicts that higher initial viral doses lead to higher peaks of
within-host titer. Second, the model also shows that with lower initial doses, the timing of the peak
titer is delayed. In other words, it takes longer for the virus population to build up within the host.
This is due, in part, to the exponential growth of the virus and, in part, to the action of the immune
system that inhibits viral growth.

Our model-fitting and model comparison approach suggests strong differences between the
appropriateness of our different model structures. Models B2 and B3 had indistinguishable model

fits (Table 1, Figure S4), but model B3 included an additional parameter, assuming logistic growth
of the virus population (i.e., a carrying capacity within the host). This is a classic case of over-fitting,
however, because the within-sample prediction for the model with more parameters was
indistinguishable from the simpler model. Therefore, we conclude that model B2 is more
parsimonious.

Table 2. Parameter estimates (median and 95% credible intervals) from the most parsimonious model, B2.

Parameter	Description	Units	Estimate
$V(0)$ low			0.15 (0.01 – 0.96)
$V(0)$ medium	Initial viral densities (per dosage)	Viral DNA copy (VC)	4.02 (0.44 – 25.42)
$V(0)$ high			180.63 (26.50 – 934.05)
$Z(0)$	Initial immune component denisty	Immune component (IC)	0.65 (0.09 – 14.93)
ϕ	Viral growth rate	day^{-1}	1.71 (0.85 – 3.16)
β	Mass-action attack rate	(IC)$^{-1}$ day^{-1}	0.67 (0.04 – 2.29)
N_Z	Rate of growth that ensures return of immune system to homeostasis	(IC) day^{-1}	$N_Z = \delta Z(0)$
δ	Rate of decline that ensures return of immune system to homeostasis	day^{-1}	0.84 (0.33 – 1.82)
ψ	Immune component growth rate in response to virus	day^{-1}	0.67 (0.18 – 1.62)
γ	Half saturation constant	VC	0.16 (0.03 – 0.69)

We had trouble fitting model A1 to the data, because it fits so poorly to the low-dose treatment.
Therefore, the model ends up predicting a more or less linear decline in virus load across time, with
high variance (Figure S1). This linear effect allowed the model to fit very well to the data from the
low dose treatment, increasing its overall likelihood and improving its LOO-IC value. However,
this parameterized version of the model fails to capture basic features of the data set. In contrast,
model A2 fits to the data reasonably well (Figure S2). However, this model predicts the same peak
titer across dosage treatments, leading to a very poor fit to the data from the low-dose treatment.

Model B1 also has a decent fit to the data (Figure S3). However, with this model's estimated
parameters, the dynamics show damped oscillations towards the endemic equilibrium. This pattern
does not seem biologically realistic, as the data do not show any clear oscillations.

4. Discussion

Our study represents an initial step towards making model-based, quantitative predictions about ranavirus growth and immune system functioning within larval amphibian hosts, which complements the history of empirical work in these research areas. The experimental data with bullfrog (*Lithobates catesbeianus*) tadpoles suggests a non-monotonic relationship between within-host viral titer and time post-exposure. That is, an immune response appears to be reducing the growth of the virus over time. This is broadly consistent with the immunological studies with *Xenopus laevis* tadpoles, showing a robust interferon response [15,16]. Through model comparisons, we identified a model of exponential viral growth and a mass-action immune system response that captures key features of our data and generally fits our data quite well. Importantly, this model-based approach allows us to make predictions and to develop testable hypotheses for follow-up experiments.

The best-fitting model predicts that individuals that survive infection should exhibit a viral titer pattern that rises and falls over time. This prediction contrasts with the empirical work that uses *Xenopus*, because larval *Xenopus* almost always die from frog virus 3 (FV3) exposure. Even a pre-treatment of type III interferon, which should boost the immune response, only delays mortality in the *Xenopus* system [16]. However, it is well known that the American bullfrog is uncommon in its high resistance and tolerance to multiple diseases, including FV3 [7,24] and the amphibian chytrid fungus (*Batrachochytrium dendrobatidis*) [25,26]. Therefore, it is not surprising that we could infer a more robust immune response from this species compared to *X. laevis*. Still, our current data cannot validate that, in surviving individuals, virus loads rise and then fall, because we do not have individual host-level time series of viral load. Future studies could use various sources of eDNA (e.g., swabbing, skin scrapes, or water filtration), which are correlated with internal titers [17], to validate whether there is a rise and fall of viral load in individuals. Alternatively, it may be fruitful to experiment with larger-bodied animals that would permit sampling blood repeatedly through time.

Our model also predicts that, if a host does not die from the initial exponential growth of the virus, the immune system will decrease the virus to an endemic equilibrium (Appendix A). In other words, we should see persistent infections with very low levels of virus in some infected bullfrog tadpoles. It is worth noting that this model prediction is not relevant to quiescent virus persisting in particular tissues, such as that found in peritoneal leukocytes in adult *Xenopus* [27]. Rather, the persistence in our model is due to a dynamic balance between viral replication and host immune responses. In any case, our model predicts low-level infections that would be detectable beyond 60 days from initial exposure time and should be actively replicating. That said, there are also several reasons that an endemic equilibrium may not hold. First, demographic stochasticity in the virus population could cause low-level infections to fade out [6,28]. Second, variability in the immune response over time (e.g. waning immunity) could cause fluctuations in viral titer. Third, the costs of continued immune responses could lead to host death even with low levels of infection [29].

Several of the tested model structures predict full viral clearance at equilibrium, but none of these models fit as well to the observed dynamics in bullfrog tadpoles. Future studies that collect time-series data and that have some long-term exposures (e.g., greater than 50 days) will improve our ability to distinguish between viral clearance and viral persistence. Furthermore, because we did not have simultaneous empirical data from the larval immune system, our model fitting algorithm was agnostic to the model's prediction of immune system dynamics. If future studies simultaneously capture time-series data on viral titer and measures of larval immune system, we would have much more power to distinguish between different models of within-host dynamics.

There are key features of the data set that our deterministic modeling scheme does not fully capture, and these are areas for future model development. First, the main process that our model ignores is host mortality due to viral growth. For instance, our current model predicts that immunocompromised individuals would show exponential viral growth, and in reality, this likely

lead to death. Future versions of our model could include stochasticity, following Kennedy and
Dwyer [6], such that demographic stochasticity in the viral population could lead to the virus
exceeding a threshold growth rate or titer that causes host death. Then, we could analyze the
probability of this occurring over many realizations of the stochastic model. This would allow us to
explicitly evaluate how initial exposure dosage affects the probability of host mortality, from a
within-host, mechanistic framework. Relatedly, the model also does not fit well to the initial data at
high viral dosage. It predicts a higher initial viral dosage than the data would suggest. This is a
peculiar issue; however, it could be the case that if we added stochasticity in viral growth, and/or
stochasticity in the initial immune response, we may better capture these initial dynamics.

Viral titers within a host can be important determinants of the infection dynamics among hosts [3],
affecting both the duration of infections and their propensity to be transmitted. Many dose-response
experiments, for instance, show that hosts exposed to higher doses of ranaviruses, and so presumably
harbor larger viral populations, are more likely to die and die more rapidly [30–32]. More intensely
infected individuals may also shed more virus (e.g., [17]) and thus be more infectious. Similar effects
of the intensity of infection are common in other host-virus systems [33,34], as well as for other
amphibian pathogens (e.g., *B. dendrobatidis*; [35]). Pathogen titers or intensity of infection can be a
natural way in which to link within and between-host dynamics [3,36,37]. Within-host models of
host-virus interactions may therefore be a fruitful way of integrating the myriad effects that are
known to influence the outcome of ranavirus infections.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1, Supplement1.docx,
which includes Figures S1 – S4.

Author Contributions: Conceptualization, JRM, ALG, JLB.; methodology, JRM, ALG, JLB; software, JRM;
validation, JRM; formal analysis, JRM; investigation, JRM, ALG, JLB; data curation, JLB; writing—original draft
preparation, JRM, JLB; writing—review and editing, JRM, ALG, JLB; visualization, JRM; supervision, JRM, ALG,
JLB; project administration, JRM, ALG, JLB.

Funding: JRM was funded under the State of Arizona Technology and Research Initiative Fund (TRIF),
administered by the Arizona Board of Regents, through Northern Arizona University. JLB was supported by
National Science Foundation grant 1754474. Data collection were supported by Association of Zoo & Aquariums
CGF grant 14-1263 and a Zoological Medicine & Wildlife Health Research Grant from the American Association
of Zoo Veterinarians to JLB. ALG was supported by the Canada Research Chairs Program and the Natural
Sciences and Engineering Research Council (NSERC) of Canada.

Acknowledgments:

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the
study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to
publish the results.

Appendix A

The coexistence equilibrium of model B2, which analytically expresses the amount of virus and host
immune components at the stable equilibrium with a persistent infection, is as follows:

$$Z^* = \frac{\phi}{\beta} \qquad (A1.1)$$

$$V^* = \frac{\frac{\gamma}{\psi}\left(\delta - \frac{N_Z}{Z^*}\right)}{\left(1 - \frac{\delta}{\psi} + \frac{N_Z}{\psi Z^*}\right)} \qquad (A1.2)$$

References

1. Hawley, D.M.; Altizer, S.M. Disease ecology meets ecological immunology: Understanding the links between organismal immunity and infection dynamics in natural populations. *Funct. Ecol.* **2011**, *25*, 48–60.

2. Ezenwa, V.O.; Jolles, A.E. From host immunity to pathogen invasion: the effects of helminth coinfection on the dynamics of microparasites. *Integr. Comp. Biol.* **2011**, *51*, 540–51.

3. Mideo, N.; Alizon, S.; Day, T. Linking within- and between-host dynamics in the evolutionary epidemiology of infectious diseases. *Trends Ecol. Evol.* **2008**, *23*, 511–7.

4. Day, T.; Alizon, S.; Mideo, N. Bridging scales in the evolution of infectious disease life histories: Theory. *Evolution (N. Y).* **2011**, *65*, 3448–3461.

5. Alizon, S.; van Baalen, M. Acute or Chronic? Within-Host Models with Immune Dynamics, Infection Outcome, and Parasite Evolution. *Am. Nat.* **2008**, *172*, E244–E256.

6. Kennedy, D. a.; Dukic, V.; Dwyer, G. Pathogen Growth in Insect Hosts: Inferring the Importance of Different Mechanisms Using Stochastic Models and Response-Time Data. *Am. Nat.* **2014**, *184*, 407–423.

7. Hoverman, J.; Gray, M.; Miller, D. Anuran susceptibilities to ranaviruses: role of species identity, exposure route, and a novel virus isolate. *Dis. Aquat. Organ.* **2010**, *89*, 97–107.

8. Duffus, A.L.J.; Waltzek, T.B.; Stöhr, A.C.; Allender, M.C.; Gotesman, M.; Whittington, R.J.; Hick, P.; Hines, M.K.; Marschang, R.E. Distribution and Host Range of Ranaviruses. In *Ranaviruses*; 2015; pp. 9–57 ISBN 978-3-319-13754-4.

9. Hoverman, J.T.; Gray, M.J.; Haislip, N. a; Miller, D.L. Phylogeny, life history, and ecology contribute to differences in amphibian susceptibility to ranaviruses. *Ecohealth* **2011**, *8*, 301–19.

10. Haislip, N. a; Gray, M.J.; Hoverman, J.T.; Miller, D.L. Development and disease: how susceptibility to an emerging pathogen changes through anuran development. *PLoS One* **2011**, *6*, e22307.

11. Warne, R.W.; Crespi, E.J.; Brunner, J.L. Escape from the pond: Stress and developmental responses to ranavirus infection in wood frog tadpoles. *Funct. Ecol.* **2011**, *25*, 139–146.

12. Brunner, J.L.; Storfer, A.; Gray, M.J.; Hoverman, J.T. Ranavirus Ecology and Evolution: From Epidemiology to Extinction. In *Ranaviruses*; 2015; pp. 71–104.

13. De Jesús Andino, F.; Chen, G.; Li, Z.; Grayfer, L.; Robert, J. Susceptibility of Xenopus laevis tadpoles to infection by the ranavirus Frog-Virus 3 correlates with a reduced and delayed innate immune response in comparison with adult frogs. *Virology* **2012**, *432*, 435–43.

14. Wendel, E.S.; Yaparla, A.; Koubourli, D. V.; Grayfer, L. Amphibian (Xenopus laevis) tadpoles and adult frogs mount distinct interferon responses to the Frog Virus 3 ranavirus. *Virology* **2017**, *503*, 12–20.

15. Grayfer, L.; De Jesus Andino, F.; Robert, J. The Amphibian (Xenopus laevis) Type I Interferon Response

to Frog Virus 3: New Insight into Ranavirus Pathogenicity. *J. Virol.* **2014**, *88*, 5766–5777.

16.   Grayfer, L.; De Jesús Andino, F.; Robert, J. Prominent Amphibian (Xenopus laevis) Tadpole Type III
Interferon Response to the Frog Virus 3 Ranavirus. *J. Virol.* **2015**, *89*, 5072–5082.

17.   Brunner, J.L.; Olson, A.D.; Rice, J.G.; Meiners, S.E.; Le Sage, M.J.; Cundiff, J.A.; Goldberg, C.S.; Pessier,
A.P. Ranavirus infection dynamics and shedding in American bullfrogs: consequences for spread and
detection in trade. *In Review.*

18.   Gosner, K.L. A Simplified Table for Staging Anuran Embryos Larvae with Notes on Identification.
Herpetologica **1960**, *16*, 183–190.

19.   Stilwell, N.; Whittington, R.J.; Hick, P.; Becker, J.A.; Ariel, E.; van Beurden, S.; Vendramin, N.; Olesen,
N.; Waltz Partial validation of a TaqMan real-time quantitative PCR for the detection of ranaviruses. *Dis.*
Aquat. Organ. **2018**, *128*, 105–116.

20.   Menten, L.; Michaelis, M.I. Die kinetik der invertinwirkung. *Biochem Z* **1913**, *49*, 5.

21.   Rosenbaum, B.; Rall, B.C. Fitting functional responses: Direct parameter estimation by simulating
differential equations. *Methods Ecol. Evol.* **2018**, *9*, 2076–2090.

22.   Carpenter, B.; Gelman, A.; Hoffman, M.D.; Lee, D.; Goodrich, B.; Betancourt, M.; Brubaker, M.; Guo, J.;
Li, P.; Riddell, A. *Stan* : A Probabilistic Programming Language. *J. Stat. Softw.* **2017**, *76*.

23.   Gelman, A. Inference and monitoring convergence. In *Markov Chain Monte Carlo in Practice*; Gilks, W.,
Richardson, S., Spiegelhalter, D., Eds.; CRC Press, 1996; pp. 131–143.

24.   Majji, S.; Lapatra, S.; Long, S.M.; Sample, R.; Bryan, L.; Sinning, A.; Chinchar, V.G. Rana catesbeiana
virus Z (RCV-Z): a novel pathogenic ranavirus. *Dis. Aquat. Organ.* **2006**, *73*, 1–11.

25.   Daszak, P.; Strieby, A.; Cunningham, A.A.; Longcore, J.E.; Brown, C.C.; Porter, A.D. Experimental
Evidence That the Bullfrog (Rana Catesbeiana) Is a Potential Carrier of Chytridiomycosis, an Emerging
Fungal Disease of Amphibians. *Herpetol. J.* **2004**, *14*, 201–207.

26.   Peterson, A.C.; McKenzie, V.J. Investigating differences across host species and scales to explain the
distribution of the amphibian pathogen Batrachochytrium dendrobatidis. *PLoS One* **2014**, *9*.

27.   Morales, H.D.; Abramowitz, L.; Gertz, J.; Sowa, J.; Vogel, A.; Robert, J. Innate immune responses and
permissiveness to ranavirus infection of peritoneal leukocytes in the frog Xenopus laevis. *J Virol* **2010**,
84, 4912–4922.

28.   Kennedy, D.A.; Dwyer, G. Effects of multiple sources of genetic drift on pathogen variation within hosts.
PLoS Biol. **2018**, *16*, 1–17.

29.   Hanssen, S.A.; Hasselquist, D.; Folstad, I.; Erikstad, K.E. Costs of immunity: Immune responsiveness
reduces survival in a vertebrate. *Proc. R. Soc. B Biol. Sci.* **2004**, *271*, 925–930.

30. Plumb, J.A.; Zilberg, D. The lethal dose of largemouth bass virus in juvenile largemouth bass and the
comparative susceptibility of striped bass. *J Aquat Anim Heal.* **1999**, *11*, 246–252.

31. Brunner, J.L.; Richards, K.; Collins, J.P. Dose and host characteristics influence virulence of ranavirus
infections. *Oecologia* **2005**, *144*, 399–406.

32. Duffus, A.L.J.; Nichols, R.A.; Garner, T.W.J. Experimental evidence in support of single host
maintenance of a multihost pathogen. *Ecosphere* **2014**, *5*, article 142.

33. Cattadori, I.M.; Albert, R.; Boag, B. Variation in host susceptibility and infectiousness generated by co-
infection: the myxoma-Trichostrongylus retortaeformis case in wild rabbits. *J R Soc Interface* **2007**, *4*, 831–
840.

34. Lowen, A.C.; Mubarka, S.; Steel, J.; Palese, P. Influenza Virus Transmission Is Dependent on Relative
Humidity and Temperature. *PLoS Pathog.* **2007**, *3*, e151.

35. Wilber, M.Q.; Weinstein, S.B.; Briggs, C.J. Detecting and quantifying parasite-induced host mortality
from intensity data: method comparisons and limitations. *Int J Parasitol* **2016**, *46*, 59–66.

36. Gog, J.R.; Pellis, L.; Wood, J.L.N.; McLean, A.R.; Arinaminpathy, N.; Lloyd-Smith, J.O. Seven challenges
in modeling pathogen dynamics within-host and across scales. *Epidemics* **2014**, *10*, 45–48.

37. Metcalf, C.J.; Graham, A.L.; Martinez-Bakker, M.; Childs, D.Z. Opportunities and challenges of Integral
Projection Models for modelling host-parasite dynamics. *J Anim Ecol* **2016**, *85*, 343–355.

402