1　**Title: BCL6 is required for the development of functionally responsive IgM+**

2　**GC-independent Memory B Cells**

Authors: Gretchen Harms Pritchard[1], Akshay T. Krishnamurty[1], Lauren Rodda[1],

Christopher Thouvenel[2], Derek J. Bangs[1], Courtney E. McDougal[1], David J.

Rawlings[2], Marion Pepper[1]*

Affiliations: [1]Department of Immunology, University of Washington School of

Medicine, Seattle, WA, 98109; USA; [2]Center for Immunity and Immunotherapies,

Seattle Children's Research Institute, Seattle, WA 98101, USA

22　*Corresponding author. mpepper@uw.edu

Abstract:

Humoral immunity depends upon long-lived, antibody-secreting plasma cells and memory B cells (MBCs). MBCs exhibit significant phenotypic and functional heterogeneity shaped by interactions with CD4+ T cells. It is currently unclear how specific CD4+ T cell interactions with B cells influence specific MBC subset generation. We used genetic ablation and antibody depletion to dissect key CD4+ T cell/B cell receptor-ligand pair interactions to define the critical signals that govern the development of specific MBC populations. While it has previously been suggested that germinal center (GC) experience is required for the development of CD73+CD80+ MBCs that rapidly form secondary plasmablasts, highly functional IgM+ MBCs differentiate in a BCL6 and CD4+ T cell-dependent, but GC-independent manner. BCL6 upregulation, in the presence or absence of a GC, can therefore serve as a predictor of long-lived, functional MBCs.

One Sentence Summary: Differential requirements of BCL6 and Tfh cells lead to the generation of functionally distinct memory B cell populations.

Keywords: Humoral immunity, Memory B cell, Malaria, BCL6, Germinal Center, Somatic Hypermutation

INTRODUCTION

Humoral immunity is mediated by long-lived plasma cells (LLPCs) and memory B cells (MBCs) differentiated from B cells activated through B cell receptor (BCR) binding to cognate antigens (Inoue and Kurosaki, 2024; Plotkin, 2010). Pre-existing antibodies secreted by LLPCs can provide immediate defense against invading pathogens while MBCs maintain their BCR expression and rapidly differentiate into antibody-secreting plasmablasts (PBs) upon rechallenge (Akkaya et al., 2020). Phenotypically and functionally diverse MBC populations can form in response to immunization or infection in mice and humans (Callahan et al., 2024; Krishnamurty et al., 2016; Price et al., 2021; Seifert et al., 2015; Weisel et al., 2016; Yates et al., 2013; Zuccarino-Catania et al., 2014). We previously demonstrated in a murine model of infection that both IgG+ and IgM+ CD73+CD80+ MBCs rapidly form PBs in response to rechallenge, with CD73+CD80+ IgM+ MBCs (referred to as DP IgM+) responding faster than CD73+CD80+ IgG+ MBCs (referred to as DP IgG+ or DP swIg+) (Krishnamurty et al., 2016). A third population of expanded CD73-CD80- IgD+ MBCs, are more naïve-like and thought to predominantly contribute to a subsequent GC response that can repopulate the MBC pool (Callahan et al., 2024; Dogan et al., 2009; Krishnamurty et al., 2016; Mesin et al., 2020; Zuccarino-Catania et al., 2014). It is currently unclear how these distinct functionally responsive MBC populations form and how the transcription factor BCL6 may be involved.

The current paradigm of MBC formation suggests that activated B cells undergo iterative and complex interactions with CD4+ T cells that influence their fate. CD4+ T cells and B cells first engage at the T-B border and interfollicular regions of the secondary lymphoid organs. Some of these activated B cells subsequently form low-affinity GC-independent MBCs (Taylor et al., 2012b), whereas other activated B cells continue to interact with CD4+ Tfh cells in the GC (GC Tfh), where BCL6 is expressed by both B cell and Tfh cell populations (Vinuesa et al., 2016). Within the GC, class switch recombination (CSR), BCR somatic hypermutation (SHM), and testing by CD4+ T cells leads to affinity

maturation and the downregulation of BCL6 and eventual exit of diversified class-
switched MBC and LLPC populations (Papa and Vinuesa, 2018). The distinct
signals that lead to the differentiation of specific MBC populations, particularly the
DP IgM+ MBCs that have undergone SHM but not CSR, are unknown. Using
various knockout mice and antibody-mediated temporal deletion, we tested how
specific CD4+ T cell interactions and transcriptional programs can influence MBC
fate. Our results demonstrate that both DP MBC populations require intrinsic
expression of BCL6, but only the majority of the DP swIg+ MBCs require GC Tfh
cells whereas an expanded population of DP IgM+ MBCs can form in the
absence of GC Tfh cells.
96
RESULTS and DISCUSSION
B cell-intrinsic BCL6 expression is required for the development of *P.ch*-
specific DP MBCs
As the hallmark transcription factor of the GC, BCL6 regulates many
aspects of GC B cell program, such as localization, DNA repair, and inhibition of
a plasma cell fate (Laidlaw and Cyster, 2021). To test the requirement of BCL6
for the development of DP MBC populations, we set up chimeras using
congenically disparate WT and B cell conditional BCL6 deficient (BCL6BKO)
bone marrow (see Methods) in order to create a system in which BCL6-sufficient
and -deficient B cells could respond to infection in the same inflammatory
environment. WT mice were infected with 10^6 *P.ch* infected red blood cells
(iRBCs) and the B cell response was examined at various time points thereafter.
Previously published techniques were used to examine rare antigen-specific B
cells specific for the carboxy terminus of the blood stage expressed merozoite
surface protein 1 (MSP1) (Krishnamurty et al., 2016). After infection with *P.ch*,
MSP1-specific BCL6BKO did not form GC B cells while MSP1-specific WT cells
did (Fig. S1A), confirming the important role for BCL6 in GC B cell generation
(Dent et al., 1997; Fukuda et al., 1997; Ye et al., 1997). The complete loss of GC

B cells among the BCL6BKO population was reflected in a decrease of the total
number of MSP1-specific B cells as well as MBCs (Fig. S1B).

Among the MBC populations at memory time points >40 days post-
infection, BCL6 deficiency resulted in a loss of IgM+ MBCs and swIg+ MBCs
(reduced by ~50 and ~98%, respectively), compared to WT MSP1-specific MBCs
(Fig. 1A). Analysis of the expression of CD73 and CD80 on each subset further
revealed a selective loss of the DP cells that accounted for a significant portion of
the IgM+ and swIg+ MBCs that were lacking in the BCL6BKO population (both
reduced by ~99%) (Fig. 1B). We also tested to see if cre toxicity (Becher et al.,
2018; Schmidt-Supprian and Rajewsky, 2007) was responsible for the defect that
we saw in the transgenic B cells, rather than loss of BCL6, by infecting WT mice
or Mb1cre+ mice with *P.ch.* We saw that the GC compartment was unaffected,
neither were DP MBC populations (Fig. S1C,D). Intrinsic BCL6 expression is
therefore critical for the formation of the GC and DP MBCs.

BCL6-dependent DP MBCs are required for rapid PB formation

As previously mentioned, BCL6 controls many aspects of the GC B cell
program including the negative regulation of several molecules associated with
MBC phenotype including CD80 and PD-L2 (Niu et al., 2003; Peng et al., 2019).
Therefore, while the preceding data demonstrated that BCL6BKO B cells do not
form DP MBCs, it was possible that we had altered the phenotype but not the
function of the MBCs. It was also possible that DP MBCs were not the only
source of novel antibody secreting B220+CD138+ PBs (Krishnamurty et al.,
2016). Thus, we compared functional responsiveness of WT and BCL6BKO cells
to rechallenge. WT:BCL6BKO bone marrow chimeric mice were infected, allowed
to form memory and subsequently rechallenged with *P.ch* at similar memory time
points that we had previously seen functional PB responses to rechallenge
(Krishnamurty et al., 2016). As expected, prior to rechallenge, WT B cells
consisted of two populations of memory cells: B220+CD138- MBCs and
B220^{low}CD138+ LLPCs. LLPCs did not form in the BCL6BKO population as

expected based on their GC-dependence (Fig. 1C) (Weisel et al., 2016). Similar

to our previous results, WT cells formed an expanded B220+CD138+ PB

population and an expanded LLPC population 3 days after rechallenge (Fig. 1C)

(Krishnamurty et al., 2016). Yet both populations were absent in the BCL6BKO

cells (Fig. 1C). B cell-intrinsic expression of BCL6 is therefore required for the

formation of DP MBCs and LLPCs, and in the absence of these populations, no

PBs are produced shortly after rechallenge with *P.ch*, again highlighting the

importance of the DP MBC populations in a secondary response to infection.

CD4+ T cells induce BCL6 expression in early activated B cells

While BCL6 is considered a "lineage defining" transcription factor of the GC, it is

expressed during multiple phases of B cell activation. Elegant studies by Okada

and colleagues using immunofluorescent microscopy of BCL6-YFP reporting B

cells demonstrated that antigen-activated B cells initially up-regulate BCL6 in a T

cell-dependent manner in the outer region of the follicle, before the GC has

formed (Kitano et al., 2011). Moreover, studies from the Haberman laboratory

identified perifollicular activated cells that express intermediate levels of BCL6

along with IRF4, and demonstrated that the cells that further upregulate BCL6

enter the GC whereas the cells that further upregulate IRF4 presumably form PBs

(Zhang et al., 2017). We therefore sought to better understand the kinetics of

BCL6 expression after infection, after excluding PBs from our analysis. WT mice

were infected with 10^6 *P.ch* infected red blood cells (iRBCs) and the infected-

induced B cell response was examined at various time points thereafter.

MSP1-specific cells did not express BCL6 until day 6 post-infection, at

which point IgD-BCL6+ cells began to accumulate (Fig. 2A). The majority of the

IgD- BCL6-expressing cells were IgM+ six days post-infection yet the proportion

of IgM+ BCL6+ cells decreased steadily and by day 20 post-infection BCL6+ cells

were almost exclusively swIg+ (Fig. S2A). During this early activation period,

CD73 and CD80 were both upregulated on IgD-BCL6- cells (Fig. 2B), suggesting

the emergence of memory precursor cells (Kuo et al., 2007; Laidlaw et al., 2017).

We next asked whether early MSP1+ B cell BCL6 expression is dependent
on CD4+ T cells, and if the early BCL6 expression drives the development of the
subsequent DP MBC formation. To answer these questions, we first initiated
treatment of mice with a CD4+ T cell-depleting antibody (clone GK1.5) one day
prior to infection and examined the B cell response at day 7 post-infection.
Depletion of CD4+ T cells prior to infection resulted in no MSP1 specific B cell
BCL6 expression at day 7 whereas ~25% of the CD138- B cells from the
untreated controls did (Fig. 3A). Additionally, while CD4+ T cell depletion
beginning on day 3 also impaired BCL6 expression, depletion at day 5 post-
infection did not affect BCL6 expression at day 7 post-infection (Fig. 3A).
Together with Figure 2A, these data demonstrate that BCL6 is upregulated by
day 6 post-infection, and interactions with CD4+ T cells within the first 3 days of
infection are essential for B cell upregulation of BCL6. Yet, when CD4+ T cells
are depleted beginning at day 5 post-infection, BCL6 expression is maintained,
indicating that delivery of the T cell signals inducing BCL6 upregulation have
been initiated by this time point.

In order to determine whether the CD4+ T cell dependent expression of
BCL6+ occurring within the first week of infection was sufficient for the
development of DP MBCs, or if CD4+ T cell interactions that occurred later than
day 7 also contributed to the B cell response, we depleted CD4+ T cells at
various time points of infection, and examined whether or not this led to distinct
memory populations. Depletion of CD4+ T cells prior to infection resulted in very
few IgM+ (mean=16.5, Fig. 3B) or swIg+ (mean=7, Fig. 3C) DP MBCs, reflecting
a >98% reduction compared to untreated controls. These data demonstrate that
CD4+ T cells are essential for the development of DP MBCs. While there was
also a reduction in the number of DP MBCs in mice that had CD4+ T cells
depleted beginning at all time points ≥ day 7 post-infection, the percentage of
cells displaying CD73 and CD80 was largely intact, with no significant decrease
in mice that received CD4+ T cell depleting antibody beginning at either day 14 or
day 21, in both IgM+ (Fig. 3B) and swIg+ (Fig. 3C) MBC populations. Together,

these data indicate that early after infection, CD4+ T cells deliver signals to B cells that initiate BCL6 upregulation and the subsequent development of DP MBCs, yet CD4+ T cell maintenance is required for the magnitude of the DP MBC populations.

GC Tfh cells are not required for the development of somatically hypermutated DP IgM+ MBCs

Our data demonstrated that early interactions with T cells leads to the generation of BCL6-expressing B cells. We next asked what type of T cells are important for the development of DP MBC populations, specifically whether GC Tfh cells are required. Using a transgenic strain of *Plasmodium yoelii* that expresses the GP66 glycoprotein from LCMV (*P.y-GP66*), thus enabling us to track an antigen-specific T cell response, we have previously reported that by day 4 post-infection an expanded population of CXCR5 expressing T cells was detectible. By day 8 post-infection a subset of these cells had begun to upregulate PD-1, indicative of the emergence of a GC Tfh cell population (Arroyo and Pepper, 2020). Because B cell activation and upregulation of BCL6 followed similar kinetics to T cell activation and GC Tfh generation, we next sought to understand whether the early T cell signals were provided by GC Tfh cells. In order to test this, we generated mice with *T cell* intrinsic ablation of BCL6 (BCL6TKO) (see Methods), which is essential for the formation of GC Tfh cells (Hollister et al., 2013).

At 8 days post-infection with *P.ch*, there were equal numbers of CD38+GL7+ activated B cells between the WT and BCL6TKO mice (Fig. 4A). Furthermore, BCL6 was upregulated in B cells from both WT and BCL6TKO mice, although to a slightly diminished degree in the BCL6TKO cells (Fig. 4B). The data indicate that non- GC Tfh CD4+ T cells are able to provide the early signals leading B cell activation and BCL6 upregulation. To confirm that these early T cell signals can be delivered outside the context of a GC, we performed histology on spleens from WT and BCL6TKO mice at days 0 (i.e. naïve) and 6-8

post-infection with *P.ch* (Fig. 4C). In order to determine whether or not T cells had

entered the B cell follicle we quantified size and T cell density of the early foci of

BCL6+ cells surrounded by IgD+ follicular B cells. While both WT and BCL6TKO

spleens contained early clusters within the follicle, they were smaller in the

BCL6TKO mice and, as expected, contained a paucity of CD3+ cells (Fig. 4C).

Together, these data demonstrate that while BCL6 expression within the T cell

compartment is necessary for the development of a CD4+ GC Tfh cell population,

B cells can become activated, express BCL6, and begin cluster coalescence in

the absence of such cells.

Previous studies have described B cell clustering in the absence of T cells;

however, these were in response to T-independent antigens and resulted in B

cells that had not undergone SHM (de Vinuesa et al., 2000; Lentz and Manser,

2001). Therefore, we next analyzed the DP MBC response to determine if the B

cells in the BCL6TKO mice were able to undergo SHM despite the absence of

GC Tfh cells. By day 28 post-infection with *P.ch*, there were far fewer MSP1-

specific B cells in BCL6TKO mice, and specifically, the GC B cell population was

abolished as expected (Fig. S3A). Among MBCs, the DP IgM+ MBC population

was intact yet there was a pronounced loss of DP swIg+ MBC in BCL6TKO mice

compared to WT mice (Fig. 4D). We next performed BCR sequencing of sorted

MBCs from *P.ch* infected WT and BCL6TKO mice and saw that BCRs from DP

IgM+ MBCs from both WT and BCL6TKO had undergone SHM. However, there

was a slight reduction in the level of SHM in the DP IgM+ MBCs from the

BCL6TKO mice (mean number of mutations = 2.05) compared to the DP IgM+

MBCs from the WT mice (mean number of mutations = 2.78) (Fig. 4E). We were

able to sort too few MSP1-specific DP swIg+ cells from the BCL6TKO mice

(n=10) to make conclusions about this population (Fig. S3C). Together, these

data demonstrate that the population of DP IgM+ MBCs can arise and undergo

SHM in the absence of GC Tfh cells.

To address whether DP IgM+ MBCs can also develop independently of GC

Tfh cells in other inflammatory settings or whether this phenotype was unique to

P.ch, we examined the formation of PE-specific MBCs in mice immunized with
PE protein in adjuvant. As seen with *P.ch* infection, the total number of PE-
specific DP IgM+ MBCs was not affected by the loss of the GC Tfh compartment
whereas the DP swIg+ MBCs were severely diminished (Fig. S3D). Together,
these data demonstrate that while both DP IgM+ and DP swIg+ MBCs have
undergone SHM, albeit to varying degrees, only the DP swIg+ MBCs require the
presence of GC Tfh cells in order to expand. DP IgM+ MBCs can expand and
undergo SHM outside of the GC reaction.

274   The GC is not required for rapid secondary PB formation

275   Having established that the expanded DP IgM+, but not DP swIg+, MBCs
can develop independently from the GC reaction, we next wanted to know
whether these cells are able to form secondary PBs. Our previous experiments
had compared the ability of different MBC populations to respond to secondary
challenge within individual mice, however genetic ablation of the GC Tfh
compartment necessitated comparing populations of cells in different groups of
mice. When we assessed parasitemia in WT and BCL6TKO we found that there
were no significant differences in parasitemia through day 30 post-infection,
consistent with previous results (data not shown) (Pérez-Mazliah et al., 2017).
However, to ensure that the PBs formed were induced by secondary infection
rather than any sub-patent parasites that remained in the blood at undetectable
levels, we treated WT and BCL6TKO mice with the antimalarial drug atovaquone
prior to rechallenge. At 3 days post-secondary challenge we examined secondary
PB formation. There was no significant difference between the ability of the WT
and the BCL6TKO mice to mount a secondary IgM+ PB response (Fig. 4F),
demonstrating the intact functionality of the GC Tfh-independent DP IgM+ MBCs.
We also confirmed the ability of the GC Tfh deficient mice to mount a secondary
PB response using the PE immunization setting and saw similar results (Fig.
S3E).

Our data provide evidence that prior to entering the follicle and establishing

GC formation, CD4+ T cells deliver activation signals that are sufficient to induce

BCL6 upregulation. In the WT:BCL6BKO mixed chimeras, BCL6 deletion in the B

cells results in ablation of both DP MBC populations. However, in mice without

GC Tfh cells (i.e. BCL6TKO mice), the DP IgM+ MBC population is intact while

the DP swIg+ MBC population is reduced, but not absent. Together, these data

demonstrate that development of both of these populations is dependent on BCL6

and is induced prior to GC formation, yet the population of mutated DP MBCs that

retain IgM expression bypasses the GC in its development.

The current model for the role of the GC in MBC development suggests

that only low affinity, unmutated DN MBCs can be generated independently of the

GC; in contrast, high affinity, mutated DP MBCs are dependent on the GC. Our

approach of ablating BCL6 in either the B cell compartment or the T cell

compartment enabled us to decouple the effects of intrinsic BCL6 expression and

GC experience on the development of MBC populations. These data suggest that

BCL6 has an essential function(s) in the development of MBC responses,

independent of its role in GC formation. Indeed, BCL6 has myriad functions

including regulation of B cell localization through repression of EBI-2, regulation of

additional co-stimulatory molecule expression and protection from DNA-damage

induced apoptosis (Kerfoot et al., 2011; Klein and Dalla-Favera, 2008; Pereira et

al., 2009; Ranuncolo et al., 2007). Without this protection, B cells are not able to

withstand DNA damage, such as that incurred during SHM and isotype class

switch recombination (CSR). The GC is often considered to be the sole location of

both of these functions (De Silva and Klein, 2015), yet in keeping with the work of

others, our results demonstrate that both can take place outside of the GC. For

example, Shlomchik and colleagues have reported that SHM can occur at the T

zone-red pulp border (William et al., 2002), and somatically hypermutated swIg+

PBs are formed in extrafollicular sites during infection with *Salmonella*

typhirmurium or *Ehrlichia muris* (Di Niro et al., 2015; Trivedi et al., 2019).

The studies presented here give us further insight into the intricacies of
endogenous MBC development after infection and protein immunization and
challenge current paradigms of MBC development.
326
MATERIALS AND METHODS

Mice

C57BL/6   (WT), B6.SJL-Ptprc[a] Pepc[b]/BoyJ (CD45.1+), and
B6.Cg-Tg(Cd4-cre)1Cwi/BfluJ (CD4-Cre+), mice were purchased from The
Jackson Laboratory. CD45.1+ mice were crossed to WT mice to generate
CD45.1+CD45.2+ mice. Mb1Cre+ mice were provided by Dr. Michael Reth (Max
Planck Institute of Immunobiology and Epigenetics) and crossed to Bcl6flx/flx
mice, provided by Dr. Alexander Dent (Indiana University), to generate Mb1-
Cre+Bcl6flx/flx (BCL6BKO) mice. CD4-Cre+ mice were crossed to Bcl6flx/flx mice
to generate CD4-Cre+ Bcl6flx/flx (BCL6TKO) mice. For experiments with
BCL6TKO mice, Bcl6flx/flx Cre- littermates were used at WT controls. All mice
were maintained/bred under specific pathogen-free conditions at the University of
Washington. All experiments were performed in accordance with the University of
Washington Institutional Care and Use Committee guidelines.
341

Infections, drug treatment and parasitemia analysis

Plasmodium chabaudi chabaudi (AS) parasites were maintained as frozen blood
stocks and passaged through donor mice. Primary mouse infections were
initiated by intraperitoneal (i.p.) injection of 1×10^6 iRBCs from donor mice.
Secondary mouse infections were performed using a dose of 1×10^7 iRBCs
injected intravenously (i.v.). When indicated, mice were treated i.p. with
14.4mg/kg atovaquone resuspended in DMSO (Sigma) on days 7-11. When
indicated, mice were i.p. injected with 250ug CD4 depleting antibody clone GK1.5
(BioXcell) beginning at the indicated time points and continued semi-weekly
through the duration of the experiment. Parasitemia was measured by flow

cytometry by fixing a drop of blood with 0.025% glutaraldehyde and staining with

Ter119 FITC, CD45 APC, Hoechst33342, and CD71 PE. For memory PE

experiments, mice were injected at the base of the tail with 15ug PE in CFA. For

secondary PE responses, previously immunized mice were given 60ug PE in

CFA i.p. (Pape et al., 2011).

Bone Marrow Chimeras

Bone marrow cells were collected from the tibia, femur, humerus, and sternum

and labeled with anti-Thy1.2 (30-H12, eBioscience) and anti-NK1.1 (PK136,

eBioscience). Cells were resuspended and incubated with low toxicity rabbit

complement (Cedarlane Laboratories). After complement lysis, cells were

washed with media containing 10% fetal calf serum. Recipient mice were lethally

irradiated (1000 rads) and injected intravenously with $5x10^6$ total bone marrow

cells with congenically disparate WT cells mixed with BCL6BKO cells mixed in

equal portions and provided with antibiotic (Enrofloxacin)-treated water for 8

weeks. Prior to infection, chimerism was determined analyzing the ratio of

CD45.1 and CD45.2 expression among B220+ cells in the blood. In order to

adjust for differences in initial chimerism, this ratio was then applied to the

calculations obtained after infection.

Tetramers

Purified recombinant His-tagged C-terminal MSP1 protein (amino acids 4960 to

5301) (Ndungu et al., 2009) was biotinylated and tetramerized with streptavidin-

PE or streptavidin-APC (Prozyme), as previously described (Krishnamurty et al.,

2016). Decoy reagent to exclude non-specific binding to the MSP1-PE tetramer

was made by conjugating SA-PE to AF647 using an AF647 protein labeling kit

(ThermoFisher), washing and removing any unbound AF647, and incubating with

an excess of an irrelevant biotinylated His-tagged Class II-associated invariant

chain peptide (Krishnamurty et al., 2016; Taylor et al., 2012a). Decoy reagent to

exclude non-specific binding to the MSP1-APC tetramer was made by

conjugating SA-APC to DyLight 755 using a DyLight 755 antibody labeling kit

(ThermoFisher), washing and removing any unbound DyLight 755, and

incubating with an excess of irrelevant biotinylated His-tagged Class II-associated

invariant chain peptide.

Mouse Cell Enrichment and Flow Cytometry

Splenic single-cell suspensions were prepared by mashing spleens and passing

through 100um Nitex mesh (Amazon.com). For B cell enrichment, cells were

resuspended in 200uL in PBS containing 2% FBS and Fc block (2.4G2) and first

incubated with Decoy tetramer at a concentration of 10nM at room temperature

for 20 min. MSP1-PE tetramer or MSP1-APC tetramer was added at a

concentration of 10nM and incubated on ice for 30 min. Cells were washed,

incubated with anti-PE or anti-APC magnetic beads for 30 min on ice, and

enriched for using magnetized LS columns (Miltenyi Biotec). All bound cells were

stained with surface antibodies followed by fixation and intracellular antibody

staining when needed (Table 1). Cell counts were determined using Accucheck

cell counting beads. All cells were run on the LSRII, Fortessa, Canto RUO or

Symphony (BD) and analyzed using FlowJo software (Treestar).

Bulk BCR Sequencing and Analysis

cDNA was generated from bulk sorted cells using a Template Switching RT

Enzyme Mix (NEB) and the manufacturer's protocol for Smart-seq at double

volume per reaction. To account for larger cell numbers greater than 1000, a

longer 17 base barcoded Template Switch Oligo (TSO) was used, and a shorter 8

base barcoded TSO was used for counts less than 1000. All reactions used a

poly-T primer that shared the sequence of the TSOs and allowed for subsequent

smart-seq amplification of all cDNA. A 1/3 portion of the barcoded cDNA was

used in a qPCR reaction to determine the optimal number of cycles before

plateau for each reaction. The rest of the barcoded cDNA was then amplified

accordingly. To amplify BCRs, an Illumina sequencing method was used

(manuscript in preparation as Thouvenel et al). Briefly, a multiplex PCR was performed on each bulk sample using pooled constant region primers for IgM, IgG, IgA, IgD, IgK, and IgL and sample-barcoded forward primers for the universal template switch region. Samples were then pooled and library prepped using enzymatic fragmentation and ligation with an anchored multiplex PCR approach (Zheng et al., 2014) to retain sample and transcript barcodes. Indexed library preps were run on an Illumina NovaSeq X at 300 cycles. Individual samples were demultiplexed by sample-barcoded sequences. UMI-tools (Smith et al., 2017) was used to create whitelists of transcript barcodes for each sample after manually determining the barcode counts from the knee in the rank abundance plots. TRUST4 (Song et al., 2021) was run for each sample along with the barcode whitelist to create BCR contigs. Contigs were submitted to IMGT/HighV-Quest (GENE-DB Version 3.1.42) for annotation and nucleotide mutation counts were parsed using an in-house python script from the output tables. Sequences were filtered by length so that all included at least the CDR1 to the end of the J-gene. Sequences for each sample were collapsed by V-gene, J-gene, nucleotide junction and amino acid mutations, and plotted in R (Version 4.4.1, http://www.r-project.org/) using the tidyverse (2.0.0, (Wickham et al., 2019) and ggplot2 (3.5.1).

Histology

Spleens were fixed in 4% PFA for 2 hours at 4°C, washed three times for 10 min in 1X PBS and sunk in 30% sucrose overnight before freezing in OCT. 7μm cryosections were dried for 1 hour and rehydrated in 1X PBS with 1% BSA for 10 min. Sections were stained in antibody overnight at 4°C, washed for 10min in 1X PBS with 1% BSA and stained in secondary antibody for 2 hours at room temperature. All stains were done in 1X PBS with 2% normal mouse serum (Jackson Immunoresearch), 2% normal rat serum (Jackson Immunoresearch) and 1% BSA. Antibodies used were anti-IgD ef450 (11-26c, eBioscience), anti-CD3 Alexa488 (17A2, BioLegend), anti-CD35 biotin (8C12, BD Pharmingen),

anti-BCL6 Alexa647 (K112-91, BD Biosciences) and streptavidin Cy3 (Jackson Immunoresearch). Immunofluoresence microscopy images were captured with a Nikon Eclipse Ti microscope and analyzed with NIS-Elements AR.

Statistical Analysis

Two-tailed Student's *t* tests were applied to determine the statistical significance of the differences between individual groups. One-way ANOVA or two-way ANOVA was used to determine the statistical significance of the differences between multiple groups. All analyses were done with Prism (Graphpad) software. The p-values were considered significant when $p < 0.05$ (*), $p < 0.01$ (**), and $p < 0.001$ (***).

List of Supplementary Materials

Figure S1. Heterogeneous B cells populations are dependent on intrinsic expression of BCL6**.**

Figure S2. BCL6 is expressed in both IgM+ and swIg+ B cells

Figure S3. GC B cells do not develop in BCL6TKO mice and are not required for the DP IgM+ MBC response after PE immunization.

Table1. List of antibodies used

References

Akkaya, M., K. Kwak, and S.K. Pierce. 2020. B cell memory: building two walls of protection against pathogens. *Nat Rev Immunol* 20:229-238.

Arroyo, E.N., and M. Pepper. 2020. B cells are sufficient to prime the dominant CD4+ Tfh response to Plasmodium infection. *J Exp Med* 217:

Becher, B., A. Waisman, and L.F. Lu. 2018. Conditional Gene-Targeting in Mice: Problems and Solutions. *Immunity* 48:835-836.

Callahan, D., S. Smita, S. Joachim, K. Hoehn, S. Kleinstein, F. Weisel, M. Chikina, and M. Shlomchik. 2024. Memory B cell subsets have divergent developmental origins that are coupled to distinct imprinted epigenetic states. *Nat Immunol*

De Silva, N.S., and U. Klein. 2015. Dynamics of B cells in germinal centres. *Nat*
Rev Immunol 15:137-148.
de Vinuesa, C.G., M.C. Cook, J. Ball, M. Drew, Y. Sunners, M. Cascalho, M.
Wabl, G.G. Klaus, and I.C. MacLennan. 2000. Germinal centers without T
cells. *J Exp Med* 191:485-494.
Dent, A.L., A.L. Shaffer, X. Yu, D. Allman, and L.M. Staudt. 1997. Control of
inflammation, cytokine expression, and germinal center formation by BCL-
6. *Science* 276:589-592.
Di Niro, R., S.J. Lee, J.A. Vander Heiden, R.A. Elsner, N. Trivedi, J.M. Bannock,
N.T. Gupta, S.H. Kleinstein, F. Vigneault, T.J. Gilbert, E. Meffre, S.J.
McSorley, and M.J. Shlomchik. 2015. Salmonella Infection Drives
Promiscuous B Cell Activation Followed by Extrafollicular Affinity
Maturation. *Immunity* 43:120-131.
Dogan, I., B. Bertocci, V. Vilmont, F. Delbos, J. Mégret, S. Storck, C.A. Reynaud,
and J.C. Weill. 2009. Multiple layers of B cell memory with different effector
functions. *Nat Immunol* 10:1292-1299.
Fukuda, T., T. Yoshida, S. Okada, M. Hatano, T. Miki, K. Ishibashi, S. Okabe, H.
Koseki, S. Hirosawa, M. Taniguchi, N. Miyasaka, and T. Tokuhisa. 1997.
Disruption of the Bcl6 gene results in an impaired germinal center
formation. *J Exp Med* 186:439-448.
Hollister, K., S. Kusam, H. Wu, N. Clegg, A. Mondal, D.V. Sawant, and A.L. Dent.
2013. Insights into the role of Bcl6 in follicular Th cells using a new
conditional mutant mouse model. *J Immunol* 191:3705-3711.
Inoue, T., and T. Kurosaki. 2024. Memory B cells. *Nat Rev Immunol* 24:5-17.
Kerfoot, S.M., G. Yaari, J.R. Patel, K.L. Johnson, D.G. Gonzalez, S.H. Kleinstein,
and A.M. Haberman. 2011. Germinal center B cell and T follicular helper
cell development initiates in the interfollicular zone. *Immunity* 34:947-960.
Kitano, M., S. Moriyama, Y. Ando, M. Hikida, Y. Mori, T. Kurosaki, and T. Okada.
2011. Bcl6 protein expression shapes pre-germinal center B cell dynamics
and follicular helper T cell heterogeneity. *Immunity* 34:961-972.
Klein, U., and R. Dalla-Favera. 2008. Germinal centres: role in B-cell physiology
and malignancy. *Nat Rev Immunol* 8:22-33.
Krishnamurty, A.T., C.D. Thouvenel, S. Portugal, G.J. Keitany, K.S. Kim, A.
Holder, P.D. Crompton, D.J. Rawlings, and M. Pepper. 2016. Somatically
Hypermutated Plasmodium-Specific IgM(+) Memory B Cells Are Rapid,
Plastic, Early Responders upon Malaria Rechallenge. *Immunity* 45:402-
414.
Kuo, T.C., A.L. Shaffer, J. Haddad, Y.S. Choi, L.M. Staudt, and K. Calame. 2007.
Repression of BCL-6 is required for the formation of human memory B
cells in vitro. *J Exp Med* 204:819-830.
Laidlaw, B.J., and J.G. Cyster. 2021. Transcriptional regulation of memory B cell
differentiation. *Nat Rev Immunol* 21:209-220.
Laidlaw, B.J., T.H. Schmidt, J.A. Green, C.D. Allen, T. Okada, and J.G. Cyster.
2017. The Eph-related tyrosine kinase ligand Ephrin-B1 marks germinal
center and memory precursor B cells. *J Exp Med* 214:639-649.

Lentz, V.M., and T. Manser. 2001. Cutting edge: germinal centers can be induced in the absence of T cells. *J Immunol* 167:15-20.

Mesin, L., A. Schiepers, J. Ersching, A. Barbulescu, C.B. Cavazzoni, A. Angelini, T. Okada, T. Kurosaki, and G.D. Victora. 2020. Restricted Clonality and Limited Germinal Center Reentry Characterize Memory B Cell Reactivation by Boosting. *Cell* 180:92-106.e111.

Ndungu, F.M., E.T. Cadman, J. Coulcher, E. Nduati, E. Couper, D.W. Macdonald, D. Ng, and J. Langhorne. 2009. Functional memory B cells and long-lived plasma cells are generated after a single Plasmodium chabaudi infection in mice. *PLoS Pathog* 5:e1000690.

Niu, H., G. Cattoretti, and R. Dalla-Favera. 2003. BCL6 controls the expression of the B7-1/CD80 costimulatory receptor in germinal center B cells. *J Exp Med* 198:211-221.

Papa, I., and C.G. Vinuesa. 2018. Synaptic Interactions in Germinal Centers. *Front Immunol* 9:1858.

Pape, K.A., J.J. Taylor, R.W. Maul, P.J. Gearhart, and M.K. Jenkins. 2011. Different B cell populations mediate early and late memory during an endogenous immune response. *Science* 331:1203-1207.

Peng, C., Q. Hu, F. Yang, H. Zhang, F. Li, and C. Huang. 2019. BCL6-Mediated Silencing of PD-1 Ligands in Germinal Center B Cells Maintains Follicular T Cell Population. *J Immunol* 202:704-713.

Pereira, J.P., L.M. Kelly, Y. Xu, and J.G. Cyster. 2009. EBI2 mediates B cell segregation between the outer and centre follicle. *Nature* 460:1122-1126.

Plotkin, S.A. 2010. Correlates of protection induced by vaccination. *Clin Vaccine Immunol* 17:1055-1065.

Price, M.J., C.D. Scharer, A.K. Kania, T.D. Randall, and J.M. Boss. 2021. Conserved Epigenetic Programming and Enhanced Heme Metabolism Drive Memory B Cell Reactivation. *J Immunol* 206:1493-1504.

Pérez-Mazliah, D., M.P. Nguyen, C. Hosking, S. McLaughlin, M.D. Lewis, I. Tumwine, P. Levy, and J. Langhorne. 2017. Follicular Helper T Cells are Essential for the Elimination of Plasmodium Infection. *EBioMedicine* 24:216-230.

Ranuncolo, S.M., J.M. Polo, J. Dierov, M. Singer, T. Kuo, J. Greally, R. Green, M. Carroll, and A. Melnick. 2007. Bcl-6 mediates the germinal center B cell phenotype and lymphomagenesis through transcriptional repression of the DNA-damage sensor ATR. *Nat Immunol* 8:705-714.

Schmidt-Supprian, M., and K. Rajewsky. 2007. Vagaries of conditional gene targeting. *Nat Immunol* 8:665-668.

Seifert, M., M. Przekopowitz, S. Taudien, A. Lollies, V. Ronge, B. Drees, M. Lindemann, U. Hillen, H. Engler, B.B. Singer, and R. Küppers. 2015. Functional capacities of human IgM memory B cells in early inflammatory responses and secondary germinal center reactions. *Proc Natl Acad Sci U S A* 112:E546-555.

Smith, T., A. Heger, and I. Sudbery. 2017. UMI-tools: modeling sequencing errors
in Unique Molecular Identifiers to improve quantification accuracy. *Genome*
Res 27:491-499.
Song, L., D. Cohen, Z. Ouyang, Y. Cao, X. Hu, and X.S. Liu. 2021. TRUST4:
immune repertoire reconstruction from bulk and single-cell RNA-seq data.
Nat Methods 18:627-630.
Taylor, J.J., R.J. Martinez, P.J. Titcombe, L.O. Barsness, S.R. Thomas, N. Zhang,
S.D. Katzman, M.K. Jenkins, and D.L. Mueller. 2012a. Deletion and anergy
of polyclonal B cells specific for ubiquitous membrane-bound self-antigen.
J Exp Med 209:2065-2077.
Taylor, J.J., K.A. Pape, and M.K. Jenkins. 2012b. A germinal center-independent
pathway generates unswitched memory B cells early in the primary
response. *J Exp Med* 209:597-606.
Trivedi, N., F. Weisel, S. Smita, S. Joachim, M. Kader, A. Radhakrishnan, C.
Clouser, A.M. Rosenfeld, M. Chikina, F. Vigneault, U. Hershberg, N. Ismail,
and M.J. Shlomchik. 2019. Liver Is a Generative Site for the B Cell
Response to Ehrlichia muris. *Immunity* 51:1088-1101.e1085.
Vinuesa, C.G., M.A. Linterman, D. Yu, and I.C. MacLennan. 2016. Follicular
Helper T Cells. *Annu Rev Immunol* 34:335-368.
Weisel, F.J., G.V. Zuccarino-Catania, M. Chikina, and M.J. Shlomchik. 2016. A
Temporal Switch in the Germinal Center Determines Differential Output of
Memory B and Plasma Cells. *Immunity* 44:116-130.
Wickham, H., M. Averick, J. Bryan, W. Chang, L. D'Agostino McGowan, R.
François, G. Grolemund, A. Hayes, L. Henry, J. Hester, M. Kuhn, T. Lin
Pedersen, E. Miller, S. Milton Bache, K. Müller, J. Ooms, D. Robinson,
D.P. Seidel, V. Spinu, K. Takahashi, D. Vaughan, C. Wilke, K. Woo, and H.
Yutani. 2019. Welcome to the Tidyverse. *The Journal of Open Source*
Software
William, J., C. Euler, S. Christensen, and M.J. Shlomchik. 2002. Evolution of
autoantibody responses via somatic hypermutation outside of germinal
centers. *Science* 297:2066-2070.
Yates, J.L., R. Racine, K.M. McBride, and G.M. Winslow. 2013. T cell-dependent
IgM memory B cells generated during bacterial infection are required for
IgG responses to antigen challenge. *J Immunol* 191:1240-1249.
Ye, B.H., G. Cattoretti, Q. Shen, J. Zhang, N. Hawe, R. de Waard, C. Leung, M.
Nouri-Shirazi, A. Orazi, R.S. Chaganti, P. Rothman, A.M. Stall, P.P.
Pandolfi, and R. Dalla-Favera. 1997. The BCL-6 proto-oncogene controls
germinal-centre formation and Th2-type inflammation. *Nat Genet* 16:161-
170.
Zhang, T.T., D.G. Gonzalez, C.M. Cote, S.M. Kerfoot, S. Deng, Y. Cheng, M.
Magari, and A.M. Haberman. 2017. Germinal center B cell development
has distinctly regulated stages completed by disengagement from T cell
help. *Elife* 6:
Zheng, Z., M. Liebers, B. Zhelyazkova, Y. Cao, D. Panditi, K.D. Lynch, J. Chen,
H.E. Robinson, H.S. Shim, J. Chmielecki, W. Pao, J.A. Engelman, A.J.

Iafrate, and L.P. Le. 2014. Anchored multiplex PCR for targeted next-
generation sequencing. *Nat Med* 20:1479-1484.
Zuccarino-Catania, G.V., S. Sadanand, F.J. Weisel, M.M. Tomayko, H. Meng,
S.H. Kleinstein, K.L. Good-Jacobson, and M.J. Shlomchik. 2014. CD80
and PD-L2 define functionally distinct memory B cell subsets that are
independent of antibody isotype. *Nat Immunol* 15:631-637.

Acknowledgements: The authors would like to thank Brian Hondowicz and Brian
Johnson for technical assistance. We would also like to thank Dr. Michael Reth
(Max Planck Institute of Immunobiology and Epigenetics) for providing the
Mb1Cre+ mice, and Dr. Alexander Dent (Indiana University) for providing the
Bcl6flx/flx mice. The authors have no conflicting financial interests.

Funding: MP was supported by NIH RO1 A1-118803 and the Burroughs
Wellcome Fund. GHP was supported by NIH R21 AI-166022-02.

Author contributions: GHP conceptualized, performed and analyzed
experiments, prepared figures, wrote and edited the manuscript, ATK
conceptualized, performed and analyzed experiments, LR performed and
analyzed experiments and prepared figures, CT and DJB analyzed experiments
and prepared figures, CEM performed experiments, DJR contributed to
experimental design and analyses, MP conceptualized and analyzed data from
experiments and wrote and edited the manuscript.

FIGURE LEGENDS

Figure 1. B cell intrinsic expression of BCL6 is required for IgM+ and swIg+ MBCs. Chimeras were made using congenically disparate WT (CD45.1/.2+) and BCL6BKO (CD45.2+) bone marrow mixed in equal proportions. After ≥8 weeks reconstitution, mice were infected with *P. chabaudi* and MSP1-specific B cell responses in the spleen were assessed at memory time points >40 days post-infection (A,B). MSP1-specific MBCs were assessed by isotype (A), and CD73 and CD80 expression (B). Data are pooled from 5 experiments with 1-5 mice per experiment. At memory time points (>40 days post-infection), mice were either unchallenged (memory) or rechallenged intravenously with 1×10^7 iRBCs and MSP1-specific B cell responses in the spleen were analyzed by B220 and CD138 expression 3 days later (C). Data are pooled from 3 experiments with 1-3 mice per group in each experiment.

Figure 2. BCL6 is expressed in B cells after infection with *P. chabaudi*. WT mice were infected with *P. chabaudi* and MSP1-specific B cell responses were assessed at the indicated time points. Gating strategy and quantification of CD138-BCL6+IgD- cells (A). Gating strategy and quantification of CD73+CD80+ among BCL6-IgD- cells (B). Data are pooled from 9 independent experiments with 10-31 mice per time point.

Figure 3. Early BCL6 expression is dependent on CD4+ T cells and required for DP MBC development. WT mice were treated with GK1.5 at the indicated time points during *P. chabaudi* infection and BCL6+IgD- cells were quantified at 7 days post-infection (A). Data are pooled from 2 independent experiments with 1-3 mice per group in each experiment. WT mice were treated with GK1.5 at the indicated time points during *P. chabaudi* infection and CD73 and CD80 expression was assessed on IgM+ (B) and swIg+ (C) GL7-IgD- cells. Data are pooled from 3 independent experiments with a total of 6-7 mice per time point.

Figure 4. GC Tfh cells are not required for early B cell activation or the development of DP IgM+ MBCs. WT and BCL6TKO mice were infected with *P. chabaudi* and activation (CD38+GL7+ (A)) and BCL6 expression (B) was assessed at d8 post-infection. WT and BCL6TKO mice were infected with *P. chabaudi* and spleens prepared for histological analysis on 0 (n= 2 mice per genotype) and 6-8 (n = 3 mice per genotype) post-infection. The size and density of CD3+ cells in BCL6+ clusters from sections were quantified from d6-8 infected mice (C). Data are representative of 2 experiments and 3-30 follicles per mouse, 11-60 follicles per timepoint. Immunofluorescence microscopy of splenic B cell follicles. Scale bar is 50µm. WT and BCL6TKO mice were infected with *P. chabaudi.* At 28 days post-infection, CD73 and CD80 expression was assessed on IgM+ and swIg+ MBCs (D). Data are pooled from 3 individual experiments with 2-4 mice in each group. MSP1-specific DP IgM+ MBCs from 2 memory WT and 2 memory BCL6TKO mice were sorted and BCRs were sequenced and analyzed for the number of mutations among unique sequences (WT n=87, BCL6TKO n=44) compared to germline sequences (E). WT and BCL6TKO mice were infected with *P. chabaudi* and treated with atovaquone daily for 5 days starting at day 7 post infection. At ≥28 days post infection, mice were rechallenged with *P. chabaudi* and the IgM+ secondary plasmablast response was quantified 3 days later (F). Data are pooled from 2 independent experiments with 2-3 mice per group in each experiment.

Figure S1. Heterogeneous B cell populations are dependent on intrinsic expression of BCL6. WT:BCL6BKO chimeric mice were infected with *P. chabaudi* and MSP1-specific GC B cells (A) and MBCs (B) were quantified at days 12 and >40 post-infection, respectively. WT and Mb1Cre+ mice were infected with *P. chabaudi* and MSP1-specific GC B cells (C) and DP MBCs (D) were analyzed ≥28 days later.

Figure S2. BCL6 is expressed in both IgM+ and swIg+ B cells. Isotype was

assessed among BCL6+IgD- cells at the indicated time points (A).

Figure S3. GC B cells do not develop in BCL6TKO mice and are not required

for the DP IgM+ MBC response after PE immunization. WT and BCL6TKO

mice were infected with *P. chabaudi* and total MSP1-specific (A) and GC (B) B

cells were quantified at day 28 post-infection. At 28 days post-infection with *P.*

chabaudi DP IgG+ MBCs were sorted, BCRs were sequenced (WT n=141,

BCL6TKO n=10), and mutation counts were determined by comparing to

germline (C). WT and BCL6TKO mice were immunized with PE and PE-specific

MBC isotypes were analyzed for CD73 and CD80 expression (D). WT and

BCL6TKO mice were immunized with PE and allowed to form memory, then

rechallenged with PE. PE-specific plasmablast responses were quantified 3 days

later (E).

Fig. 1

Fig. 2

A.

MSP1-specific CD138- B cells

B.

BCL6-IgD- B cells

Fig. 3

Fig. 4