High-throughput genotyping of a full voltage-gated sodium

channel gene via genomic DNA using target capture

sequencing and analytical pipeline MoNaS to discover

novel insecticide resistance mutations

Kentaro Itokawa[1], Koji Yatsu[1], Tsuyoshi Sekizuka[1], Yoshihide Maekawa[2], Osamu
Komagata[2,3], Masaaki Sugiura[4], Tomonori Sasaki[5], Takashi Tomita[2], Makoto Kuroda[1],
Kyoko Sawabe[2], and Shinji Kasai[2*]
1. Pathogen Genomics Center, National Institute of Infectious Diseases, Japan
2. Department of Medical Entomology, National Institute of Infectious Diseases, Japan
3. Antimicrobial Resistance Research Center, National Institute of Infectious Diseases,
Japan
4. Global Research and Development Department, Fumakilla Limited, Japan
5. Research and Development Department, Fumakilla Limited, Japan
14
15　*To whom correspondence should be addressed: kasacin@nih.go.jp
16

Abstract

Insects' voltage-gated sodium channel (VGSC) is the primary target site of pyrethroid
insecticides. Various amino acid substitutions in the VGSC protein are known to confer
insecticide resistance and are selected under insecticide pressure. In the genome, the
VGSC gene consists of more than 30 exons sparsely distributed across a large genomic
region, which often exceeds 100 kbp. Due to this complex genomic structure of gene *VGSC*,
it is usually challenging to genotype full coding nucleotide sequences (CDSs) of *VGSC* from
individual genomic DNA (gDNA). In this study, we designed biotinylated oligonucleotide
probes via annotated CDSs of *VGSC* of Asian tiger mosquito, *Aedes albopictus*. The probe
set effectively concentrated (>80,000-fold) all targeted regions of gene *VGSC* from pooled
barcoded Illumina libraries each constructed from individual *A. albopictus* gDNAs. The
probe set also captured all homologous *VGSC* CDSs except tiny exons from the gDNA of
other Culicinae mosquitos, *A. aegypti* and *Culex pipiens* complex, with comparable
efficiency by virtue of the high conservation of *VGSC* at the nucleotide level. Furthermore,
we developed an automated bioinformatic analysis pipeline to genotype *VGSC* after capture
sequencing—MoNaS (Mosquito Na$^+$ channel mutation Search)—which conducts mapping
of reads, variant calling, and variant annotation for nonsynonymous mutations. The
proposed method and our bioinformatic tool should facilitate the discovery of novel amino
acid variants conferring insecticide resistance on VGSC and population genetics studies on
resistance alleles with respect to the origin, selection, and migration of both clinically and
agriculturally important insect pests.

Key words: Targeted enrichment, Variant analysis, Voltage Gated Sodium Channel,
Insecticide resistance, Mosquitos

                                   Introduction

Synthetic pyrethroids are currently the most frequently used insecticides for the control of

clinically important mosquitos. The mode of action of pyrethroids is inhibition of the voltage-

gated sodium channel (VGSC) in the nervous system of insects. Various amino acid (aa)

substitutions in VGSC are known to alter its susceptibility to pyrethroids, thereby causing

resistance known as knockdown resistance (_kdr_) (Dong et al., 2014; Rinkevich, Du, & Dong,

2013). VGSC is highly conserved among insects, and some resistance-conferring aa

substitutions occur parallelly in different species. Therefore, it is relatively straightforward to

infer the effect of certain aa substitutions seen in one species if the influence of those

substitutions has already been elucidated in other species. Nonetheless, sequence analysis

of an entire coding sequence (CDS) of the _VGSC_ gene from genomic DNA (gDNA) is usually

more complicated, because in the genome, _VGSC_ genes typically consist of many (>30)

small exons sparsely distributed across a large genomic region, which often exceeds 100

kbp (Davies, Field, Usherwood, & Williamson, 2007). Although direct sequencing of PCR-

amplified genomic fragments is employed to genotype _VGSC_ in many studies, these

analyses often target only the regions where the resistance-conferring substitutions are

frequently found (e.g., IIS5–6). Such a bias may lower the chance of discovering novel

resistance mutations outside the region investigated.

Aedes albopictus, or Asian tiger mosquito, is a medically important mosquito species

ubiquitously present on most continents on Earth. This species is capable of transmitting

important viral diseases including Dengue, Chikungunya, and Zika. Although the

transmission potential of _A. albopictus_ for the dengue virus is believed to be limited as

compared to the major vector species (_A. aegypti)_, _A. albopictus_ has caused outbreaks in

Japan in 2014, where _A. aegypti_ is absent (Kutsuna et al., 2015). The _kdr_ substitution in _A._

albopictus had not been reported until the F1534C allele was discovered in Singapore 2009

(S Kasai et al., 2011). Since this discovery, identical and other _kdr_ substitutions at the same

aa position, F1534S/L, has been reported from various geographic locations worldwide (H.

Chen et al., 2016; Marcombe, Farajollahi, Healy, Clark, & Fonseca, 2014; Xu et al., 2016).

More recently, we also discovered the new _kdr_ substitution V1016G in _A. albopictus_ by

extending the region of the search for mutations (Shinji Kasai et al., 2019).

The next-generation sequencing (NGS) technology has revolutionized almost all fields of

biology by reducing the cost and time of DNA sequencing by orders of magnitude. The

recent _Anopheles gambiae_ 1000 Genomes project, Ag1000G, has uncovered a number of

previously unknown nonsynonymous mutations in the _VGSC_ gene in _A. gambiae_ and _A._

coluzzii (Clarkson et al., 2018), some of which have been suspected to cause resistance

directly or indirectly. The study also showed that even neutral variations within or flanking

79	the *VGSC* locus represent valuable information to infer the origin and evolution of resistance.
80	Although whole-genome sequencing may discover novel variants of *VGSC* unequivocally,
81	this simplistic approach is still too costly to be justified when a number of wild-caught
82	individuals need to be analyzed. An enrichment approach involving hybridization of oligo
83	DNA/RNA is often employed to selectively sequence targeted genome regions (Gnirke et
84	al., 2009) for studies e.g. on genotyping of disease-related genes in humans. This
85	technology is aimed at increasing the depth of analysis and the number of samples to be
86	multiplexed within a given sequencing capacity in return for limiting the region to be
87	analyzed. In this study, we designed biotinylated oligonucleotide DNA probes from *A.*
88	*albopictus VGSC* CDSs. The probe set efficiently concentrated targeted regions from the
89	gDNA of individual *A. albopictus*. Although the probe set was designed by means of only
90	the *A. albopictus VGSC* gene, the same probe set captured most CDSs of *A. aegypti* and
91	*Culex pipiens* complex owing to the high nucleotide conservation of *VGSC*. This technology
92	allows for full-CDS analysis of the complex *VGSC* gene in a relatively low-cost and highly
93	multiplexed manner, which is expected to promote discoveries of novel resistance-
94	conferring aa substitutions both in medical and agricultural insect pests.

| 95 | <div align="center">Materials and Methods</div> |

| 96 | **Design of custom probes** |

97	The full-length *VGSC* gene (AALF000723-RA) was found in scaffold JXUM01S000562 in
98	the genome assembly of an *A. albopictus* Foshan strain, AaloF1 (X.-G. Chen et al., 2015)
99	at vectorbase.org (Giraldo-Calderón et al., 2015). Because the annotation (gene set:
100	AaloF1.2) obviously missed some CDSs, we refined the annotation by aligning shotgun-
101	sequenced NGS reads of *VGSC* cDNA (Shinji Kasai et al., 2019) using Hisat2 (Kim,
102	Langmead, & Salzberg, 2015) and the *Musca domestica* VGSC protein sequence
103	(GenBank accession No.: AAB47604) via BLASTX (Altschul, Gish, Miller, Myers, & Lipman,
104	1990). Compared to AaloF1.2, the refined annotation included three added CDSs, and four
105	extended CDSs (Table S1) among the 35 coding CDSs in total. Of those, 34 (32 + 2 mutually
106	excluding exons) CDSs exactly corresponded to the *A. aegypti VGSC* CDSs annotated by
107	Davies et al. (2007) in terms of sizes. Therefore, the numbering of exons in this paper was
108	set to be concordant with the *A. aegypti VGSC* exons described by Davies et al. (2007).
109	The additional optionally used 45 bp small exon, referred to as exon 16.5, was found in
110	cDNA data between exons 16 and 17 in the genome. We also included some exons of other
111	genes flanking *VGSC* (AALF020128, AALF020129, AALF020130, AALF000725,
112	AALF000726, AALF000727, AALF000728, and AALF000730) or a gene nested in the
113	intronic region of *VGSC* (AALF020132) during the probe design to take advantage of the

population genetic analysis in future studies. All CDS sequences, some of which contained a flanking intronic region (for tiny exons less than 120 bp in size) were submitted to the IDT website (https://sg.idtdna.com) to design 120 bp xGen Lockdown biotinylated oligonucleotide DNA probes, all overlapping pairwise by 60 bp (2× Tiling density option). By means of 15 kbp genomic regions in total, 229 probes were designed (Table S2), of which 145 target *VGSC*. Nonetheless, in the more recent contiguous assembly of the C6/36 cell line (see below), AALF020132, AALF000725, AALF000726, AALF000727, AALF000728, and AALF000730 are not located in the same assembly with the *VGSC* locus. For this reason, in this paper, we evaluate the performance of the probe set only in terms of *VGSC* CDS enrichment.

Samples

Fifty-six mosquitos belonging to species *A. albopictus* or *A. aegypti* or to the *Culex pipiens* complex—either kept in the laboratory or caught in the wild (Table 1)—served as a source of gDNA. Of those, strains Aalb-SP, Aaeg-SP, and Cpip-JPP were already known to possess haplotypes with 1534C, 989P-1016G, and 1014F aa variants, respectively (Hardstone et al., 2007; S Kasai et al., 2014; Shinji Kasai et al., 2019).

gDNA extraction

gDNA was individually extracted from the whole body of an adult or pupa using the MagExtractor Genome Kit (TOYOBO). The protocol was modified to conduct the extraction in 8-strip PCR tubes or a 96-well PCR plate as follows. The whole body of a single insect was homogenized in a PCR well containing 50 µl of the Lysis & Binding Solution and zirconia beads (ø 2 mm; Nikkato) in TissueLyser II (QIAGEN) at 25 Hz for 30 s. After that, the samples were centrifuged at $2000 \times g$ for 1 min to precipitate large debris, and each supernatant was transferred to a new well containing 50 µl of the Lysis & Binding Solution and 5 µl of DNA-binding Magnetic Beads. The solution was shaken in MicroMixer E-36 (TAITEC) at the maximum speed (2500 rpm) for 10 min, and then, on a magnetic plate, the supernatant was discarded. The beads bound to DNA were washed twice with 100 µl of the Washing Solution and twice with 75% ethanol each. Finally, DNA was eluted with 50 µl of low-TE buffer (0.1 mM EDTA, 10 mM Tris-HCl pH 8.0) by shaking in MicroMixer E-36 at the maximum speed for 10 min. The obtained DNA was quantified with the Qubit Highly Sensitive DNA Assay Kit (Invitrogen). The obtained DNA concentration ranged from 2.3 to 8.6 ng/µl for *A. albopictus*, 5.3 to 10 ng/µl for *A. aegypti*, and 7.8 to 11 ng/µl for *C. pipiens* complex mosquitos.

Library construction and hybridization capture

Illumina libraries with TruSeq barcode adapters were prepared by means of the NEBNext

Ultra II FS DNA Library Prep Kit for Illumina (NEB) on the 1/4 scale of the manufacturer-

suggested protocol. Briefly, 4 µl of the gDNA extracted above (without adjusting the

concentration) was mixed with 0.4 µl of the Enzyme Mix, 1.4 µl of Reaction Buffer, and 1.2

µl of H_2O on ice. The mixture was then incubated at 37 °C for 10 min followed by incubation

at 65 °C for 30 min. Those end-prepped DNAs were directly ligated with Illumina adapters

by the addition of 0.5 µl of TruSeq 96 dual-index adapters (Illumina) instead of adapters

supplied with the kit, 6 µl of the Ligation Master Mix, and 0.2 µl of Ligation Enhancer, with

incubation at 20 °C for 15 min. Next, the libraries were incubated at 65 °C for 30 min to

inactivate the ligase; then, all the 56 libraries were pooled together in a single 1.5 ml LoBind

tube (Eppendrof). The pooled library was purified with 1.2× SPRIselect (Beckman Coulter)

and eluted with 20 µl of low-TE buffer. A 7 µl aliquot of the pooled library was mixed with 0.8

µl of a 10 µg/µl UltraPure Salmon Sperm DNA Solution (Invitrogen) in a PCR-tube and was

concentrated by incubation at 80 °C for 10 min while the lid of the tube and thermal cycler

were opened. The concentrated library mix was hybridized, captured, and washed with the

designed oligo DNA probe set and the xGen Hybridization and Wash Kit (IDT). After that,

the streptavidin magnetic beads were subjected to PCR amplification with HiFi Kapa (Kapa

Biosystems) for 12 cycles. The amplified library was purified with 1.2× SPRIselect beads

and quantified by real-time PCR using P5 and P7 adapter primers and qPCR double

quencher   probe   (6-FAM)-5′-ACACTCTTT-(ZEN)-CCCTACACGACGCTCTTC-3′-(Iowa

Black FQ) (IDT) in the PrimeTime Gene Expression Master Mix (IDT). Serial dilutions of the

phiX library (Illumina) were used for construction of the standard curve. The quantified

library was sequenced on Illumina MiniSeq with the Mid Output Kit (Illumina) for 151 cycles

from both ends along with the libraries from other studies.

Reference genomes and annotation for *VGSC*

Although the probe sets were designed from assembly AaloF1, we chose a C6/36 cell line

genome assembly, canu_80X_arrow2.2 (Miller et al., 2018), as a reference genome of *A.*

albopictus for further bioinformatic analysis because this assembly has better contiguity and

fewer scaffolds than AaloF1 does. In the canu_80X_arrow2.2 assembly, the whole *VGSC*

gene was found in scaffolds MNAF02001058.1 and MNAF02001442.1 annotated as Gene

IDs LOC109421922 and LOC109432678, respectively, in the NCBI *Aedes albopictus*

Annotation Release 101. The two *VGSC* genes were assumed to be redundant haplotigs

due to heterogeneity in the original DNA source. To avoid dual mapping of the NGS reads,

we purged MNAF02001442 by hard-masking this entire scaffold (replacing all bases with

the "N" character) rather than MNAF02001058.1 because LOC109432678 in MNAF02001442.1 has a single frame-shifting nucleotide deletion in the thymine homopolymer track within exon 4 (TTTTTT → TTTTT), which was suspected due to an uncorrected base-calling error. LOC109421922 was defined by the number of transcriptional variants in the NCBI's annotation because *VGSC* is known to have complex alternative splicing patterns (Davies et al., 2007). Nevertheless, we simplified the *VGSC* gene model into two possible transcriptional variants to build a GFF3 annotation file for annotating aa changes. These two transcripts include all the regions of mandatory or optional CDSs but differ by the two mutually exclusive exons "19c/k" and "26d/l," where one carries exons "19c" and "26k," and the other contains exons "19d" and "26l." CDSs of all the transcriptional variants of LOC109421922 were merged via overlaps. Those merged CDSs perfectly matched AaloF1 except for LOC109421922 including exon 16.5 and except for one mutually exclusive exon "26k" whose sequence itself was found to be intact in MNAF02001058.1.

The *VGSC* gene in the chromosome level assembly of the *A. aegypti* genome, AaegL5.0 (Matthews et al., 2018), was annotated in the same manner. The whole *VGSC* gene is encoded as AAEL023266 (the NCBI *Aedes aegypti* Annotation Release 101) on chromosome 3. AAEL023266 has 13 transcripts, these CDSs were merged via overlaps as in the canu_80X_arrow2.2 assembly of *A. albopictus*. AAEL023266 appeared to be lacking an exon corresponding to exon 16.5, whereas we found its sequence between exons 16 and 17. AAEL023266, however, contains an additional exon between exons 11 and 12. The 21 bp small exon was assumed to correspond to exon "12" in the *Anopheles gambiae* genome described by Davies et al. (2007) and is situated within the intracellular loop between domains I and II. We found the sequence homologous to this exon also in the two *A. albopictus* genome assemblies, AaloF1 and canu_80X_arrow2.2; thus, we had failed to include this exon in the probe design. The complete *VGSC* sequence was also found in scaffold NIGP01000811 and was assumed to be a redundant haplotig. This scaffold was purged from the assembly by hard-masking. In *C. quinquefasciatus* genome assembly Cpip_J2 (Arensburger et al., 2010), the *VGSC* gene is located in scaffold supercont3.182. The *VGSC* gene in supercont3.182, however, contains shorter exon 13 which was truncated by scaffolding gap and lacks the entire exon 14. Complete exons 13 and 14 were found in another scaffold, supercont3.1170, which contains an incomplete *VGSC* gene probably as a haplotig. We fused contig AAWU01037504.1 containing exons 13 and 14 of *VGSC* from supercont3.1170 into supercont3.182 to restore the complete coding sequence of the *VGSC* gene, thereby creating supercont3.182_2. The *VGSC* in supercont3.182 (and supercont3.182_2) still contained *kdr* aa substitutions, L932F and I936V, as already

reported by Davies et al. (2007) plus unusual frame-shifting deletions in exons 26l and 32.

For these reasons, supercont3.182_2 was further polished by means of the *consensus*

module in BCFtools (Danecek & McCarthy, 2017) with the "-H 1" option using the variant

information for Cpip-JNA-01 in the VCF file generated as described below, thereby finally

resulting in supercont3.182_3. The latter scaffold was added to the genome assembly, and

the original scaffolds supercont3.182 and supercont3.1170 were purged by hard-masking.

We were not able to find an exon corresponding to "exon 16.5" in *A. albopictus* and *A.*

aegypti.

Bioinformatic analysis

The FASTQ data were mapped to the reference using *BWA mem* (v.0.7.17) (Li & Durbin,

2009) with default options. The resultant BAM files were sorted by the *sort* program from

the SAMtools suite (v.1.9) (Li et al., 2009), and we removed PCR duplicates by the *rmdup*

programs from the SAMtools. Variant calling was performed on the resulting BAM files of

each species in the FreeBayes software (v.1.2.0) (Garrison & Marth, 2012) with default

options. The variant annotation (for aa changes) was conducted with the *csq* program from

the BCFtools suite (v.1.9) (Danecek & McCarthy, 2017) with options "*-l -p a*". Finally, the

discovered aa changes were mapped onto the corresponding position in the *Musca*

domestica VGSC protein sequence (GenBank accession No.: AAB47604). Those

bioinformatic processes (Fig. 1) were automated in pipeline tools *MoNaS* (Mosquito Na$^+$

channel mutation Search; https://github.com/ItokawaK/MoNaS) written in the Python3 script

language.

For estimating the level of enrichment, five sets of random data on whole-genome shotgun

paired-end reads (150 bp × 2, 300 ± 50 bp insert length, 1 million read pairs) from each

reference genomic assembly were simulated in the *wgsim* software

(https://github.com/lh3/wgsim). The *multicov* program from the Bedtools suite (v.2.27.1)

(Quinlan & Hall, 2010) was applied to calculate the number of reads overlapping with any

targeted CDS regions. Nucleotide identities of exons were calculated using *Muscle* (Edgar,

2004) and BioPython's *Phylo* package (Talevich, Invergo, Cock, & Chapman, 2012). *R*

(v.3.5.1) (R_Development_Core_Team, 2014) and the *ggplot2* package (Wickham, 2016)

were utilized for summarizing and visualizing the data.

248   Results

For the 56 mosquito gDNA samples, 4.9 million demultiplexed read pairs (150 bp PE) in

total were obtained after a single run of Illumina MiniSeq. From those samples, 0.040–0.17,

0.050–0.12, and 0.063–0.10 million read pairs were obtained from each individual mosquito

of species *A. albopictus* and *A. aegypti* and *C. pipiens* complex, respectively. In *A.*

albopictus, 44% of all reads on average overlapped with any of the *VGSC* CDSs under

study (Reads mapped per kilobase exon per million "sequenced" reads: RPKM = 6.5×10^4),

which was approximately 8.3×10^4-fold enrichment compared to simulated random data on

whole-genome shotgun sequencing (Fig. 2). Although the probe set was designed on the

basis of the *A. albopictus* genomic sequence, the same probe set captured *VGSC* CDSs

from the gDNA of *A. aegypti* and *C. pipiens* complex at an on-target rate similar to that of *A.*

albopictus (Fig. 2).

Fig. 3 shows a distribution of median and minimum sequencing depths within each CDS

in each individual sample after PCR duplicates were removed. In *A. albopictus*, most

nucleotides in all exons were covered deeply with minimum bias. In *A. aegypti* and *C.*

quinquefasciatus, however, some exons were covered at low depth partly or entirely. In

particular, exons 2 and 16.5 were covered at nearly or absolutely zero depth.

Fig. 4 presents a distribution of the allele balance in genotypes containing single or multiple

nucleotide variants (SNVs or MNVs). The ratio of the first allele in a heterozygous genotype

was distributed mostly around 50%, which was substantially different from the homozygous

genotype (near 100%) except for one SNV or MNV site in exon 32 of the *A. albopictus* gene

located in the GGT (Gly) trinucleotide tandem repeats variable in length near the C terminus

of VGSC, where accurate calling of the genotype is difficult.

Aa substitutions identified at the end of the MoNaS pipeline are listed in Table 2. Of

those, F1534C in Aalb-SP (Shinji Kasai et al., 2019)*,* S989P and V1016G in Aaeg-SP (S

Kasai et al., 2014), and L1014F in Cpip-JPP (Hardstone et al., 2007), all previously known

to exist in those strains, were recalled correctly. In the Aaeg-Mex strain, V410L, V1016I,

and F1534C variants, which are known as *kdr* (Brengues et al. 2003; Haddi et al. 2017;

Kawada et al. 2009), were detected. Other aa substitutions detected—C749*Y (*in

mosquito aa coordinates because there was no corresponding aa in *M. domestica*),

A2023T and G2046E in *A. albopictus*, S723T in *A. aegypti*, and K109R, Y319F, T1632S,

E1633D, E1856D, G2051A, and A2055V in *C. pipiens* complex—are not known for their

effects on insecticide susceptibility.

281   Discussion

In this study, we conducted targeted enrichment sequencing of *VGSC* CDSs from the

gDNA of mosquitos using hybridization capture probes. The result of the experiment is quite

promising: most nucleotides of *VGSC* CDSs were covered at sufficient read depths even in

samples with less than a 30 Mbp (0.1 million read-pairs) sequencing effort.

Even though the probe set was designed on the basis of the *A. albopictus* genome

sequence only, it successfully enriched *VGSC* CDSs from the gDNA of two other Culicinae

mosquito species, *A. aegypti* and *C. pipiens* complex, owing to the high sequence homology

among *VGSC* orthologs at the nucleotide level. Although the estimated entire efficiency of
enrichment was lower by approximately 50% in *C. pipiens* samples than in *A. albopictus*
samples, the on-target ratio was still comparable or rather higher in *C. pipiens* (Fig. 2). This
result can be explained by the much smaller genome size of *C. pipiens* complex (579 Mb in
the CpipJ2 assembly) as compared to *A. albopictus* (2.25 Gbp in the C6/36 assembly).
Applicability of a single probe set to multi species (e.g., the same genus or family) is clearly
advantageous because this obviates the need to prepare each custom probe sets specific
for each single species and may enable capture even in species lacking prior genome
information. Nonetheless, the evolutionary distance will limit the range of species that one
probe set can be applied. In this study, the mapping results on each exon indicated that
capture efficiency decreased in some exons (Fig. 3). This empirical observations in this
suggested that less than 87.5% in identity or less than 60 bp in size for the homology track
of targets could decrease the efficiency of capture (Fig. 5). Especially, our probe set failed
to capture two optionally used exons, 2 and 16.5, in *A. aegypti* and *C. pipiens* complex,
which are among the smallest exons targeted (Fig. 3). It is assumed that those tiny exons
alone do not provide enough thermostability for probe–target DNA duplex during the capture.
Given that the probes for those small exons contain flanking intronic sequences of *A.*
albopictus genome as well, those flanking sequences may have provided a length of the
homologous track sufficient to capture sequences from *A. albopictus*. Although it is
straightforward to optimize our probe set further at least for the two species of mosquito
simply by adding species specific probes for those exons and flanking intronic regions, small
exons in general will be a major challenge when a probe set is aimed to be used for a group
of species rather than specific targets because the homology in an intronic region will decay
more rapidly than that in an exonic region during speciation. We also missed another exon
corresponding to "exon 12" in *Anopheles gambiae* described by Davies et al. (2007) during
probe design (see Materials and Methods). Such tiny and rarely used exons may be difficult
to annotate without high-quality high-throughput RNA sequencing data. Nevertheless, in
mosquitoes, all such tinyl exons are actually situated on the N-terminal intracellular loop or
the intracellular loop between domains I and domain II in VGSC, where no resistance-
associated mutation has been found so far (Dong et al., 2014). Therefore, it is not clear
whether ignoring those small exons of *VGSC* from analysis does have serious consequence
for insecticide resistance research.
The process of genotyping *VGSC* carried out in this study was automated in MoNaS. This
program sequentially runs tools conducting mapping of NGS reads to a reference, sorting,
removal of PCR duplicates, indexing for BAM files, variant calling, variant annotation, and
finally integration of these results across multiple samples into a single table with conversion

of the aa coordinates to those corresponding to the *M. domestica* VGSC protein. The
automation in MoNaS allows researchers to process raw NGS reads of many samples via
a simple command line operation without expert knowledge of the bioinformatics field.
MoNaS can be run locally with appropriate genome reference data or executed online as a
Web service (https://gph.niid.go.jp/monas).
330

Acknowledgements

This research was partially supported by the Research Program on Emerging and Re-
emerging Infectious Diseases (JP19fk0108067) from Japan Agency for Medical Research
and Development (AMED) and by Japan Initiative for Global Research Network on
Infectious Diseases (J-GRID) from the Ministry of Education, Culture, Sports, Science and
Technology of Japan and AMED.

Authors' contributions

KI and SK designed the study. KI and OK designed the capture probes. KI and TT refined
the annotation of the *VGSC* gene. KI conducted all the experiments. KI, TSe, KY, and MK
contributed to the bioinformatic analysis and development of MoNaS. SK, YM, MS ans TSa
contributed maintaining the laboratory colony or collected specimens of mosquitoes in the
field. KI drafted this manuscript. All the coauthors critically revised the manuscript and
approved it for publication.
345

Reference

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment
search tool. *Journal of Molecular Biology*, *215*(3), 403–410. doi:10.1016/S0022-
2836(05)80360-2
Arensburger, P., Megy, K., Waterhouse, R. M., Abrudan, J., Amedeo, P., Antelo, B., …
Atkinson, P. W. (2010). Sequencing of Culex quinquefasciatus establishes a platform for
mosquito comparative genomics. *Science*, *330*(6000), 86–88.
doi:10.1126/science.1191864
Brengues, C., Hawkes, N. J., Chandre, F., Mccarroll, L., Duchon, S., Guillet, P., … Hemingway,
J. (2003). Pyrethroid and DDT cross-resistance in Aedes aegypti is correlated with novel
mutations in the voltage-gated sodium channel gene. *Medical and Veterinary Entomology*,
17(1), 87–94. doi:10.1046/j.1365-2915.2003.00412.x
Chen, H., Li, K., Wang, X., Yang, X., Lin, Y., Cai, F., … Ma, Y. (2016). First identification of kdr
allele F1534S in VGSC gene and its association with resistance to pyrethroid insecticides

in Aedes albopictus populations from Haikou City, Hainan Island, China. *Infectious*

Diseases of Poverty, *5*(1), 31. doi:10.1186/s40249-016-0125-x

Chen, X.-G., Jiang, X., Gu, J., Xu, M., Wu, Y., Deng, Y., … James, A. A. (2015). Genome

sequence of the Asian Tiger mosquito, *Aedes albopictus* , reveals insights into its biology,

genetics, and evolution. *Proceedings of the National Academy of Sciences*, *112*(44),

E5907–E5915. doi:10.1073/pnas.1516410112

Clarkson, C. S., Miles, A., Harding, N. J., Weetman, D., Kwiatkowski, D., Donnelly, M., &

Consortium, T. <I>Anopheles gambiae</I> 1000 G. (2018). The genetic

architecture of target-site resistance to pyrethroid insecticides in the African malaria

vectors Anopheles gambiae and Anopheles coluzzii. *BioRxiv*, 323980. doi:10.1101/323980

Danecek, P., & McCarthy, S. A. (2017). BCFtools/csq: haplotype-aware variant consequences.

Bioinformatics, *33*(13), 2037–2039. doi:10.1093/bioinformatics/btx100

Davies, T. G. E., Field, L. M., Usherwood, P. N. R., & Williamson, M. S. (2007). A comparative

study of voltage-gated sodium channels in the Insecta: implications for pyrethroid

resistance in Anopheline and other Neopteran species. *Insect Molecular Biology*, *16*(3),

361–75. doi:10.1111/j.1365-2583.2007.00733.x

Dong, K., Du, Y., Rinkevich, F., Nomura, Y., Xu, P., Wang, L., … Zhorov, B. S. (2014, July).

Molecular biology of insect sodium channels and pyrethroid resistance. *Insect*

Biochemistry and Molecular Biology. NIH Public Access. doi:10.1016/j.ibmb.2014.03.012

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res*, *32*(5), 1792–1797. doi:10.1093/nar/gkh340

Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read

sequencing. Retrieved from http://arxiv.org/abs/1207.3907

Giraldo-Calderón, G. I., Emrich, S. J., MacCallum, R. M., Maslen, G., Dialynas, E., Topalis, P.,

… Lawson, D. (2015). VectorBase: an updated bioinformatics resource for invertebrate

vectors and other organisms related with human diseases. *Nucleic Acids Research*,

43(D1), D707–D713. doi:10.1093/nar/gku1117

Gnirke, A., Melnikov, A., Maguire, J., Rogov, P., LeProust, E. M., Brockman, W., … Nusbaum,

C. (2009). Solution hybrid selection with ultra-long oligonucleotides for massively parallel

targeted sequencing. *Nature Biotechnology*, *27*(2), 182–189. doi:10.1038/nbt.1523

Haddi, K., Tomé, H. V. V., Du, Y., Valbon, W. R., Nomura, Y., Martins, G. F., … Oliveira, E. E.

(2017). Detection of a new pyrethroid resistance mutation (V410L) in the sodium channel

of Aedes aegypti: a potential challenge for mosquito control. *Scientific Reports*, *7*(1),

46549. doi:10.1038/srep46549

Hardstone, M., Leichter, C., Harrington, L., Kasai, S., Tomita, T., & Scott, J. (2007). Cytochrome

P450 monooxygenase-mediated permethrin resistance confers limited and larval specific

cross-resistance in the southern house mosquito, Culex pipiens quinquefasciatus.

Pesticide Biochemistry and Physiology, *89*(3), 175–184. doi:10.1016/j.pestbp.2007.06.006

Kasai, S., Caputo, B., Tsunoda, T., Cuong, T. C., Maekawa, Y., Lam-Phua, S. G., … Tomita, T.

(2019). First detection of a Vssc allele V1016G conferring a high level of insecticide

resistance in Aedes albopictus collected from Europe (Italy) and Asia (Vietnam), 2016: a

new emerging threat to controlling arboviral diseases. *Eurosurveillance*, *24*(5), 1700847.

doi:10.2807/1560-7917.ES.2019.24.5.1700847

Kasai, S., Komagata, O., Itokawa, K., Shono, T., Ng, L. C., Kobayashi, M., & Tomita, T. (2014).

Mechanisms of pyrethroid resistance in the dengue mosquito vector, Aedes aegypti: target

site insensitivity, penetration, and metabolism. *PLoS Negl Trop Dis*, *8*(6), e2948.

doi:10.1371/journal.pntd.0002948

Kasai, S., Ng, L. C., Lam-Phua, S. G., Tang, C. S., Itokawa, K., Komagata, O., … Tomita, T.

(2011). First detection of a putative knockdown resistance gene in major mosquito vector,

Aedes albopictus. *Jpn J Infect Dis*, *64*(3), 217–221. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/21617306

Kawada, H., Higa, Y., Komagata, O., Kasai, S., Tomita, T., Thi Yen, N., … Takagi, M. (2009).

Widespread Distribution of a Newly Found Point Mutation in Voltage-Gated Sodium

Channel in Pyrethroid-Resistant Aedes aegypti Populations in Vietnam. *PLoS Negl Trop*

Dis, *3*(10), e527. doi:10.1371/journal.pntd.0000527

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low memory

requirements. *Nature Methods*. doi:10.1038/nmeth.3317

Kutsuna, S., Kato, Y., Moi, M. L., Kotaki, A., Ota, M., Shinohara, K., … Ohmagari, N. (2015).

Autochthonous Dengue Fever, Tokyo, Japan, 2014. *Emerging Infectious Diseases*, *21*(3),

517–520. doi:10.3201/eid2103.141662

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics*, *25*(14), 1754–1760. doi:10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R. (2009). The

Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

doi:10.1093/bioinformatics/btp352

Marcombe, S., Farajollahi, A., Healy, S. P., Clark, G. G., & Fonseca, D. M. (2014). Insecticide

Resistance Status of United States Populations of Aedes albopictus and Mechanisms

Involved. *PLoS ONE*, *9*(7), e101992. doi:10.1371/journal.pone.0101992

Matthews, B. J., Dudchenko, O., Kingan, S. B., Koren, S., Antoshechkin, I., Crawford, J. E., …

Vosshall, L. B. (2018). Improved reference genome of Aedes aegypti informs arbovirus

vector control. *Nature*, *563*(7732), 501–507. doi:10.1038/s41586-018-0692-z

Miller, J. R., Koren, S., Dilley, K. A., Puri, V., Brown, D. M., Harkins, D. M., … Shabman, R. S.

(2018). Analysis of the Aedes albopictus C6/36 genome provides insight into cell line utility

for viral propagation. *GigaScience*, *7*(3). doi:10.1093/gigascience/gix135

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics*, *26*(6), 841–842. doi:10.1093/bioinformatics/btq033

R_Development_Core_Team. (2014). R: A language and environment for statistical computing.

R Foundation for Statistical Computing, Vienna, Austria. Retrieved from http://www.r-

project.org

Rinkevich, F. D., Du, Y., & Dong, K. (2013). Diversity and convergence of sodium channel

mutations involved in resistance to pyrethroids. *Pesticide Biochemistry and Physiology*,

106(3), 93–100. doi:10.1016/j.pestbp.2013.02.007

Saavedra-Rodriguez, K., Urdaneta-Marquez, L., Rajatileka, S., Moulton, M., Flores, A. E.,

Fernandez-Salas, I., … Black, W. C. th. (2007). A mutation in the voltage-gated sodium

channel gene associated with pyrethroid resistance in Latin American Aedes aegypti.

Insect Mol Biol, *16*(6), 785–798. doi:IMB774 [pii]10.1111/j.1365-2583.2007.00774.x

Srisawat, R., Komalamisra, N., Eshita, Y., Zheng, M., Ono, K., Itoh, T. Q., … Rongsriyam, Y.

(2010). Point mutations in domain II of the voltage-gated sodium channel gene in

deltamethrin-resistant Aedes aegypti (Diptera: Culicidae). *Applied Entomology and*

Zoology, *45*(2), 275–282. doi:10.1303/aez.2010.275

Talevich, E., Invergo, B. M., Cock, P. J., & Chapman, B. A. (2012). Bio.Phylo: A unified toolkit

for processing, analyzing and visualizing phylogenetic trees in Biopython. *BMC*

Bioinformatics, *13*(1), 209. doi:10.1186/1471-2105-13-209

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Retrieved from http://ggplot2.org

Xu, J., Bonizzoni, M., Zhong, D., Zhou, G., Cai, S., Li, Y., … Chen, X.-G. (2016). Multi-country

Survey Revealed Prevalent and Novel F1534S Mutation in Voltage-Gated Sodium

Channel (VGSC) Gene in Aedes albopictus. *PLOS Neglected Tropical Diseases*, *10*(5),

e0004696. doi:10.1371/journal.pntd.0004696

Data Accessibility

- NGS read data: DRR167897– DRR167952

- MoNaS source code: https://github.com/ItokawaK/MoNaS

Table 1 Mosquito samples used in this study

ID	Species	Lab. Colony / Wild	num	Description	DDBJ Accession no.
Aalb-SP	*A. albopictus*	Lab. Colony	8	Originated from Singapore in 2016. This strain is known to possess 1534C *kdr* variant (Kasai et al., 2019).	DRR167925–932
Aalb-Viet	*A. albopictus*	Lab. Colony	2	Originated from Hanoi, Viet Nam in 2016.	DRR167935–936
Aalb-Okayama	*A. albopictus*	Lab. Colony	2	Originated from Okayama, Japan in 2015.	DRR167923–924
Aalb-Toyama	*A. albopictus*	Lab. Colony	2	Originated from Tokyo, Japan in 2015.	DRR167933–934
Aalb-Ishigaki	*A. albopictus*	Lab. Colony	2	Originated from Ishigaki-zima, Okinawa, Japan in 2016.	DRR167921–922
Aalb-Yona	*A. albopictus*	Wild	8	Caught wild from Yonaguni-zima, Okinawa, Japan in 2017.	DRR167937–944
Aaeg-Mex	*A. aegypti*	Lab. Colony	16	Originated from Monterrey, Mexico in 2008.	DRR167897–912
Aaeg-SP	*A. aegypti*	Lab. Colony	8	Originated from Singapore in 2009. This strain is known to possesses 989P−1016G kdr haplotype (Kasai 2014).	DRR167913–920
Cpip-JNA	*C. quinquefasciatus*	Lab. Colony	2	This strain was selected for *CYP9M10* genotype (Itokawa et al., 2011) from JHB strain originated from Johannesburg, South Africa in 2001 (Arensburger et al., 2010).	DRR167945–946
Cpip-JPP	*C. quinquefasciatus*	Lab. Colony	2	Originated from Saudi Arabia, selected by permethrin for 20 generations (Amin et al., 1989). This strain is known to possesses 1014F *kdr* variant (Hardstone et al., 2007) .	DRR167947–948
Cpip-Ryo	*C. pipiens pallens*	Lab. Colony	2	Originated from Kanagawa, Japan in 2015.	DRR167951–952
Cpip-JP_mix	*C. pipiens* form *molestus*	Lab. Colony	2	Mixed from several lab. colonies originated from different places of Japan in 2003–2004.	DRR167949–950

Table 2 Detected amino-acid substitutions

Species	Population	n	Amino-acid substitutions (num. of homozygous, heterozygous individuals)
A. albopictus	Aalb-SP	8	F1534C**(8,0); A2023T(8,0)
	Aalb-Viet	2	A2023T(0,1)
	Aalb-Okayama	2	C749*Y(0,1); A2023T(0,1); G2046E(0,1)
	Aalb-Toyama	2	A2023T(0,1)
	Aalb-Ishigaki	2	A2023T(1,0)
	Aalb-Yona	8	A2023T(0,2)
A. aegypti	Aaeg-Mex	16	V410L**(1,7); S723T(1,7); V1016I**(1,7); F1534C**(16,0)
	Aaeg-SP	8	S989P**(8,0); V1016G**(8,0)
C. quinquefasciatus	Cpip-JNA	2	K109R(0,1); T1632S(0,1); E1633D(0,1); G2051A(0,1); A2055V(0,1)
C. quinquefasciatus	Cpip-JPP	2	R261K(2,0); L1014F**(2,0)
C. pipiens pallens	Cpip-Ryo	2	Y319F(2,0); T1632S(0,2); E1633D(0,2)
C. pipiens form *molestus*	Cpip-JP_mix	2	Y319F(2,0); L1014F**(2,0); T1632S(2,0); T1633D(2,0); E1856D(2,0)

The amino-acid coordination is *M. domestica* except C749Y with asterisk (*) since there was no corresponding amino-acid in *M. domestica* (genbank id: AAB47604). Double asterisks (**) indicate known *kdr* substitutions conferring pyrethroid resistance. Variants seen on the Gly repeats near the C-terminal are omitted.

Fig. 1 Analytical pipeline MoNaS

   A diagram for analytical pipeline MoNaS. MoNaS executes several bioinformatic tools to call variants and aa substitutions. Finally, a custom script converts species aa coordinates to the standard housefly aa coordinates, tells whether each aa substitution is among the known listed known *kdr* substitutions and creates a human-readable table from Variant Call Format (VCF).

Fig. 2 The NGS reads are enriched in targeted *VGSC* exons by capture

　A distribution of the numbers of reads per kilobase of *VGSC* exons per million sequenced reads (on a logarithmic scale) in all individual samples. Legend: "unrmdup", raw mapped reads, "rmdup", reads after removal of possible PCR duplicates, "sim", simulated whole genome shotgun reads from each genome. Species names: "Aalb", *A. albopictus*, "Aaeg", *A. aegypti*, "Cpip", *C. pipiens* complex. The values associated with each arrow indicate size of a fold-change.

Fig. 3 Coverage of targeted *VGSC* exons

Distribution of median (MED) and minimum (MIN) depths of nucleotides (with 1 and on a logarithmic scale) within each exon and each individual sample after PCR duplicates were removed. Exons labeled with "●" contained nucleotide sites with relatively low coverage.

Fig. 4 The allele balance in genotypes containing a variant

The distribution of the balance in read depth for each allele at heterozygous or homozygous genotypes containing either SNVs or MNVs allele(s). The balance was calculated as [read depth of the first allele in "GT" info] / [total depth] in the VCF format. Gray points in *A. albopictus* are at the genotype at GGT (Gly) trinucleotide repeats in exon32.

Fig. 5 Length and conservation of the *VGSC* exons (CDSs) targeted

Length (on a logarithmic scale) and percentage identity to *A. albopictus* of each exon in *A. aegypti* and *C. pipiens* complex. Red exon names are those with low-coverage nucleotides in Fig. 3. The green area represents >60 bp length and >87.5% identity.