Sphingomyelin is critical in organizing phosphoinositide dynamics during phago-cytic uptake of *Mycobacterium tuberculosis*

Patrick Niekamp, Gaelen Guzman, Hans Leier, Ali Rashidfarrokhi, Veronica Richina, Joost Holthius, Fikadu G. Tafesse[1]

[1]Department of Molecular Microbiology & Immunology, Oregon Health & Science University (OHSU), Portland, OR 97239, USA

*Correspondence: tafesse@ohsu.edu

Abstract

Tuberculosis, the pathological result of infection by *Mycobacterium tuberculosis* (*M.tb*), represents a massive and persistent global health burden with vast overrepresentation in regions with poor access to healthcare. This pathogen survives intracellularly within lung macrophages, and has been shown to extensively reprogram the host lipid compartment to the effect of enhanced survival and immune evasion. Sphingolipids are a class of lipid with extensive functions in a wide variety of cellular contexts, including apoptosis, autophagy, stress response, and inflammation. There remains a significant knowledge gap surrounding the lipid-mediated interactions between *M.tb* and its host. Our investigation aims to explore the mechanisms through which *M.tb* harnesses host-derived sphingolipids to gain entry to the host and to survive intracellularly. To this end, we use chemical and genetic perturbations of the sphingolipid *de novo* synthesis pathway and fluorescence microscopy to show that sphingolipids are required for efficient phagocytosis of *M.tb*. We furthermore show that the dynamics of phosphoinositides at the phagosomal synapse are perturbed upon blockade of sphingolipid synthesis. Finally, we demonstrate that perturbing the homeostasis of a specific sphingolipid (sphingomyelin) through both enrichment

and depletion at the cell surface reduces *M.tb* uptake. This work represents a first step in

building comprehensive knowledge of the mechanisms of host-pathogen interactions in

M.tb, and demonstrates a cohort of methodologies that may be applied to a variety of

pathogenic contexts.

Introduction

Phagocytosis is critical for clearing dead and effete cells, and serves as a first line of

defense against pathogens such as bacteria, parasites, and fungi. Phagocytosis culmi-

nates in the engulfment and internalization of target particles, which are subsequently

degraded via passage through the endocytic pathway. This cellular process is distinct

from other forms of extracellular uptake, including endocytosis and micropinocytosis, in

that phagocytosis is necessarily initiated by the engagement of receptor to multi-valent

ligand on particles >0.5um in diameter (Levin, Grinstein, & Schlam, 2015). While all eu-

karyotic cells are capable of phagocytic uptake, the bulk of phagocytosis in mammals is

performed by professional phagocytes with a common myeloid precursor: monocytes,

macrophages, dendritic cells, neutrophils, and mast cells (Stuart & Ezekowitz, 2005). In

the context of pathogenic clearance, phagocytosis results in robust inflammatory re-

sponse and presentation of antigen to the adaptive immune response (Levin et al., 2015).

Some pathogens, however, thrive within the phagosomal compartment – of particular

note is *Mycobacterium tuberculosis* (*M.tb*). This bacterium is the etiologic agent of tuber-

culosis, an ancient and devastating disease. This pathogen is referenced in the earliest

records of human history, and currently remains a massive global health burden – in 2018,

M.tb was responsible for an estimated 1.3 million deaths among HIV-negative individuals,

and an additional 300,000 deaths among HIV-positive individuals (Daniel, 2006; *Global*

tuberculosis report 2018., 2018). A highly-specialized intracellular pathogen, *M.tb* en-

gages phagocytic receptors on alveolar macrophages and dendritic cells to establish in-

fection. It thensubverts host immune responses to evade lysosomal degradation and rep-

licate within a modified phagosomal compartment (Cambier, Falkow, & Ramakrishnan,

2014).

As *M.tb* must interact with host membranes and lipids at nearly all stages of its infec-

tive lifecycle, it is critical to develop an understanding of the specific mechanisms of host-

pathogen interactions at the lipid level. Research suggests that *M.tb* induces substantial

lipidomic reprogramming in its host. Notably, this pathogen has been shown to interfere

with Ca^{2+}-induced phagosomal maturation by interfering with the activity of sphingosine

kinase, which produces the potent signaling molecule sphingosine-1-phosphate (Malik et

al., 2003; Thompson et al., 2005). However, the specific functional roles played by host

lipid species at the site of phagocytic uptake of *M.tb* have remained relatively unstudied.

In particular, the structural and signaling functions played by sphingolipids at the cell

membrane and within the phagosome during *M.tb* infection are virtually unknown.

Sphingolipids are a complex family of natural lipids that contain a sphingoid base.

Recent research has shown that sphingolipids mediate functions beyond membrane bar-

rier components; these lipids accumulate in lipid rafts and act as secondary messengers

within the cell (El Alwani, Wu, Obeid, & Hannun, 2006; Hla & Dannenberg, 2012; Taha,

Mullen, & Obeid, 2006). These functions are relevant to many disparate biological pro-

cesses that span from apoptosis and autophagy to stress response and inflammation

(Hakomori & Igarashi, 1995; Hannun & Obeid, 2002; Snider, Alexa Orr Gandy, & Obeid,

2010). Of particular interest to the study of phagocytosis, sphingomyelin is enriched to

the exoplasmic leaflet, and has been shown to contribute to the formation of cholesterol-

enriched lipid rafts – its synthesis is mediated by sphingomyelin synthase-1 in the Golgi

network and sphingomyelin synthase-2 in lysosomes, and it is degraded at the cell mem-

brane by sphingomyelinase (García-Arribas, Alonso, & Goñi, 2016; Hla & Dannenberg,

2012; van Meer & de Kroon, 2011). Additionally, sphingosine 1-phosphate (S1P), pro-

duced by sphingosine kinase, acts through various S1P receptors to activate a wide range

of signaling processes; these S1PRs have been shown to activate several factors related

to phagocytosis, including Rac, phospholipase C, and PI_3K (Snider et al., 2010). As such,

this investigation endeavors to elucidate the roles played by this dynamic lipid class in the

phagocytic uptake of *M.tb*.

To determine the functions of sphingolipids in phagocytosis, we chemically and ge-

netically manipulated various stages of the sphingolipid metabolic pathway in multiple

lines of immunologically-relevant phagocytes of murine and human origin (macrophages,

dendritic cells, and monocyte-derived macrophages), performed uptake assays with the

intracellular pathogen *M.tb*, and visualized the progression of phagocytosis using fluores-

cence microscopy. We first demonstrate that the efficiency of phagocytic uptake of *M.tb*

is significantly impaired when bulk sphingolipid synthesis is inhibited in murine and human

host cells. In characterizing this observation, we show that the initial stages of phagocy-

tosis (particle binding and recognition) are largely unaffected by defects in sphingolipid

synthesis, but that sphingolipid-deficient cells are defective in the propagation of phago-

cytic signaling beginning at intermediate stages of phagocytosis (engulfment, and inter-

nalization). We observe that cell devoid of sphingolipids fail to convert phosphatidylinosi-

tol 4,5-*bis*phosphate to phosphatidylinositol 3,4,5-*tris*phosphate – producing an overall

defect in proper phagosome formation. Next, we show that genetically blocking the syn-

thesis of sphingomyelin at the cell membrane confers a defect in the phagocytic uptake

of *M.tb*. These findings highlight the diversity of functions that sphingolipids play during

this immunologically significant phase of the *Mycobacterium tuberculosis* infectious lifecy-

cle.

Results

Chemical inhibition and genomic editing effectively blocks sphingolipid synthesis in phag-

ocytes

Cellular proportions of sphingolipids are maintained primarily through a central *de*

novo synthesis pathway and secondarily through salvage pathways. The rate-limiting first

step of sphingolipid synthesis is mediated by the serine palmitoyltransferase (SPT) com-

plex in the ER to produce the sphingoid base from serine and palmitoyl-CoA. Just down-

stream, ceramide is produced by ceramide synthase (CerS). Ceramid is frequently con-

sidered the central hub of sphingolipid metabolism, and is converted to higher-order

sphingolipids (including sphingosine, sphingomyelin, and galactosylceramide) within the

Golgi and at the plasma membrane (Holthuis & Menon, 2014). These species are further

modified via phosphorylation, glycosylation, and sulfidation to produce a dynamic popu-

lation of dozens of distinct lipid species in constant flux (Gault, Obeid, & Hannun, 2010).

The sphingolipid synthesis pathway, and the resulting proportions of its products, can be

manipulated using an array of genetic and chemical tools.

The chemical inhibitor myriocin (Myr) blocks the activity of the SPT complex, pre-

venting the synthesis of all lipid species with the sphingoid base; similarly, the inhibitor

Fumonisin B1 (FB1) blocks ceramide synthase activity to produce a defect in the synthe-

sis of all high order sphingolipids. We confirmed the potency of these inhibitors in three

phagocytosis-competent cell lines through radioactive labeling with 3-L-(^{14}C)-serine. Total

lipids were extracted and analyzed using thin layer chromatography (TLC) and autoradi-

ography. For THP-1 and U937 human monocyte-derived macrophages, RAW 264.7 mu-

rine macrophages, and DC2.4 murine dendritic cells, three-day treatment with Myr and

FB1 (at 15uM and 5uM, respectively) resulted in significantly reduced levels of (^{14}C)-

sphingomyelin – demonstrating the capacity of these drugs to block sphingolipid synthe-

sis (Fig. 1A and B, and data not shown).

To orthogonally investigate the effects of chemical inhibition of sphingolipid syn-

thesis, we prepared genetic knockouts of Sptlc2 (one of the two primary SPT complex

members) in RAW 264.7 and DC2.4 cells using CRISPR/Cas9 genomic editing. We ver-

ified knockdown using Western blotting (Fig1C), and then assessed the capacity of these

cells to synthesize sphingolipids using TLC and autoradiography (Fig. 1D+E). To further

characterize of the effects of Sptlc2 knockout, we performed lipidomics analysis of RAW

264.7 macrophages using liquid chromatography coupled with tandem mass spectrome-

try. We found all observed sphingolipid species to be depleted in Sptlc2-/- cells relative

to wildtype levels, whereas other lipid species were unaffected (Fig. 1F).

Finally, to characterize the effects of myriocin and FB1 treatment, as well as Sptlc2

knockout, on cell survival and growth, we performed proliferation and survival assays. We

found that there was no significant difference in morphology, proliferation, or survival

(Supp. Fig. 1 and data not shown). We subsequently found that RAW 264.7 Sptlc2-/- cells

display normal membrane trafficking through in pulse-chase assessment of [^{35}S]-methio-

nine/cysteine incorporation and protein secretion (Supp. Fig. 1B).

Sphingolipid depletion results in impaired phagocytic uptake of M.tb

To investigate the role of sphingolipids during phagocytic uptake of *M.tb*, we used

Myr and FB1 to block sphingolipid synthesis in THP-1, RAW 264.7, and DC2.4 cells, in-

fected with an mCherry-expressing strain *of M.tb,* and quantified treated versus untreated

cells. Briefly, cells were infected at a multiplicity of infection (MOI) of 10 for two hours,

after which cells were fixed and stained with 1) Alexa Fluor 488-conjugated phalloidin to

visualize F-actin at cell boundary and 2) DAPI to visualize nuclei for cell count. Fluores-

cence microscopy was used to quantify uptake efficiency. Efficiency was calculated

through automated counting of cell number (by nuclei) and internalized bacteria (red flu-

orescent particles within green cell boundary).We report uptake efficiency as the ratio of

total number of internalized *M.tb* particles against the number of identified nuclei, and

normalized the uptake rate of treated cells to that of untreated cells. Both THP-1 and

U937 monocyte-derived macrophages treated with Myr or FB1 showed a significant im-

pairment in phagocytosis compared to untreated cells, with a reduction in uptake of [60

and 40% respectively] (Fig. 2A-D). RAW 264.7 macrophages and DC 2.4 dendritic cells

both displayed similar reductions in phagocytic uptake following Myr and FB1 treatment

with decreases of 80% (Fig. 2E-H).

We next assessed the capacity of Sptlc2-/- RAW 264.7 and DC 2.4 cells to inter-

nalize *M.tb* using a similar paradigm: two-hour infection with mCherry-expressing *M.tb*,

followed by fixation and staining with Alexa Fluor 488-conjugated phalloidin and DAPI.

Fluorescence microscopy was performed and analyzed as above. In agreement with our

previous results, we find that in both RAW 264.7 and DC 2.4 cells, Sptlc2 knockout results

in a significant decrease in *M.tb* uptake compared to wildtype (Fig. 2 I-L). Together with

the results of chemical inhibition, these results suggest that the maintenance of total cel-

lular sphingolipid levels through sphingolipid synthesis is required for efficient uptake of

M.tb across all of our model cell lines*.*

Sphingolipid depleted cells recognize and bind pathogen-particles, but are defective in

proper phagosome formation

Phagocytosis is strictly regulated through a signaling network that involves the dy-

namic activation and recruitment of both enzymes (such as GTPases, kinases, and phos-

phatases) and lipids – particularly several phosphatidylinositol species. As a target parti-

cle is engaged at the cell membrane, the aggregation of ligand-bound phagocytic receptor

initiates a sequence of phosphorylation that activates the Rho GTPases RAC1 and

Cdc45, which subsequently activate phosphatidylinositol 5-kinase. This kinase catalyzes

a localized increase in phosphatidylinositol 4,5-*bis*phosphate ($PI_{(4,5)}P_2$). High $PI_{(4,5)}P_2$ lev-

els initiate actin assembly at the base of the nascent phagosome, driving the membrane

into a leading edge that works around the target particle. As this leading edge progresses

away from the base of the nascent phagosome, local $PI_{(4,5)}P_2$ levels at the base are re-

duced via three mechanisms. First, it may be converted through phosphorylation by phos-

phatidylinositol 3-kinase (PI_3K) to form phosphatidylinositol 3,4,5-*tris*phosphate

($PI_{(3,4,5)}P_3$),which initiates actin disassembly and recycling at the trailing edge. In parallel,

$PI_{(4,5)}P_2$ is degraded by phosopholipase C-gamma (PLCγ) into the secondary messengers

diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP_3), which activate innate immune

response components such as NFκB. Finally, $PI_{(4,5)}P_2$ is also degraded to $PI_{(4)}P$ by phos-

phatases OCRL and INPP5B. Evidence suggests that the trailing wave of $PI_{(4,5)}P_2$ degra-

dation following phagosome assembly is critical for the complete phagocytic uptake of a

target particle (Levin et al., 2015). To date, however, there have been no reports connect-

ing phosphoinositide signaling and sphingolipids during phagocytosis.

To assess the roles played by sphingolipids during this signaling cascade, we chal-

lenged wildtype and Sptlc2-knockout RAW 264.7 cells with Alexa594-conjugated zymo-

san A particles. Zymosan A is an inert bead composed of fungal β-glucan that is engaged

by phagocytic receptors to stimulate particle engulfment in macrophages and dendritic

cells (Underhill, 2003). Using fluorescence microscopy and several transfected fusion

protein biosensors, we quantified the spatiotemporal enrichment of activated GTPases

Rac1 and Cdc42, $PI_{(4,5)}P_2$, $PI_{(3,4,5)}P_3$, and F-actin at the phagocytic cup (Schlam et al.,

2015).

First, wildtype and Sptlc2-deficient RAW 264.7 macrophages were transfected

with PAK(PBD)-YFP – this fusion construct binds activated Rac1/Cdc42 GTPases

through the p21-binding domain of the Protein Activated Kinase (Hoppe & Swanson,

2004; Schlam et al., 2015; Srinivasan et al., 2003). Cells were then challenged with zy-

mosan A and the localization of the reporter construct was quantified through live-cell

fluorescence imaging. Phagocytic uptake was partitioned into five stages (*Start*, *Recog-*

nition, *Engulfment*, *Internalization*, and *End*) and the localized enrichments of biosensor

in at least four cells were quantified for each stage. We show that both wildtype and

Spltc2-deficient cells display an increase in Rac1/Cdc42 activity at the pseudopod after

binding zymosan A particles (Fig. 3A). As the wildtype cells fully engulf particles, there is

a concomitant increase in YFP signal intensity, followed by a sharp decline in YFP signal

at the phagosome as the bead is fully internalized and Rac1/Cdc42 activity is terminated.

In contrast, over the span of imaging, Sptlc2-deficient cells fail to fully engulf or internalize

particles, and correspondingly, there is a defect in the aggregation of YFP signal. By

quantifying the degree of enrichment at each stage, we observe a marked overall reduc-

tion in the spatiotemporal activity of the Rac1 and Cdc42 GTPases in the Sptlc2 knockout

cells. (Fig. 3B).

To characterize subsequent stages of the phagocytic signaling cascade, we visu-

alized the dynamics of $PI_{(4,5)}P_2$ at the phagosome using the PH(PLCδ)-GFP reporter con-

struct (Schlam et al., 2015) – which binds to $PI_{(4,5)}P_2$ through the PH domain of phospho-

lipase-C *delta*. As above, wildtype and Sptlc2-deficient RAW 264.7 macrophages were

transfected with the reporter construct, challenged with zymosan A, and visualized using

fluorescence live-cell imaging. As with the local increases of Rac1 and Cdc42 activity at

the pseudopod, both the wildtype and the Sptlc2 knockout cells display an enrichment of

$PI_{(4,5)}P_2$ upon particle recognition. However, $PI_{(4,5)}P_2$ enrichment peaks as wildtype cells

begin engulfing particles, whereas Sptlc2-deficient cells never reach a significant enrich-

ment of reporter signal at the nascent phagosome and again fail to internalize the particles

(Fig. 3C). Quantification again shows a statistically significant gap in $PI_{(4,5)}P_2$ aggregation

at the phagosomal cup, further suggesting an impairment of phagocytic signaling follow-

ing particle recognition (Fig. 3D).

Next, we visualized $PI_{(3,4,5)}P_3$ at the site of particle engagement of wildtype and

Sptlc2 knockout RAW 264.7 macrophages by transfection with the PH(AKT)-GFP fluo-

rescent biosensor. This reporter construct is composed of the PH domain of the protein

kinase Akt and GFP (Marshall et al., 2001). As this phosphatidylinositol species is re-

quired for actin disassembly at the trailing edge of pseudopod progression, we also used

the F-actin biosensor LifeAct to visualize actin dynamics in the developing phagosome.

Unlike with Rac1/Cdc42 activity and $PI_{(4,5)}P_2$ localized concentration, Sptlc2 knockout

cells never attain the degree of $PI_{(3,4,5)}P_3$ enrichment that wildtype cells quickly reach upon

particle recognition (Fig. 3E and F). Accordingly, imaging with LifeAct demonstrates that

F-actin cycling is aberrant in Sptlc2 knockout cells (Fig. 3G and H).

Together, these data suggest that sphingolipids play a critical role in the proper

maturation of the phagosome, but are dispensable components of the early stages of

particle binding and recognition.

Perturbation of sphingomyelin levels at exoplasmic leaflet inhibits M.tb *uptake*

Because sphingomyelin is particularly enriched to the outer leaflet of the plasma

membrane, we hoped to investigate the role that this specific lipid species plays during

the phagocytic uptake of *M.tb*. First, to determine the local enrichment of sphingomyelin

at the phagocytic synapse, we briefly infected THP-1 macrophages with zymosan A par-

ticles, and stained sphingomyelin at the exoplasmic leaflet using purified GFP-lysenin.

The protein lysenin is a pore-forming toxin derived from the earthworm *Eisenia foetida*

and binds specifically to sphingomyelin – and GFP-conjugated lysenin, purified from bac-

terial expression, serves as a functional staining tool for visualizing localized concentra-

tions of sphingomyelin (Kinoshita et al., 2017; Yamaji et al., 1998). We find that after 10

minutes of exposure to zymosan, sphingomyelin is enriched at the phagocytic cup upon

receptor binding (data not shown). These data corroborate prior findings that sphingomy-

elin is enriched to domains such as lipid rafts and caveolae, and suggest that this specific

sphingolipid species plays an important role in efficient phagocytosis (Kinoshita, Suzuki,

Murata, & Matsumori, 2018).

To further characterize the role of sphingomyelin at the cell surface during *M.tb*

infection, we endeavored to manipulate the levels of sphingomyelin at the cell surface

and quantify uptake through fluorescence microscopy. Sphingomyelin is synthesized

along the secretory pathway by sphingomyelin synthase 1 (Sgms1), and within lysosomal

compartments by Sgms2. We isolated clonal knockouts of the Sgms1 gene in U937

monocytes using CRISPR/Cas-9 genomic editing. As above, we infected these and

wildtype cells for 2 hours and quantified uptake through fluorescence microscopy. We

found that Sgms1-deficient cells are impaired in their capacity to internalize *M.tb* to a

similar degree as cells depleted of all sphingolipids (Fig. 4A and B). Thus, we conclude

that sphingomyelin plays a significant role in the phagocytic uptake of *M.tb*, and that this

lipid species contributes significantly to our observed results on the effect of total sphin-

golipid depletion reported above. Further efforts to characterize the effects of sphingomy-

elin-specific depletion are ongoing.

Discussion

In this study, we report that the capacity of *M.tb* to invade host cells by engagement

of phagocytic receptors is significantly dependent on the presence of host sphingolipids

at the cell membrane. This result corroborates our prior observations on the uptake of

Candida albicans (Tafesse et al., 2015). In that publication, we demonstrated that sphin-

golipid depletion inhibits the phagocytic clearance of this fungal pathogen, and that sphin-

golipid-depleted mice are more susceptible to death by infection due to systemic candid-

iasis. The observations we report here similarly agree with prior reports that sphingolipids

– particularly ceramide – are required for defense against infection by *Pseudomonas ae-*

ruginosa. This process occurs through the formation of sphingolipid-enriched membrane

platforms on epithelial cells that enable the clustering of receptors that mediate innate

immune response to pathogens (Grassmé et al., 2003).

Using a total of four human and murine phagocytic cell models (THP-1, U937,

RAW 264.7, and DC 2.4), we demonstrate that chemical and genetic tools may be used

to deplete total levels of bulk sphingolipids, and that these sphingolipid-deficient phago-

cytes are incapable of taking up *Mycobacterium tuberculosis.* This phenotype is mirrored

in the genetic blockade of sphingomyelin synthesis at the cell surface via CRISPR/Cas9

deletion of Sphingomyelin synthase-1 in U937 human monocytes. These results demon-

strate the importance of sphingolipids in the context of this cellular process. However, it

is notable that sphingolipid depletion does not totally ablate phagocytic uptake.

There are several possible explanations to this observation. First, the sphingolipid

salvage pathway of these cells remains intact – any sphingolipids in the growth medium

may be taken up to a degree that provides cells with sufficient quantities of sphingolipids

to eventually phagocytose *M.tb* particles, despite a reduction in total efficiency. Alterna-

tively, the limited rate of uptake we observe may be due to limitations in visualization and

quantification; bacteria which may appear to be internalized may in fact not actually be

fully within the host cell. In particular, we note that there is overlap between bacterial

fluorescent signal and plasma membrane in approximately 50% of U937 Sgms1-/- cells,

whereas this is the case for only 20% of bacteria in control cells (data not shown). The

incorporation of an antibiotic treatment during infection may clarify this possibility. In

agreement with this possibility, Hauck and colleagues have used treatment of ectopic

acid sphingomyelinase enzyme to deplete cells of sphingomyelin at the exoplasmic leaflet

to show that the internalization of N. gonorrhoeae was reduced, but that its binding to the

cell surface was not affected (Hauck et al., 2000).

In addition to the initial observations of defective internalization of *M.tb* by sphin-

golipid-deficient hosts, we present a possible mechanism through which sphingolipids

enable the propagation of phagocytic signaling by coordinating the dynamics of phospho-

inositide turnover. Upon engagement of phagocytic receptor to cognate ligand, Rho

GTPases Rac1 and Cdc42 are activated and initiate the localized production of $PI_{(4,5)}P_2$.

This local increase in the concentration of $PI_{(4,5)}P_2$ initiates F-actin polymerization at the

base of the forming phagocytic cup, which drives pseudopodia around the particle. Con-

comitantly, the proper progression of the pseudopod (and the subsequent full engulfment

of the particle) necessitates F-actin disassembly and recycling; this process is driven by

the conversion of $PI_{(4,5)}P_2$ to $PI_{(3,4,5)}P_3$ via PI3K (Schlam et al., 2015). Because we observe

defects in the propagation of this signaling pathway upon depletion of sphingolipids, we

believe that sphingolipids, sphingomyelin in particular, are responsible for coordinating

the necessary interactions between kinases and phosphoinositide substrates. One can-

didate of interest for further investigation is PI3K, which may require interaction or mem-

brane tethering with one or more sphingolipids at the site of phagocytic receptor engage-

ment..

This study represents, to our knowledge, the first evidence for sphingolipids as

regulators of phosphoinositide turnover during phagocytosis. However, prior work has

shown that sphingolipids play roles in phosphoinositide signaling in other contexts. For

example, phosphatidylinositol 4-phosphate (PI$_4$P) has been implicated in the transfer of

ceramides from the ER to the Golgi by interacting with the ceramide transfer protein CERT

(Hanada et al., 2003). Additionally, Capasso and colleagues showed that sphingomyelin

production triggers dephosphorylation of PI$_4$P at the trans-Golgi as a negative feedback

mechanism, suggesting regulatory interplay that maintains cellular levels of sphingomy-

elin (Capasso et al., 2017).

Collectively, this study demonstrates that sphingolipids are indispensable for the

phagocytosis of *Mycobacterium tuberculosis*. In the context of infection with this patho-

gen, phagocytosis represents a double-edged sword. As phagocytosis represents the in-

vasive mechanism of entry for this pathogen, it may be of interest to investigate mecha-

nisms of inhibiting the interactions between host phagocytes and *M.tb.* However, phago-

cytosis also represents a significant mechanism through which professional antigen pre-

senting cells activate and hone the adaptive immune response – without the capacity to

efficiently sense and probe the internal environment, host organisms may be at greater

risk of death by systemic infection, as was observed in the context of *C. albicans* (Tafesse

et al., 2015). Despite this conundrum, it is essential to understand the mechanisms

through which pathogens harness host factors to initiate infection. This study represents

an important step in understanding the mechanisms of pathogenesis in *Mycobacterium*

tuberculosis infection; we hope that such work contributes to the development of new

therapeutic strategies to fight this ancient pathogen.

Materials and Methods

Reagents. Myriocin and Fumonisin B1 were purchased from Cayman Chemical Com-

pany. 3-L-[14C]serine with a specific activity of ??? was purchased from ???. Phorbol 12-

myristate 13-acetate (PMA) was purchased from Abcam?. 4% Paraformaldehyde was

from Fisher Scientific. Alexa Fluor® 488 Phalloidin and Zymosan A (S. cerevisiae) Bi-

oParticles Alexa Fluor 594 conjugate were purchased from Thermo Fisher. Rabbit poly-

clonal antibody against phospho-PI3 Kinase p85 (Tyr458)/p55 (Tyr199) (p-PI3K) (#4228)

was purchased from Cell Signaling Technology. Mouse Monoclonal antibody against

GAPDH (ab8245) was purchased from Abcam. Anti-rabbit IgG, HRP-linked antibody and

anti-mouse IgG, HRP-linked antibody were purchased from Cell Signaling.

Plasmids

Active (GTP-bound) Rac1/Cdc42 was detected with PAK(PBD)-YFP, a plasmid encoding

the PBD of PAK fused to YFP (Hoppe 2004.). $PI_{(4,5)}P_2$ was detected with PH(PLCδ)-GFP,

a plasmid encoding the PH domain of PLCδ fused to GFP (kindly provided by Grinstein).

$PI_{(3,4,5)}P_3$ levels were monitored by PH(AKT)-GFP, a plasmid encoding the PH domain of

AKT fused to GFP (Feng 2014). F-actin was monitored by LifeAct-GFP, a F-actin staining

peptide fused to GFP (Riedl 2008).

The plasmids used for generation of lentivirus consists of the pCDH-CMV(-)-sgRNA plas-

mid, the psPAX2 plasmid (packaging) and pMD2.G plasmid (envelope). pCW-Cas9 len-

tivirus was kindly provided by David Sabatini (Whitehead Insititute for Biomedical Re-

search)

Cell culture and drug treatment

THP-1, U937, Raw 264.7 and DC 2.4 cells were routinely cultured in RPMI medium sup-

plemented with 10% FBS (Seradigm) and 1% Pen Strep (Gibco). HeLa cells were cul-

tured in DMEM supplemented with 10% FBS and 1% Pen Strep. All cell lines were cul-

tured at 37 °C and 5% CO_2.

THP-1 and U937 cells were differentiated to macrophages by 50ng/ml PMA for 1 day in

RPMI medium supplemented with 10% FBS, after which the cells were treated with 5µM

myriocin or fumonisin B1 for 3 days in RPMI supplemented with 10% FBS and 1%PS.

Radio-active labeling and TLC

$500x10^3$ THP-1, U937, and RAW cells were seeded into a well of a 6-well plate, differen-

tiated and treated as described above (Cell culture and drug treatment). After 3 days of

drug treatment the cells were labeled with 2µCi/ml ??? of 3-L-[^{14}C]-serine for 4 hours in

Opti-MEM supplemented with the appropriate drug concentration of myriocin and FB1

and incubated at 37°C. Cells were washed two times with PBS and lipid extraction was

done following Bligh and Dyer method (Bligh and Dyer 1959). The methanol/chloroform-

lipid extracts were dried by nitrogen gas. Dried lipids were re-dissolved in a few drops of

chloroform/methanol (1:2, vol/vol) and loaded on a TLC plate. Lipids were separated by

developing the TLC plate first in acetone and then in a mixture of chloroform, methanol

and 25% ammonia solution (50:25:6, vol/vol/vol). Radiolabeled lipids were detected on a

Phosphor-Imager (Company?).

Generation of Lentivirus. (Fikadu)

Doxycycline inducible Cas9 expressing plasmid, pCW-Cas9, was kindly provided by Da-

vid Sabatini (Whitehead Institute for Biomedical Research). Lentivirus was produced by

co-transfection of HEK 293T cells with the pCDH-sgRNA/pCW-Cas9 and psPAX2 and

pMD2.G. Transfection was performed with Lipofectamine 3000 (Thermo Fisher) accord-

ing to manufacturer's instructions. Cells were cultured in DMEM supplemented with

10%FBS and the growth medium was replaced after 6 hours. 48 hours after transfection,

lentivirus-containing supernatant was harvested, centrifuge for 5mins at 1250 rpm and

filtered through a 0.45μm filter.

Generation of stable Raw264.7 inducible Cas9 cell lines.

Raw264.7 cells were infected with lentivirus (pCW-Cas9) encoding Cas9 cDNA, and were

cultured in media containing 7 μg/mL of puromycin (Sigma Aldrich).

Generation of CRISPR/Cas9-mediated Sptlc2-/- Raw264.7 cell line. (Fikadu)

Potential target sequences for CRISPR interference were found with the rules outlined in

(Mali 2013). The following seed sequences (CRISPR target sequences) preceding the

PAM motif that were found in the exon of Sptlc2 gene were used: Sptlc2 #1 GAAC-

GGCTGCGTCAAGAAC; Sptlc2 #2: AGCAGCACCGCCACCGTCG Potential off-target

effects of the seed sequence were confirmed using the NCBI Mus musculus Nucleotide

BLAST. Generation of CRISPR/Cas9-mediated Sptlc2-knockout Raw264.7 cell line was

performed as described in (Tafesse 2014). Briefly, CRISPR gBlock was designed to clone

into the restriction enzymatic site NheI/BamHI of pCDH-CMV(-) (SBI; CD515B-1) as fol-

lows: cacagtcagacagtgactcaGTGTCACAgctagcTTTCCCATGATTCCTTCATATTT-

GCATA-

TACGATACAAGGCTGTTAGAGAGATAATTAGAATTAATTTGACTGTAAACACAAAGA

TATTAGTACAAAATACGTGACGTAGAAAGTAATAATTTCTTGGGTAGTTTGCAG-

TTTTAAAATTATGTTTTAAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTT

CGATTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACAC-

CGnnnnnnnnnnnnnnnnnnn-

nGTTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAA

AAGTGGCACCGAGTCGGTGCTTTTTTTggatccTGTGCACAgtcagtcacagtcagtctac   (n:

CRISPR target sequences). The gBlock was then digested using the restriction enzymes

NheI and BamHI and ligated into pCDH-CMV(-) vector that was linearized by digesting

with the same restriction enzyme.

The Cas9-inducible cells were infected with lentivirus carrying pCDH-CMV(-)-sgRNA, and

were cultured in media containing 250 µg/mL of hygromycin B (Life Technology). To in-

duce expression of Cas9, cells were treated with 2 µg/mL of doxycycline (Clontech) for

3–5 days. Clonal selection was performed by single cell dilution on 96-well plates. The

individual colonies were collected and the expression of Sptlc2 was examined by western-

blotting using Sptlc2 antibody. Or TLC?

U937 SGMS1 -/- cell lines

U937 SGSM1 -/- cells were kindly provided by Sebastian Virreira Winter (Max Plack Insi-

tute) and the CRISPR/Cas9-mediated knock out cell lines were established as described

in Winter et al., 2016 (Winter 2016).

Culturing of *Mycobacterium tuberculosis*

Culturing and infection of *Mycobacterium tuberculosis (M.tb)* was conducted in a Bi-osafety-level 3 laboratory following general safety guidelines. *M.tb* strain H37R, express-ing constitutively mCherry, was cultured at 37°C in 7H9 Broth medium supplemented with 50µg/ml hygromycin B.

Infection of macrophages with *M.tb*

For infection of THP-1 and U937 cells, the cells were seeded into a SensoPlate™ 96-Well Glass-Bottom Plates (Greiner Bio-One) at a density of $15x10^3$ cells per well in RPMI supplemented with 10% FBS, 1% PS and 50ng/ml PMA at 37°C. 1 day after differentiation THP-1 cells were treated for 3 days with 5µM myriocin or fumonisin B1. Infection with *M.tb* was performed after 3 days of differentiation of U937 cells and 3 days of drug treat-ment of differentiated THP-1 cells. THP-1 and U937 cells were infected with a multiplicity of infection (MOI) of 50 in RPMI supplemented with 10% FBS for 4 hours at 37°C.

For infection of Raw264.7 and DC2.4 cells, $20x10^3$ cells / well were seeded 24 hours prior infection into a SensoPlate™ 96-Well Glass-Bottom Plates (Greiner Bio-One) and cul-tured in RPMI 10%FBS, 1%PS at 37°C. Raw264.7 and DC2.4 cells were infected with a MOI of 10 in RPMI supplemented with 10% FBS for 4 hours at 37°C.

After 4 hours *M.tb* containing supernatant was aspirated and cells were washed one time with PBS and then fixed with 4% PFA over night at 4°C.

Microscopy

Spinning disk confocal (SDC) imaging was performed on a Yokogawa CSU-W1 SDC microscope (Nikon, Tokyo, Japan) using a 20x Plan Apo VC (NA 0.75) objective and a 60x Plan Apo VC (NA 1.4) oil immersion objective equipped with a Andor's Zyla sCMOS

camera under the control of the Nikon acquisition software and a stage motorized in

X,Y. Live cell imaging was performed at 37°C and 5% CO_2 using a multi-position imag-

ing with an automated focusing at each time point. Representative images were ana-

lyzed and prepared using the FIJI software.

Further quantitative imaging was performed on a Keyence BZ-X700 all in one fluores-

cence microscope (Keyence Corporation, Osaka, Japan) using a 40x Plan Apo oil im-

mersion objective, and controlled using the Keyence BZ-X Viewer software. Image anal-

ysis was performed using Keyence BZ-X Analyzer software and FIJI.

Phagocytosis assay

Fixed cells were washed twice with PBS and incubated in PBS for 15 min in Permaebili-

zation Buffer (0.1% Triton X-100 and 1%BSA). Cells were stained with Phalloidin-Alexa

488 (Thermo Fisher) at a final concentration of 33nM for 60 min and washed one time

with PBS. After the Phalloidin-staining, cells were stained with 1xDAPI for 10 mins and

washed 3 times with PBS. THP-1 cells were then imaged using the SDC microscope

(Nikon) with a 20x objective and the U937, Raw264.7 and DC2.4 cells were imaged with

the Keyence BZ-X700 with a 20x objective. The parameters for imaging were kept the

same for each sample. For image analysis either the Nikon or Keyence BZ-X Analyzer

software was used. In brief: The outline of the cells was determined via the A488 signal

(Phalloidin-staining) and the number and area of mCherry (*M.tb*) signal in the A488 sig-

nal was detected by the program. In a second round the number of cells was deter-

mined by counting the nuclei via the DAPI staining. Number of bacteria / area of

mCherry-signal inside the cell was divided by the number of cells analyzed to identify

the phagocytosis efficiency.

Live imaging of phagocytosis

For transient transfection, Raw264.7 cells were seeded into a SensoPlate™ 96-Well

Glass-Bottom Plates at a cell number of 10x10^3 cells/well 24 hours prior transfection.

Transfection was conducted with Lipofectamine 3000 (Thermo Fisher) according to

manufacturer's instructions. Briefly, 100ng of plasmid DNA and 0.2µl of P3000 reagent

were mixed in 5µl Opti-MEM and 0.33µl Lipofectamine 3000 was mixed in 5µl Opti-

MEM. Both reagent dilutions were mixed together and allowed to sit for 15 min. The mix

was then added directly to the appropriate well and mixed by swirling the plate gently.

For the imaging, the growth medium was aspirated and replaced with RPMI without

phenol red containing Zymosan A Bioparticles A595 (Thermo Fisher) at an MOI of 20.

The 96-well plate were centrifuged at 1000rpm for 1 mins. The cells were imaged with

the SDC microscope (Nikon) and 3-4 different spots per well were imaged at the same

time for 60 min with capture intervals of 15 sec. The parameters for imaging were kept

the same for each sample.

Live imaging analysis

Out of all cells imaged, those which showed to recognize a zymosan particle and main-

tained a heathy phenotype throughout the whole imaging process were selected for fur-

ther analysis. Imaging analysis was performed using FIJI software (Schindelin et al.,).

The parameters for image processing were kept constant when comparing different

data sets. The area of the phagosome was selected, cropped and opened in a new win-

dow. For the original image, the threshold was set automatically on the GFP-channel

and for the threshold of the GFP-phagosome area, the threshold of the whole cell was

multiplied by 1.5 to ensure that only an increase in signal is counted (GFP-threshold cell

x 1.5 = GFP-threshold phagosome area). The threshold area at different time points

was measured by the analyze-measure function of FIJI and the results were exported

into Excel (Microsoft). The values for the GFP-threshold of the phagosome area were

divided by the values for the GFP-threshold of the whole cell at the corresponding time

points.

Immunoblotting

Differentiated THP-1 cells treated with myriocin or left untreated were infected with *M.tb*

at a MOI of 20 for 10 and 20 mins or left uninfected. And, differentiated U937 WT and

SGSM1 -/- cells were infected with *M.tb* at a MOI of 20 for 5 and 10 mins or left uninfected.

Afterwards, the washed cells were lysed and post-nuclear fractions were subjected to

12% SDS-PAGE and separated by electrophoresis, before being transferred to methanol-

activated Polyvinylidene difluoride membranes. Membranes were blocked in Tris-buff-

ered saline (TBS) supplemented with 0.1% Tween 20 (TBS-T) and 5% bovine serum

albumin (BSA) for 30 min. Primary antibodies against p-PI3K and GAPDH were used in

3% BSA TBS-T at 1:1000 dilutions and incubated over night or 60 min, respectively. HRP-

conjugated antibodies were used at a 1:5000 dilution with an incubation time of 60 min.

Membranes exposed to ECL western blotting substrate (GE Life Science) were visualized

using the ImageQuant LAS 4000. Brightness/Contrast were adjusted globally across the

entire image using the FIJI software. Immunoblots were cropped for presentation; see

Supplementary Fig. 5 for uncropped images.

References

Cambier, C. J., Falkow, S., & Ramakrishnan, L. (2014). Host evasion and exploitation

schemes of Mycobacterium tuberculosis. *Cell*, *159*(7), 1497–1509.

https://doi.org/10.1016/j.cell.2014.11.024

Daniel, T. M. (2006). The history of tuberculosis. *Respiratory Medicine*, *100*(11), 1862–

1870. https://doi.org/10.1016/j.rmed.2006.08.006

El Alwani, M., Wu, B. X., Obeid, L. M., & Hannun, Y. A. (2006). Bioactive sphingolipids

in the modulation of the inflammatory response. *Pharmacology and Therapeutics*,

112(1), 171–183. https://doi.org/10.1016/j.pharmthera.2006.04.004

García-Arribas, A. B., Alonso, A., & Goñi, F. M. (2016). Cholesterol interactions with

ceramide and sphingomyelin. *Chemistry and Physics of Lipids*, *199*, 26–34.

https://doi.org/10.1016/j.chemphyslip.2016.04.002

Gault, C., Obeid, L., & Hannun, Y. (2010). An overview of sphingolipid metabolism: from

synthesis to breakdown Introduction to Sphingolipid Metabolism. *Adv Exp Med Biol*,

688, 1–23. https://doi.org/10.1007/978-1-4419-6741-1_1

Global tuberculosis report 2018. (2018). Geneva.

Grassmé, H., Jendrossek, V., Riehle, A., von Kürthy, G., Berger, J., Schwarz, H., …

Gulbins, E. (2003). Host defense against Pseudomonas aeruginosa requires

ceramide-rich membrane rafts. *Nature Medicine*, *9*, 322. Retrieved from

https://doi.org/10.1038/nm823

Hakomori, S. I., & Igarashi, Y. (1995). Functional role of glycosphingolipids in cell

recognition and signaling. *Journal of Biochemistry*, *118*(6), 1091–1103.

https://doi.org/10.1093/oxfordjournals.jbchem.a124992

Hannun, Y. A., & Obeid, L. M. (2002). The ceramide-centric universe of lipid-mediated

cell regulation: Stress encounters of the lipid kind. *Journal of Biological Chemistry*,

277(29), 25847–25850. https://doi.org/10.1074/jbc.R200008200

Hla, T., & Dannenberg, A. J. (2012). Sphingolipid Signaling in Metabolic Disorders. *Cell*

Metabolism, *16*(4), 420–434. https://doi.org/10.1016/j.cmet.2012.06.017

Holthuis, J. C. M., & Menon, A. K. (2014). Lipid landscapes and pipelines in membrane

homeostasis. *Nature*, *510*(7503), 48–57. https://doi.org/10.1038/nature13474

Hoppe, A. D., & Swanson, J. A. (2004). Cdc42, Rac1, and Rac2 Display Distinct

Patterns of Activation during Phagocytosis. *Molecular Biology of the Cell*, *15*(8),

3509–3519. https://doi.org/10.1091/mbc.e03-11-0847

Kinoshita, M., Suzuki, K. G. N., Matsumori, N., Takada, M., Ano, H., Morigaki, K., …

Murata, M. (2017). Raft-based sphingomyelin interactions revealed by new

fluorescent sphingomyelin analogs. *Journal of Cell Biology*, *216*(4), 1183–1204.

https://doi.org/10.1083/jcb.201607086

Kinoshita, M., Suzuki, K. G. N., Murata, M., & Matsumori, N. (2018). Evidence of lipid

rafts based on the partition and dynamic behavior of sphingomyelins. *Chemistry*

and Physics of Lipids, *215*(May), 84–95.

https://doi.org/10.1016/j.chemphyslip.2018.07.002

Levin, R., Grinstein, S., & Schlam, D. (2015). Phosphoinositides in phagocytosis and

macropinocytosis. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell*

Biology of Lipids, *1851*(6), 805–823. https://doi.org/10.1016/j.bbalip.2014.09.005

Malik, Z. A., Thompson, C. R., Hashimi, S., Porter, B., Iyer, S. S., & Kusner, D. J.

(2003). Cutting Edge: Mycobacterium tuberculosis Blocks Ca2+ Signaling and

Phagosome Maturation in Human Macrophages Via Specific Inhibition of Sphingosine Kinase. *The Journal of Immunology*, *170*(6), 2811–2815. https://doi.org/10.4049/jimmunol.170.6.2811

Marshall, J. G., Booth, J. W., Stambolic, V., Mak, T., Balla, T., Schreiber, A. D., … Grinstein, S. (2001). Restricted accumulation of phosphatidylinositol 3-kinase products in a plasmalemmal subdomain during Fcγ receptor-mediated phagocytosis. *Journal of Cell Biology*, *153*(7), 1369–1380. https://doi.org/10.1083/jcb.153.7.1369

Schlam, D., Bagshaw, R. D., Freeman, S. A., Collins, R. F., Pawson, T., Fairn, G. D., & Grinstein, S. (2015). Phosphoinositide 3-kinase enables phagocytosis of large particles by terminating actin assembly through Rac/Cdc42 GTPase-activating proteins. *Nature Communications*, *6*(1), 8623. https://doi.org/10.1038/ncomms9623

Snider, A. J., Alexa Orr Gandy, K., & Obeid, L. M. (2010). Sphingosine kinase: Role in regulation of bioactive sphingolipid mediators in inflammation. *Biochimie*, *92*(6), 707–715. https://doi.org/10.1016/j.biochi.2010.02.008

Srinivasan, S., Wang, F., Glavas, S., Ott, A., Hofmann, F., Aktories, K., … Bourne, H. R. (2003). Rac and Cdc42 play distinct roles in regulating PI(3,4,5)P3 and polarity during neutrophil chemotaxis. *Journal of Cell Biology*, *160*(3), 375–385. https://doi.org/10.1083/jcb.200208179

Stuart, L. M., & Ezekowitz, R. A. B. (2005). Phagocytosis : Elegant Complexity Review, *22*, 539–550. https://doi.org/10.1016/j.immuni.2005.05.002

Tafesse, F. G., Rashidfarrokhi, A., Schmidt, F. I., Freinkman, E., Dougan, S., Dougan, M., … Ploegh, H. L. (2015). Disruption of Sphingolipid Biosynthesis Blocks

Phagocytosis of Candida albicans. *PLoS Pathogens*, *11*(10), 1–27.

https://doi.org/10.1371/journal.ppat.1005188

Taha, T. A., Mullen, T. D., & Obeid, L. M. (2006). A house divided: Ceramide,

sphingosine, and sphingosine-1-phosphate in programmed cell death. *Biochimica*

et Biophysica Acta - Biomembranes, *1758*(12), 2027–2036.

https://doi.org/10.1016/j.bbamem.2006.10.018

Thompson, C. R., Iyer, S. S., Melrose, N., VanOosten, R., Johnson, K., Pitson, S. M., …

Kusner, D. J. (2005). Sphingosine Kinase 1 (SK1) Is Recruited to Nascent

Phagosomes in Human Macrophages: Inhibition of SK1 Translocation by

Mycobacterium tuberculosis. *The Journal of Immunology*, *174*(6), 3551–3561.

https://doi.org/10.4049/jimmunol.174.6.3551

Underhill, D. M. (2003). Macrophage recognition of zymosan particles. *Journal of*

Endotoxin Research, *9*(3), 176–180. https://doi.org/10.1179/096805103125001586

van Meer, G., & de Kroon, A. I. P. M. (2011). Lipid map of the mammalian cell. *Journal*

of Cell Science, *124*(1), 5–8. https://doi.org/10.1242/jcs.071233

Yamaji, A., Sekizawa, Y., Emoto, K., Sakuraba, H., Inoue, K., Kobayashi, H., & Umeda,

M. (1998). Lysenin, a novel sphingomyelin-specific binding protein. *The Journal of*

Biological Chemistry, *273*(9), 5300–5306. Retrieved from

http://www.ncbi.nlm.nih.gov/pubmed/9478988

TLC: THP-1   TLC: RAW264.7   TLC: RAW264.7

WB

612   Figure 1.

615   Figure 2.