Sphingomyelin biosynthesis is critical in organizing phosphoinositide turnover during

phagocytic uptake of *Mycobacterium tuberculosis*

Patrick Niekamp[,1,2], Gaelen Guzman[*,1], Hans Leier[1], Ali Rashidfarrokhi[3], Veronica Richina[4],*

Fabian Pott[1], Joost CM Holthuis[2], Fikadu G. Tafesse[1#]

[*]These authors contributed equally.

[1]Department of Molecular Microbiology and Immunology, Oregon Health & Science University,

3181 SW Sam Jackson Park Road, Portland, OR 97239.

[2]Biology & Chemistry Department, University of Osnabrück, Fachbereich Biologie/Chemie,

Barbarastrasse 13, 49076 Osnabrück, GERMANY. [3]Department of Pathology and Department of

Medicine, New York University Langone, 550 First Avenue, MSB 594, New York, NY 10016.

[4]Department of Biomedicine University Basel Hebelstrasse 20 4031 Basel Switzerland.

[#]Correspondence: tafesse@ohsu.edu

Abstract

Phagocytosis by alveolar macrophages is the obligate first step of infection by *Mycobacterium*

tuberculosis, yet the mechanisms of this potential vulnerability in the bacterial life cycle are

incompletely understood. Here we show that sphingolipids, a lipid class enriched in the outer

leaflet of plasma membranes, are required for *M. tuberculosis* to efficiently enter and infect

phagocytes. Genetic knockout or inhibition of serine palmitoyltransferase or ceramide synthases,

key enzymes that catalyze the *de novo* sphingolipid biosynthesis, greatly reduces *M. tuberculosis*

uptake by diverse phagocytic cell types without affecting other forms of endocytosis. This

phenotype is caused by multiple defects in the cellular processes that are critical for the

formation of a phagocytic cup. While the clustering of innate immune receptors such as Dectin-1

is normal, the exclusion of the regulatory phosphatase CD45 from the receptor engagement site

at the phagocytic synapse is defective in cells deficient in sphingolipid biosynthesis. We also

show that the turnover of phosphoinositides and activation of small GTPases at the phagosomal

synapse are impaired upon blockade of sphingolipid synthesis. Human monocytes deficient in

sphingomyelin biosynthesis show reduced *M. tuberculosis* uptake comparable to that of cells

depleted of total sphingolipids, indicating that sphingomyelin is specifically required for

phagocytosis. Together, these data identify the sphingomyelin biosynthetic pathway as a critical

cellular process mediating the earliest stage of *M. tuberculosis* invasion and establish the

significance of this lipid in innate immune responses.

Introduction

The etiologic agent behind an estimated 10 million new cases of tuberculosis and 1.6 million deaths in 2017 alone, *Mycobacterium tuberculosis* (Mtb) has established itself as a highly effective and insidious pathogen[1,2]. Mtb invades alveolar immune cells through phagocytosis, and wields an extensive repertoire of effector molecules that operate to dampen the immune response, reprogram the host lipidome, and promote long-term intracellular survival[3,4]. As an intracellular pathogen, Mtb necessarily interacts with its host along numerous lipid interfaces throughout its infectious lifecycle. Understanding the biological roles of host lipids during infection may illuminate novel routes of therapeutic strategies to prevent the onset and spread of tuberculosis[5].

In most settings, phagocytosis is critical for clearing dead and effete cells, and serves as a first line of defense against microbial pathogens. This process culminates in the engulfment and internalization of target particles, which are subsequently degraded following convergence of the phagosome with the lysosome[6]. In the context of pathogen clearance, uptake by professional phagocytes (primarily monocytes, macrophages, and dendritic cells) results in a robust inflammatory response and antigen presentation that may contribute to an adaptive immune response[7]. For example, previous work shows that phagocytic uptake of *Candida albicans* is essential for the clearance of infection *in vivo*[8]. In contrast, Mtb is highly adept at subverting its own degradation to survive within phagosome-derived compartments through an array of mechanisms. As one example, this pathogen inhibits the Ca^{2+}-dependent fusion of the phagosome to the lysosome by disrupting the activity of sphingosine kinase[6,9]. Cholesterol is reported to be required for Mtb entry[10]. Despite extensive literature on the host-pathogen interactions of Mtb, the specific functional roles played by host lipid species at the site of initial phagocytic uptake of Mtb have remained relatively unstudied. In particular, structural and signaling functions

played by sphingolipids at the cell membrane and within the phagosome during Mtb infection are

unknown.

Sphingolipids are a complex family of membrane lipids that contain a sphingoid base. A

species of particular note, sphingomyelin constitutes a significant fraction of the total lipid

composition of the exoplasmic leaflet of the plasma membrane and plays critical roles in the

organizing membrane dynamics at the cell surface[11,12]. A growing body of literature shows that

sphingolipids mediate functions beyond existing as membrane barrier components: members of

this family accumulate in specialized lipid domains and act as secondary messengers within the

cell[13–15]. These signaling functions are relevant to many biological processes that span

from apoptosis and autophagy to stress response and inflammation in many (if not all) mammalian

cell types[16–18]. The cellular proportions of ceramide and spingosine-1-phosphate have been shown

to act as a rheostat that determines cell survival under stress[19]. Notably, it has been shown that the

biosynthetic flow of sphingolipids through the Golgi creates a regulatory feedback loop that

controls the levels of phosphatidylinositol-4-phosphate, although interactions outside of the *trans*-

Golgi network are unknown[20].

Prior work has investigated the roles played by sphingolipids during the phagocytosis of

other particles – contradictory reports describe the effect of sphingolipid depletion during

phagocytosis. We have previously reported that cellular sphingolipids are critical for the *in vivo*

clearance of *Candida albicans* infection through phagocytosis by macrophages[8]. In contrast,

depletion of cellular sphingolipids reportedly enhances phagocytic uptake of opsonized particles[21].

Together, these results suggest that the roles played by sphingolipids during phagocytosis may be

pathogen or particle specific, based on the specific ligands decorating a particle[22]. However, the

role of sphingolipids during entry of Mtb remained poorly understood.

Here, we employed chemical and genetic tools to manipulate levels of sphingolipids in

mammalian phagocytes, and infected them using an mCherry-expressing reporter strain of H37Rv

Mtb. We first show that host sphingolipid depletion reduces the capacity of Mtb to invade host

cells through phagocytosis – but that other mechanisms of endocytosis are unaffected. We next

show that sphingolipid depletion resulted in the impairment of the regulatory phosphatase CD45

displacement by phagocytic receptors at the site of particle engagement. Furthermore, we

demonstrate that blockade of sphingolipid biosynthesis results in defects of phagocytic signaling,

including an unsuccessful activation and recruitment of the small GTPases Rac1 and CD42 to the

phagocytic cup. This in turn resulted in only transient and futile turnover of phosphoinositidyl-4,5-

*bis*phosphate, a critical process for actin remodeling during phagosome formation. Finally, we

show that specifically inhibiting the *de novo* synthesis of sphingomyelin results in a robust

phenocopy of total blockade of sphingolipid biosynthesis. Together, these data demonstrate that

intact sphingomyelin biosynthesis is required for the earliest stages of *M. tuberculosis* invasion

and highlights the importance of this host lipid in innate immune responses against this pathogen.

Results

Manipulating sphingolipid biosynthesis in mammalian phagocytes using genetic and chemical tools

To define the role of the sphingolipid biosynthetic pathway in phagocytic uptake of Mtb, we sought to inhibit sphingolipid biosynthesis in four different phagocytic cell lines: the murine macrophage cell line RAW 246.7, the murine dendritic cell line DC2.4, the human monocyte cell lines THP-1 and U937. We first used a chemical approach to inhibition using the compounds myriocin (Myr) and fumonisin B1 (FB1). As depicted in Figure 1A, the initial and rate limiting step of sphingolipid synthesis is the production of the sphingoid base through the condensation of serine and palmitoyl-CoA, and is performed by the enzymatic complex serine palmitoyltransferase (SPT)[23]. Previously we reported that genetic ablation of Sptlc2 results in blockade of *de novo* synthesis of all sphingolipid, as does treatment with the atypical amino acid Myr[8]. The fungal toxin FB1 acts several steps downstream in the sphingolipid biosynthetic pathway by blocking the activity of ceramide synthase[8]. As the central nexus of sphingolipid synthesis, ceramide is converted into many classes of higher sphingolipids - including sphingomyelin, glycosphingolipids, and galactosylceramides[19,24].

To verify the reported effects of myriocin and fumonisin B1 and Sptlc2 knockout, we used thin layer chromatography to visualize the incorporation of 3-L-[^{14}C]-serine into sphingolipid species in RAW 246.7, THP-1, DC2.4, and U937 cells following 3-day treatment with Myr and FB1 (Figure 1B and D and Supplementary Figure 1). Across all cell lines, treatment with these compounds resulted in significant reduction of sphingolipid content, without disrupting phosphatidylserine content – demonstrating the specificity and potency of these compounds (Figure 1E and F).

To further examine the lipid composition of cells deficient in sphingolipid biosynthesis,

we performed mass spectrometry-based lipidomic analysis of RAW 264.7 macrophages following

treatment with chemical inhibitors and Sptlc2 knockout. We examined cellular levels of

sphingosine, ceramide, sphingomyelin, and glucosylceramide. As expected, both myriocin and

Sptlc2 knockout result in significant reduction of all sphingolipid species (Figure 1 G-J). In FB1

treated cells, we observe reduction of all sphingolipid species downstream of ceramide synthesis

(the point of inhibition along the biosynthetic pathway), but we observe a concordant increase in

sphingosine (synthesized upstream of ceramide), as previously reported (Figure 1G-J)[25].

We then sought to verify the mutation in Sptllc2 gene or inhibitor treatments do not have a

pleiotropic effect. To this end, we carefully examined for any defect in the overall viability and

morphology of the cell. Despite a reduced level of sphingolipids, these cells grow normally with

no apparent defect in morphology or viability (Supplementary Figure 2). Notably, we observe

that myriocin and FB1 treatment, as well as Sptlc2 knockout, do not result in the total ablation of

cellular sphingolipid content. This is at least partially attributable to the activity of the

sphingolipid salvage pathways, through which cells may acquire sphingolipids from additives in

media – primarily serum. It is likely that cells deficient in sphingolipid biosynthesis acquire

sphingolipids from media at only the minimal level to allow cell growth, as their overall

sphingolipid content is significantly lower than in wildtype cells (Figure 1D-J). This is consistent

with our previous report that shows the Sptlc2[-/-] DC2.4 cells do not harbor any significant defect

with regards to cell survival, morphology, and membrane trafficking [8]. As described below, the when

grown in the presence of serum, the primary phenotype following blockade of sphingolipid

biosynthesis is a reduced rate of phagocytic uptake of pathogens such as Mtb.

Cells deficient in sphingolipid biosynthesis display reduced capacity to phagocytose Mtb.

To investigate the role of sphingolipids during phagocytic uptake of Mtb, we used Myr and FB1 to block sphingolipid synthesis in RAW 264.7, THP-1, DC2.4, and U937 cells. We then infected cells with an mCherry-expressing strain of Mtb to quantify rates of phagocytosis. Figure 2A depicts the experimental and analysis workflow. Briefly, cells were infected at a multiplicity of infection (MOI) of 10 for two hours, after which cells were fixed and stained with DyLight 488-conjugated phalloidin and with DAPI (to visualize F-actin at cell boundary and nuclei, respectively). Fluorescence microscopy was used to quantify uptake efficiency through automated counting of cell number (by nuclei) and internalized bacteria (red fluorescent particles within green cell boundary). We report uptake efficiency as the ratio of the total number of internalized Mtb particles against the number of identified nuclei, and normalized the uptake rate of treated cells to that of untreated cells to address variability between biological replicates. In all cell models analyzed, we observed a statistically significant reduction in Mtb uptake of approximately 50% following treatment with myriocin, and approximately 40% for FB1 (Figure 2B-I). Similarly, the knockout of Sptlc2 resulted in approximately 50% of wildtype uptake in both RAW 264.7 and DC2.4 cells (Figure 3A-D). These results suggest that the maintenance of total cellular sphingolipid levels through active sphingolipid biosynthesis is required for efficient phagocytic uptake of Mtb across four model cell lines and irrespective of the mechanism of inhibition.

We next sought to assess whether blocking sphingolipid biosynthesis affects the phagocytosis of Zymosan A, a model particle composed of β-glucan chains derived from Saccharomyces cerevisiae. Zymosan A particles engage the C-type Lectin receptor Dectin-1, expressed on professional phagocytes[8,26,27]. Dectin-1 is one of several pathogen recognition receptors that engage Mtb and initiate phagocytic signaling to facilitate host cell entry[28,29]. We

thus hypothesized that inhibiting sphingolipid biosynthesis in host cells would result in a reduced

uptake rate for this Zymosan A, reflecting that of Mtb. We found that Sptlc2-deficient RAW 264.7

cells are reduced in their capacity to uptake Zymosan A to a similar degree as to that of Mtb uptake

(Figure 3E and F). We thus used Zymosan A as a model particulate to gain further insight regarding

the mechanism in which sphingolipids are involved in phagocytosis.

Sphingolipid-depleted cells are permissive to virus that utilizes phagocytosis-like cellular
processes.

Are the effects of sphingolipid depletion restricted to phagocytic uptake? To answer this

question, we performed parallel infections of RAW 264.7 wildtype and Sptlc2 knockout cells using

a fluorescent reporter strain of Herpes Simplex Virus-1 (HSV) encoding a fusion of viral protein

VP26 and mCherry[30]. HSV is an enveloped virus that gains entry to host cells through the

phagocytosis-like process of macropinocytosis, and the viral receptors have been shown to be

nectin-1 (also known as herpesviruses entry mediator)[31,32]. It has been shown that HSV entry is

dependent on cholesterol and dynamin2[33]. At 12-hours post-infection, we visualized and quantified

rates of infection using high-content imaging as described above. Because the VP26-mCherry

reporter construct localizes to the nucleus, we did not stain cells using phalloidin and instead used

the co-localization of DAPI with mCherry as a measure of infection. We observe that there is no

significant difference in the infection rates of wildtype and Sptlc2$^{-/-}$ cells, suggesting that

sphingolipids play no significant role in the entry of HSV-1 (Figure 3G and H). These data

demonstrate that endocytic pathways are functional in sphingolipid-deficient cells, suggesting a

specific role for sphingolipids in phagocytosis of bacterial pathogens such as Mtb by immune

cells.

The exclusion of the regulatory phosphatase CD45 from the site of particle engagement is

impaired in Sptlc2$^{-/-}$ cells

It has been shown that the engagement of phagocytic receptor to ligand initiates the

accumulation of pathogen recognition receptors to the phagocytic synapse, which results in the

physical expulsion of inhibitory phosphatases such as CD45 from the phagocytic synapse and the

formation of a lipid raft-like domain[34]. Prior work has shown that the several intracellular

pathogens, including Mtb, induce their uptake in a cholesterol-dependent manner[5,35]. Because

sphingolipids such as sphingomyelin have been shown to play a role in the formation of both lipid

raft and cholesterol-rich domains, we hypothesized that cellular sphingolipid content is required

for the aggregation of the pathogen recognition receptor Dectin-1 to the site of particle

engagement, and the concomitant exclusion of CD45 from the contact site.

To address this hypothesis, we transiently transfected wildtype and Sptlc2$^{-/-}$ RAW 264.7

macrophages with a GFP-tagged Dectin-1 construct, and assessed the co-localization of this

fluorescent protein with the inhibitory phosphatase CD45 upon treatment with non-fluorescent

Zymosan A particles two minutes post-incubation. In both wildtype and Sptlc2 knockouts, we find

that Dectin-1 aggregates at the site of contact with Zymosan A within the first two minutes. As

expected, in wildtype cells the inhibitory phosphatase CD45 is efficiently displaced by Dectin-1.

To this end, the co-localization of CD45 with Dectin-1 in wildtype cells is significantly reduced at

the site of particle engagement (Figure 4A and B). In contrast, the co-localization of Dectin-1-GFP

and CD45 in Sptlc2 knockout cells is only minimally reduced (Figure 4A and B). These data

suggest that sphingolipid biosynthesis is required to mediate the proper displacement of CD45, a

step that is critical for the initiation of phagocytic synapse formation.

To control against the possibility that deficient sphingolipid biosynthesis results in altered

surface expression of CD45 and Dectin-1, we used flow cytometry to compare surface expression

of these proteins between wildtype and Sptlc2$^{-/-}$ RAW 264.7 cells. We probed cells for endogenous

surface protein using APC-conjugated anti-CD45 antibody and FITC-conjugated anti-Dectin-1

antibody. We found agreement in overall fluorescent signal for both wildtype and Sptlc2 knockout

cells for both surface markers (Figure 4C-F). These results suggest that the levels of both CD45

and Dectin-1 at the cell surface are unaffected by blocking the biosynthesis of sphingolipids.

Activation of Rho GTPases Rac1 and Cdc42 are defective in Sptlc2$^{-/-}$ cells.

As we describe above, the initial stages of phagosomal synapse formation are impaired in

Sptlc2$^{-/-}$ cells. Downstream of receptor activation, phagocytosis is strictly regulated through a

signaling network that involves the dynamic activation and recruitment of both enzymes (such as

GTPases, kinases, and phosphatases) and lipids – particularly several phosphatidylinositol species.

To determine whether blocking sphingolipid biosynthesis perturbs downstream phagocytic

signaling events following particle engagement, we transfected wildtype and Sptlc2$^{-/-}$ cells with

Rac1/Cdc42 fluorescent biosensors to examine the activation and localization of these proteins

during phagocytosis.

As depicted in Figure 5A, the Rho GTPases Rac1 and Cdc42 are activated via

phosphorylation following engagement of phagocytic receptors to pathogenic particles such as

Mtb[36]. To visualize the activation of these GTPases, we transfected wildtype and Sptlc2$^{-/-}$ RAW

264.7 macrophages with a fluorescent biosensor consisting of the fusion of YFP to the p21-binding

domain of the p21-activated kinase (PBD(PAK)-YFP) – this biosensor enables the visualization

of localized GTPase activation[37,38]. Twenty-four hours after transfection, we infected cells using

Zymosan A particles at an MOI of 10 and performed live imaging to visualize the distribution of

PBD(PAK)-YFP in these cells. In wildtype cells, we observe increased localization of fluorescent

signal to the site of particle contact – corresponding to the expected pattern of Rac1/Cdc42

activation (Figure 5B and C, and Supplementary Movie1). In Sptlc2$^{-/-}$ cells on the other hand, we

observed only a transient (and futile) activation which resulted in significantly reduced

accumulation of a sustained fluorescent signal in Sptlc2$^{-/-}$ cells, suggesting that these cells are

defective in the activation of Rho GTPases following particle binding (Figure 5B and C, and

Supplementary Movie2).

Phosphatidylinositol turnover is impaired in Sptlc2 knockout cells.

Next, we visualized the turnover of phosphatidylinositols at the phagocytic synapse. Rho

GTPase activation initiates the conversion of phosphatidylinositol-4-phosphate (PIP) to

phosphatidylinositol-4,5-*bis*phosphate (PIP2) at the base of the phagocytic synapse. The

accumulation of PIP2 is essential to initiate the polymerization of actin filaments that drive the

leading edges of the growing phagosome around the particle[38]. Following the leading edge of PIP2

production, the enzyme phosphatidylinositol 3-kinase (PI3K) converts PIP2 into

phosphatidylinositol-3,4,5-*tris*phosphate (PIP3), which initiates a trailing edge of actin

disassembly at the base of the phagosomal cup (Figure 5A)[7]. To date, there have been no reports

connecting phosphoinositide signaling to cellular sphingolipid biosynthesis during phagocytosis.

However, sphingolipid and phosphoinositide levels within the Golgi regulate each other in a

homeostatic feedback circuit: buildup of sphingomyelin in the Golgi initiates dephosphorylation

of PI4P, which subsequently reduces the import of cholesterol sphingolipid into the Golgi[20,39].

To visualize turnover of these signaling molecules, we transfected RAW 264.7 wildtype

and Sptlc2 knockout cells with biosensors that specifically bind PIP2 and PIP3, and used live

imaging to examine their dynamics during phagocytois of Alexa55 labeled Zymosan A particles.

The first such biosensor we used consisted of the fusion of GFP to the plekstrin homology (PH)

domain of phospholipase C-delta (PLCδ), which selectively binds to PIP2 (PH(PLCδ)-GFP)[7]. We

show an enrichment of GFP signal to the site of particle engagement as wildtype cells synthesize

PIP2 following particle recognition, and that this synthesis peaks with the engulfment of the

particle (Figure 5D and E and Supplemental Movie 3). In Sptlc2[-/-] cells, however, we observe only

a transient (and futile) accumulation of GFP signal and fails to accumulate to the same degree as

in wildtype cells – and thus appear fail to synthesize PIP2 (Figure 5D and E and Supplemental

Movie 4).

We then transfected RAW 264.7 cells with a PIP3 biosensor consisting of the PH domain

of Akt fused to GFP (PH(AKT)-GFP)[40]. We visualized the localization of biosensor signal to show

that wildtype cells begin synthesizing PIP3 early in phagocytosis, and that this synthesis strongly

peaks as a particle is fully engulfed (Figure 5F and G and Supplementary Movie 5). However,

Sptlc2 knockout cells fail to produce a trailing wave of PIP3, and thus never reach the levels of

synthesis observed in wildtype cells (Figure 5F and G and Supplementary Movie 6).

Together, these results support the hypothesis that active sphingolipid biosynthesis is

required for the initiation of the phagocytic signaling cascade, as well as for the transduction of

pro-phagocytic signaling.

Sphingomyelin biosynthesis is required for efficient Mtb uptake

Which sphingolipid is most essential for enabling phagocytic signaling? Because

sphingomyelin is highly enriched to the exoplasmic leaflet of the plasma membranes, we

hypothesized that this lipid species is responsible for reduced Mtb uptake in cells depleted of total

sphingolipids[11,24,41]. Sphingomyelin constitutes a major branch of the sphingolipid biosynthetic

pathway, and the majority of its synthesis is catalyzed by the enzyme sphingomyelin synthase 1

(SGMS1) in the Golgi − as such, cells may be depleted of this specific lipid species through

CRISPR/Cas9-mediated knockout of this gene. Two SGMS1 CRISPR/Cas9 knockout clones of

U937 cells that were isolated as previously described[42]. Using these SGMS1 knockouts, we

performed Mtb uptake assays as described above. We found that SGMS1$^{-/-}$ cells are reduced in

their capacity to uptake Mtb to a similar degree as to that of cells deficient in the biosynthesis of

all sphingolipids (Figure 6A and B, and Figure 2A and B).

To address the possibility that other sphingolipids beyond sphingomyelin play roles in the

phagocytosis of Mtb, we used CRISPR-Cas9 to knock out UDP-glucose ceramide

glycosyltransferase (UGCG) in THP-1 cells (Figure 6C and D). UGCG is singularly responsible

for catalyzing the synthesis of glucosylceramide − which is in turn the precursor of complex

glycosphingolipids, a branch of sphingolipid biosynthesis comprising over 400 known lipid

species[43]. If any of these species play a significant role in the phagocytic uptake of Mtb, we

hypothesized that the inhibition of glycosphingolipid synthesis would result in a decrease in Mtb

internalization. We found that two UGCG$^{-/-}$ clones displayed no change in the rate of Mtb

phagocytosis, as compared to wildtype THP-1 cells (Figure 6C and D). Together, these data

suggest that sphingomyelin is particularly essential for the phagocytic uptake of Mtb.

Discussion

In this study, we report that the capacity of Mtb to invade host cells by engagement of

phagocytic receptors is significantly dependent on active sphingolipid biosynthesis pathway. Our

central findings are thus: First, the blockade of total sphingolipid synthesis reduces Mtb uptake by

approximately 50%. Critically, we show that blocking sphingolipid biosynthesis has no effect on

the macropinocytosis of Herpes Simplex Virus – suggesting that these cells remain otherwise

susceptible to infection via other routes of cellular uptake. We next show that blocking

sphingolipid synthesis results in defects in the exclusion of the regulatory phosphatase CD45 at

the phagocytic synapse. Furthermore, we show that several critical stages of phagocytic signaling

involving GTPase activation and phosphoinositide turnover are perturbed upon depletion of

sphingolipid synthesis, resulting in defective phagosome formation. And finally, we show that the

blockade of sphingomyelin synthesis produces a phenocopy of total sphingolipid depletion –

suggesting that sphingomyelin specifically is an essential lipid factor during Mtb uptake.

These data support a model in which sphingolipids are essential for successful activation

of signaling events upon binding of pathogen-particles such as Mtb – and are thereby critical in

initiating phagocytic synapse. Sphingolipids are instrumental in the accumulation of activated

pathogen recognition receptor at the site of particle binding. This receptor accumulation is in turn

essential for activating the phagocytic signaling cascade – including the activation of Rho GTPases

and the turnover of phosphoinositides. Without active sphingolipid biosynthesis, there is an overall

defect in phagosome formation and a reduction in phagocytic uptake of Mtb – though we find it

notable that the lack of sphingolipid biosynthesis does not completely ablate the invasive

phagocytosis of Mtb. We postulate that this may be partially due to the activity of the sphingolipid

salvage pathways, which allow cells to acquire high-order sphingolipids from the surrounding

Discussion

In this study, we report that the capacity of Mtb to invade host cells by engagement of

phagocytic receptors is significantly dependent on active sphingolipid biosynthesis pathway. Our

central findings are thus: First, the blockade of total sphingolipid synthesis reduces Mtb uptake by

approximately 50%. Critically, we show that blocking sphingolipid biosynthesis has no effect on

the macropinocytosis of Herpes Simplex Virus – suggesting that these cells remain otherwise

susceptible to infection via other routes of cellular uptake. We next show that blocking

sphingolipid synthesis results in defects in the exclusion of the regulatory phosphatase CD45 at

the phagocytic synapse. Furthermore, we show that several critical stages of phagocytic signaling

involving GTPase activation and phosphoinositide turnover are perturbed upon depletion of

sphingolipid synthesis, resulting in defective phagosome formation. And finally, we show that the

blockade of sphingomyelin synthesis produces a phenocopy of total sphingolipid depletion –

suggesting that sphingomyelin specifically is an essential lipid factor during Mtb uptake.

These data support a model in which sphingolipids are essential for successful activation

of signaling events upon binding of pathogen-particles such as Mtb – and are thereby critical in

initiating phagocytic synapse. Sphingolipids are instrumental in the accumulation of activated

pathogen recognition receptor at the site of particle binding. This receptor accumulation is in turn

essential for activating the phagocytic signaling cascade – including the activation of Rho GTPases

and the turnover of phosphoinositides. Without active sphingolipid biosynthesis, there is an overall

defect in phagosome formation and a reduction in phagocytic uptake of Mtb – though we find it

notable that the lack of sphingolipid biosynthesis does not completely ablate the invasive

phagocytosis of Mtb. We postulate that this may be partially due to the activity of the sphingolipid

salvage pathways, which allow cells to acquire high-order sphingolipids from the surrounding

environment (e.g. fetal calf serum in growth media) and incorporate these species into the cellular

lipid pool. Ongoing work seeks to address this hypothesis.

This study represents the first evidence for sphingolipids as regulators of signaling during

phagocytosis of particles such as Zymosan and Mtb. We have previously reported that active

sphingolipid biosynthesis is critical for the phagocytic clearance of *Candida albicans* both *in vitro*

and *in vivo* – though we found no insight into the mechanism behind this defect in uptake. Further,

a recent report suggests that treatment with FB1 results in reduced entry of *M. smegmatis* by

J774A.1 cells (whereas there was no effect on the entry of *E. coli*)[44]. Adding an additional layer of

complexity, another study suggests that blocking sphingolipid synthesis via treatment with FB1

results in enhanced phagocytosis of opsonized particles[22]. We postulate that these data suggest a

paradigm in which sphingolipids serve differing roles during the phagocytic uptake of different

particles: enhancing the internalization of some, and limiting that of others.

Together, we show the dependency of Mtb on the sphingolipid biosynthetic pathway to

efficiently enter and invade the host cells. In this context, phagocytosis represents a double-edged

sword: while this immune process is critical for the clearance of many pathogenic particles, Mtb

harnesses phagocytosis to gain entry to host cells and establish long-term survival. This study

represents an important step towards understanding the underlying mechanisms of cellular

processes mediating the earliest stage of Mtb invasion and contributes to the development of new

strategies to fight this ancient pathogen.

Acknowledgements

This work is supported through the National Institutes of Health (3R21AI124225-01A1)

and the Medical Research Foundation. We would like to thank the labs of Dr. David Underhill

(Cedars-Sinai Medical Center, Los Angeles, CA), Dr. Sergio Grinstein (The Hospital for Sick

Children Research Institute, Toronto, ON), Dr. Sebastian Winter (Max Planck Institute of

Biochemistry, Munich, Germany), Dr. Tim Nice (Oregon Health and Science University,

Portland, OR), and Dr. David Johnson (Oregon Health and Science University, Portland, OR) for

kindly providing reagents, cell lines, viruses, and plasmid constructs essential to the completion

of this study. We would also like to thank the members of the OHSU Cores for Advanced Light

Microscopy and Flow Cytometry for assistance and training in data collection and analysis – in

particular, we would like to thank Dr. Stefanie Kaech Petrie, Matthew Lewis, and Sara

Christensen. We would finally like to thank Madeleine Faucher, Marie Foss, and Jossef Osborn

who kindly and patiently assisted in the editing and preparation of this work.

Conflicts of interest

The authors have no conflicts of interest to declare.

Materials and Methods

Reagents.

Myriocin and FB1 were purchased from Cayman Chemical Company. 3-L-[^{14}C]-serine

was purchased from American Radioactivity. Phorbol 12-myristate 13-acetate (PMA) was

purchased from BioLegend. 4% paraformaldehyde was from Fisher Scientific. DyLight® 488

Phalloidin was purchased from Cell Signaling Technology, and Alexa Fluor 647-conjugated

Zymosan A (*S. cerevisiae*) BioParticles were purchased from Thermo Fisher. Mouse Monoclonal

antibody against GAPDH (ab8245) was purchased from Abcam. Anti-rabbit IgG HRP-linked

antibody and anti-mouse IgG HRP-linked antibody were purchased from Cell Signaling. APC-

conjugated, rat monoclonal antibody against CD45 antibody (Clone 30-F11) was purchased from

Biolegend. FITC-conjugated, rat monoclonal antibody against Dectin-1 (Clone 2A11) was

purchased from AbD Serotec. Zombie Aqua™ Fixable Viability Kit (part #77143) was purchased

from BioLegend.

Plasmids

The plasmids used for generation of lentivirus consists of the pCDH-CMV(-)-sgRNA plasmid, the

psPAX2 (packaging) and pMD2.G (envelope) plasmids. pCW-Cas9 lentivirus was purchased from

Addgene. The pMXsIP-Dectin-1-GFP was kindly provided by the lab of Dr. David Underhill

(Cedars-Sinai Medical Center, Los Angeles, CA)[45]. Active (GTP-bound) Rac1/Cdc42 was

detected with PAK(PBD)-YFP, a plasmid encoding the PBD of PAK fused to YFP[37] was obtained

from Addgene. PI(4,5)P2 was detected with PH(PLCδ)-GFP, a plasmid encoding the PH domain

of PLCδ fused to GFP. PI(3,4,5)P3 levels were monitored by PH(AKT)-GFP, a plasmid encoding

the PH domain of AKT fused to GFP[46]. These two fusion constructs were kindly provided by the

lab of Dr. Sergio Grinstein (The Hospital for Sick Children Research Institute, Toronto, ON).

Cell culture and small molecule inhibitor treatment

THP-1, U937, and DC2.4 cells were routinely cultured in RPMI medium supplemented

with 10% FBS (Seradigm) and 1% Pen Strep (Gibco). RAW 264.7 cells were cultured in DMEM

supplemented with 10% FBS and 1% PenStrep. All cell lines were cultured at 37 °C and 5% CO_2.

In plating adherent cells (RAW 264.7 and DC2.4), cells were lifted using 0.25% trypsin solution

(Gibco) and counted using hemocytometer, with Trypan Blue solution (Gibco) for counterstain.

THP-1 and U937 cells were differentiated to macrophages by 50ng/ml PMA for 1 day in

RPMI medium supplemented with 10% FBS, after which the cells were treated with 5μM myriocin

or FB1 for 3 days in RPMI supplemented with 10% FBS and 1%Pen Strep. RAW 264.7 and DC2.4

cells were treated similarly with myriocin and FB1 – however, 24-hours before infection, cells

were lifted and counted, and seeded at appropriate density in 96-well plates or on coverslips.

Biological replicates represent independent treatments with chemical inhibitors and infections on

separate days.

Metabolic labeling and thin layer chromatography

$5x10^5$ cells (wildtype, knockout, or inhibitor-treated) were seeded into a 6-well plate. Cells

were labeled with 1μCi/ml of 3-L-[^{14}C]-serine for 4 hours in Opti-MEM supplemented with the

appropriate inhibitors. Cells were then washed two times with PBS and lipid extraction was done

following the Bligh and Dyer method[47]. The methanol/chloroform-lipid extracts were dried by

nitrogen gas. Dried lipids were re-dissolved in several drops of chloroform/methanol (1:2, vol/vol)

and loaded on a TLC plate. Lipids were separated by developing the TLC plate first in acetone and

then in a mixture of chloroform, methanol and 25% ammonia solution (50:25:6, vol/vol/vol).

Radiolabeled lipids were detected on a Storm 825 Phosphor-Imager (GE Healthcare).

Generation of lentiviruses

Lentivirus production was performed as described previously[48]. Briefly, lentiviruses were

produced by co-transfection of HEK 293T cells with the lentiviral vectors that contain our gene of

interest (pCDH-sgRNA or pCW-Cas9) and the packaging plasmids (psPAX2 and pMD2.G).

Transfection was performed with Lipofectamine 3000 (Thermo Fisher) according to

manufacturer's instructions. Cells were cultured in DMEM supplemented with 10%FBS and the

growth medium was replaced after 6 hours. 48 hours after transfection, lentivirus-containing

supernatants were harvested, centrifuge for 5min at 1250 rpm and filtered through a 0.45µm filter.

Generation of CRISPR/Cas9-mediated knockout cell lines

CRISPR/Cas9-mediated genome-editing for genes involved in the sphingolipid pathway

were performed as described previously[8]. Briefly, to generate Sptlc2$^{-/-}$ knockout RAW 264.7 cells

were infected with lentivirus (pCW-Cas9) encoding Cas9 cDNA, and were cultured in media

containing 2 µg/mL of puromycin (Sigma Aldrich). Potential target sequences for CRISPR

interference were found with the rules outlined in Mali et al. 2013[49]. The following seed sequences

(CRISPR target sequences) preceding the PAM motif that were found in the exon of Sptlc2 gene

were used: Sptlc2 #1 GAACGGCTGCGTCAAGAAC; Sptlc2 #2:

AGCAGCACCGCCACCGTCG; UGCG GCTGTGGCTGATGCATTTCA. Potential off-target

effects of the seed sequence were evaluated using the NCBI Mus musculus Nucleotide BLAST.

Generation of CRISPR/Cas9-mediated Sptlc2-knockout RAW 264.7 cell line and UGCG-

knockout THP-1 cell line were performed as previously described[48]. Briefly, CRISPR gBlock was

designed to clone into the restriction enzymatic site NheI/BamHI of pCDH-CMV(-) (SBI;

CD515B-1) as follows:

cacagtcagacagtgactcaGTGTCACAgctagcTTTCCCATGATTCCTTCATATTTGCATAT

ACGATACAAGGCTGTTAGAGAGATAATTAGAATTAATTTGACTGTAAACACAAAGA

TATTAGTACAAAATACGTGACGTAGAAAGTAATAATTTCTTGGGTAGTTTGCAGTTT

TAAAATTATGTTTTAAAATGGACTATCATATGCTTACCGTAACTTGAAAGTATTTCG

ATTTCTTGGCTTTATATATCTTGTGGAAAGGACGAAACACCGnnnnnnnnnnnnnnnnnnnnG

TTTTAGAGCTAGAAATAGCAAGTTAAAATAAGGCTAGTCCGTTATCAACTTGAAAAA

GTGGCACCGAGTCGGTGCTTTTTTTggatccTGTGCACAgtcagtcacagtcagtctac (n: CRISPR

target sequences).

The gBlock was then digested using the restriction enzymes NheI and BamHI and ligated

into pCDH-CMV(-) vector that was linearized by digesting with the same restriction enzyme.

The Cas9-inducible cells were infected with lentivirus carrying pCDH-CMV(-)-sgRNA,

and were cultured in media containing 250 μg/mL of hygromycin B (Life Technology). To induce

expression of Cas9, cells were treated with 1 μg/mL of doxycycline (Clontech) for 3–5 days.

Clonal selection was performed by single cell dilution on 96-well plates. The individual colonies

were collected and the expression of Sptlc2 was examined by western-blotting using Sptlc2

antibody.

U937 SGSM1$^{-/-}$ cells were kindly provided by Sebastian Virreira Winter (Max Planck

Institute of Biochemistry, Munich, Germany) and the CRISPR/Cas9-mediated knock out cell lines

were established as described in Winter et al., 2016[42].

Culturing of Mtb

Culturing and infection of Mtb was conducted in a Biosafety-level 3 laboratory following general safety guidelines. Mtb strain H37R, expressing constitutively mCherry were obtained from the Fortune lab (Ragon Institute, Cambridge, MA). It was cultured at 37°C in 7H9 Broth medium supplemented with 50µg/ml hygromycin B. The density of the bacteria at the time of infection is between OD600 of 0.6 - 0.8.

Mtb uptake assay

For infection of THP-1 and U937 cells, the cells were seeded into a SensoPlate™ 96-Well Glass-Bottom Plates (Greiner Bio-One) at a density of $1.5x10^4$ cells per well in RPMI supplemented with 10% FBS, 1% PenStrep and 50ng/ml PMA at 37°C. One day after differentiation, cells were treated for 3 days with 5µM myriocin or 15µM FB1. These cells were then infected with a multiplicity of infection (MOI) of 10 in RPMI supplemented with 10% FBS for 2 hours at 37°C.

For infection of RAW 264.7 and DC2.4 cells, $2.0x10^4$ cells per well were seeded into a SensoPlate™ 96-Well Glass-Bottom Plates (Greiner Bio-One). 24 hour later, the cells were were infected with an MOI of 10 in RPMI supplemented with 10% FBS for 2 hours at 37°C. Cells were washed with PBS, and then fixed with 4% PFA over night at 4°C.

Phagocytosis measurement assay

Phagocytosis assay was performed as previously described[50]. Briefly, fixed cells were washed twice with PBS and incubated in PBS for 15 min in Permeabilization Buffer (0.1% Triton

X-100 and 1%BSA). Cells were stained with Phalloidin-Alexa 488 (Thermo Fisher) at a final

concentration of 33nM for 60 min and washed one time with PBS. After the Phalloidin-staining,

cells were stained with DAPI for 10 mins and washed 3 times with PBS. Cells were then imaged

using the Keyence BZ-X700 with a PlanFluor 20x objective. The parameters for imaging were

kept the same for each sample. For image analysis the Keyence BZ-X Analyzer software was used.

In brief: The outline of the cells was determined via the Alexa488 signal (Phalloidin staining) and

the number and area of mCherry (Mtb) signal in the Alexa488 signal was detected by the program.

In a second round the number of cells was determined by counting the nuclei via the DAPI staining.

Number of bacteria (mCherry-signal) inside the cell was divided by the number of cells (DAPI

signal) to identify the phagocytosis efficiency.

Live imaging of phagocytosis

For transient transfection, RAW 264.7 cells were seeded into a SensoPlate™ 96-Well

glass-bottom plates at a cell number of 1.0×10^4 cells/well 24 hours prior transfection. Transfection

was conducted with Lipofectamine 3000 according to manufacturer's instructions. For imaging,

the growth medium was aspirated and replaced with RPMI without phenol red containing Zymosan

A Bioparticles Alexa595 at an MOI of 10. The 96-well plates were centrifuged at 1000rpm for one

minute. The cells were imaged with the SDC microscope (Nikon) and 3-4 different spots per well

were imaged at the same time for 60 min with capture intervals of 15 sec. The parameters for

imaging were kept the same for each sample.

Live image analysis (quantification)

Of live cell images, individual images were selected for further analysis if they captured a

cell bound to a Zymosan particle while maintaining a healthy morphology throughout the imaging

time-course. Imaging analysis was performed using FIJI software[51]. The parameters for image

processing were kept constant when comparing different data sets. The area of the phagosome was

selected, cropped and opened in a new window. The threshold at different time points was

measured by the analyze-measure function of FIJI. The values for the GFP-threshold of the

phagosome area were divided by the values for the GFP-threshold of the whole cell at the

corresponding time points.

Dectin-1 and CD45 displacement assay

RAW 264.7 WT and Sptlc2$^{-/-}$ cells were seeded on glass coverslips in 12-well plates and

grown to 70-90% confluency overnight. Cells were transfected with pMXsIP Dectin-1-GFP

using Lipofectamine 3000 (Invitrogen) according to the manufacturer's instructions. After 24

hrs, cells were washed three times with ice-cold PBS, kept on ice for 5 min, and inoculated with

Alexa Fluor 594-conjugated Zymosan in chilled serum-free DMEM at a ratio of 10 beads per

cell. After centrifugation for 5 min at 250 x g and 4°C, cells were incubated for 5 min at 37°C,

washed three times with ice-cold PBS to remove unbound Zymosan, then fixed for 15 min with

4% paraformaldehyde.

Images were collected using an LSM880 confocal microscope using a PLAN APO 63x oil-

immersion objective and AIRY SCAN. Image analysis was performed using FIJI. Briefly, a

region-of-interest was defined around the Zymosan particle, and Pearson's correlation coefficient

between Dectin-1-GFP signal and anti-CD45-APC signal was calculated using the colocalization

function.

Flow cytometry

RAW 264.7 wildtype and Sptlc2$^{-/-}$ cells were rinsed with PBS and lifted from the flask using treatment with trypsin. Fc receptor was blocked with CD16/32 antibody (1:100) and cells were stained with ZombieAqua Live/Dead (1:200). Cells were then stained with either CD45-APC (1:200), or Dectin-1-FITC (1:50) for 30 min. Finally, cells were fixed for 20 mins in 4% paraformaldehyde and analyzed using an LSR-II flow cytometer. Cells independently stained with ZombieAqua, CD45-APC, and Dectin-1-FITC were used for compensation. Analysis was performed using FlowJo software.

Herpes Simplex Virus-1 infection

HSV-1-mCherry reporter virus was kindly provided by the lab of Dr. David Johnson (Oregon Health and Science University, Portland, Oregon). Twenty-four hours prior to infection, 1×10^4 wildtype and Sptlc2 knockout RAW 264.7 cells were seeded on glass coverslips in a 12-well plate. Cells were equally infected at an MOI ~1, and incubated for 12 hours at 37°C and 5% CO_2. Coverslips were fixed for 20 mins in 4% paraformaldehyde, stained with DAPI and then imaged on a Keyence BZ-X700. Infection rates were calculated by dividing the mCherry expressing cells with the total number of cells (DAPI signal).

References

1. Daniel, T. M. The history of tuberculosis. *Respir. Med.* **100**, 1862–1870 (2006).

2. *Global tuberculosis report 2018.* (2018).

3. Korb, V., Chuturgoon, A. & Moodley, D. Mycobacterium tuberculosis: Manipulator of Protective Immunity. *Int. J. Mol. Sci.* **17**, 131 (2016).

4. Ouimet, M. *et al.* Mycobacterium tuberculosis induces the miR-33 locus to reprogram autophagy and host lipid metabolism. *Nat. Immunol.* **17**, 677–686 (2016).

5. Fine-Coulson, K., Reaves, B. J., Karls, R. K. & Quinn, F. D. The Role of Lipid Raft Aggregation in the Infection of Type II Pneumocytes by Mycobacterium tuberculosis. *PLoS One* **7**, (2012).

6. Thompson, C. R. *et al.* Sphingosine Kinase 1 (SK1) Is Recruited to Nascent Phagosomes in Human Macrophages: Inhibition of SK1 Translocation by Mycobacterium tuberculosis. *J. Immunol.* **174**, 3551–3561 (2005).

7. Levin, R., Grinstein, S. & Schlam, D. Phosphoinositides in phagocytosis and macropinocytosis. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids* **1851**, 805–823 (2015).

8. Tafesse, F. G. *et al.* Disruption of Sphingolipid Biosynthesis Blocks Phagocytosis of Candida albicans. *PLoS Pathog.* **11**, 1–27 (2015).

9. Malik, Z. A. *et al.* Cutting Edge: Mycobacterium tuberculosis Blocks Ca2+ Signaling and Phagosome Maturation in Human Macrophages Via Specific Inhibition of Sphingosine Kinase. *J. Immunol.* **170**, 2811–2815 (2003).

10. Gatfield, J. & Pieters, J. Essential role for cholesterol in entry of mycobacteria into macrophages. *Science* **288**, 1647–50 (2000).

11.   Kinoshita, M., Suzuki, K. G. N., Murata, M. & Matsumori, N. Evidence of lipid rafts

based on the partition and dynamic behavior of sphingomyelins. *Chem. Phys. Lipids* **215**,

84–95 (2018).

12.   García-Arribas, A. B., Alonso, A. & Goñi, F. M. Cholesterol interactions with ceramide

and sphingomyelin. *Chem. Phys. Lipids* **199**, 26–34 (2016).

13.   El Alwani, M., Wu, B. X., Obeid, L. M. & Hannun, Y. A. Bioactive sphingolipids in the

modulation of the inflammatory response. *Pharmacol. Ther.* **112**, 171–183 (2006).

14.   Hla, T. & Dannenberg, A. J. Sphingolipid Signaling in Metabolic Disorders. *Cell Metab.*

16, 420–434 (2012).

15.   Taha, T. A., Mullen, T. D. & Obeid, L. M. A house divided: Ceramide, sphingosine, and

sphingosine-1-phosphate in programmed cell death. *Biochim. Biophys. Acta - Biomembr.*

1758, 2027–2036 (2006).

16.   Hakomori, S. I. & Igarashi, Y. Functional role of glycosphingolipids in cell recognition

and signaling. *J. Biochem.* **118**, 1091–1103 (1995).

17.   Hannun, Y. A. & Obeid, L. M. The ceramide-centric universe of lipid-mediated cell

regulation: Stress encounters of the lipid kind. *J. Biol. Chem.* **277**, 25847–25850 (2002).

18.   Snider, A. J., Alexa Orr Gandy, K. & Obeid, L. M. Sphingosine kinase: Role in regulation

of bioactive sphingolipid mediators in inflammation. *Biochimie* **92**, 707–715 (2010).

19.   Leier, H. C., Messer, W. B. & Tafesse, F. G. Lipids and pathogenic flaviviruses : An

intimate union. 1–7 (2018). doi:https://doi.org/10.1371/journal.ppat.1006952

20.   Capasso, S. *et al.* Sphingolipid metabolic flow controls phosphoinositide turnover at the

trans -Golgi network. *EMBO J.* **36**, 1736–1754 (2017).

21.   Hinkovska-Galcheva, V., Boxer, L., Mansfield, P. J., Schreiber, A. D. & Shayman, J. A.

Enhanced phagocytosis through inhibition of de novo ceramide synthesis. *J. Biol. Chem.*

278, 974–982 (2003).

22.   Hinkovska-Galcheva, V., Boxer, L., Mansfield, P. J., Schreiber, A. D. & Shayman, J. A.

Enhanced Phagocytosis through Inhibition of de Novo Ceramide Synthesis. *J. Biol. Chem.*

278, 974–982 (2003).

23.   Hanada, K. Serine palmitoyltransferase, a key enzyme of sphingolipid metabolism.

Biochim. Biophys. Acta **1632**, 16–30 (2003).

24.   Gault, C., Obeid, L. & Hannun, Y. An overview of sphingolipid metabolism: from

synthesis to breakdown Introduction to Sphingolipid Metabolism. *Adv Exp Med Biol* **688**,

1–23 (2010).

25.   Meivar-Levy, I., Sabanay, H., Bershadsky, A. D. & Futerman, A. H. The Role of

Sphingolipids in the Maintenance of Fibroblast Morphology. *J. Biol. Chem.* **272**, 1558–

1564 (2002).

26.   Underhill, D. M. Macrophage recognition of zymosan particles. *J. Endotoxin Res.* **9**, 176–

180 (2003).

27.   Mansour, M. K. *et al.* Dectin-1 Activation Controls Maturation of β-1,3-Glucan-

containing Phagosomes. *J. Biol. Chem.* **288**, 16043–16054 (2013).

28.   Rothfuchs, A. G. *et al.* Dectin-1 Interaction with Mycobacterium tuberculosis Leads to

Enhanced IL-12p40 Production by Splenic Dendritic Cells. *J. Immunol.* **179**, 3463–3471

(2007).

29.   Wagener, M., Hoving, J. C., Ndlovu, H. & Marakalala, M. J. Dectin-1-Syk-CARD9

Signaling Pathway in TB Immunity. *Front. Immunol.* **9**, (2018).

30.   Wisner, T. W., Sugimoto, K., Howard, P. W., Kawaguchi, Y. & Johnson, D. C.

Anterograde Transport of Herpes Simplex Virus Capsids in Neurons by both Separate and

Married Mechanisms. *J. Virol.* **85**, 5919–5928 (2011).

31.   Tiwari, V. & Shukla, D. Nonprofessional Phagocytosis Can Facilitate Herpesvirus Entry

into Ocular Cells. *Clin. Dev. Immunol.* **2012**, 1–8 (2012).

32.   Clement, C. *et al.* A novel role for phagocytosis-like uptake in herpes simplex virus entry.

J. Cell Biol. **174**, 1009–1021 (2006).

33.   Gianni, T., Gatta, V. & Campadelli-Fiume, G. V 3-integrin routes herpes simplex virus to

an entry pathway dependent on cholesterol-rich lipid rafts and dynamin2. *Proc. Natl.*

Acad. Sci. **107**, 22260–22265 (2010).

34.   Goodridge, H. S. *et al.* Activation of the innate immune receptor Dectin-1 upon formation

of a 'phagocytic synapse'. *Nature* **472**, 471–475 (2011).

35.   Grassmé, H. *et al.* Host defense against Pseudomonas aeruginosa requires ceramide-rich

membrane rafts. *Nat. Med.* **9**, 322 (2003).

36.   Mao, Y. & Finnemann, S. C. Regulation of phagocytosis by Rho GTPases. *Small*

GTPases **6**, 89–99 (2015).

37.   Hoppe, A. D. & Swanson, J. A. Cdc42, Rac1, and Rac2 Display Distinct Patterns of

Activation during Phagocytosis. *Mol. Biol. Cell* **15**, 3509–3519 (2004).

38.   Schlam, D. *et al.* Phosphoinositide 3-kinase enables phagocytosis of large particles by

terminating actin assembly through Rac/Cdc42 GTPase-activating proteins. *Nat. Commun.*

6, 8623 (2015).

39.   Hanada, K. *et al.* Molecular machinery for non-vesicular trafficking of ceramide. *Nature*

426, 803–809 (2003).

40.   Marshall, J. G. *et al.* Restricted Accumulation of Phosphatidylinositol 3-Kinase Products

in a Plasmalemmal Subdomain during Fcγ Receptor-Mediated Phagocytosis. *J. Cell Biol.*

153, 1369–1380 (2001).

41.   Subbaiah, P. V. & Sargis, R. M. Sphingomyelin: A natural modulator of membrane

homeostasis and inflammation. *Med. Hypotheses* **57**, 135–138 (2001).

42.   Virreira Winter, S., Zychlinsky, A. & Bardoel, B. W. Genome-wide CRISPR screen

reveals novel host factors required for Staphylococcus aureus α-hemolysin-mediated

toxicity. *Sci. Rep.* **6**, 24242 (2016).

43.   Wegner, M. S., Gruber, L., Mattjus, P., Geisslinger, G. & Grösch, S. The UDP-glucose

ceramide glycosyltransferase (UGCG) and the link to multidrug resistance protein 1

(MDR1). *BMC Cancer* **18**, 1–10 (2018).

44.   Viswanathan, G., Jafurulla, M., Kumar, G. A., Raghunand, T. R. & Chattopadhyay, A.

Macrophage sphingolipids are essential for the entry of mycobacteria. *Chem. Phys. Lipids*

213, 25–31 (2018).

45.   Goodridge, H. S. *et al.* Activation of the innate immune receptor Dectin-1 upon formation

of a ˜ Phagocytic synapse-™. *Nature* **472**, 471–475 (2011).

46.   Feng, S. *et al.* A Rapidly Reversible Chemical Dimerizer System to Study Lipid Signaling

in Living Cells. *Angew. Chemie Int. Ed.* **53**, 6720–6723 (2014).

47.   BLIGH, E. G. & DYER, W. J. A rapid method of total lipid extraction and purification.

Can. J. Biochem. Physiol. **37**, 911–7 (1959).

48.   Tafesse, F. G. *et al.* GPR107, a G-protein-coupled Receptor Essential for Intoxication by

Pseudomonas aeruginosa Exotoxin A, Localizes to the Golgi and Is Cleaved by Furin. *J.*

Biol. Chem. **289**, 24005–24018 (2014).

49.   Mali, P. *et al.* RNA-guided human genome engineering via Cas9. *Science* **339**, 823–6

(2013).

50.   Rashidfarrokhi, A., Richina, V. & Tafesse, F. G. Visualizing the Early Stages of

661   Phagocytosis. *J. Vis. Exp.* (2017). doi:10.3791/54646

662   51.   Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*

663   *Methods* **9**, 676–682 (2012).

Figure 1: Manipulation of cellular sphingolipid levels using genetics and chemical tools. (A) Simplified overview of sphingolipid biosynthetic pathway. Genes that are deleted using CRISPR/Cas9 system are indicated in blue and the inhibitors that are used in this study are shown in red. (B) Model phagocyte cell lines used in this study, respective species background, and genetic knockouts available. (C) Western blot probed using anti-Sptlc2 antibody in wildtype and CRISPR/Cas9 knockout DC2.4 and RAW 264.7 cells (upper), and probed against calnexin as loading control (lower). (D) Control, mutant and inhibitor-treated cells were labeled with the sphingolipid precursor 3-L-[14C]-serine, and total lipids were extracted and analyzed by thin layer chromatography (TLC) and autoradiography. (E, F) Quantification of the [14C]-SM and [14C]-PS signals from [14C]-serine labeling of RAW264.7 and THP-1 cells shown in *D*. (G-J) Total lipids were extracted from control, Sptlc2⁻/⁻ and inhibitor-treated cells and lipid profiling was performed using LC/MS. Levels of sphingosine (G), ceramide (H), sphingomyelin (I) and glucosylceramide (GlucCer) are shown. All graphs display SD of three independent experiments, and an unpaired t-test was used to analyze the significance of the data. *$p<0.05$, **$p<0.01$, *** $p < 0.001$.

Figure 2: Sphingolipid biosynthesis is required for efficient phagocytosis of Mtb. (A) Overview of the phagocytosis experiment. (B, D) Representative images of Mtb-infected RAW 264.7 (B) and DC2.4 (D) cells. (C, E) Quantification of Mtb uptake in RAW 264.7 (C) and DC2.4 (E) cells. (E, H) Representative images of infected THP-1 (F) and U937 (H) monocyte-derived macrophages. 2×10^4 THP-1 and U937 cells were plated in 96-well plate and were differentiated using 80ng/ml PMA for one day, and then treated for three days with myriocin or FB1. (G, I) Quantification of Mtb uptake in THP-1 (G) and U937 (I) cells. Three independent experiments were performed each in triplicate with at least 1,000 cells quantified per replicate. Mtb uptake was defined using total number of bacteria (mCherry signal) within cellular periphery (Phalloidin-Alexa488 signal) and cellular counts were defined by nuclear stain. Data are mean ± SD. ***p<0.001, two-tailed unpaired t-test. Data are mean ± SD. **p<0.01, ***p<0.001, two-tailed unpaired t-test.

Figure 3: Sphingolipid biosynthesis is required for efficient phagocytosis of Mtb and Zymosan A, but not for HSV entry. (A) Representative images of wildtype and Sptlc2$^{-/-}$ RAW 264.7 (A) and DC2.4 (C) cells infected with mCherry-expressing Mtb. (B, D) Quantification of Mtb uptake for experiments performed as in *A* and *C*, respectively. (E) Representative images of wildtype and Sptlc2$^{-/-}$ RAW 264.7 cells infected with Alexa fluor 594-conjugated Zymosan A particles. (F) Quantification of Zymosan A uptake uptake performed as in *E*. (G) Microscopy images of cells that were infected with HSV-mCherry (MOI of 10). At 12 h post infection, cells were fixed, stained with DAPI and analyzed by microscopy. (H) Quantification of experiments performed as described *G*. Three independent experiments were performed each in triplicate with at least 1,000 cells quantified per replicate. Data are mean ± SD. ***p<0.001, two-tailed unpaired t-test. Data are mean ± SD. **p<0.01, ***p<0.001, ns = not significant, two-tailed unpaired t-test.

Figure 4: Exclusion of CD45 by Dectin-1 from the cup formation site is impaired in Sptlc2$^{-/-}$ cells.
(A) Confocal microscopy of GFP-tagged Dectin-1-expressing RAW264.7 macrophages stimulated with
Zymosan A for 2 min and stained for CD45 (red). Zymosan particles are shown in star (*) at the
phagocytic cup. (B) Colocalization measurement of CD45 and Dectin-1 at the site of cup formation as
presented by Pearson's colocalization coefficient. (C, E) Flow cytometry of wildtype and Sptlc2$^{-/-}$
RAW264.7 cells that were stained for CD45 and Dectin-1at the cell surface. (D, F) Graphs showing
quantification of cell surface expression of the receptors described in *C* and *E*, respectively. All data are
representative of 3 independent experiments performed in triplicate. Data are mean ± SD. ns = not
significant, two-tailed unpaired t-test.

Figure 5. Sphingolipids are essential for efficient recruitment of Rac1/Cdc42 and phosphoinositide turnover at the phagocytic cup. (A) Schematic representation of the small GTPase Rac1/Cdc42 and phosphoinositide dynamics during phagocytosis. (B) Confocal images of wild type and Sptlc2$^{-/-}$ RAW264.7 cells transfected with PAK(PBD)–GFP, a biosensor for active Rac1/Cdc42, during phagocytosis. Cells were incubated with Alexa fluor 594-conjugated Zymosan A particles, and live imaging was performed. Images captured at 30-second intervals are shown. (C) A graph showing quantitation of the PAK(PBD) fluorescence intensity in the course of formation of a phagocytic cup (shown in arrow) as performed in *B*. (D, F) Time-lapse confocal microscopy images of wildtype and Sptlc2$^{-/-}$ RAW 264.7 cells expressing PH(PLCδ)-GFP, the PI(4,5)P$_2$ biosensor (D) or PH(AKT)-GFP, the (PI3,4,5)P$_3$ biosensor (F) after infection with Alexa flour 594 conjugated Zymosan A particles at MOI of 10. Representative images that are captured at 30-second intervals are shown. (E, G) Accumulation of GFP signal to site of particle binding was quantified by the ratio of GPF signal intensity at phagosome versus GFP signal across entire cell at the indicted time points for experiments performed as *D* and *F*. n= 3 independent experiments and the mean GFP intensity profiles of 10 - 20 cells are represented. Data are mean ± SD. *p<0.05, **p<0.01, two-tailed unpaired t-test, ns = not significant

Figure 6: SGMS1^{-/-} cells, but not UGCG^{-/-}, are defective in phagocytic uptake of Mtb.
(A, C) Representative images of wildtype and two clonal isolates of SGMS1 CRISPR/Cas9 knockout
U937 cells (A) and THP-1 cells deficient in UGCG (Glucosylceramide synthase) gene infected with
mCherry-expressing Mtb. Cells were activated with 80nM PMA for 24 h before Mtb infection. (B, D)
Quantification of Mtb uptake for experiments described as in *A* and *C*, respectively. Three independent
experiments were performed each in triplicate with at least 1,000 cells quantified per replicate. Data are
mean ± SD. ***$p<0.001$, ns = not significant, two-tailed unpaired t-test.

740
741

742
743 **Supplemental Figure 1: Genetic and small molecule inhibitors mediated depletion of sphingolipid in**
744 **phagocytes.** (A, B) Control, mutant and inhibitor-treated cells were labeled with the sphingolipid
745 precursor 3-L-[^{14}C]-serine, and total lipids were extracted and analyzed by thin layer chromatography and
746 autoradiography. All data are representative of 3 independent experiments.

747
748

749
750 **Supplemental Figure 2: Depletion of sphingolipids has no effect on cell morphology, growth rate or**
751 **cell division.** (A) Microscopy images of control and Sptlc2$^{-/-}$ RAW 264.7 macrophages, showing no
752 morphological difference between the mutant and control cells. (B) The cell division of the control and
753 Sptlc2$^{-/-}$ RAW 264.7 macrophages at different time points was measured and presented.

754
755
756
757