Structural basis of the differential binding of engineered knottins to integrins αVβ3 and α5β1

Johannes F. Van Agthoven [1,2,3], Hengameh Shams [4], Frank V. Cochran [5], José L. Alonso [1,2,3], James R. Kintzing [5], Kiavash Garakani [4], Brian D. Adair [1,2,3], Jian-Ping Xiong [1,2,3], Mohammad R. K. Mofrad [4], Jennifer R. Cochran [5] and M. Amin Arnaout [1,2,3*]

From the [1]Leukocyte Biology and Inflammation Program, [2]Structural Biology Program, [3]Division of Nephrology/Department of Medicine, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA, 02129, USA; [4] Departments of Bioengineering and Mechanical Engineering, University of California, Berkeley, CA, 94720, USA. [5] Departments of Bioengineering and Chemical Engineering, Stanford University, Stanford, CA, 94305, USA.

* Lead and corresponding author: EM: aarnaout1@mgh.harvard.edu

Highlights

- Knottin 2.5F binds both αVβ3 and α5β1 whereas knottin 2.5D is αVβ3-specific.
- RGD loop is conformationally flexible in apo-2.5F but constrained in 2.5D.
- The mode of interaction of each ligand with the integrin is structurally defined.
- Promiscuous binding of 2.5F may require focused motional freedom of the RGD loop.

Abstract

Targeting both integrins αVβ3 and α5β1 simultaneously appears to be more effective in cancer therapy than targeting each one alone. The structural requirements for bispecific binding of ligand to integrins has not been fully elucidated. RGD-containing knottin 2.5F binds selectively to αVβ3 and α5β1, whereas knottin 2.5D is αVβ3-specific. To elucidate the structural basis of this selectivity, we determined the structures of 2.5F and 2.5D as apo-proteins and in complex with αVβ3, and compared their interactions with integrins using molecular dynamics simulations. These studies show that 2.5D engages αVβ3 by an induced fit, but conformational selection of a flexible RGD loop accounts for high affinity selective binding of 2.5F to both integrins. The contrasting binding of the highly flexible low affinity linear RGD peptides to multiple integrins, suggests that a "Goldilocks zone" of conformational flexibility of the RGD loop in 2.5F underlies its selective binding promiscuity to integrins.

Keywords:

Keywords: Integrins; Receptors/Structure-Function; Protein crystallography; NMR; Molecular dynamics; Cancer therapy

Introduction

Heterodimeric α/β integrins comprise a large family of divalent cation-dependent adhesion receptors that mediate cell-cell and cell-matrix interactions, which underlie their essential roles in normal metazoan physiology but also in contributing to many diseases including pathologic thrombosis, inflammation, autoimmunity and cancer (Raab-Westphal et al., 2017). In response to cell-activating stimuli, intracellular signals are generated that rapidly convert integrins into a ligand-competent state, a process termed inside-out signaling (Arnaout et al., 2007). Physiologic ligands, prototyped by the Arg-Gly-Asp (RGD) sequence motif, then bind the integrin head (formed of the α-subunit propeller and β-subunit βA domains) (Xiong et al., 2002). A carboxylate group of the ligand Asp makes an electrostatic contact with a Mg^{2+} ion coordinated at a *metal-ion-dependent adhesion site* (MIDAS) of the βA domain, and the ligand Arg inserts into a pocket in the α-subunit propeller. Ligand binding induces tertiary changes in βA that are converted to quaternary changes in the ectodomain, thus forging links of the integrin cytoplasmic tails with the actin cytoskeleton to regulate cell function, a process termed outside-in signaling (Friedland et al., 2009).

Therapeutic targeting of integrins has generally focused on development of peptides or small molecules that primarily target a single integrin (Kapp et al., 2017), an approach that has been effective in platelets, where integrin αIIbβ3 is most abundant (Coller and Shattil, 2008). However, in other cell types expressing multiple integrins, high selectivity for a single integrin may promote upregulation of another integrin sharing the same ligand, leading to reduced effectiveness, drug resistance, or even paradoxical effects. This scenario may be particularly relevant in cancer cells, where primary targeting of αVβ3 with cilengitide failed to prolong survival of patients with glioblastoma (Mason, 2015), likely related to unfavorable pharmacokinetics, enhanced α5β1-mediated cell migration (Caswell et al., 2008; Christoforides et al., 2012) and agonist-like behavior (Reynolds et al., 2009). Recent studies also showed superiority of targeting αVβ3 plus α5β1 as compared to αVβ3 alone in noninvasive *in vivo* imaging of brain cancer in mice (Moore et al., 2013).

The engineered 3.5kDa miniproteins knottins 2.5D and 2.5F bind with nanomolar affinity to αVβ3 (2.5D) or to both αVβ3 and α5β1 (2.5F) (Kimura et al., 2009a). 2.5D and 2.5F only differ in four residues: two on either side of the RGD motif (Fig. 1A). In this report, we determined the solution structures of 2.5F and 2.5D and their crystal structures in complex with αVβ3. Our results show that the 2.5F and 2.5D use different binding modes to interact with αVβ3 that are critically dependent on the degree of conformational flexibility of the respective RGD loop backbone. These data suggest that flexibility of the RGD loop in 2.5F is just sufficient to allow it to bind both integrins by adopting conformations to fit both binding sites but not so large, as in linear RGD peptides, that the entropic cost of stabilizing the loop in a single conformation will compromise its high-affinity binding.

Results

Integrin binding to 2.5D and 2.5F. We measured binding of 2.5D-Fc or 2.5F-Fc fusion proteins to K562 cells stably expressing recombinant αVβ3 (αVβ3-K562) and to K562 cells, which constitutively express $\alpha_5\beta_1$ integrin. 2.5D bound to αVβ3-K562 cells with nanomolar affinity (1.4 $\pm$ 0.4 nM, Fig. 1B, D), but exhibited no measurable binding affinity to K562 cells (Fig. 1B, D), over the same range, as was also true for the scrambled knottin FN-RDG2 (where the RGD motif is replaced with RDG (Kimura et al., 2009b)). In contrast, 2.5F bound both αVβ3-K562 and K562 with similar affinities (6.9 $\pm$ 1.3nM and 9.2 $\pm$ 1.4 nM, respectively) (Fig. 1C, D). Both 2.5D and 2.5F also bound U87MG glioblastoma cells (1.7 $\pm$ 0.6 nM and 5.9 $\pm$ 1.3 nM, respectively) (Fig. 1B- D), which express high levels of αVβ3 (Dumont et al., 2009).

NMR structures of 2.5D and 2.5F. To begin to elucidate the structural basis for selectivity of knottin/integrin binding, we first determined the solution structures of 2.5D and 2.5F by NMR (Table 1). As expected, both knottins assumed the same compact structure held together by three disulfide bonds, typical of the cysteine inhibitor family (Fig. 2A, B). However, structure of the engineered RGD-containing loop flanked by prolines 3 and 11 was drastically different in the two knottins (Fig. 2A, B). In 2.5D, this loop maintains a nearly single packed conformation (Fig. 2A, C, D), but is flexible in 2.5F (Fig. 2B-D).

X-ray structures of integrin-bound 2.5D and 2.5F. Knottins 2.5D and 2.5F were each soaked into preformed αVβ3 ectodomain crystals in presence of 1 mM Mn^{2+} and the crystal structure of the respective complex was determined as previously described (Van Agthoven et al., 2014; Xiong et al., 2002). Simulated annealing composite omit maps showed clear ligand density, and allowed complete tracing of the knottin macromolecule (Fig. 3, Table 2, Supplemental Fig.1) with Real Space Cross Correlations (RSCCs) for RGD in both knottins of 0.93-0.97, suggesting almost full occupancy of the ligand. The RGD motif of each ligand inserts into the crevice between the propeller and βA domains and contacts both in an identical manner (Fig. 3A-F). The Arg[6] guanidinium of each knottin contacts α_V-Asp[218] of the propeller, with a carboxylate from Asp[8] contacting the MIDAS Mn^{2+}. In the αVβ3/2.5D structure, 2.5D residues Ala[10], Pro[11], Pro[28], Asn[29] and Phe[31] form additional van der Waals contacts with βA (Fig. 3G). In the αVβ3/2.5F structure, 2.5F-Arg[4] hydrogen bonds βA-Asn[313] and contacts the ADMIDAS metal ion indirectly through a chloride ion (Fig. 3F). Other interactions include van der Waals contacts of 2.5F residue Pro[10], Pro[11], and Phe[31] with βA (Fig. 3H). These interactions, which bury a surface area of 654.6 $Å^2$ for αVβ3/2.5F and 606.2 $Å^2$ for αVβ3/2.5D structures account for the high affinity binding of each ligand to αVβ3.

As with binding of the natural ligand FN10 or the partial agonist cilengitide to αVβ3 (Van Agthoven et al., 2014; Xiong et al., 2002), binding of 2.5D or 2.5F induced a 3.7Å inward movement of the α1 helix of the βA domain towards the MIDAS Mn^{2+}, and restructuring of the F/α7 loop (Supplemental Fig. 2A), confirming that both knottins are partial agonists. The shape of the CD loop of the βTD in both knottin/integrin structures was also comparable to the one published for αVβ3/wtFN10 (Van Agthoven et al., 2014) (Supplemental Fig. 2 B-D). However, whereas the crystal structure of the pure antagonist hFN10 bound to αVβ3 showed a h-bond between $β_3$-Glu[319] of βA and β3-Ser[674] of βTD and a visible glycan at Asn[711], both features were absent in the αVβ3/2.5F and αVβ3/2.5D structures (Supplemental Fig. 2 B-D).

Conformations of the RGD-containing loops of 2.5D and 2.5F bound to αVβ3. In contrast to the major differences in conformation of the RGD-containing loops of 2.5D and 2.5F (Fig. 2), the two loops were largely superposable in the integrin-bound state (Fig. 4A), with a root mean square deviation (r.m.s.d.) of $0.62\pm0.27Å^2$ (mean$\pm$sd). Superposing the crystal structures of the integrin-bound loops on the respective NMR structure of the lowest energy state showed dramatic differences in the RGD-containing loop of 2.5D (Fig. 4B). The r.m.s.d. of this loop in 2.5D between the integrin-bound and solution states is $3.52\pm1.01Å^2$, which is significantly higher than its narrow r.m.s.d. in solution that is maintained in all 20 conformers (0.67 ± 0.22 $Å^2$, Figure 2D, Supplemental Fig. 3). The apo-protein state is stabilized by a 4-residue type I β-turn spanning Arg[6] to Trp[9], with hydrogen bonds involving the carbonyl oxygens and amide nitrogens of Trp[9] and Arg[6], and maintains a distance of ~6.4 Å between the β carbons of Arg[6] and Asp[8] ($C_{β-β}^{RD}$)(Fig. 4C). The importance of a Trp residue immediately after RGD in forming a β-turn was previously noted (Park et al., 2002). When 2.5D is bound to αVβ3, the β-turn unfolds, with Trp[9] moving from the solvent-exposed state to form an internal van der Waals bond with Gly[5] (Fig. 4D), thus extending the $C_{β-β}^{RD}$ distance to 9Å (Fig. 4E). In contrast, the r.m.s.d. of the RGD-containing loop of 2.5F between the bound and solution structure is $3.13\pm1.64Å^2$, comparable to its r.m.s.d. in solution (2.76 ± 1.0 $Å^2$) (Fig. 4F), with conformational flexibility of the RGD loop backbone reflected in $C_{β-β}^{RD}$ varying from 4.1Å in the lowest energy state to 8.8Å in conformer #14 (Fig. 4G, Supplemental Fig. 3) that approaches the 9.1Å $C_{β-β}^{RD}$ found in αVβ3/2.5F (Fig. 4H) or RGD-bound α5β1 ($C_{β-β}^{RD}$=8.8Å) structures (Fig. 4 I).

MD simulations of knottin binding to αVβ3 and α5β1. MD simulation was used to characterize the early stages in binding of 2.5F and 2.5D to αVβ3 and α5β1. The lowest energy NMR structures of 2.5F and 2.5D were docked onto the αVβ3 and α5β1 crystal structures, resulting in four protein complexes. In each complex, the knottin was moved 9Å away from the integrin surface allowing several water layers to form between the two before simulation was initiated. Over a 500-ns run, 2.5F associated with both αVβ3 (135±37 kcal/mole) and α5β1 (81±51 kcal/mole) (Fig. 5A, B, Supplemental Movies 1 and 2), recapitulating the cell-based data (Fig. 1 B-D). The first dual contact of 2.5F with αVβ3 was detected at 0.020 ns of simulation by 2.5F-Arg[6] hydrogen-bonding β3-Thr[212] and salt-bridging β3-Asp[150], and 2.5F-Asp[8] salt-bridging β3-Arg[214] (Fig 5C). The first dual contact of 2.5F with α5β1 was detected at 0.35 ns, but the interaction stabilized at 1.25 ns through a salt-bridge between 2.5F-Asp[8] and β1-Lys[182] (Fig 5D).

As expected, 2.5D also bound αVβ3 effectively (164±37 kcal/mole) (Fig. 5A, B, Supplemental Movie 3). The low energy of interaction between 2.5D-Trp[9] and 2.5D-Gly[5] suggested that surrounding residues in the binding pocket of αVβ3 are involved in the conformational switch of 2.5D-Trp[9] from the solvent to the buried state. Consistently, over the course of the αVβ3/2.5D simulation, 2.5D-Trp[9] first formed an S-π bond with β3-Met[180] at 0.38 ns after the start of simulation, which was reinforced at 0.96 ns via an electrostatic interaction between 2.5D-Asp[8] and β3-Arg[214] (Fig. 5E). In contrast 2.5D rapidly diffused away from α5β1 after 1.1 ns of interaction (7±19 kcal/mole) (Fig. 5A, B), unable to sustain the initial binding of 2.5D-Arg[6] to β1-Gln[221] and β1-Asp[227] at 0.01 ns (not shown) with additional contacts to the integrin (Fig. 5F, Supplemental Movie 4).

Binding of 2.5D and 2.5F to native and mutant cellular αVβ3. To assess the contribution of the early contacts of 2.5D makes with αVβ3 on binding energy (Fig. 5E), we replaced β3-Met[180] with alanine (β3-Met[180] has no homolog in β1) and β3-Arg[214] with glycine (the equivalent residue in β1). MD simulations showed that the Met[180]/Arg[214]-Ala-Gly αVβ3 mutant (αVβ3[**]) sustained a significant loss in binding energy (25±19%) to 2.5D, but maintained the energy of interaction (111±19%) with 2.5F (Fig. 6A, B). To validate the MD data, we quantified the binding of Alexa-647-labeled 2.5F and 2.5D to wild-type αVβ3 and αVβ3[**], each transiently expressed in HEK293 cells. The double mutation reduced surface expression of αVβ3[**] by ~50% compared to wild-type αVβ3 (Fig. 6C). When binding of each labeled ligand to the integrin was corrected for the degree of receptor expression, binding of 2.5F to αVβ3[**] was minimally affected (85±9% of binding to wt- αVβ3, Fig. 6D), but binding of 2.5D to αVβ3[**] was markedly reduced (28±1% of binding to wt- αVβ3) (Fig. 6E).

Discussion

The present studies show that specific recognition of αVβ3 by 2.5D requires high structural plasticity of the RGD-containing loop, revealed by comparing the conformational changes in loop backbone in structures of the apo-protein and αVβ3/2.5D complex. These comparisons also reveal a pronounced induced fit binding mechanism upon complex formation with αVβ3, which also resembles the well-known interactions between antibodies and antigens (Wilson and Stanfield, 1994). These features were not observed in binding of 2.5F to αVβ3, where the RGD-containing loop of the apo-protein is flexible, with some conformers having a $C_{\beta-\beta}^{RD}$ distance comparable to that found in the αVβ3-bound state, suggesting that 2.5F binds αVβ3 by conformation selection.

MD simulations elucidated the structural basis of the induced fit that underlies binding of 2.5D to αVβ3. The RGD-containing loop in the apo-protein is stabilized by a Type I β-turn, yielding a $C_{\beta-\beta}^{RD}$ distance of ~6.4Å, which extends to the optimal 9Å distance as a result of the switch of 2.5D-Trp[9] from a solvent to a buried state. This switch appears to be driven by an early contacts with β3-Met[180] and β3-Arg[214], and is later influenced by the surrounding hydrophilic environment (β3-Tyr[166], β3-Arg[214]), as 2.5D-Asp[8] coordinates the metal ion at MIDAS. Substitution of β3-Met[180] to Ala and β3-Arg[214] to Gly (as in β1) resulted in a major loss in binding energy of 2.5D to αVβ3. This was confirmed by assessing knottin binding to HEK293 transiently expressing wild-type αVβ3 or αVβ3**. Since α5β1 lacks the equivalent Met and Arg residues, the induced fit mechanism cannot proceed, accounting for the lack of binding of 2.5D

to α5β1. The $C_{\beta-\beta}^{RD}$ distance offered by some conformers of the flexible RGD-containing loop in 2.5F also accounts for its high affinity binding to α5β1.

The differences between 2.5F and 2.5D in adapting to the ligand-binding pocket in α5β1 likely relates to the RGD flanking residues of each ligand. Notably, 2.5F harbors two more prolines Pro^5 and Pro^{10} in the RGD-containing loop (R$\underline{P}^5$RGDN$\underline{P}^{10}$P) when compared to that in 2.5D (Q$\underline{G}^5$RGDW$\underline{A}^{10}$P)(Fig. 1A)(Kimura et al., 2009b). These trans-isomers of the prolines likely introduce local rigidity through their pyrrolidine ring, limiting the backbone dihedral angle to ~ 90°, thus restraining the RGD-containing loop from adopting a β-turn fold, as in 2.5D (Krieger et al., 2005). Previous studies showed that the highly flexible linear RGD peptides have low affinity and are not specific for a particular integrin (Ruoslahti and Pierschbacher, 1987). Decreasing structural flexibility of the RGD loop by cyclization favors high affinity binding to integrins (Bogdanowich-Knipp et al., 1999), as it decreases the entropy term of the Gibbs free energy. However, our study shows that rigidifying the RGD loop limits the ability of 2.5D to bind certain integrins. The critical proline residues in the RGD-containing loop of 2.5F achieves a fine equilibrium between stability and flexibility of the RGD loop enabling a focused motional freedom (Krieger et al., 2005; Pabon and Camacho, 2017), where the RGD loop in 2.5F is flexible enough to bind both αVβ3 and α5β1 without excessive entropic contribution, which would hamper high affinity binding.

Eight of the 24 known mammalian integrins including αVβ3 and α5β1 bind to an RGD motif present in a host of natural ligands (Takada et al., 2007). High affinity peptidomimetics or RGD-like small molecules targeting single integrins have been developed. However, most of these continue to have residual but significant affinity to other RGD-binding integrins, and the selectivity of newer versions of these ligand-mimetics have not yet been fully explored in cell-based systems and at the high concentrations likely needed *in vivo* (Kapp et al., 2017). Given the ability of cancer cells to utilize both αVβ3 and α5β1 integrins for growth and metastasis, small molecules with multispecificity have been developed for potential applications in cancer therapy (Hatley et al., 2018; Sheldrake and Patterson, 2014). Despite these successes, development of multifunctional small molecule integrin antagonists that maintain high affinity and suitable pharmacokinetic properties remains a challenge (Nero et al., 2014). The engineered knottin miniproteins 2.5D and 2.5F have a number of advantages over small molecules and short peptides including exceptional structural stability, high affinity, and specificity to tumor-associated integrins (Kimura et al., 2009b; Kwan et al., 2017). Their amenability to large-scale synthesis provides a manufacturing advantage over monoclonal antibodies. These features highlight knottins as promising candidates to bridge the gap between small molecule drugs and monoclonal antibodies.

Acknowledgements. This work was supported by NIH grants DK088327 and DK48549 (to MAA), DK101628 (to JVA), from the National Institutes of Diabetes, Digestive and Kidney diseases (NIDDK) of the National Institutes of Health; the Stanford Bio-X Interdisciplinary Initiatives Program, the Stanford Cancer Institute and the Stanford Maternal & Child Health Research Institute (JRC); National Science Foundation (NSF) grant CBET-0829205 (to MRKM). This work used resources of the Extreme Science and Engineering Discovery Environment (XSEDE) awarded to MRKM, which is supported by the National Science

Foundation (NSF) (grant ACI-1053575). JRK was supported by a National Science Foundation Graduate Research Fellowship and an ARCS Scholar Award.

Author contribution. JVA, JLA, JRK, BDA and JRC designed and performed the ligand-binding studies. FVC performed the NMR studies. JVA collected the x-ray diffraction data and refined the crystal structures with JPX. HS, KG and MRKM generated the molecular dynamic simulation data. All authors interpreted the data. MAA conceived and oversaw the project and wrote the manuscript with input from all authors.

Declaration of Interests
JRC is a cofounder of xCella Biosciences. The other authors declare no competing interests.

References

Adams, P.D., Afonine, P.V., Bunkoczi, G., Chen, V.B., Davis, I.W., Echols, N., Headd, J.J., Hung, L.W., Kapral, G.J., Grosse-Kunstleve, R.W.*, et al.* (2010). PHENIX: a comprehensive Python-based system for macromolecular structure solution. Acta Crystallogr D Biol Crystallogr *66*, 213-221.

Arnaout, M.A., Goodman, S.L., and Xiong, J.P. (2007). Structure and mechanics of integrin-based cell adhesion. Curr Opin Cell Biol *19*, 495-507.

Blystone, S.D., Graham, I.L., Lindberg, F.P., and Brown, E.J. (1994). Integrin alpha v beta 3 differentially regulates adhesive and phagocytic functions of the fibronectin receptor alpha 5 beta 1. J Cell Biol *127*, 1129-1137.

Bogdanowich-Knipp, S.J., Jois, D.S., and Siahaan, T.J. (1999). The effect of conformation on the solution stability of linear vs. cyclic RGD peptides. J Pept Res *53*, 523-529.

Brooks, B.R., Brooks, C.L., 3rd, Mackerell, A.D., Jr., Nilsson, L., Petrella, R.J., Roux, B., Won, Y., Archontis, G., Bartels, C., Boresch, S.*, et al.* (2009). CHARMM: the biomolecular simulation program. J Comput Chem *30*, 1545-1614.

Caswell, P.T., Chan, M., Lindsay, A.J., McCaffrey, M.W., Boettiger, D., and Norman, J.C. (2008). Rab-coupling protein coordinates recycling of alpha5beta1 integrin and EGFR1 to promote cell migration in 3D microenvironments. J Cell Biol *183*, 143-155.

Christoforides, C., Rainero, E., Brown, K.K., Norman, J.C., and Toker, A. (2012). PKD controls alphavbeta3 integrin recycling and tumor cell invasive migration through its substrate Rabaptin-5. Dev Cell *23*, 560-572.

Coller, B.S., and Shattil, S.J. (2008). The GPIIb/IIIa (integrin alphaIIbbeta3) odyssey: a technology-driven saga of a receptor with twists, turns, and even a bend. Blood *112*, 3011-3025.

Dumont, R.A., Hildebrandt, I., Su, H., Haubner, R., Reischl, G., Czernin, J.G., Mischel, P.S., and Weber, W.A. (2009). Noninvasive imaging of alphaVbeta3 function as a predictor of the antimigratory and antiproliferative effects of dasatinib. Cancer Res *69*, 3173-3179.

Emsley, P., and Cowtan, K. (2004). Coot: model-building tools for molecular graphics. Acta Crystallogr D Biol Crystallogr *60*, 2126-2132.

Friedland, J.C., Lee, M.H., and Boettiger, D. (2009). Mechanically activated integrin switch controls alpha5beta1 function. Science *323*, 642-644.

Guntert, P., Mumenthaler, C., and Wuthrich, K. (1997). Torsion angle dynamics for NMR structure calculation with the new program DYANA. J Mol Biol *273*, 283-298.

Gupta, V., Gylling, A., Alonso, J.L., Sugimori, T., Ianakiev, P., Xiong, J.P., and Arnaout, M.A. (2007). The beta-tail domain (betaTD) regulates physiologic ligand binding to integrin CD11b/CD18. Blood *109*, 3513-3520.

Hatley, R.J.D., Macdonald, S.J.F., Slack, R.J., Le, J., Ludbrook, S.B., and Lukey, P.T. (2018). An alphav-RGD Integrin Inhibitor Toolbox: Drug Discovery Insight, Challenges and Opportunities. Angew Chem Int Ed Engl *57*, 3298-3321.

Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: visual molecular dynamics. J Mol Graph *14*, 33-38, 27-38.

Kapp, T.G., Rechenmacher, F., Neubauer, S., Maltsev, O.V., Cavalcanti-Adam, E.A., Zarka, R., Reuning, U., Notni, J., Wester, H.J., Mas-Moruno, C.*, et al.* (2017). A Comprehensive Evaluation of the Activity and Selectivity Profile of Ligands for RGD-binding Integrins. Sci Rep *7*, 39805.

Kimura, R.H., Cheng, Z., Gambhir, S.S., and Cochran, J.R. (2009a). Engineered knottin peptides: a new class of agents for imaging integrin expression in living subjects. Cancer Res *69*, 2435-2442.

Kimura, R.H., Levin, A.M., Cochran, F.V., and Cochran, J.R. (2009b). Engineered cystine knot peptides that bind alphavbeta3, alphavbeta5, and alpha5beta1 integrins with low-nanomolar affinity. Proteins *77*, 359-369.

Krieger, F., Moglich, A., and Kiefhaber, T. (2005). Effect of proline and glycine residues on dynamics and barriers of loop formation in polypeptide chains. J Am Chem Soc *127*, 3346-3352.

Kryshtafovych, A., Moult, J., Bales, P., Bazan, J.F., Biasini, M., Burgin, A., Chen, C., Cochran, F.V., Craig, T.K., Das, R.*, et al.* (2014). Challenging the state of the art in protein structure prediction: Highlights of experimental target structures for the 10th Critical Assessment of Techniques for Protein Structure Prediction Experiment CASP10. Proteins *82 Suppl 2*, 26-42.

Kwan, B.H., Zhu, E.F., Tzeng, A., Sugito, H.R., Eltahir, A.A., Ma, B., Delaney, M.K., Murphy, P.A., Kauke, M.J., Angelini, A.*, et al.* (2017). Integrin-targeted cancer immunotherapy elicits protective adaptive immune responses. J Exp Med *214*, 1679-1690.

Mason, W.P. (2015). End of the road: confounding results of the CORE trial terminate the arduous journey of cilengitide for glioblastoma. Neuro Oncol *17*, 634-635.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J. (2007). Phaser crystallographic software. J Appl Crystallogr *40*, 658-674.

Mehta, R.J., Diefenbach, B., Brown, A., Cullen, E., Jonczyk, A., Gussow, D., Luckenbach, G.A., and Goodman, S.L. (1998). Transmembrane-truncated alphavbeta3 integrin retains high affinity for ligand binding: evidence for an 'inside-out' suppressor? Biochem J *330 (Pt 2)*, 861-869.

Moore, S.J., Hayden Gephart, M.G., Bergen, J.M., Su, Y.S., Rayburn, H., Scott, M.P., and Cochran, J.R. (2013). Engineered knottin peptide enables noninvasive optical imaging of intracranial medulloblastoma. Proc Natl Acad Sci U S A *110*, 14598-14603.

Nero, T.L., Morton, C.J., Holien, J.K., Wielens, J., and Parker, M.W. (2014). Oncogenic protein interfaces: small molecules, big challenges. Nat Rev Cancer *14*, 248-262.

Otwinowski, Z., and Minor, W. (1997). Processing of X-ray diffraction data collected in oscillation mode. Methods Enzymol *276*, 307-326.

Pabon, N.A., and Camacho, C.J. (2017). Probing protein flexibility reveals a mechanism for selective promiscuity. Elife *6*.

Park, H.S., Kim, C., and Kang, Y.K. (2002). Preferred conformations of RGDX tetrapeptides to inhibit the binding of fibrinogen to platelets. Biopolymers *63*, 298-313.

Phillips, J.C., Braun, R., Wang, W., Gumbart, J., Tajkhorshid, E., Villa, E., Chipot, C., Skeel, R.D., Kale, L., and Schulten, K. (2005). Scalable molecular dynamics with NAMD. J Comput Chem *26*, 1781-1802.

Raab-Westphal, S., Marshall, J.F., and Goodman, S.L. (2017). Integrins as Therapeutic Targets: Successes and Cancers. Cancers (Basel) *9*.

Reynolds, A.R., Hart, I.R., Watson, A.R., Welti, J.C., Silva, R.G., Robinson, S.D., Da Violante, G., Gourlaouen, M., Salih, M., Jones, M.C.*, et al.* (2009). Stimulation of tumor growth and angiogenesis by low concentrations of RGD-mimetic integrin inhibitors. Nat Med *15*, 392-400.

Ruoslahti, E., and Pierschbacher, M.D. (1987). New perspectives in cell adhesion: RGD and integrins. Science *238*, 491-497.

Sheldrake, H.M., and Patterson, L.H. (2014). Strategies to inhibit tumor associated integrin receptors: rationale for dual and multi-antagonists. J Med Chem *57*, 6301-6315.

Shen, Y., Delaglio, F., Cornilescu, G., and Bax, A. (2009). TALOS+: a hybrid method for predicting protein backbone torsion angles from NMR chemical shifts. J Biomol NMR *44*, 213-223.

Takada, Y., Ye, X., and Simon, S. (2007). The integrins. Genome Biol *8*, 215.

Van Agthoven, J.F., Xiong, J.P., Alonso, J.L., Rui, X., Adair, B.D., Goodman, S.L., and Arnaout, M.A. (2014). Structural basis for pure antagonism of integrin alphaVbeta3 by a high-affinity form of fibronectin. Nat Struct Mol Biol *21*, 383-388.

Wilson, I.A., and Stanfield, R.L. (1994). Antibody-antigen interactions: new structures and new conformational changes. Curr Opin Struct Biol *4*, 857-867.

Xiong, J.P., Mahalingham, B., Alonso, J.L., Borrelli, L.A., Rui, X., Anand, S., Hyman, B.T., Rysiok, T., Muller-Pompalla, D., Goodman, S.L.*, et al.* (2009). Crystal structure of the complete integrin alphaVbeta3 ectodomain plus an alpha/beta transmembrane fragment. J Cell Biol *186*, 589-600.

Xiong, J.P., Stehle, T., Diefenbach, B., Zhang, R., Dunker, R., Scott, D.L., Joachimiak, A., Goodman, S.L., and Arnaout, M.A. (2001). Crystal structure of the extracellular segment of integrin alpha Vbeta3. Science *294*, 339-345.

Xiong, J.P., Stehle, T., Zhang, R., Joachimiak, A., Frech, M., Goodman, S.L., and Arnaout, M.A. (2002). Crystal structure of the extracellular segment of integrin alpha Vbeta3 in complex with an Arg-Gly-Asp ligand. Science *296*, 151-155.

Table 1. NMR structural statistics for 2.5D and 2.5F 20-conformer ensembles

	2.5D	2.5F
PDB Code	2M7T	6MM4
NMR-derived restraints		
Interproton total	462	340
short-range, $|i-j| \leq 1$	248	223
medium-range, $1 < |i-j| < 5$	78	48
long-range, $|i-j| \geq 5$	136	69
Dihedral angles	32	35
Hydrogen bonds	16	16
Disulfide bonds	18	18
Ramachandran statistics		
Residues in		
most favored regions (%)	86.3	87.5
additional allowed regions (%)	13.7	11.4
generously allowed regions (%)	0.0	0.5
disallowed regions (%)	0.0	0.7
RMSD statistics (residues 1-33)		
Average backbone RMSD to mean (Å)	0.26 +/- 0.06	1.04 +/- 0.28
Average heavy atom RMSD to mean (Å)	0.48 +/- 0.09	1.52 +/- 0.3

Table 2. Data collection and refinement statistics.

Data collection	αVβ3/2.5D	αVβ3/2.5F
PDB Code	6MSL	6MSU
Beamline	ID19 at APS	ID19 at APS
Space group	P3$_2$21	P3$_2$21
Unit cell dimensions (Å, °)	$a=b=129.8$, $c=305.9$; $\alpha=\beta=90$, $\gamma=120$	$a=b=129.9$, $c=305.9$; $\alpha=\beta=90$, $\gamma=120$
Resolution range (Å)	50.0-3.1	50.0-3.1
Wavelength (Å)	0.97921	0.97921
Total reflections	1,919,957	945,125
Unique reflections	54,885 (5,389)*	54,732 (5,385)
Completeness	100 (100)	100 (100)
Redundancy	6.9 (7.0)	6.2 (6.3)
Molecules in asymmetric unit	1	1
Average I/σ	14.1 (1.7)	15.3 (2.0)
R_{merge} (%)	14.9 (169.4)	12.7 (131.8)
R_{meas} (%)	16.2 (183.2)	13.9 (143.9)
R_{pim} (%)	6.2 (68.6)	5.5 (57.1)
Wilson B-factor	65.6	62.7
cc1/2	0.97 (0.52)	0.98 (0.61)
Refinement statistics		
Resolution range (Å)	49.2-3.1	49.3-3.1
R_{factor} (%)	22.7 (30.2)	22.9 (31.8)
R_{free} (%)#	26.8 (29.2)	27.6 (34.6)
No. of atoms	13,132	13,158
Protein	12,713	12,714
Water	4	4
Mn^{2+}	8	8
Glc-NAc	407	431
Chloride	0	1
Average B-factor for all atoms (Å^2)	62.3	61.4
r.m.s. deviations		
Bond lengths (Å)	0.007	0.010
Bond angles (°)	1.10	1.63
Ramachandran plot		
Most favored (%)	89.9	90.2
Allowed regions (%)	9.8	9.5
Outliers (%)	0.2	0.2
Clashscore (%)	8.8	8.1
Rotamer outliers (%)	0	0

* Values in parentheses are for the highest resolution shell (0.1Å)

R_{free} was calculated with 5% of the data

Figure Legends

Figure 1. Primary sequence and binding properties of knottins 2.5D and 2.5F. (A) The primary structure of EETI-II and of the engineered knottins 2.5D and 2.5F, where the 6-residue trypsin-binding loop in native knottin is replaced with an 11-residue loop (amino acids 2-12) containing the RGD motif. (B, C) Dose-response curves showing binding of Fc fusions of 2.5D (B), 2.5F (C) or a scrambled (scr) knottin (B, C) to native integrins expressed on U87MG, αVβ3 expressed on transfected K562 cells (αVβ3-K562), and native α5β1 on K562 cells. Points display the mean and standard deviation for triplicate determinations. (D) Equilibrium binding constant (K_d) values from the binding data shown in B, C, along with the standard error derived from the fitted curves. The K_d value derived from the curve fit of 2.5D-Fc binding to K562 in Fig. 1B is not reliable as determined by the P value for the parameter, and thus not reported in 1D. nd, not determined.

A.

```
EETI-II   GCP-----RILMRCKQDSDCLAGCVCGPNGFCG
2.5D      GCPQGRGDWAPTSCKQDSDCLAGCVCGPNGFCG
2.5F      GCPRPRGDNPPLTCKQDSDCLAGCVCGPNGFCG
          1   5    10    15     20  25  30
```

B. 2.5D

C. 2.5F

D.

Ligand	K562 (mean±se)	αVβ3-K562 (mean±se)	U87MG (mean±se)
2.5D	nd	1.4 (±0.4) nM	1.7 ±0.6 nM
2.5F	6.9 ±1.3 nM	9.2 ±1.3 nM	5.9 ±1.3 nM

Ligand	RMSD RGD loop (4-10)	RMSD Core domain (14-33)	RMSD Overall
2.5F	2.76 Å (±1.00)	0.35 Å (±0.18)	1.14 Å (±1.29)
2.5D	0.67 Å (±0.22)	0.22 Å (±0.12)	0.35 Å (±0.26)

Figure 2. NMR structures and analysis of 2.5D and 2.5F. (A, B) Solution NMR structures of knottins 2.5D (A) and 2.5F (B) represented as the 20 lowest energy conformers. Ribbon diagrams of the lowest energy conformer of 2.5D and 2.5F are shown to the right. The RGD loop is in cyan with the integrin RGD binding sequence in sticks. The core domain is blue with disulfide bonds in yellow sticks. The 3_{10} helix is represented in red cartoon. (C) RMSD of solution structure of 2.5D and 2.5F plotted versus residue number. The RGD loop (residue 4-10) and core domain (residue 14-33) are indicated. (D) Table summarizing the RMSD values per domain shown as mean±S.D.: core domain (residue 14-33) and RGD loop (residue 4-10).

Figure 3. Crystal structures of αVβ3 bound to knottins 2.5D or 2.5F. (A, B) Ribbon diagrams of αVβ3 (αV is in blue and β3 in green) bound to 2.5D (brown in A, C, E and G) or 2.5F (purple, in B, D, F and H). (C, D) Ribbon diagrams of the αVβ3 head bound to 2.5D (C) and 2.5F (D). The propeller is in blue and βA domain in green. Mn^{2+} ions at LIMBS (gray), MIDAS (cyan) and ADMIDAS (magenta) are shown as spheres (also in E-H). (E, F) Ribbon diagrams showing key electrostatic and hydrogen bond interactions and metal ion coordination in the structure of αVβ3/2.5D (E) and αVβ3/2.5F (F). 2.5F-R4 hydrogen bonds with β3-N313. Additionally, 2.5F contacts the ADMIDAS ion through a chloride (Cl) ion represented as a blue sphere. Water molecules are shown as small red spheres. (G, H) Solvent accessible surface view of the integrin/ligand interface showing residues in 2.5D (A10, P11, P28, N29, and F31) and in 2.5F (N9, P10, P11 and F31) forming van der Waals (≤4Å) contacts with the βA domain.

Figure 4. Structural comparisons of 2.5D and 2.5F in solution and integrin-bound. (A) Ribbon diagram of the superposed crystal structures of 2.5D (brown) and 2.5F (purple) in the integrin bound state. (B, F) Crystal structures of 2.5D (brown) and 2.5F (purple) superimposed on lowest energy model #1 of apo 2.5D (B, tale) and apo 2.5F (F, green). (C, G) Stick diagrams of the RGD loop in apo 2.5D (tale) and apo 2.5F (green) of lowest energy NMR #1 and of NMR#14. The 2.5D apo structure shows a β hairpin Type I turn in the RGD loop. Cβ-Cβ distances is stable among 2.5D conformers but is variable in case of 2.5F. (D) Superposition of GRGDW (residues 5-9) of 2.5D in the apo (tale) and bound states (brown). The red arrow shows conformational change of 2.5D-W9 from solvent-exposed state in the apo form to buried state in the crystal structure. (E, H) Crystal structures of the integrin-bound RGD loops of 2.5D (E) and 2.5F (H). (I) Stick diagram of the RGD loop bound to α5β1 crystal structure (PDB code 4WK4). The Cβ-Cβ distances between R6 and D8 in C, G, E, H, and R and D in I are shown.

Figure 5. MD simulations of 2.5D/2.5F binding to αVβ3/α5β1. Knottins were initially separated by 9Å. (A) Energy over time and (B) bar graph (mean± SE) showing absolute value of binding energies between 2.5D and 2.5F to αVβ3 and α5β1 averaged over 500 ns of simulation time. Difference between 2.5D/α5β1 and 2.5D/αVβ3, 2.5F/αVβ3, 2.5F/α5β1 is significant at p<2.2e^{-16}. (C, D) Selected residues in structures of MD simulation of 2.5F binding to αVβ3 at t=0.020 ns (C) and to α5β1 at t=1.250 ns (D). (E, F) Selected residues in structures of MD simulation of 2.5D binding to αVβ3 at t=0.960 ns (E), and to α5β1 at t=1.250 ns (F). Distances (dotted lines) are indicated. The head segment of the respective integrin is displayed.

Figure 6. Interaction of 2.5D and 2.5F with αVβ3 and α5β1. (A, B) MD simulations of 2.5D and 2.5F binding to integrins. (A) Energy over time and (B) absolute value of energies (histograms (mean $\pm$ SD) for 160 ns of the binding of 2.5D and 2.5F to wild type αVβ3 and αVβ3** double mutant (β3-M180A and β3-R214G) simulations. Differences among the groups are significant at $p < 2.2e^{-16}$ except for the difference between αVβ3/2.5F and αVβ3**/2.5F where the p value is $1.892e^{-06}$. (C-E) Histograms (mean $\pm$ SE, n=4 independent experiments) showing binding of integrin antibody or Alexa 647-labelled 2.5F or 2.5D (each at 65 nM) to transiently transfected HEK293T-αVβ3 and double mutant β3-M180A and β3-R214G αVβ3** in 1 mM Ca^{2+}/Mg^{2+} as determined by FACS analysis. (C) Binding of αVβ3 heterodimer specific antibody LM609 detected by APC-labeled goat anti-mouse Fc-specific antibody. (D) Binding of Alexa647-2.5F to αVβ3 and αVβ3**. (E) Binding of Alexa647-2.5D to αVβ3 and αVβ3**. Binding of the knottin to wild-type αVβ3 in D and E was set to 100%.

Star Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
LM609	Sigma Aldrich	MAB1976
APC-labeled goat anti-mouse Fc-specific antibody		
Bacterial and Virus Strains		
BacMagic baculovirus DNA (derived from AvNPV genome)	Novagen	72350-3
Bl21-DE3	New England Biolabs	C2527
Chemicals, Peptides, and Recombinant Proteins		
Peptide 2.5F	(Kimura et al., 2009b)	N/A
Peptide 2.5D	(Kimura et al., 2009b)	N/A
Restriction Enzymes	New England Biolabs	
Cell culture reagents	Invitrogen Corp (San Diego), Fisher Scientific (Hampton, NH)	
PBS	Boston BioProducts	Cat#BM-220
Hepes	Boston BioProducts	Cat#BBH-85
Alexa Fluor 647 dye	Invitrogen Corp (San Diego)	Cat#A20173
Alexa Fluor 488 dye	Invitrogen Corp (San Diego)	Cat#A10235
Dulbecco's Modified Eagle Medium (DMEM)	Invitrogen	Cat#11965126
Iscove's Modified Dulbecco's Medium (IMDM)	Invitrogen	Cat#12440061
Dulbecco's Phosphate Buffer Saline (DPBS)		Cat#A1285801
Fetal Bovine Serum (FBS)	Invitrogen	Cat#16000036
Cell Dissociation Buffer, Enzyme Free PBS	Invitrogen	Cat#13151013
Lipofectamine 2000	Invitrogen	Cat#11668027
Ni-NTA agarose	Qiagen	Cat#30210
Protein A Sepharose	Sigma	Cat#P3391
Tris	Sigma	Cat#252859
$MgCl_2$	Sigma	Cat#M8266
$CaCl_2$	Sigma	Cat#449709

MnCl$_2$	Sigma	Cat#244589
L-glutamine	Invitrogen	Cat#21051024
Penicillin/Streptavidin	Invitrogen	Cat#15140122
IPTG	Sigma	Cat#6758
Bovine Serum Albumin	Sigma	Cat#A2058
Critical Commercial Assays		
QuikChange Lightning	Agilent	Cat#210515
Deposited Data		
2.5F (apo) structure	This paper	PDB: 6MM4
2.5D (apo) structure	(Kryshtafovych et al., 2014)	PDB: 2M7T
2.5F-αVβ3 structure	This paper	PDB: 6MSL
2.5D-αVβ3 structure	This paper	PDB: 6MSU
Experimental Models: Cell Lines		
U87MG glioblastoma cells	ATCC (Manassas, VA)	Cat#HTB-14
K562 leukemia cells	Blystone et. al, 1994	SUNY Upstate Medical University
HEK293T	ATCC (Manassas, VA)	Cat#CRL-3216
Oligonucleotides		
β3-M180A-F gaaaacccctgctatgatgcgaagaccacctgcttgcc	This paper	N/A
β3-M180A-R ggcaagcaggtggtcttcgcatcatagcaggggtttc	This paper	N/A
β3-R214G-F gtgaagaagcagagtgtgtcacggaaccgagat	This paper	N/A
β3-R214G-R atctcggttccgtgacacactctgcttcttcac	This paper	N/A
Recombinant DNA		
pcDNA3-αV	(Gupta et al., 2007)	N/A
pcDNA3-β3		N/A
pBacPAK8-αV	(Mehta et al., 1998)	N/A
pBacPAK8-β3	(Mehta et al., 1998)	N/A
pET-32-2.5d	(Kryshtafovych et al., 2014)	N/A
pET-32-2.5F	This paper	N/A
pADD2 shuttle vector	(Moore et al., 2013)	N/A

Software and Algorithms		
SigmaPlot	Systat software	https://systatsoftware.com
FlowJo	TreeStar Inc.	https://www.flowjo.com
HKL2000	(Otwinowski and Minor, 1997)	http://www.hkl-xray.com
Phaser		https://www.phenix-online.org
Coot	(Emsley and Cowtan, 2004)	https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/
Phenix	(Adams et al., 2010)	https://www.phenix-online.org
PyMol	Schrödinger	https://pymol.org/2/
Haddock		http://www.bonvinlab.org/software/haddock2.2/
Charmm27	(Brooks et al., 2009)	https://www.charmm.org/charmm/?CFID=a0742b75-0ba2-4feb-94e3-a1fc7391a45a&CFTOKEN=0
NAMD	(Phillips et al., 2005)	http://www.ks.uiuc.edu/Research/namd/
VMD	(Humphrey et al., 1996)	http://www.ks.uiuc.edu/Research/vmd/
CYANA	(Guntert et al., 1997)	http://www.cyana.org
YASARA		http://www.yasara.org
TALOS+	(Shen et al., 2009)	https://spin.niddk.nih.gov/NMRPipe/talos/

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, M. Amin Arnaout (aarnaout1@mgh.harvard.edu).

METHOD DETAILS

Peptide Synthesis. Knottins 2.5D, 2.5F and the scrambled FN-RDG2 (where RGD motif is replaced with RDG) were prepared as previously described (Kimura et al., 2009b). Briefly, the linear 33-amino acid peptides starting with Gly1 (Figure 1A) were made by solid-phase peptide synthesis on a CS Bio (Menlo Park, CA) instrument using standard 9-fluorenylmethyloxycarbonyl chemistry. Knottin peptides were folded by promoting disulfide bond formation in oxidizing buffer at room temperature with gentle rocking overnight. Folded knottins were purified by reversed-phase HPLC, where they appeared as a sharp peak with a shorter retention time than unfolded or misfolded precursors. The molecular masses of folded knottins were determined by matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry (Stanford Protein and Nucleic Acid Facility). Folded 2.5F and 2.5D (2 mg/mL) were incubated with an amine-reactive succinimidyl ester derivative of Alexa Fluor 647 carboxylic acid in 0.1 M Hepes, pH 8.0, at a 5:1 dye/peptide molar ratio for 1 h at room temperature and then at 4 °C overnight. The free dye was removed by dialysis and buffer exchange into phosphate buffered saline (PBS).

Plasmids, mutagenesis, protein expression and purification. Human αVβ3 ectodomain was expressed in insect cells and purified as described (Mehta et al., 1998). The genetic sequence for knottins 2.5F, 2.5D or FN-RDG2 starting with Gly1 and ending with Gly33 (Figure 1A) was fused to the fragment crystallizable (Fc) region of mouse IgG2a in the pADD2 shuttle vector as described (Moore et al., 2013). The knottin Fc fusion proteins were expressed in human embryonic kidney (HEK293) cells following the manufacturer's protocols in the FreeStyle MAX 293 Expression System (Invitrogen). Secreted knottin-Fc fusion proteins were purified using Protein A Sepharose (Sigma) followed by size exclusion chromatography (Superdex 75 column; GE Life Sciences). Purified knottin-Fc fusion proteins were bivalent homo-dimers of the expected molecular weight of ~60 kDa (Moore et al., 2013).

Cell lines, cell culture and transfection. U87MG glioblastoma cells, K562 leukemia cells, and HEK293T embryonic kidney cells were obtained from American Type Culture Collection (Manassas, VA); integrin-transfected K562 cells (Blystone et al., 1994) were provided by Scott Blystone (SUNY Upstate Medical University). U87MG cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% Fetal Bovine Serum (FBS) and 1% penicillin-streptomycin (P/S). K562 cells were grown in liquid culture in IMEM Iscove's Modified Dulbecco's Medium (IMDM) supplemented with 10% FBS and 1% P/S. HEK293T cells were cultured in DMEM supplemented with 10% FCS, 2 mM l-glutamine, 1 mM sodium pyruvate, penicillin, and streptomycin, and were transiently co-transfected with pcDNA3 plasmids encoding full-length wild-type αVβ3, αVβ3** (αVβ3 with β3-M180A and β3-R214G) using Lipofectamine 2000 reagent (Invitrogen) according to the manufacturer's protocol.

Binding assays. 2.5D-Fc and 2.5F-Fc fusions were labeled with the succinimidyl ester derivative of Alexa Fluor 488 (Invitrogen) according to the manufacturer's protocol. Free dye was removed by dialysis and buffer exchange into PBS. U87MG cells were detached using Enzyme-Free Cell Dissociation Buffer (Gibco); K562 cells were grown in suspension. 4×10^4 cells then incubated with varying concentrations (0.01 – 500 nM) of Alexa Fluor 488 labeled 2.5D- or 2.5F-Fc fusion proteins in 25 mM Tris pH 7.4, 150 mM NaCl, 2 mM CaCl$_2$, 1 mM MgCl$_2$, 1 mM MnCl$_2$, and 0.1% bovine serum albumin (BSA) for 3 hours at 4 °C, to minimize internalization. Cells were

pelleted and washed twice with 800 µL of PBSA (phosphate buffered saline containing 0.1% bovine serum albumin) and the fluorescence of remaining surface-bound protein was measured using flow cytometry using a Guava EasyCyte 8HT instrument (EMD Millipore). Resulting data were evaluated using FlowJo software (TreeStar Inc.) and equilibrium dissociation constants (Kd) were determined in SigmaPlot (Systat Software, San Jose, CA) using a least-square fit to a logistic curve. Kd values are determined from the fitted parameter, and the standard error, t and p values for the parameter calculated using the reduced χ^2 method. The validity of the Kd value was assessed with a p value cutoff of 0.05.

Wild type αVβ3 and mutated αVβ3** transiently transfected HEK293T cells were gently trypsinized and washed in DPBS buffer. Cells were re-suspended in complete culture medium, incubated for one hour at 37˚C and subsequently washed in 1 mM Ca^{2+}/Mg^{2+}, 0.1% bovine serum albumin-supplemented Hepes buffered saline pH 7.4 (binding buffer). 5×10^6 cells were incubated with Alexa647-labeled 2.5F or 2.5D (50 nM) in binding buffer for 30 min, at 25 °C then washed, re-suspended, fixed in 1% paraformaldehyde and analyzed by flow cytometry in a LSRII flow cytometer (BD). Anti-$\alpha_V\beta_3$ antibody LM609 (20 µg/ml) was used to normalize $\alpha_V\beta_3$ cell surface expression in a separate set of tubes. Transfected HEK293T cells were stained with LM609 for 30 min at 4˚C. After washing the excess antibody, APC-labeled goat anti-mouse Fc-specific antibody (10 µg/ml) was added for 30 min at 4°C, and the stained cells were washed, fixed and expression analyzed by flow cytometry as described above. Alexa-647 labeled knottin 2.5F or 2.5D binding to αVβ3 and αVβ3** was measured in mean fluorescence intensity (MFI) units, normalized according to LM609 binding and expressed as percentage of 2.5F or 2.5D binding to αVβ3.

NMR A codon-optimized DNA sequence was prepared by assembly PCR and cloned into the pET-32 vector to express a protein product in *E. coli* containing a thioredoxin and His-tag fusion protein separated by a TEV protease site. Uniform [15]N- and [13]C-labeling was achieved by IPTG-induced expression in BL21-DE3 cells in M9 minimal media containing [15]NH_4Cl and [13]C-glucose. Cell lysis was followed by initial purification by His-tag capture with Ni-NTA. The thioredoxin fusion protein portion was removed with TEV protease to provide the exact 33-residue peptide sequence. Disulfide bond formation to fold the peptides was performed using the previously reported redox buffer (Kimura et al., 2009b). Final purification by RP-HPLC and characterization by ESI mass spectrometry confirmed folded engineered EETI-II peptides. NMR samples were prepared using sodium phosphate buffer, pH 6 containing 10% D_2O. Standard multidimensional NMR datasets were acquired for backbone and side chain resonance assignments, along with [13]C (aliphatic and aromatic) and [15]N 3D-NOESY datasets for NOE-derived distance restraints. Dihedral angle restraints were derived from backbone assignments using TALOS+. 3D structure calculations were performed using the CYANA automated NOE assignment and simulated annealing algorithms. Initial structures calculated using standard automated NOE assignments led to convergence of 20 lowest energy structures, which were consistent with the expected knottin disulfide pattern. Further improved 3D structures were calculated by including disulfide bond restraints and hydrogen bond restraints determined by cross-hydrogen bond scalar couplings identified from long-range HNCO datasets. Final structures were refined by restrained molecular dynamics in explicit solvent using the YASARA package.

Crystallography, structure determination and refinement. The αVβ3 ectodomain was crystallized at 4 °C by vapor diffusion using the hanging drop method as previously described (Xiong et al., 2009; Xiong et al., 2001; Xiong et al., 2002). Knottins 2.5F or 2.D (5 mM) was soaked into $\alpha_v\beta_3$ crystals in the crystallization well solution containing 1 mM Mn^{2+} for 2–3 weeks. Crystals were harvested in 12% PEG 3500 (polyethylene glycol, molecular weight 3500) in 100 mM sodium acetate, pH 4.5, 800 mM NaCl plus 2 mM Mn^{2+} and 2.5F or 2.5D (at 5 mM), cryoprotected by the addition of glycerol in 2% increments up to a 24% final concentration and then flash frozen in liquid nitrogen. Diffraction data from cryo-cooled crystals were collected on the ID19 beamline fitted with a CCD detector at the APS Facility (Chicago, IL). Data were indexed, integrated and scaled with the HKL2000 (Otwinowski and Minor, 1997) program. Phases were determined by molecular replacement using PHASER (McCoy et al., 2007), with the structures $\alpha_V\beta_3$ ectodomain (PDB ID 4MMX). Composite simulated annealing omit maps were then generated using the program Phenix Composite Omit Map package by turning on the simulated Cartesian annealing option with a temperature of 5,000K. The knottin structure was traced by the extra density using PDB 2IT7 and introducing the engineered mutations using Coot (Emsley and Cowtan, 2004). The resulting models were refined with the 1.10.1 version of Phenix (Adams et al., 2010) using simulated annealing, TLS, positional and individual temperature-factor refinement and default restrains. Several cycles of refinement and model building using Coot were applied to refine the structures of αVβ3/2.5D, αVβ3/2.5F (Table 2), with automatic optimization of X-ray and stereochemistry and additional Ramachandran restrains in the last cycles. A-weighted 2*fo-fc* electron density map was generated from the final models and structure factors using Phenix. All structural illustrations were prepared with the PyMol software (Schrödinger).

Docking and Initial Configurations. The NMR structures of 2.5F (PDB: 6MM4) and 2.5D (PDB ID: 2M7T) were docked onto the crystal structures of the integrin αVβ3 (PDB ID: 4MMZ) and α5β1 (PDB ID: 4WJK) headpieces using the expert interface of the HADDOCK webserver. Four HADDOCK docking runs, between integrin α5β1 and 2.5D, integrin α5β1 and 2.5F, integrin αVβ3 and 2.5D, and integrin αVβ3 and 2.5F were performed. As docking inputs, the RGD sequence was specified as the active site of 2.5D and 2.5F. Residues α5-Glu[221], α5-Asp[227], and Mg^{2+} ion at the MIDAS site were specified as the active site of integrin α5β1, while residues αV-Asp[150], αV-D[218], and the MIDAS Mn^{2+} ion, were specified as the active site of integrin αVβ3. To preserve the ion coordination of the MIDAS, ADMIDAS, and LIMBS ions during the docking run, unambiguous distance restraints between the coordinating groups and the MIDAS, ADMIDAS, and LIMBS ions were fed into HADDOCK. Furthermore, integrins αVβ3 and α5β1 were specified as non-flexible in HADDOCK to prevent integrin backbone movement during docking. The RGD sequence of the knottins were specified as semi-flexible, while the non-RGD sequence was specified as non-flexible. Upon completion of the docking runs, the top solution from each generated cluster was analyzed. The best solution was then selected from these solutions based on maximal engagement of the specified active site residues.

Molecular Dynamics Simulations To simulate the interaction of 2.5D and 2.5F with integrins αVβ3 and α5β1, we performed Molecular Dynamics simulations using the top solution from each of the four HADDOCK structures. To setup the separated simulations, the axis between the center of mass of the integrin βA domain and the knottin variants was determined in the four HADDOCK complexes. Knottin was then separated from the integrin by 9Å along this axis,

allowing water molecules to populate the space in between the knottin and the integrin upon solvation of the structure. We also performed equilibration simulations on the available crystal structures of αVβ3 in complex with 2.5D and 2.5F to be used as references. All structures were solvated and then ionized at a combined KCl concentration of 0.15 M. Structures were subsequently minimized for 100,000 steps and equilibrated for 0.5 ns using the NAMD molecular dynamics package (Phillips et al., 2005) and CHARMM27 force field (Brooks et al., 2009). Upon minimization, each of the four generated complexes ran for 500 ns. All simulations ran at an initial temperature of 310 K using the Nose-Hoover thermostat, and pressure was maintained at 1 atm using the Langevin piston. All equilibration and production run simulations were performed using a time step of 2 fs. Electrostatics of the system were determined using the Particle mesh Ewald (PME) method. van der Waals (VDW) interactions were modeled using a switching function to smoothly reduce the VDW force to zero at the cutoff distance of 1.2 nm. Simulations were then analyzed using Visual Molecular Dynamics (VMD) (Humphrey et al., 1996).

Accession numbers. Coordinates for the NMR structures of knottins 2.5D and 2.5F and the crystal structures of αVβ3/2.5D and αVβ3/2.5F the have been deposited in the PDB with ID codes 2M7T, 6MM4, 6MSL and 6MSU, respectively.

QUANTITATION AND STATISTICAL ANALYSIS.
Quantitation and statistical analysis are reported in the Method Details section.

DATA AND SOFTWARE AVAILABILITY
All software and libraries used are reported in the Method Details section, together with the Key Resources Table.