EPIP: MHC-I epitope prediction integrating mass spectrometry derived motifs and tissue-specific expression profiles

Weipeng Hu[*1,2,3], Si Qiu[*2,3], Youping Li[2,3,4], Xinxin Lin[2,3], Le Zhang[2,3,4], Haitao Xiang[2], Xing Han[2,3], Lei Chen[2,3], Sha Li[2,3], Wenhui Li[2,3], Zhe Ren[2], GuiXue Hou[2], Zhilong Lin[2], Jianliang Lu[2], Geng Liu[2,3], Bo Li@[2,3,5], Leo J Lee@[2,3,6]

1. BGI-Shenzhen, Shenzhen 518083, China
2. BGI-GenoImmune, Wuhan 4300794, China
3. BGI Education Center, University of Chinese Academy of Sciences, Shenzhen 518083, China
4. Guangdong Enterprise Key Laboratory of Human Disease Genomics, Shenzhen 518083, China
5. Department of Electrical and Computer Engineering, Donnelly Centre for Cellular and Biomolecular Research, University of Toronto, Toronto, Ontario M5S 3G4, Canada

Correspondence: ljlee@psi.toronto.edu, libo@genomics.cn

Abstract

Background: Accurate prediction of epitopes presented by human leukocyte antigen (HLA) is

crucial for personalized cancer immunotherapies targeting T cell epitopes. Mass spectrometry

(MS) profiling of eluted HLA ligands, which provides high-throughput measurements of HLA

associated peptides *in vivo*, can be used to faithfully model the presentation of epitopes on the

cell surface. In addition, gene expression profiles measured by RNA-seq data in a specific

cell/tissue type can significantly improve the performance of epitope presentation prediction.

However, although large amount of high-quality MS data of HLA-bound peptides is being

generated in recent years, few provide matching RNA-seq data, which makes incorporating gene

expression into epitope prediction difficult.

Methods: We collected publicly available HLA peptidome and matching RNA-seq data of 34

cell lines derived from various sources. We built position score specific matrixes (PSSMs) for 21

HLA-I alleles based on these MS data, then used logistic regression (LR) to model the

relationship among PSSM score, gene expression and peptide length to predict whether a peptide

could be presented in each of the cell line. We further built a universal LR model, termed

Epitope Presentation Integrated Prediction (EPIP), based on more than 180,000 unique HLA

ligands collected from public sources and ~3,000 HLA ligands generated by ourselves, to predict

epitope presentation for 66 common HLA-I alleles.

Results: When evaluating EPIP on large, independent HLA eluted ligand datasets, it performed

substantially better than other popular methods, including MixMHCpred (v2.0), NetMHCpan

(v4.0), and MHCflurry (v1.2.2), with an average 0.1% positive predictive value (PPV) of 52.01%,

compared to 37.24%, 36.96%, 24.90% and 23.76% achieved by MixMHCpred, NetMHCpan-4.0

(EL), NetMHCpan-4.0 (BA) and MHCflurry, respectively. It is also comparable to EDGE, a

recent deep learning-based model that is not publicly available, on predicting epitope

presentation and selecting immunogenic cancer neoantigens. However, the simplicity and

flexibility of EPIP makes it easier to be applied in diverse situations, and we demonstrated this

by generating MS data for the HCC4006 cell line and adding the support of HLA-A*33:03 to

EPIP. EPIP is publicly available at < http://epip.genomics.cn/>.

Conclusions: we have developed an easy to use, publicly available epitope prediction tool, EPIP,

that incorporates information from both MS and RNA-seq data, and demonstrated its superior

performance over existing public methods.

Keywords: T cell epitope, MHC-I, HLA peptidome, Eluted ligand, Mass spectrometry, RNA-

seq, Neoantigen, Cancer immunotherapy

Background

T cell epitopes, which are short peptides presented by major histocompatibility complex (MHC)

molecules on the cell surface and recognized by T-cell receptors (TCRs), lie at the heart of the

human immune system that can remove infected and malignant cells. With the rapid progress of

cancer immunotherapies, the mechanistic understanding, computational prediction and clinical

manipulation of T cell epitopes have gathered renewed and widespread interests [1–4]. It is well

known that peptide binding to MHC molecules is the most selective step in the antigen

presentation pathway [5], and intensive computational efforts have been exerted to predict this

binding process based on peptide sequences (allele-specific models) or peptide and the

corresponding human leukocyte antigen (HLA) sequences (pan-specific models). Traditionally,

these models are trained on data accumulated from various types of *in vitro* binding assays,

which can provide quantitative binding affinity measurements between pre-selected peptides and

HLA-I or HLA-II molecules. However, the requirement of synthesizing peptides beforehand

limited the ability of these assays to perform unbiased, high-throughput screening of the vast

peptide space, and it also did not account for the antigen-loading process *in vivo* [6]. The recent

advance of mass spectrometry (MS) profiling of HLA ligands, mainly for HLA-I alleles at the

moment, overcomes most of these limitations by providing *in vivo* measurements of peptides

presented by MHC molecules on the cell surface in a high-throughput manner, although the

resulting measurements are only qualitative (binary).

The value of MS data as contributing to the ultimate prediction of peptide immunogenicity has

been increasingly recognized [6,7], and as a result, a number of recently developed peptide-MHC

binding prediction methods incorporated some form of MS data, in addition to the traditional

binding assay data [5,8]. For example, NetMHCpan-4.0 trained an ensemble of two hidden layer

neural networks on both affinity and MS data, and was able to make predictions on both types of

outputs. MHCflurry adopted a deep learning framework containing both locally connected and

fully connected layers and was mainly trained on affinity data, but used MS data at the model

selection stage, while MS data may play an even bigger role in its future releases. Other methods

chose to rely on MS data alone [7,9,10], such as MixMHCpred [9,10], which built allele-specific

position score specific matrixes (PSSMs) after deconvolving mixed-allele MS data, and the very

recent EDGE method [7], which used a comprehensive, deep learning model that pooled together

large sets of mono-allele and mixed-allele MS data as well as matching RNA-seq data. With the

increasing amount of MS data becoming available, it is our view that it might be better to build

separate models for binding affinity (based on binding assay data) and epitope presentation

(based on MS data) since the underlying data generation processes are different. These two

related pieces of information could be further combined at a later stage to contribute to the

prediction of immunogenicity. The presentation of a peptide by MHC not only depends on

information contained within the protein sequences but is also modulated by other factors such as

protein abundance, localization and turnover [11,12]. Among these, the corresponding gene

expression levels can be handily profiled by RNA-seq and have been shown to significantly

improve prediction accuracy, first by the MS IntrinsicEC method on mono-allele cell lines [12],

then more comprehensively by EDGE [7]. Unfortunately, both methods are not publicly

available.

We set out to develop an effective, flexible and publicly available HLA-I epitope prediction

method that takes advantage of large sets of MS and RNA-seq data. We first collected public MS

and matching RNA-seq datasets for 21 HLA-I alleles, which frequently appear in the European

or Asian population. By carefully examining the contributions of PSSM scores, gene expression

levels and peptide length features to epitope presentation, we were able to effectively model

these variables with a universal logistic regression (LR) model. We then trained such a LR model,

termed Epitope Presentation Integrated Prediction (EPIP), to support 66 common HLA-I alleles.

In the remainder of this article, we first describe our design rationale in detail, followed by

comprehensive evaluation of EPIP on large, independent MS and immunogenic datasets as well

as a user case showing its extendibility, before concluding with discussions on future

improvements and research directions. We believe EPIP is a timely contribution to the research

community of cancer immunotherapies.

Methods

Collection of MS and RNA-seq data

We collected matching MS and RNA-seq data from 16 mono-allele HLA-A and HLA-B cell

lines [12] and 18 mixed-allele cell lines [13] to build the EPIP model. We also collected MS data

of 15 mono-allele HLA-C cell lines from Di Marco et al [14], deconvolved MS data of 26 more

HLA alleles from SysteMHC [15] and a small portion of mixed-allele MS data released with

EDGE [7] to expand the allele coverage of EPIP. To perform independent evaluation of EPIP,

we further downloaded MS data from Trolle et al [16] and Bassani-Sternberg et al [11], in

addition to test data reserved from Pearson et al [13] and Bulik-Sullivan et al [7]. Detailed

information and usage of all collected MS data are provided in Table S1.

MS profiling of HCC4006 HLA-I peptidome

HLA-I peptidome samples for HCC4006 were prepared according to Bassani-Sternberg et al [11].

They were then analyzed by LC-MS/MS to obtain peptide MS spectra, and full details of the

experimental procedure are provided in the Supplementary Material. We employed the MS-GF+

search engine version 2018.07.17 [17] to search the peak lists against the UniProt databases

(161,521 entries for human as of December 2017) and a file containing 245 frequently observed

contaminants such as human keratins, bovine serum proteins, and proteases. N-terminal

acetylation (42.01 Da) and methionine oxidation (15.99 Da) were set as variable modifications.

The enzyme specificity was set as unspecific. The initial allowed mass deviation of the precursor

ion deviation was set to 10 ppm. Possible peptide lengths were restricted to 8 to 15. Possible

precursor charges were restricted to 2 to 5. Range of allowed isotope peak errors was restricted

to -1 to 2. Percolator version 3.02.0 was applied as a post-processing step for result filtering [18].

The q value cutoff was set to 0.01. From the "pout.tab" output file produced by percolator, hits to

the contaminants were eliminated.

Data and features of EPIP models

From the matching MS and RNA-seq data collected from public sources [12,13], we first built

separate LR models, EPIP_s, to predict epitope presentation for 21 alleles and tissues. This

included all 16 mono-allele data profiled in Abelin et al [12], as well as five more alleles (HLA-

A*11:01, HLA-A*32:01, HLA-B*15:01, HLA-B*40:01, HLA-B*07:02) selected from Pearson

et al [13], which frequently appear in the Chinese population [19]. All RNA-seq datasets were

downloaded and re-processed with the same pipeline to ensure consistency. Specifically, raw

fastq files were cleaned by fastp (version 0.18.0) [20] and aligned to human genome assembly

GRCh38 and its associated transcriptome (Ensembl Release 92) with STAR (version 2.5.3a) [21];

gene expression (transcripts per million or TPM) was quantified by RSEM (version 1.3.0) [22]

based on the alignment.

EPIP_s uses three features to model epitope presentation: peptide motifs as modelled by PSSM,

peptide lengths and the corresponding gene expression levels. For mono-allele MS data , we built

PSSMs directly, as described in Liu et al [23]. For mixed-allele MS data, we first performed

motif deconvolution by GibbsCluster [24] before building PSSMs. Peptides from deconvolved

motifs that matched the HLA motifs of interest generated from SysteMHC data were retrieved.

For each allele, length-specific PSSMs were built for those that have at least 100 peptides, up to

the most abundant three lengths (9-11-mers or 8-10-mers, depending on the allele, details in

Table S1), and this allowed us to make use of 74%-96% of MS profiled peptides across different

alleles. To fully utilize the training data and avoid overfitting, we used the stacking method [25]

to build PSSM (see supplementary materials, Fig S1). For each HLA allele, we also obtained an

empirical probability mass function (pmf) to depict the length distribution of 8-15-mers based on

the allele-specific MS data, and the peptide length feature is the corresponding probability in the

empirical pmf of peptide length distribution. Expression values of a peptide was calculated by

summing up the expression levels of all transcripts containing the peptide (in TPM) before

transformed by log2(TPM+1). For the mono-allele cell lines in Abelin et al [12], transcript

expression levels were taken as the average of the four cell lines with RNA-seq data available;

similarly, for the mixed-allele cell lines in Pearson et al [13] , transcript expression levels were

averaged over the 10 cell lines with RNA-seq data available. All three features were further

normalized to be zero mean, unit variance Gaussian before using as inputs to a LR model with

weak L2 regularization implemented in scikit-learn 0.19.0 [26].

MS profiled peptides can only provide a positive training set for our model. To generate the

negative set, we randomly sampled the human peptidome to obtain peptides of the matching

length and remove those that overlap with the positive set. Adopting the same strategy as in

NetMHCpan 4.0, we used more negative peptides than positive ones to train EPIP_s. For

example, if the positive set contains 9-11-mers with the most abundant 9-mers having a size of n,

we then generate a negative set with $10n$ peptides for each length ($30n$ in total).

To build the full EPIP model for the above 21 alleles from different cell lines, we pooled all the

PSSM scores, peptide length distributions and corresponding gene expression levels together to

train a single LR model with the same training data as EPIP_s. Five-fold cross validation was

used to evaluate EPIP during training and to compare EPIP and EPIP_s, while the full training

set was used to build the final EPIP model and it was evaluated on independent test data. To

expand the number of HLA alleles that EPIP can support, mono-allele HLA-C MS data from Di

Marco et al [14], deconvolved MS data from SysteMHC [15] and some mixed-allele MS data

released by EDGE [7] were used to learn PSSMs for these additional HLA alleles. For the mono-

allele HLA-C MS data from Di Marco et al, we deconvolved the unfiltered list of peptides to

remove the B35 and C04 motifs through GibbsCluster [24] before using them to train PSSMs.

We further generated additional MS-data from the HCC4006 cell line so that EPIP can support

HLA-A*33:03, a high frequency allele with 8.03% prevalence in the Chinese population [19].

Immunogenicity evaluation

The immunogenic dataset used in this paper is the same as in Bulik-Sullivan et al. [7], which

contains 2,023 assayed mutations from 17 patients with annotated HLA information from four

previous studies [27–30]. In three of the four studies [27,28,30] , mutations were tested using

25mer tandem minigene (TMG) assays. For each mutation in these datasets, we used EPIP,

MixMHCpred 2.0, NetMHCpan4.0-EL, NetMHCpan4.0-BA, MHCflurry 1.2.2 to make

predictions with respect to the HLA alleles of each patient for overlapping 8-11mers that harbor

the mutation, and the score of the mutation is taken as the best score among these peptides (EPIP:

highest presentation probability; MixMHCpred 2.0: highest predicted score; NetMHCpan4.0-EL

and NetMHCpn4.0-BA: lowest rank score; MHCflurry 1.2.2: minimum binding affinity). We

then ranked the mutations according to the best scores from each software. For the fourth study

[29] , we ranked mutations by taking the best scores from each software across all mutation-

spanning peptides tested in the tetramer assays. To compare EPIP with EDGE on predicting

immunogenic peptides, we excluded the peptides whose immunogenicity was undetermined as

described in Bulik-Sullivan et al. We retrieved a total of 31 immunogenic epitopes from Tran et

al., 216 Gros et al. and Stronen et al., which were not provided by Bulik-Sullivan et al and

slightly different from the number 29 reported in Bulik-Sullivan et al. Since the epitope

prediction scores by EDGE was not available, we directly compared its immunogenic prediction

results with EPIP.

Results

Construction and analysis of EPIP models

After collecting matching MS and RNA-seq data from public sources, we first developed

separate LR models for each of the 21 tissues and alleles (termed EPIP_s) to predict epitope

presentation by combining motif, peptide length and gene expression features. Full details of the

datasets and features are described in Methods, and the overall process is sketched in Figure 1A.

According to previous studies[12,31–33], about one in a thousand peptides generated from the

human proteome can be presented by MHC. Therefore, we further generated 999-fold decoy

peptides that are not in the training set to evaluate our models. Similar to Abelin et al [12], the

main evaluation criterion used throughout the paper is 0.1% positive predictive value (PPV),

which measures the proportion of truly presented peptides in the top-scoring 0.1% ones found by

our model. The 0.1% PPV is both a more realistic measure for immunopeptidome prediction and

a more stringent one than many other popular metrics, such as ROC-AUC and F1 scores, for this

highly biased classification task.

We used PSSM, a simple model that can learn patterns in biological sequences, to model peptide

motifs since it has previously been widely used in modelling MHC binding peptides [34–36], is

easy to visualize, and seems to have adequate representational power. It is also a data efficient

model due to its simplicity. To show this, we used subsampling to build PSSM models for

different lengths of peptides to predict epitope presentation and tested their performances (0.1%

PPV) on various alleles. As shown in Figure 1B, the performances of PSSM models reached

saturation when having 100-200 peptides in the training set. Therefore, we selected 100 as a

cutoff to build length-specific PSSM models for up to three most abundant lengths in each allele

(9-11-mers or 8-10-mers, depending on the allele, details in Table S1), and the model would also

be restricted to make predictions on these peptide lengths.

Although it is reasonable to assume that the relationship among motif, peptide length and gene

expression is a general one across different tissues (cell lines) and alleles so that data from these

different sources could be pooled together to train a general LR model (as outlined in Figure 1C),

building EPIP_s first allowed us to rigorously analyze and quantify this. We first examined the

distribution of optimized model parameters (weights and biases) among these 21 EPIP_s models.

As shown in Figure 2A, the corresponding model parameters distributed quite tightly. Since the

features have been normalized prior to model training, the relative magnitudes of model

parameters also displayed the contribution of each feature, where PSSM played the most

important role, followed by expression levels and peptide lengths (the average weights of PSSM,

expression level and peptide lengths are 4.30, 1.37, 0.37, respectively). Furthermore, when

displaying these 21 sets of weights and biases as hive plots (Figure 2B), they share very similar

inter-relationships and seem to be mostly differ by a scaling factor, which is likely caused by

somewhat different influences of L2 regularization in each model. This provided us with a strong

motivation to build a universal LR model, EPIP, to account for the contribution of PSSM,

peptide length and gene expression across these 21 tissues and alleles, to compare against the

EPIP_s models, which will be described next.

EPIP and EPIP_s achieved similar performance on epitope presentation prediction

Based on five-fold cross validation, we performed comprehensive evaluations between EPIP and

EPIP_s in terms of 0.1% PPV. As shown in Figure 3A, when evaluating across different alleles

originating from the same cell line, the performances of EPIP and EPIP_s are quite similar. The

mean 0.1% PPV of EPIP is slightly better (0.5741 vs 0.5692), likely due to the more training

data available to EPIP, but such a difference is not statistically significant ($P = 0.1385$, paired t-

test), and both substantially outperformed predictions based on PSSM scores alone (the mean

0.1% PPV of PSSM is 0.4421).We also performed similar analyses for the same HLA alleles

across different cell lines with diverse tissue origins and thus diverse gene expression profiles

(Figure 3B). When comparing 0.1% PPV of EPIP and EPIP_s, EPIP is not worse than EPIP_s

overall, and even notably better for some cell lines (Figure 3C-F), also likely due to the larger

training set of EPIP. Based on these comparisons, we concluded that developing a universal LR

model, EPIP, to account for the relationship among these three variables is a valid approach. This

seemly simple strategy has substantial practical values, since the EPIP model trained on the

selected 21 HLA alleles and cell lines from two different tissue origins can now be

straightforwardly extended to previously unseen HLA alleles and tissues without adjusting the

model parameters of LR, or requiring extra RNA-seq data for training. We did so for 44 more

HLA alleles by incorporating public mono-allele [14] and deconvolved mixed allele [7,15] MS

data. We also generated our own MS data to demonstrate as a user case, after evaluating EPIP

against other methods.

EPIP significantly outperforms other existing methods

We evaluated EPIP against MixMHCpred (v2.0), NetMHCpan-4.0 (EL), NetMHCpan-4.0 (BA),

and MHCflurry (v1.2.2) on independent validation MS datasets collected from public sources,

which contained 29 MS datasets and was not used in the training of EPIP. EPIP significantly

outperformed other methods with mean 0.1% PPV of 52.01%, compared to 37.24%, 36.96%,

24.90% and 23.76% achieved by MixMHCpred, NetMHCpan(EL), NetMHCpan (BA) and

MHCflurry, with highly significant p-values (P = 2.819e-11, 3.960e-11, 1.738e-14, 6.826e-14,

respectively). Since other methods in the comparison do not make use of gene expression values,

we further combined different gene expression value cutoffs with their predictions. The

advantage of EPIP persists and the differences are still highly significant, as shown in Figure 4A.

Full comparisons for each validation dataset are provided in Figure S2. The recently developed

EDGE method has also been shown to excel at epitope presentation prediction comparing to

existing methods, but we were not able to do a fair comparison since it is not publicly available.

Nevertheless, we will discuss this further with some qualitative comparisons in the Discussion

section to show that the performances of EPIP and EDGE are comparable. Overall, the results on

independent MS validation sets showed the clear advantage of EPIP in predicting epitope

presentation.

EPIP is valuable to immunogenicity prediction

For cancer immunotherapies, as well as other clinical applications, the ultimate task is usually to

predict the immunogenicity of an epitope. While this remains an elusive goal in general [37], it

has recently been shown that accurate prediction of epitope presentation can significantly

improve immunogenicity predictions [7]. Therefore, we downloaded the same dataset as used in

Bulik-Sullivan et al [7], which was collected from four previous publications [27–30] that

consisted of 26 SNVs and 31 neoantigens with pre-existing T-cell responses among 2,023

assayed single nucleotide variants (SNVs) from 17 patients, to evaluate our method against

others. We first asked if the increased accuracy of EPIP over other publicly available methods on

epitope presentation could translate into better immunogenicity prediction. To quantify this, we

plotted the precision recall curves (PRCs) by ranking all SNVs scored by different methods,

where the score of a SNV is the maximum score of all possible epitopes generated by the SNV.

EPIP's area under PRC (AUPRC) was more than twice that of other methods, and kept the

advantages when we applied expression value threshold of 1 or 2 TPM to other methods'

predictions (Figure 4B and Figure S3). We also compared EPIP with EDGE at the level of

ranking mutations and minimal epitopes (the recognized 8-11-mers overlapping the mutation) as

originally done in Bulik-Sullivan et al [7]. EPIP is very comparably to EDGE on these two

evaluations, especially for the top 10 and top 20 ranked mutations and epitopes. When

prioritizing the mutations, the number of immunogenic SNVs ranked in the top 20, 10, and 5

were 18 (69.23%), 16 (61.54%) and 11 (42.31%) for EPIP, and 19 (73.08%), 18 (69.23%) and 12

(46.15%) for EDGE (Figure 4C). When prioritizing the minimal epitopes, EPIP ranked 15

(48.39%), 12 (38.71%) and 5 (16.13%) CD8+ recognized epitopes in the top 20, 10 and 5.

Therefore, despite being a much simpler model trained with less data comparing to EDGE, EPIP

is also quite valuable for immunogenicity prediction in neoantigen-based cancer

immunotherapies and is a significant improvement over other publicly available methods.

EPIP can be extended to support more alleles by incorporating new MS data

Due to its simplicity and modular structure, EPIP can be straightforwardly extended to support

more HLA alleles by incorporating new MS data. An allele that we are particularly interested in

is HLA-A*33:03, which appears often in the Chinese population but with very limited MS or

binding affinity data prior to this publication (more details in the supplementary material). We

first performed MS profiling of eluted HLA ligands on the HCC4006 lung cancer cell line, which

is homozygous on all three HLA alleles (HLA-A*33:03, HLA-B*44:03 and HLA-C*07:06,

respectively). Upon deconvolution of MS data by GibbsCluster and removing the trash cluster,

we were only able to obtain two clusters, likely due to the low expression of HLA-C alleles. We

were able to assign one of the clusters to HLA-B*44:03 by comparing to the known motif of

HLA-B*44:03 obtained from mono-allelic HLA-B*44:03 cell line [12]. We believe the other

cluster should belong to HLA-A*33:03, since the motif is quite similar to the A33:03 motif

predicted by NetMHCpan-4.0 in its pan-specific mode (although binding affinity has not been

directly profiled for this allele) and ~80% of the peptides in this cluster has binding affinity <

500nm to HLA-A*33:03 as predicted by NetMHCpan-4.0. To further check on this, we also

performed t-SNE [38] analysis for the peptides in this cluster, together with peptides of HLA-

*A02:01 and HLA-A*11:01 in EPIP's training set. As shown in Figure S4, peptides from this

cluster is much closer to those from HLA-A*11:01 than HLA-A*02:01, consistent with the fact

that HLA-A*33:03 and HLA-A*11:01 belong to the same HLA supertype while HLA-A*02:01

belongs to a different supertype. Based on these evidences, we confidently assigned this cluster

to HLA-A*33:03. By inputting these HLA-A*33:03-specific peptides to EPIP, where a dedicated

module has been built to accept new allele-specific MS data, it can learn the corresponding

PSSM and length distribution and add HLA-A*33:03 to its supported alleles. The overall work

flow is illustrated in Figure 5.

Discussions

Besides making use of recently available, large-scale MS immunopeptidome data, our method

clearly demonstrated the advantage of taking gene expression levels into account in a principled

way when modelling epitope presentation, but the use of gene expression levels could present its

own challenges. Even when discovering neoantigens for cancer patients where RNA-seq is

routinely preformed, this information could be missing or inaccurate. In this case, since it is well-

known that gene expression levels are largely preserved in the same tissue [39], one could make

use of the gene expression levels quantified in a different sample from a similar tissue origin. For

example, we showed that when replacing the expression levels of train_sample55 with HELA,

which are both originated from ovary tumors, the performance of EPIP on HLA-A*2402 only

dropped a little and still significantly outperformed other public methods (Figure S6). In other

applications of epitope prediction, such as those originating from pathogens, the definition of

gene expression levels could be less clear, but there have been efforts to do this by more

sophisticated RNA-seq analysis methods. Although the effect of gene expression on pathogen

epitopes is less clear, several studies had shown the importance of gene expression and sufficient

protein supplement in virus epitopes presentation in infected human cells. While the 2B2-CTL

failed to cytotoxic HPV-16-positive cervical cancer cell lines, transfer HPV E6 protein cDNA

into those cancer cell lines could enhance endogenous E6 mRNA expression level and overcome

the failure [40,41]. Since virus employ the human cell machinery to produce their protein, the

virus epitopes may share the same processing disciplines with human epitopes. As more

sophisticated RNA-seq analysis methods now enable researchers to analyses the transcriptome of

virus infected cells and calculate the TPM/RPKM of specific virus proteins, EPIP could be used

to predict virus epitopes in these cells. [42]

The incorporation of gene expression levels in EPIP is similar in principle to what has been done

by EDGE [7]. Although EDGE is not publicly available, nor is it straightforward to reproduce

their work, some comparisons with EDGE on epitope presentation prediction would be

informative. To achieve this, we used the same evaluation criteria as in EDGE to compare

against other affinity-based methods, i.e., to further reduce the positive to negative ratio, or

prevalence, to 1:2,500 and 1:10,000 and checked the PPVs of EPIP at 40% recall. Although it is

difficult to justify the optimal threshold of prevalence for neoantigen prediction, further reducing

this threshold does have the side effect of enlarging the performance gap between EPIP and other

methods. As shown in Figure S5, EPIP reached a more than 10-fold improvement over

MHCflurry with 1:10,000 prevalence, similar to what has been achieved by EDGE. Combined

with the fact and EPIP and EDGE performed similarly on the same immunogenicity test data, we

concluded that the performances of EPIP and EDGE should be quite comparable. Since EPIP is a

much simpler model comparing to EDGE, this also implies that the performance gain of EDGE

is to a larger degree contributed by its large-scale training data than the more sophisticated deep

learning framework.

One clear future work for EPIP is to add more allele support as more MS data becomes available,

both from public sources and our own experiments. Enabled by EPIP's simple and modular

structure, doing so is quite straightforward as demonstrated by adding the support of HLA-

A*33:03, and we will keep doing so and update our web tool over time. We chose to adopt

simple machine learning techniques, i.e., PSSMs and LR, for this initial version of EPIP for easy

analysis and development. More recent deep learning techniques for sequence analysis, mostly

developed in the field of natural language processing [43,44], can make better use of large

datasets and model different contributing factors more effectively, and could lead to more

powerful variable length, pan-specific models. This is another future research direction of EPIP.

Moreover, immunogenicity prediction in a clinical setting depends on more than epitope

presentation [45–47], and we believe it is important to account for the extra factors contributing

to immunogenicity as well. To that end, community efforts to generate more immunogenicity

data and improved experimental techniques to profile TCR and peptide-MHC interactions [48,49]

might lead to larger datasets and deeper biological understanding that can eventually enable

accurate predictions of immunogenicity.

Conclusion

In this paper, we have developed a user-friendly, publicly available epitope prediction tool, EPIP,

that incorporates information from both MS and RNA-seq data. It significantly outperforms

other publicly available tools and can be easily extended to support more HLA alleles when new

MS data becomes available. We believe EPIP is a timely contribution to the community to

advance cancer immunotherapies and beyond.

List of abbreviations

HLA: human leukocyte antigen

MS: Mass spectrometry

PSSMs: position score specific matrixes

LR: logistic regression

EPIP: Epitope Presentation Integrated Prediction

MHC: major histocompatibility complex

TCRs: T-cell receptors

PMF: probability mass function

PPV: positive predictive value

TMG: tandem minigene

PRCs: precision recall curves

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The MS data of HCC4006 generated during the current study are available in the CNGB

Nucleotide Sequence Archive (CNSA: https://db.cngb.org).

Competing interests

The authors declare that they have no competing interests

Funding

This project is supported by National Natural Science Foundation of China (grant numbers

81702826 and 81772910), Science, Technology and Innovation Commission of Shenzhen

Municipality (grant no. JCYJ20170303151334808) and Shenzhen Municipal Government of

China (grant no. 20170731162715261)

Authors' contributions

SQ, LJL, WH, BL designed the project. WH, XH, GL, JL implemented the EPIP model. WH and

YL collected the data for training and evaluation of the model. XL, LC, SL cultured the

HCC4006 cell line. XL, LC, SL, GH, ZL, YR generated the MS data of HCC4006. LZ, YL, WH,

WL, HX, ZR analyzed the MS data of HCC4006. LJL, WH, SQ, YL, XL, LZ, BL wrote the

manuscript. LJL, SQ, BL supervised the project. All authors read and approved the final

manuscript.

Acknowledgements

We wish to thank four anonymous reviewers whose comments greatly improved the quality of

this work.

Reference

1.   Sahin U, Derhovanessian E, Miller M, Kloke BP, Simon P, Löwer M, et al. Personalized

RNA mutanome vaccines mobilize poly-specific therapeutic immunity against cancer.

Nature. 2017;547: 222–226. doi:10.1038/nature23003

2.   Ott PA, Hu Z, Keskin DB, Shukla SA, Sun J, Bozym DJ, et al. An immunogenic personal

neoantigen vaccine for patients with melanoma. Nature. 2017. pp. 217–221.

doi:10.1038/nature22991

3.   Hilf N, Kuttruff-Coqui S, Frenzel K, Bukur V, Stevanović S, Gouttefangeas C, et al.

Actively personalized vaccination trial for newly diagnosed glioblastoma. Nature. 2018.

doi:10.1038/s41586-018-0810-y

4.   Keskin DB, Anandappa AJ, Sun J, Tirosh I, Mathewson ND, Li S, et al. Neoantigen

vaccine generates intratumoral T cell responses in phase Ib glioblastoma trial. Nature.

2018. doi:10.1038/s41586-018-0792-9

5.   Jurtz V, Paul S, Andreatta M, Marcatili P, Peters B, Nielsen M. NetMHCpan-4.0:

Improved Peptide–MHC Class I Interaction Predictions Integrating Eluted Ligand and

Peptide Binding Affinity Data. J Immunol. 2017; ji1700893.

doi:10.4049/jimmunol.1700893

6.   Gfeller D, Bassani-Sternberg M. Predicting antigen presentation-What could we learn

from a million peptides? Front Immunol. 2018;9: 1–17. doi:10.3389/fimmu.2018.01716

7.   Bulik-Sullivan B, Busby J, Palmer CD, Davis MJ, Murphy T, Clark A, et al. Deep

learning using tumor HLA peptide mass spectrometry datasets improves neoantigen

identification. Nat Biotechnol 2018. 2018. doi:10.1038/nbt.4313

8.   O'Donnell TJ, Rubinsteyn A, Bonsack M, Riemer AB, Laserson U, Hammerbacher J.

MHCflurry: Open-Source Class I MHC Binding Affinity Prediction. Cell Syst. 2018;7:

129-132.e4. doi:10.1016/j.cels.2018.05.014

9.   Bassani-Sternberg M, Chong C, Guillaume P, Solleder M, Pak H, Gannon PO, et al.

Deciphering HLA motifs across HLA peptidomes correctly predicts neo-antigens and

identifies allostery in HLA specificity. bioRxiv. 2017; 1–30. doi:10.1101/098780

10.   Gfeller D, Guillaume P, Michaux J, Pak H-S, Daniel RT, Racle J, et al. Peptide length

distribution and multiple specificity in naturally presented HLA-I ligands. bioRxiv. 2018;

1–36. doi:10.1101/335661

11.   Bassani-Sternberg M, Pletscher-Frankild S, Jensen LJ, Mann M. Mass Spectrometry of

Human Leukocyte Antigen Class I Peptidomes Reveals Strong Effects of Protein

Abundance and Turnover on Antigen Presentation. Mol Cell Proteomics. 2015;14: 658–

673. doi:10.1074/mcp.M114.042812

12.   Abelin JG, Keskin DB, Sarkizova S, Hartigan CR, Zhang W, Sidney J, et al. Mass

Spectrometry Profiling of HLA-Associated Peptidomes in Mono-allelic Cells Enables

More Accurate Epitope Prediction. Immunity. 2017;46: 315–326.

doi:10.1016/j.immuni.2017.02.007

13.   Pearson H, Daouda T, Granados DP, Durette C, Bonneil E, Courcelles M, et al. MHC

class I – associated peptides derive from selective regions of the human genome. J Clin

Invest. 2016;126: 1–12. doi:10.1172/JCI91302

14.   Di Marco M, Schuster H, Backert L, Ghosh M, Rammensee H-G, Stevanović S. Unveiling

the Peptide Motifs of HLA-C and HLA-G from Naturally Presented Peptides and

Generation of Binding Prediction Matrices. J Immunol. 2017; ji1700938.

doi:10.4049/jimmunol.1700938

15.   Shao W, Pedrioli PGA, Wolski W, Scurtescu C, Schmid E, Vizcaíno JA, et al. The

SysteMHC Atlas project. Nucleic Acids Res. 2017; 1–11. doi:10.1093/nar/gkx664

16.   Trolle T, McMurtrey CP, Sidney J, Bardet W, Osborn SC, Kaever T, et al. The Length

Distribution of Class I–Restricted T Cell Epitopes Is Determined by Both Peptide Supply

and MHC Allele–Specific Binding Preference. J Immunol. 2016;196: 1480–1487.

doi:10.4049/jimmunol.1501721

17.   Kim S, Pevzner PA. MS-GF+ makes progress towards a universal database search tool for

proteomics. Nat Commun. 2014. doi:10.1038/ncomms6277

18.   The M, MacCoss MJ, Noble WS, Käll L. Fast and Accurate Protein False Discovery Rates

on Large-Scale Proteomics Data Sets with Percolator 3.0. J Am Soc Mass Spectrom. 2016.

doi:10.1007/s13361-016-1460-7

19.   Zhou F, Cao H, Zuo X, Zhang T, Zhang X, Liu X, et al. Deep sequencing of the MHC

region in the Chinese population contributes to studies of complex disease. Nat Genet.

2016. doi:10.1038/ng.3576

20.   Chen S, Zhou Y, Chen Y, Gu J. Fastp: An ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018. doi:10.1093/bioinformatics/bty560

21.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: Ultrafast

universal RNA-seq aligner. Bioinformatics. 2013. doi:10.1093/bioinformatics/bts635

22.   Li B, Dewey CN. RSEM: Accurate transcript quantification from RNA-seq data with or

without a reference genome. Bioinformatics: The Impact of Accurate Quantification on

Proteomic and Genetic Analysis and Research. 2014. doi:10.1201/b16589

23.   Liu G, Li D, Li Z, Qiu S, Li W, Chao C, et al. PSSMHCpan: a novel PSSM based

software for predicting class I peptide-HLA binding affinity. Gigascience. 2017; 1–26.

doi:10.1093/gigascience/gix017

24.   Andreatta M, Alvarez B, Nielsen M. GibbsCluster: Unsupervised clustering and alignment

of peptide sequences. Nucleic Acids Res. 2017;45: W458–W463. doi:10.1093/nar/gkx248

25.   Breiman L. Stacked regressions. Mach Learn. 1996;24: 49–64. doi:10.1007/BF00117832

26.   Pedregosa FABIANPEDREGOSA F, Alexandre Gramfort N, Michel V, Thirion

BERTRANDTHIRION B, Grisel O, Blondel M, et al. Scikitlearn: Machine Learning in

Python Gaël Varoquaux. J Mach Learn Res. 2011. doi:10.1007/s13398-014-0173-7.2

27.   Gros A, Parkhurst MR, Tran E, Pasetto A, Robbins PF, Ilyas S, et al. Prospective

identification of neoantigen-specific lymphocytes in the peripheral blood of melanoma

patients. Nat Med. 2016;22: 433–438. doi:10.1038/nm.4051

28.   Tran E, Ahmadzadeh M, Lu YC, Gros A, Turcotte S, Robbins PF, et al. Immunogenicity

of somatic mutations in human gastrointestinal cancers. Science (80-). 2015;350: 1387–

1390. doi:10.1126/science.aad1253

29.   Strønen E, Toebes M, Kelderman S, Van Buuren MM, Yang W, Van Rooij N, et al.

Targeting of cancer neoantigens with donor-derived T cell receptor repertoires. Science

(80-). 2016. doi:10.1126/science.aaf2288

30.   Zacharakis N, Chinnasamy H, Black M, Xu H, Lu YC, Zheng Z, et al. Immune

recognition of somatic mutations leading to complete durable regression in metastatic

breast cancer. Nat Med. 2018;24: 724–730. doi:10.1038/s41591-018-0040-8

31.   Hunt DF, Henderson RA, Shabanowitz J, Sakaguchi K, Michel H, Sevilir N, et al.

Characterization of peptides bound to the class I MHC molecule HLA-A2.1 by mass

spectrometry. Science (80-). 1992. doi:10.1126/science.1546328

32.   Rammensee HG, Friede T, Stevanović S. MHC ligands and peptide motifs: first listing.

Immunogenetics. 1995. doi:10.1007/BF00172063

33.   Vita R, Overton JA, Greenbaum JA, Ponomarenko J, Clark JD, Cantrell JR, et al. The

immune epitope database (IEDB) 3.0. Nucleic Acids Res. 2015;43: D405–D412.

doi:10.1093/nar/gku938

34.   Reche PA, Glutting JP, Reinherz EL. Prediction of MHC class I binding peptides using

profile motifs. Hum Immunol. 2002;63: 701–709. doi:10.1016/S0198-8859(02)00432-9

35.   Zhang H, Lund O, Nielsen M. The PickPocket method for predicting binding specificities

for receptors based on receptor pocket similarities: Application to MHC-peptide binding.

Bioinformatics. 2009. doi:10.1093/bioinformatics/btp137

36.   Schubert B, Kohlbacher O. Designing string-of-beads vaccines with optimal spacers.

582   Genome Med. 2016. doi:10.1186/s13073-016-0263-6

583   37.   The problem with neoantigen prediction. Nature Biotechnology. 2017. p. 97.

584   doi:10.1038/nbt.3800

585   38.   Van Der Maaten L, Hinton G. Visualizing Data using t-SNE. J Mach Learn Res. 2008.

586   doi:10.1007/s10479-011-0841-3

587   39.   Brawand D, Soumillon M, Necsulea A, Julien P, Csárdi G, Harrigan P, et al. The

588   evolution of gene expression levels in mammalian organs. Nature. 2011.

589   doi:10.1038/nature10532

590   40.   Evans M, Borysiewicz LK, Evans AS, Rowe M, Jones M, Gileadi U, et al. Antigen

591   Processing Defects in Cervical Carcinomas Limit the Presentation of a CTL Epitope from

592   Human Papillomavirus 16 E6. J Immunol. 2001;167: 5420–5428.

593   doi:10.4049/jimmunol.167.9.5420

594   41.   Morishima S, Akatsuka Y, Nawa A, Kondo E, Kiyono T, Torikai H, et al. Identification of

595   an HLA-A24-restricted cytotoxic T lymphocyte epitope from human papillomavirus type-

596   16 E6: The combined effects of bortezomib and interferon-γ on the presentation of a

597   cryptic epitope. Int J Cancer. 2007;120: 594–604. doi:10.1002/ijc.22312

598   42.   Peng RJ, Han BW, Cai QQ, Zuo XY, Xia T, Chen JR, et al. Genomic and transcriptomic

599   landscapes of Epstein-Barr virus in extranodal natural killer T-cell lymphoma. Leukemia.

600   2019;33: 1451–1462. doi:10.1038/s41375-018-0324-5

601   43.   Devlin J, Chang M-W, Lee K, Toutanova K. Bert: Pre-training of deep bidirectional

602   transformers for language understanding. arXiv Prepr arXiv181004805. 2018.

44. Gehring J, Auli M, Grangier D, Yarats D, Dauphin YN. Convolutional sequence to

sequence learning. 34th International Conference on Machine Learning, ICML 2017. 2017.

45. Assarsson E, Sidney J, Oseroff C, Pasquetto V, Bui H-H, Frahm N, et al. A quantitative

analysis of the variables affecting the repertoire of T cell specificities recognized after

vaccinia virus infection. J Immunol. 2007. doi:178/12/7890 [pii]

46. Calis JJA, Maybeno M, Greenbaum JA, Weiskopf D, De Silva AD, Sette A, et al.

Properties of MHC Class I Presented Peptides That Enhance Immunogenicity. PLoS

Comput Biol. 2013. doi:10.1371/journal.pcbi.1003266

47. McGranahan N, Furness AJS, Rosenthal R, Ramskov S, Lyngaa R, Saini SK, et al. Clonal

neoantigens elicit T cell immunoreactivity and sensitivity to immune checkpoint blockade.

Science (80-). 2016;351: 1463–1469. doi:10.1126/science.aaf1490

48. Bentzen AK, Marquard AM, Lyngaa R, Saini SK, Ramskov S, Donia M, et al. Large-scale

detection of antigen-specific T cells using peptide-MHC-I multimers labeled with DNA

barcodes. Nat Biotechnol. 2016;34: 1037–1045. doi:10.1038/nbt.3662

49. Bentzen AK, Such L, Jensen KK, Marquard AM, Jessen LE, Miller NJ, et al. T cell

receptor fingerprinting enables in-depth characterization of the interactions governing

recognition of peptide–MHC complexes. Nat Publ Gr. 2018. doi:10.1038/nbt.4303

Figure legend

Fig. 1 Construction of EPIP_s and EPIP. a) Construction of allele and tissue-specific EPIP

(EPIP_s). EPIP_s was built for 21 alleles that have matching expression profile.

   b) Saturation analysis of PSSM models. The PSSM models were trained and tested on 9-

11mer peptides independently for six HLA alleles; evaluated with 0.1% PPV, the performances

of PSSM improved with increasing training data, and reached a plateau with about 100-200

peptides, depending on the specific alleles.

   c) Construction of EPIP. To build the full EPIP model, we pooled peptides from 21 alleles

encoded by PSSM, expression and length features to fit a single LR model

Fig. 2 Relationship among weights and biases of 21 EPIP_s models. a) Boxplot for feature

weights and biases of EPIP_s models: PSSMs contributed most to the prediction, followed by

expressions and peptide lengths.

   b) Hive plots demonstrating the similar inter-relationships among the feature weights and

biases of different EPIP_s models, where the negative biases have been inversed for plotting.

Fig. 3 Predictive performance comparison of EPIP and EPIP_s. a) Performance comparison of

EPIP and EPIP_s across different alleles originating from the same B cell line. The error bar

indicates the standard deviation across five-fold cross validation. The performance difference

between EPIP_s and EPIP across 21 alleles is not statistically significant ($P = 0.1385$, paired t-

test)

b) Heatmap demonstrates diverse expression profiles of different cell lines originating from

different tissues. Each row represents a gene that presents peptides in at least one cell line, a total

of 10068 genes was selected. Each column represents a cell line that used in this study, the

originated tissue is indicated below the cell line. The expression values are transformed by

log2(TPM+1).

c-f) Performance comparison of EPIP and EPIP_s for the same allele originating from

different cell lines. The error bar indicates the standard deviation across five-fold cross validation.

Fig 4. Predictive performance comparison of EPIP and other methods. a) Predictive performance

of EPIP and other publicly available methods on independent MS dataset. The evaluation dataset

was obtained from four publications that contained a total of 30 MS datasets, including 16

different tissue origins and 17 HLA alleles. The mean 0.1% PPV of these 30 datasets were

displayed. EPIP significantly outperformed other methods with different expression filtering

thresholds (TPM $> 0, > 1, > 2$), and pairwise P-values (paired t-test) for TPM > 2 were shown in

the plot.

b). Precision-recall curves of EPIP and other publicly available methods on T cell responses

data. Other methods' predictions were made with expression threshold of TPM > 0. Comparisons

of EPIP and other methods with TPM > 1 and TPM > 2 were displayed in Fig S3.

c) Predictive performance of EPIP and EDGE on immunogenic SNVs. EDGE's performance

was recalculated according to the supplementary data 3c of the original paper.

d) Predictive performance of EPIP and EDGE on minimal epitopes. EDGE's performance

was taken from Figure 4 of the original paper.

Fig 5. The workflow of adding new allele HLA-A*33:03 to EPIP's supported allele repository.

HCC4006 is homozygous on all three HLA alleles (HLA-A*33:03, HLA-B*44:03 and HLA-

C*07:06, respectively). Presented peptides from HCC4006 were identified by MS (see Methods).

The derived peptidome was then deconvolved using GibbsCluster2.0. Apart from a trash cluster

of 67 peptides, two peptide clusters were identified. One cluster matched the sequence motif of

B4403 (derived from HLA-B*44:03 mono-allelic dataset [12]), and the other matched the

sequence motif of A33:03 derived from NetMHCpan4.0 online motif reviewer. Noted that due to

the lack of training data, NetMHCpan4.0 generated the A33:03 motif based on training data from

other HLA alleles in a pan-specific mode, which could explain the slight difference between the

motif we discovered and the NetMHCpan-4.0 provided. We then inputted the HLA-A*33:03-

specific peptides to EPIP, which could learn PSSM and length distributions automatically, and

resulted in an updated version of EPIP that could support the newly added allele.

Figure 1

Figure 2

A

B

Figure 3

Figure 4

Figure 5

HCC4006
A33:03
B44:03
C07:06

MS identification

DYYDDFLQR
EAGSHIIQR
ETILKHWTR
LTMSWGNTR
ATVGYGILR
NYWDKFVKR
...

deconvolution

Group1 of 2 (1225)

Group2 of 2 (1385)

Trash (67)

A33:03

B44:03

Reference

A33:03

ALNGKIYFR
ATSPALFNR
ATVGYGILR
AYHRVFVNR
AYINHLVER
AYKSILQER
...

B44:03

ADALGEEW
AEAAASAYY
AEAATPYGI
AEAEQTLRF
AEALAQRTY
AEALATHIL
...

C07:06

EMPTY

obtain A3303
specifc petpides

Conducted by software automatically

PSSM

peptide length
distritbution

add A33:03 to
supported allele
repository

EPIP