The cellular response towards lanthanum is substrate specific and

reveals a novel route for glycerol metabolism in _Pseudomonas_

putida KT2440

Matthias Wehrmann[a], Maxime Toussaint[b,c], Jens Pfannstiel[d], Patrick Billard[b,c#], Janosch

Klebensberger[a#]

[a]_University of Stuttgart, Institute of Biochemistry and Technical Biochemistry, Department of_

Technical Biochemistry, Stuttgart, Germany

[b]_Université de Lorraine, LIEC UMR7360, Faculté des Sciences et Technologies, Vandoeuvre-_

lès-Nancy, France

[c]_CNRS, LIEC UMR7360, Faculté des Sciences et Technologies, Vandoeuvre-lès-Nancy, France_

[d]_Core Facility Hohenheim, Mass Spectrometry Module, University of Hohenheim, August von_

Hartmann-Str. 3, 70599 Stuttgart, Germany

Running title: Conditional responses of _Pseudomonas putida_ towards rare earth elements

#Address correspondence to Janosch Klebensberger (janosch.klebensberger@itb.uni-

stuttgart.de) and Patrick Billard (patrick.billard@univ-lorraine.fr)

Keywords: Lanthanides, rare earth elements, proteome, glycerol metabolism, _Pseudomonas_

putida, PQQ, PedE, PedH, GarK, dehydrogenases

Abstract

Ever since the discovery of the first rare earth element (REE)-dependent enzyme, the physiological role of lanthanides has become an emerging field of research due to the potential environmental implications and biotechnological opportunities. In *Pseudomonas putida* KT2440, the two pyrroloquinoline quinone-dependent alcohol dehydrogenases PedE and PedH are inversely produced in response to La^{3+}-availability. This REE-switch is orchestrated by a complex regulatory network including the PedR2/PedS2 two-component system and is important for efficient growth on several alcoholic volatiles. As *P. putida* is exposed to a broad variety of organic compounds in its natural soil habitat, the cellular responses towards La^{3+} during growth on various model carbon sources were investigated with a differential proteomic approach. Apart from the Ca^{2+}-dependent enzyme PedE, the differential abundance of most other identified proteins was conditional and revealed a substrate specificity. Concomitant with the proteomic changes, La^{3+} had a beneficial effect on lag-phases while causing reduced growth rates and lower optical densities in stationary phase during growth on glycerol. When these growth phenotypes were evaluated with mutant strains, a novel metabolic route for glycerol utilization was identified that seems to be functional in parallel with the main degradation pathway encoded by the *glpFKRD* operon. The newly discovered route is initiated by PedE and/or PedH, which most likely convert glycerol to glyceraldehyde. In the presence of lanthanum, glyceraldehyde seems to be further oxidized to glycerate, which, upon phosphorylation to glycerate-2-phosphate by the glycerate kinase GarK, is finally channelled into the central metabolism.

Importance

The biological role of rare earth elements has long been underestimated and research has mainly focused on methanotrophic bacteria thus far. We have recently demonstrated that *P. putida*, a plant growth promoting bacterium that thrives in the rhizosphere of various feed crops, possesses a REE-dependent alcohol dehydrogenase, but knowledge about lanthanide-dependent effects on physiological traits in non-methylotrophic bacteria is still scarce. This study demonstrates that the cellular response of *P. putida* KT2440 towards La^{3+} is mostly substrate specific and that during growth on glycerol, La^{3+} has a severe effect on several growth parameters. We provide compelling evidence that these changes are linked to glycerol oxidation by the La^{3+}-dependent alcohol dehydrogenase PedH and thereby identify a novel route for glycerol metabolism in this organism. Overall, these findings demonstrate that lanthanides can alter important physiological traits of non-methylotrophic bacteria, which might consequently influence their competitiveness during colonization of various environmental niches.

Introduction

The rhizosphere, defined as the narrow region of soil surrounding plant roots, is one of the most complex ecosystems on earth containing a multitude of organisms from different taxa including fungi, oomycetes, nematodes, protozoa, algae, viruses, archaea and arthropods as well as up to 10^8 soil dwelling bacteria per gram of fresh root (1–3). Its diversity is mainly shaped by root exudates, a complex mixture of organic compounds including carbohydrates, amino acids, or carbon acids (4, 5) and plant-, fungal-, and bacteria-derived volatiles (VOCs) such as alkenes, alcohols, terpenes or benzenoids (6, 7). As such, it is not surprising that the soil-dwelling organism *P. putida* KT2440 is equipped with a broad diversity of metabolic pathways in order to maximize its cellular fitness in different environmental niches including the rhizosphere (8–10). For efficient growth on various alcoholic VOC substrates, it uses a periplasmic oxidation system consisting of the pyrroloquinoline quinone-dependent alcohol dehydrogenases (PQQ-ADHs) PedE and PedH (11, 12). The enzymes appear to be functionally redundant but differ in their metal cofactor dependency. PedE is a Ca^{2+}-dependent enzyme, whereas PedH requires the presence of rare earth elements (REEs) of the lanthanide series (Ln^{3+}) for catalytic activity (12, 13).

Although being among the most ubiquitous metals in the earth's crust, REEs were long considered to be of no biological relevance due to their low solubility under environmental conditions (14). Indeed, the only known and characterized REE-dependent enzymes thus far belong to the family of PQQ-ADHs of methano- and methylotrophic bacteria as well as the non-methylotrophic organism *P. putida* KT2440 (12, 15–20). A characteristic aspartic acid that is additionally present in the metal coordination sphere of these enzymes is associated with Ln^{3+}-binding. Notably, this specific amino acid residue has been found in the genome of many bacteria from various origins indicating a broad distribution of Ln^{3+}-dependent enzymes (21–25). Very recently, another Ln^{3+}-binding protein, called lanmodulin, was identified in *M. extorquens* AM1 (26). This periplasmic protein, which shows structural similarities with the Ca^{2+}-binding protein calmodulin, is able to bind up to four Ln^{3+} ions per protein with picomolar affinity and changes its conformation from a largely disordered to a compact,

ordered state upon REE binding. Although its exact cellular role has yet to be established, it has been

speculated that it might play a role in Ln^{3+}-uptake. Further, homologous genes have only been

identified in the genome of some other species of *Methylobacteria* and *Bradyrhizobia*.

In addition to their functional role as metal cofactor, several studies have recently investigated the

effect of REEs on cellular physiology in different methano- and methylotrophic organisms (20, 27–

32). Some of these studies found different physiological traits to be influenced in response to Ln^{3+}-

availability including changes in metabolite cross feeding, growth rates and -yields, or biofilm

formation. It is further interesting to note that REEs have been used as micro-fertilizers, especially in

China, for over 30 years, as Ln^{3+}-supplementation can be associated with increased growth of

different food crops including rice, mungbean, maize, and coconut plants (33–37).

The aforementioned results suggest that apart from the inverse transcriptional regulation of PQQ-

ADHs, which has been described in detail for different organisms including *P. putida* (12, 19, 27, 38–

42), additional responses towards REEs exist and could depend on the specific organism and/or

environmental context. To investigate the existence of such conditional cellular responses in the non-

methylotrophic organism *P. putida* KT2440, we used a differential proteomic approach during growth

on various model carbon sources that reflect the metabolic diversity of the rhizosphere. From these

experiments, we found that the Ca^{2+}-dependent PQQ-ADH PedE was the only protein showing a

differential abundance during growth on all carbon sources tested. The vast majority of identified

proteins were differentially abundant only under one specific growth condition. During growth on

glycerol and 2-phenylethanol, which both represent substrates for PedE and PedH, a

disproportionally high number of metabolism related proteins were more abundant in the presence

of La^{3+}, while this was not the case during growth on citrate and glucose, substrates for which no

activity of these enzymes is observed. In addition, physiological characteristics, such as growth rates

and the lag-phase, of cultures could be linked to the differential activity pattern of PedE and PedH

during growth on glycerol. Based on these results, we were able to identify and reconstruct a novel

metabolic route for glycerol utilisation, which depends on PedE and/or PedH activity. This route

seems to operate in conjunction with the previously described major degradation pathway initiated

by the glycerol kinase GlpK and most likely ensures efficient growth of *P. putida* on this polyol

substrate.

Materials and Methods

Bacterial strains, plasmids and culture conditions

A complete list of all strains, plasmids, and primers used in this study can be found in **Table S1** and **Table S2**. All *Pseudomonas putida* and *Escherichia coli* strains were maintained on solidified LB medium (43). If necessary, 40 µg/mL kanamycin or 20 µg/mL 5-fluorouracil were added for maintenance and/or selection. For growth, liquid LB medium or a modified M9 salt medium (12) supplemented with 5 mM 2-phenylethanol, 25 mM succinate, 10 mM glucose, 10 mM citrate, 20 mM DL-glycerate, or 20 mM glycerol as sole source of carbon and energy was used. If not stated otherwise, precultures were grown overnight in test tubes with 3 ml M9 medium supplemented with succinate at 30°C and 180 rpm. The next day, cells were washed twice with M9 medium without supplemented C-source, and used to inoculate 200 µL of M9 medium supplemented with the desired C-source in a 96-well microtiter plate (Falcon, product no. 353047 or Sarstedt, product no. 83.3924) and incubated at 30°C and 250 rpm in a microplate reader (Xenius, Safas Monaco) or 28°C and 220 rpm in a rotary shaker (Forma, Thermo Scientific). Maximum growth rates (μ_{max}) and lag-times (λ) were estimated based on fitting the natural logarithm of the relative OD_{600} values ($\ln(N/N_0)$, with N being OD_{600} at time t) with the Richards growth model using the "grofit" package in R (44). As OD_{600} decreased directly upon begin of the experiment, $\ln(N/N_{t=3h})$ was used instead of $\ln(N/N_0)$ for better fit. Differences in lag-times, growth rates, and maximal OD_{600} during stationary phase (OD_{600}^{max}) were evaluated by statistical analysis in GraphPad PRISM using a two-tailed *t*-test ($\alpha = 0.05$, $n = 3$).

Construction of plasmids

Deletion plasmids pJOE-calA, pJOE-garK, pJOE-glp, and pMW08 were constructed as follows: the 650-bp to 1000-bp regions upstream and downstream of the *calA* (PP_2426), *garK* (PP_3178), *glpFKRD* (PP_1076 to PP_1973), or *glcDEF* (PP_3745 to PP_3747) genes were amplified from genomic DNA of *P. putida* KT2440 using primers PcalA1 to PcalA4, PgarK1 to PgarK4, Pglp1 to Pglp4, or MWH03 to

MWH06 (**Table S2**). The two up- and downstream fragments and BamHI-digested pJOE6261.2 were

then joined together using one-step isothermal assembly (45).

Strain constructions

Deletion mutant strains were constructed as previously described (46). Briefly, the integration vector

pJOE6261.2 harbouring the up- and downstream regions of the target gene(s) was transformed into

P. putida KT2440 *Δupp* (KT2440*). Kanamycin (Kan) resistant and 5-fluorouracil (5-FU) sensitive

clones were selected and one of these was incubated in LB medium at 30°C for 24 h. The cell

suspension was then plated on M9 minimal agar plates containing 25 mM succinate and 20 µg ml^{-1} 5-

FU. Clones that carried the desired gene deletion were identified by colony PCR of the 5-FU^r Kan^s

clones using primer pair PcalA1/PcalA4, PgarK1/PgarK4, Pglp1/Pglp4, or MWH03/MWH06.

Protein extraction for comparative proteome analysis

For comparative proteome analysis experiments, 50 ml M9 medium supplemented with citrate,

glucose, glycerol or 2-phenylethanol and 0 or 10 µM $LaCl_3$ were inoculated with an OD_{600} of 0.05 from

succinate precultures of strain *P. putida* KT2440* in 250 ml polycarbonate Erlenmeyer flasks and

incubated at 30°C and 180 rpm. When cell cultures reached an OD_{600} of > 0.4, cells were harvested by

centrifugation for 15 min at 6000 x g and 4°C. Cell pellets were resuspended in 1 ml sample buffer

(150 mM Tris-HCl pH 6.8; 2 % SDS; 20 mM dithiothreitol) and heated for 5 min at 95°C with gentle

shaking. Subsequently, samples were centrifuged for 15 min at 21000 x g and 4°C, and the

supernatants were stored in new reaction tubes at -20 °C. In a next step, proteins were precipitated

using chloroform-methanol (47) and pellets were resuspended in Tris-buffered (50 mM, pH 8.5) urea

(6 M). Protein concentrations were determined by the Bradford assay (48).

In-solution digest of proteins and peptide purification with C18 Stage Tips

To 25 µg protein in 60 µl Tris-buffered (50 mM, pH 8.5) urea (6 M), DTT was added to a final

concentration of 10 mM to guarantee reduction of cysteines. Samples were incubated for 30 min at

56 °C under shaking at 1000 rpm. Alkylation of cysteines was performed by adding 30 mM

iodoacetamide and incubation for 45 min at room temperature in the dark. Alkylation was stopped

by adding 50 mM DTT and samples were incubated for another 10 min at RT. 500 ng LysC protease

(Roche) in 50 mM Tris buffer (pH 8.5) was added and samples were digested overnight at 30 °C. Next,

the urea in the reaction mixture was diluted to 2 M by adding the appropriate amount of Tris buffer

(50 mM, pH 8.5). 1 µg trypsin (Roche) in Tris buffer (50 mM, pH 8.5) was added and digestion was

continued for 4 hours at 37 °C. The digest was stopped by addition of 3 µl 10% (v/v) trifluoroacetic

acid (TFA). Next, peptide mixtures were concentrated and desalted on C18 stage tips (49) and dried

under vacuum. Samples were dissolved in 20 µl 0.1% (v/v) TFA. Aliquots of 1 µl were subjected to

nanoLC-MS/MS analysis.

Mass spectrometry analysis

NanoLC-ESI-MS/MS experiments were performed on an EASY-nLC 1200 system (Thermo Fisher

Scientific) coupled to a Q-Exactive Plus mass spectrometer (Thermo Fisher Scientific) using an EASY-

Spray nanoelectrospray ion source (Thermo Fisher Scientific). Tryptic peptides were directly injected

to an EASY-Spray analytical column (2 µm, 100 Å PepMapRSLC C18, 25 cm × 75 µm, Thermo Fisher

Scientific) operated at constant temperature of 35 °C. Peptides were separated at a flow rate of 250

nL/min using a 240 min gradient with the following profile: 2% - 10% solvent B in 100 min, 10% - 22%

solvent B in 80 min, 22% - 45% solvent B in 55 min, 45% - 95% solvent B in 5 min and isocratic at 90%

solvent B for 15 min. Solvents used were 0.5 % acetic acid (solvent A) and 0.5% acetic acid in

acetonitrile/H2O (80/20, v/v, solvent B). The Q Exactive Plus was operated under the control of

XCalibur 3.0.63 software. MS spectra (m/z = 300-1600) were detected in the Orbitrap at a resolution

of 70000 (m/z = 200) using a maximum injection time (MIT) of 100 ms and an automatic gain control

(AGC) value of 1×10^{6}. Internal calibration of the Orbitrap analyzer was performed using lock-mass

ions from ambient air as described elsewhere (50). Data dependent MS/MS spectra were generated

for the 10 most abundant peptide precursors in the Orbitrap using high energy collision dissociation

(HCD) fragmentation at a resolution of 17500, a normalized collision energy of 27 and an intensity

threshold of 1.3 x 10^5. Only ions with charge states from +2 to +5 were selected for fragmentation

using an isolation width of 1.6 Da. For each MS/MS scan, the AGC was set at 5 x 10^5 and the MIT was

100 ms. Fragmented precursor ions were dynamically excluded for 30 s within a 5 ppm mass window

to avoid repeated fragmentation.

Protein quantification and data analysis

Raw files were imported into MaxQuant (51) version 1.6.0.1 for protein identification and label-free

quantification (LFQ) of proteins. Protein identification in MaxQuant was performed using the

database search engine Andromeda (52). MS spectra and MS/MS spectra were searched against *P.*

putida KT2440 protein sequence database downloaded from UniProt (53). Reversed sequences as

decoy database and common contaminant sequences were added automatically by MaxQuant. Mass

tolerances of 4.5 ppm (parts per million) for MS spectra and 20 ppm for MS/MS spectra were used.

Trypsin was specified as enzyme and two missed cleavages were allowed. Carbamidomethylation of

cysteines was set as a fixed modification and protein N-terminal acetylation and oxidation were

allowed as variable modifications. The 'match between runs' feature of MaxQuant was enabled with

a match time window of one minute and an alignment time window of 20 minutes. Peptide false

discovery rate (FDR) and protein FDR thresholds were set to 0.01.

Statistical analysis including *t*-tests and principal component analysis (PCA) were performed using

Perseus software version 1.6.0.2 (54). Matches to contaminant (e.g., keratins, trypsin) and reverse

databases identified by MaxQuant were excluded from further analysis. Proteins were considered for

LFQ (label free quantification) if they were identified by at least two peptides. First, normalized LFQ

values from MaxQuant were log2 transformed. Missing values were replaced from normal

distribution using a width of 0.2 and a downshift of 2.0. Statistical differences between two sample

groups were determined using an unpaired *t*-test and a *p*-value < 0.01 and a regulation factor > 2

(log2 fold-change > 1) were considered as significant change in protein abundance. The mass

spectrometry proteomics data will be deposited to the ProteomeXchange Consortium via the PRIDE

(55) partner repository (submitted).

Purification and activity measurement of PQQ-ADHs PedE and PedH

To measure the activity of the two PQQ-ADHs PedE and PedH, the enzymes were expressed in *E. coli*

BL21(DE3) cells using plasmids pMW09 and pMW10, and purified by affinity chromatography as

described elsewhere (12). The activities with the four substrates 2-phenylethanol, citrate, glucose

and glycerol were determined at a concentration of 10 mM using a previously described colorimetric

assay (12) with one minor modification. To represent the growth conditions, 1 µM La^{3+} instead of 1

µM Pr^{3+} was used as metal cofactor for PedH.

Funding Information

The work of Matthias Wehrmann and Janosch Klebensberger was supported by an individual

research grant from the Deutsche Forschungsgemeinschaft (DFG, KL 2340/2-1). The work of

Maxime Toussaint and Patrick Billard was supported in part by Labex Ressources21 (ANR-10-

LABX-21-01).

Acknowledgements

The authors would like to thank Prof. Bernhard Hauer for his continuous support. The

authors further declare that the research was conducted in the absence of any commercial

or financial relationships that could be construed as a potential conflict of interest.

Results

We have recently demonstrated that the two PQQ-dependent ADHs PedE and PedH are inversely regulated dependent on the presence of rare earth elements (REEs) and that a complex signalling network that includes the activity of the PedR2/PedS2 two-component system orchestrates this regulation (12, 38). In order to identify whether a global cellular response of *P. putida* KT2440 towards REEs beyond the regulation of the PQQ-ADHs exists, we used a comparative proteomic analysis during growth on four different carbon and energy sources, namely 2-phenylethanol, glycerol, glucose, and citrate.

Evaluation of proteomics data

Proteins were extracted from cells of *P. putida* by SDS to enable extraction of cytoplasmic as well as transmembrane proteins followed by label free nano-LC-MS/MS quantification. In total, 2771 proteins with at least two unique peptides and an FDR ≤ 1% were identified and quantified by our proteomics approach, corresponding to 50% of the *P. putida* KT2440 proteome. Principal component analysis revealed high reproducibility for sample replicates and distinct patterns for the different carbon sources (**Fig. S1**). The majority of proteins was increased or decreased in abundance in response to the different carbon sources. In contrast, minor differences were observed in the presence or absence of La^{3+} during growth on the same carbon source. Proteins that exhibited a 2-fold or higher change in abundance between different growth conditions and a *p*-value ≤ 0.01 were considered as differentially abundant.

Effect of lanthanum on protein abundance during growth with different substrates

According to the aforementioned criteria, 56 proteins were identified as differentially abundant comparing growing cells of *P. putida* in the presence and absence of La^{3+} with different carbon sources (**Fig. 1, Table 3, Table S3-S5**). In these studies, only the Ca^{2+}-dependent PQQ-ADH PedE (PP_2674) showed a decreased abundance in response to La^{3+} during growth on all four different

carbon and energy substrates. The Ln^{3+}-dependent PQQ-ADH PedH (PP_2679) showed an increased

abundance in response to La^{3+} during growth on glucose, glycerol, and 2-phenylethanol, whereas an

uncharacterized pentapeptide repeat containing protein (PP_2673) that is directly upstream of *pedE*

showed a decreased abundance during growth on glycerol and 2-phenylethanol (**Fig. 1**). The

remaining 53 proteins were only identified under one specific growth condition (**Table 3, Table S3-**

S5).

During growth on 2-phenylethanol and glycerol, a majority of the identified proteins was increased in

abundance (80% and 70%) in response to La^{3+} (**Table 3, Table S3**). For glucose and citrate this was

different, as most of the identified proteins were found to be less abundant in response to La^{3+} (36%

and 40% during growth on glucose and citrate; **Table S4 and Table S5**). Notably, the majority of the

identified proteins were related to metabolism according to the cluster of orthologous protein

groups (COG) database (56). To test whether the observed conditional proteomic response is linked

to PedE and/or PedH activity, we determined the corresponding enzyme activities with all four

carbon and energy sources (**Table 1**). Apart from the already known substrate 2-phenylethanol, PedE

and PedH also showed activity with glycerol, whereas no activity could be detected with citrate or

glucose.

Effect of lanthanum during growth on glycerol

From our proteomic- and biochemical data, we wondered whether PedE and PedH activity could play

a beneficial role during glycerol metabolism of *P. putida* KT2440. As the degradation pathway and

growth characteristics of this organism have been recently characterized in great detail (57, 58), we

wanted to have a closer look on the effect of La^{3+} during growth on this specific substrate. In these

experiments (**Fig. 2A, Table 2**), we consistently observed a shorter lag-phase (λ) of the cultures in

response to La^{3+} availability (10.2 ± 0.2 h vs. 17.3 ± 0.2). Additionally, the corresponding values of the

specific growth rates (μ_{max}, 0.201 ± 0.004 vs. 0.341 ± 0.010 h^{-1}) and the maximal OD_{600} in stationary

phase (OD_{600}^{max}; 0.680 ± 0.014 vs. 0.884 ± 0.004) of the cultures also differed in the presence or

absence of La^{3+}, respectively. As the purified PedH enzyme showed an 3-fold higher specific activity

towards glycerol compared to PedE *in vitro* (0.9 ± 0.1 U mg^{-1} vs. 0.3 ± 0.1 U mg^{-1}; **Table 1**), we

speculated that this increased glycerol conversion by PedH could be the underlying cause for the

observed differences in growth parameters. When subsequently a Δ*pedE* Δ*pedH* strain was analysed

for growth in the presence and absence of La^{3+} (**Fig. 2B**, **Table 2**), no significant differences in λ and

μ$_{max}$ were observed for the Δ*pedE* Δ*pedH* strain in response to La^{3+} while, although less profound,

small differences in OD$_{600}$^{max} were still detected. Further, we found that under both conditions the

double deletion strain showed a lag-phase that was undistinguishable from that of the parental strain

in the absence of La^{3+} but dramatically longer than that of the parental strain in the presence of La^{3+}

(17.5 ± 0.3 h in the absence of La^{3+} and 17.2 ± 0.4 h in the presence of 10 μM La^{3+}). Interestingly, the

growth rates under both conditions (0.276 ± 0.008 h^{-1} and 0.291 ± 0.012 h^{-1}) were significantly higher

($p < 0.01$) than those of the parental strain in presence of La^{3+} while still being significantly below ($p <$

0.05) those of the parental strain in the absence of La^{3+}.

These results implied that the two PQQ-ADHs can indeed be beneficial for growth on glycerol and

that a functionally active PedH enzyme is the underlying cause for the La^{3+}-dependent differences in

lag-times and growth rates and to some extent also for differences in OD$_{600}$^{max} of KT2440 cultures. As

PedE and PedH as well as the remaining proteins that were found to be differentially abundant in

response to La^{3+} during growth on glycerol, are not part of the described degradation pathway in *P.*

putida KT2440 (57, 58), we hypothesized that an additional metabolic route exists (**Fig. 3**). Based on

our proteomic data, this route could be initiated by the activity of PedH and the oxidation of glycerol

to glycolaldehyde. In the next steps glycolaldehyde could be oxidized to glycerate by PedH, the

aldehyde dehydrogenase AldB-II, or the aldehyde oxidase complex composed of proteins PP_3621

(IorA-II), PP_3622 and PP_3623 (AdhB). After phosphorylation by the glycerate kinase GarK,

glycerate-2-phosphate could eventually enter the central metabolism.

If such a metabolic route exists, a Δ*glpFKRD* deletion strain should still be able to grow with glycerol

as sole source of energy and carbon, whereas a Δ*pedE* Δ*pedH* Δ*glpFKRD* deletion mutant should not.

To test this scenario, the corresponding strains were constructed and characterized for their growth

phenotypes (**Fig. 4A**). We found that *P. putida* KT2440 indeed grew on glycerol independent of the

glp regulon, although growth was dramatically impaired compared to the parental strain or strain

Δ*pedE* Δ*pedH*. When PedE and PedH were additionally deleted, growth could not be observed even

after a prolonged incubation time of 5 d. This supported our hypothesis that a metabolic route for

glycerol next to the *glp* pathway exists and that this route is initiated by the PedE and PedH proteins,

most likely by the oxidation of glycerol to glyceraldehyde (**Fig. 3**). Given that the route further

proceeds via glycerate and glycerate-2-phosphate, different cellular concentrations of these

metabolites would be expected during growth on glycerol in a mutant that is not able to use the *glp*

pathway. Interestingly, a companion study to this work employed a metabolome analysis using

glycerol-growing cells of *P. putida* KT2440 and a Δ*glpK* deletion strain, which can only use the

proposed novel route via PedE and PedH (59). In their experiments, the authors observed that the

glycerate concentration measured for the Δ*glpK* strain was dramatically increased compared to the

wild type, whereas concentrations of glyceraldehyde and glyceraldehyde-3-phosphate were in the

same range for both strains. These data suggested that glycerate is indeed an intermediate during

glp-independent growth, and that the activity of downstream proteins represent the bottleneck of

the metabolic route leading to the observed accumulation. As our proteomic data indicated the

involvement of the predicted glycerate kinase GarK, we constructed a Δ*garK* deletion strain and

speculated that this strain should lack the ability to phosphorylate glycerate and would hence be

incapable of channelling glycerate-2-phosphate into the central metabolism. We indeed observed no

growth of a Δ*garK* mutant on glycerate even after incubation of up to 5 d, while strain Δ*pedE* Δ*pedH*

Δ*glpFKRD* could grow and reached OD_{600}^{max} within 72 h of incubation under the condition tested (**Fig.**

4B).

When grown on glycerol, no significant effect on growth rates and lag-times as well as only a minor,

but significant, negative effect on the OD_{600}^{max} ($p < 0.01$) was observed for the Δ*garK* deletion in the

absence of La^{3+}. In contrast, the same deletion caused a dramatic growth impairment in the presence

of La^{3+} (**Fig. 5A**, **Table 2**) and consequently the stationary phase was not reached during the 40 h of

incubation. Hence, no reliable growth parameter could be deduced from these data. It is however

obvious that the growth rate was far below the one of the parental strain in the presence of La^{3+}.

Notably, some of the most severely upregulated proteins in response to La^{3+} can either be associated

with stress responses, namely the multidrug efflux pump MexEF and the alkylhydroperoxide

reductase subunits AhpC/F, or enzymes, which play no obvious roles within the proposed metabolic

route, such as CalA, a predicted coniferyl alcohol dehydrogenase, and the glycolate oxidase GlcDEF.

Nevertheless, to test the potential metabolic influence of the latter two enzymes, we constructed

and analysed the corresponding Δ*calA* and Δ*glcDEF* mutants and tested their growth pattern with

glycerol (**Fig. 5B; Table 2**). The Δ*glcDEF* mutants showed a similar growth behaviour than the parental

strain. In contrast, strain Δ*calA* exhibited a significantly increased growth rate ($p < 0.01$) and higher

OD_{600}^{max} ($p < 0.01$) than the parental strain in the presence of La^{3+} while showing no significant

differences in OD_{600}^{max} and maximal growth rate and only minor differences ($p < 0.05$) in lag-times in

the absence of La^{3+}(**Fig. 5C; Table 2**).

Discussion

In the present study, the cellular responses of the soil bacterium *P. putida* KT2440 towards La^{3+} availability during growth on several carbon and energy sources were investigated. The only protein that showed a differential abundance independent of the substrate used for growth was the Ca^{2+}-dependent PQQ-ADH PedE. This result is in line with data from a previous study (38), which demonstrated that the La^{3+}-induced downregulation of *pedE* is dependent on the PedS2/PedR2 two-component system that, based on our current observation, seems to be functional under all tested conditions. The other two proteins that showed differential abundance under more than one culture condition (PedH, PP_2673) are both also part of the *ped* gene cluster. Notably, the carbon sources under which these proteins were identified either represent substrates of the two PQQ-ADHs PedE and PedH, or can be degraded by an enzyme depending on the same PQQ-cofactor, namely the glucose dehydrogenase Gcd. The remaining 53 proteins that showed differential abundance in response to La^{3+} were identified only during growth on one substrate, suggesting a conditional regulation. For glycerol, we provide striking evidence that the increased activity of PedH compared to PedE is the primary cause for the observed proteomic and physiological changes during growth.

Thus far, the degradation of glycerol was described to start by the uptake *via* GlpF, phosphorylation by GlpK, and subsequent GlpD-catalysed oxidation of glycerol-3-phosphate to dihydroxyacetone-3-phosphate (58). In a next step, dihydroxyacetone-3-phosphate is interconverted to glyceraldehyde-3-phosphate and enters the central metabolism. This pathway is negatively regulated by the transcriptional regulator GlpR, and the de-repression of the *glpFKRD* operon is believed to depend on the intracellular concentration of glycerol-3-phosphate, which finally impacts the lag-phase of cultures (57). As such, it was interesting to find that growth on glycerol in the presence of La^{3+} led to a shorter lag phase and lower growth rate of the parent strain, and that a ΔpedE ΔpedH mutant showed a lag-phase similar to the parent in absence of La^{3+} without any beneficial effect of La^{3+} while still growing with a higher growth rate than the parent strain in presence of La^{3+}. Further experiments revealed that a ΔglpFKRD deletion strain is still able to grow on glycerol, while a ΔpedE ΔpedH

Δ*glpFKRD* is not. With the additional observation that a Δ*garK* mutant cannot utilize glycerate, we

propose a novel route for glycerol metabolism, in which PedE and PedH catalyse the initial oxidation

of glycerol to glyceraldehyde. In the presence of La^{3+}, the route seems to proceed via a second

oxidation step to glycerate, which is subsequently converted to glycerate-2-phosphate by the activity

of GarK (**Fig. 3**). Our data suggest that the PedE/PedH-dependent route, despite being important for

efficient growth, is not the main route for glycerol metabolism, as the effect of the Δ*pedE* Δ*pedH*

deletion on the lag-phase with glycerol is far less severe than deletion of the *glpFKRD* gene cluster. It

also appears that the PedE/PedH-dependent route is less efficient than the *glp* pathway, as the

overall growth of the Δ*glpFKRD* strain is substantially impaired compared to the Δ*pedE* Δ*pedH* strain.

A possible explanation for these observations could be the formation of the toxic intermediate

glyceraldehyde, which is known for its protein crosslinking properties and the formation of

superoxide radicals due to auto-oxidation (60, 61). The observed differences in growth rates and

OD_{600}^{max} in response to La^{3+} in the parent strain would thus reflect the increased metabolic flux

towards glyceraldehyde due to the higher specific activity of PedH compared to PedE. This would

also explain the severe La^{3+}-dependent growth impairment of the Δ*garK* mutant, as one can assume

that even higher concentrations of glyceraldehyde accumulate in a mutant that cannot process

glycerate. The notion that the MexEF RND transporter proteins, which are involved in efflux of

various toxic compounds (62), and the hydroyperoxide reductase proteins AhpC and AhpF, which

have been linked to ROS detoxification in *P. putida* (63), were also more abundant in presence of La^{3+}

during growth on glycerol are supportive of such a hypothesis.

To explain the impact of La^{3+} on the lag-times of cultures, one could speculate that in addition to

glycerol-3-phosphate, also other phosphorylated derivatives, such as glycerate-2-phosphate, are able

to relieve the repression of *glpFKRD* by GlpR. However, as in the absence of La^{3+} the growth

phenotype of the Δ*garK* mutant is indistinguishable from that of the parental strain, and since the

growth rate of the parent strain in absence of La^{3+} is still significantly higher than the growth rate of

strain Δ*pedE* Δ*pedH,* we postulate that yet another metabolic route is present that contributes to

growth without affecting the lag-phase. This route could proceed via the phosphorylation of

glyceraldehyde to glyceraldehyde-3-phosphate by the activity of a so-far unknown kinase. Whether

both alternative routes to the *glp*-dependent pathway are functional in parallel or whether the

metabolic flux *via* glycerate is exclusively induced in the presence of La^{3+} is currently unknown and

would need to be tested in future studies.

Similarly, the question why proteins that cannot be associated to the predicted glycerol metabolism

pathways are among the most differentially abundant proteins in response to La^{3+} remains to be

elucidated. It is however worthwhile noting that the two enzymes CalA and GlcDEF are either known,

(GlcDEF) or predicted by the PROSITE software tool (CalA) (64) to be catalytically active on 2-hydroxy

acids. As such, activity towards pathway intermediates such as glycerate cannot be excluded at the

moment.

From our data, the La^{3+}-dependent proteomic and physiological changes during growth on glycerol

can be explained by a shift in metabolic flux resulting from differences in the specific enzyme

activities between PedH and PedE. A similar metabolic-driven interpretation can also be used to

explain the differences observed during growth on other carbon and energy sources that are known

to be substrates for PedE and PedH such as 2-phenylethanol. However, this logic clearly fails to

explain the conditional differences during growth on glucose and citrate, as they do not represent

substrates for PedE and/or PedH. Despite the fact that we currently do not know the underlying

cause for conditional regulation under these conditions, it indicates that beside the interaction with

PedH and the TCS PedS2/PedR2, additional effects of REEs in *P. putida* do exist. Such effects could

include the inhibition of protein functions by mismetallation (65, 66), changes in the physiology of

the outer membrane (67), or so far unknown REE-dependent enzymes and regulator proteins. The

latter explanation is of particular interest, as two recent studies provide strong evidence that specific

import systems that can transport Ln^{3+} into the cytoplasm of methylotrophic bacteria do exist (41,

68).

Altogether, the current study demonstrates that the utilization of REEs can influence important

physiological traits of *P. putida*, which could be highly beneficial in competitive environmental niches

such as the rhizosphere. The previously reported fertilizing effect of REEs on different food crops

could hence be the result of increased competitiveness of plant growth promoting organisms such as

P. putida during root colonization. This hypothesis is further supported by a recent study, which

found that Pseudomonads predominantly thrive on root exudates *in vivo* and are hence enriched in

the rhizosphere of *Arabidopsis thaliana* (10). As such, it will be interesting to see what future

research will add to the currently emerging theme of REEs being an important micronutrient for

methylotrophic and non-methylotrophic organisms.

Reference

1. Mendes R, Garbeva P, Raaijmakers J. 2013. The rhizosphere microbiome: significance of plant beneficial, plant pathogenic, and human pathogenic microorganisms. FEMS Microbiol Rev 37:634–663.

2. Raaijmakers J, Paulitz TC, Steinberg C, Alabouvette C, Moënne-Loccoz Y. 2009. The rhizosphere: a playground and battlefield for soilborne pathogens and beneficial microorganisms. Plant Soil 321:341–361.

3. Berg G, Roskot N, Steidle A, Eberl L, Zock A, Smalla K. 2002. Plant-Dependent Genotypic and Phenotypic Diversity of Antagonistic Rhizobacteria Isolated from Different *Verticillium* Host Plants Plants. Appl Environ Microbiol 68:3328–3338.

4. Carvalhais LC, Dennis PG, Fedoseyenko D, Hajirezaei MR, Borriss R, Von Wirén N. 2011. Root exudation of sugars, amino acids, and organic acids by maize as affected by nitrogen, phosphorus, potassium, and iron deficiency. J Plant Nutr Soil Sci 174:3–11.

5. Dakora FD, Phillips DA. 2002. Root exudates as mediators of mineral acquisition in low-nutrient environments. Plant Soil 245:35–47.

6. Schmidt R, Cordovez V, de Boer W, Raaijmakers J, Garbeva P. 2015. Volatile affairs in microbial interactions. ISME J 9:2329–2335.

7. Insam H, Seewald MSA. 2010. Volatile organic compounds (VOCs) in soils. Biol Fertil Soils 46:199–213.

8. Wackett LP. 2003. *Pseudomonas putida*—a versatile biocatalyst. Nat Biotechnol 21:136–138.

9. Nelson KE, Weinel C, Paulsen IT, Dodson RJ, Hilbert H, Martins dos Santos VAP, Fouts DE, Gill SR, Pop M, Holmes M, Brinkac L, Beanan M, DeBoy RT, Daugherty S, Kolonay J, Madupu R, Nelson W, White O, Peterson J, Khouri H, Hance I, Lee PC, Holtzapple E, Scanlan D, Tran K, Moazzez A, Utterback T, Rizzo M, Lee K, Kosack D, Moestl D, Wedler H, Lauber J, Stjepandic D, Hoheisel J, Straetz M, Heim S, Kiewitz C, Eisen J, Timmis KN, Dusterhoft A, Tummler B, Fraser CM. 2002. Complete genome sequence and comparative analysis of the metabolically versatile *Pseudomonas putida* KT2440. Environ Microbiol 4:799–808.

10.   Worsley SF, Macey MC, Newitt JT, Patrick E, Yu DW, Wilkinson B, Murrell C, Hutchings MI.

2019. Investigating the role of exudates in recruiting Streptomyces bacteria to the *Arabidopsis*

thaliana root microbiome. bioRxiv 532309.

11.   Mückschel B, Simon O, Klebensberger J, Graf N, Rosche B, Altenbuchner J, Pfannstiel J, Huber

A, Hauer B. 2012. Ethylene glycol metabolism by *Pseudomonas putida*. Appl Environ Microbiol

78:8531–9.

12.   Wehrmann M, Billard P, Martin-Meriadec A, Zegeye A, Klebensberger J. 2017. Functional Role

of Lanthanides in Enzymatic Activity and Transcriptional Regulation of Pyrroloquinoline

Quinone-Dependent Alcohol Dehydrogenases in *Pseudomonas putida* KT2440. MBio

8:e00570-17.

13.   Takeda K, Matsumura H, Ishida T, Samejima M, Igarashi K, Nakamura N, Ohno H. 2013. The

two-step electrochemical oxidation of alcohols using a novel recombinant PQQ alcohol

dehydrogenase as a catalyst for a bioanode. Bioelectrochemistry 94:75–78.

14.   Firsching FH, Brune SN. 1991. Solubility products of the trivalent rare-earth phosphates. J

Chem Eng Data 36:93–95.

15.   Pol A, Barends TRM, Dietl A, Khadem AF, Eygensteyn J, Jetten MSM, Op den Camp HJM. 2014.

Rare earth metals are essential for methanotrophic life in volcanic mudpots. Environ

Microbiol 16:255–264.

16.   Hibi Y, Asai K, Arafuka H, Hamajima M, Iwama T, Kawai K. 2011. Molecular structure of La^{3+}-

induced methanol dehydrogenase-like protein in *Methylobacterium radiotolerans*. J Biosci

Bioeng 111:547–549.

17.   Fitriyanto NA, Fushimi M, Matsunaga M, Pertiwiningrum A, Iwama T, Kawai K. 2011.

Molecular structure and gene analysis of Ce^{3+}-induced methanol dehydrogenase of

Bradyrhizobium sp. MAFF211645. J Biosci Bioeng 111:613–617.

18.   Good NM, Vu HN, Suriano CJ, Subuyuj GA, Skovran E, Martinez-Gomez NC. 2016.

Pyrroloquinoline Quinone Ethanol Dehydrogenase in *Methylobacterium extorquens* AM1

Extends Lanthanide-Dependent Metabolism to Multicarbon Substrates. J Bacteriol 198:3109–

3118.

19.   Vu HN, Subuyuj GA, Vijayakumar S, Good NM, Martinez-Gomez NC, Skovran E. 2016.

Lanthanide-Dependent Regulation of Methanol Oxidation Systems in *Methylobacterium*

extorquens AM1 and Their Contribution to Methanol Growth. J Bacteriol 198:1250–1259.

20.   Masuda S, Suzuki Y, Fujitani Y, Mitsui R, Nakagawa T, Shintani M, Tani A. 2018. Lanthanide-

Dependent Regulation of Methylotrophy in *Methylobacterium aquaticum* Strain 22A.

mSphere 3:e00462-17.

21.   Chistoserdova L. 2016. Lanthanides: New life metals? World J Microbiol Biotechnol 32:138.

22.   Keltjens JT, Pol A, Reimann J, Op den Camp HJM. 2014. PQQ-dependent methanol

dehydrogenases: rare-earth elements make a difference. Appl Microbiol Biotechnol 98:6163–

83.

23.   Sowell SM, Abraham PE, Shah M, Verberkmoes NC, Smith DP, Barofsky DF, Giovannoni SJ.

2011. Environmental proteomics of microbial plankton in a highly productive coastal

upwelling system. ISME J 5:856–865.

24.   Taubert M, Grob C, Howat AM, Burns OJ, Dixon JL, Chen Y, Murrell JC. 2015. XoxF encoding an

alternative methanol dehydrogenase is widespread in coastal marine environments. Environ

Microbiol 17:3937–3948.

25.   Lv H, Sahin N, Tani A. 2018. Isolation and genomic characterization of *Novimethylophilus*

kurashikiensis gen. nov. sp. nov., a new lanthanide-dependent methylotrophic species of

Methylophilaceae. Environ Microbiol 20:1204–1223.

26.   Cotruvo JA, Featherston ER, Mattocks JA, Ho J V, Laremore TN. 2018. Lanmodulin: A Highly

Selective Lanthanide-Binding Protein from a Lanthanide-Utilizing Bacterium. J Am Chem Soc

jacs.8b09842.

27.   Gu W, Semrau JD. 2017. Copper and cerium-regulated gene expression in *Methylosinus*

trichosporium OB3b. Appl Microbiol Biotechnol 101:8499–8516.

28.   Good NM, Walser ON, Moore RS, Suriano CJ, Huff AF, Cecilia N. 2018. Investigation of

lanthanide-dependent methylotrophy uncovers complementary roles for alcohol

dehydrogenase enzymes.

29.   Fitriyanto NA, Nakamura M, Muto S, Kato K, Yabe T, Iwama T, Kawai K, Pertiwiningrum A.

2011. Ce^{3+}-induced exopolysaccharide production by *Bradyrhizobium* sp. MAFF211645. J

Biosci Bioeng 111:146–152.

30.   Zheng Y, Huang J, Zhao F, Chistoserdova L. 2018. Physiological Effect of XoxG(4) on

Lanthanide-Dependent Methanotrophy. MBio 9:e02430-17.

31.   Krause SMB, Johnson T, Samadhi Karunaratne Y, Fu Y, Beck DAC, Chistoserdova L, Lidstrom

ME. 2017. Lanthanide-dependent cross-feeding of methane-derived carbon is linked by

microbial community interactions. Proc Natl Acad Sci 114:358–363.

32.   Akberdin IR, Collins DA, Hamilton R, Oshchepkov DY, Shukla AK, Nicora CD, Nakayasu ES,

Adkins JN, Kalyuzhnaya MG. 2018. Rare Earth Elements Alter Redox Balance in

Methylomicrobium alcaliphilum 20ZR. Front Microbiol 9:1–12.

33.   Diatloff E, Smith FW, Asher CJ. 1995. Rare earth elements and plant growth: III. Responses of

corn and mungbean to low concentrations of cerium in dilute, continuously flowing nutrient

solutions. J Plant Nutr 18:1991–2003.

34.   Diatloff E, Asher CJ, Smith FW. 1996. Rare earth elements and plant growth. Proc 8th Aust

Agron Conf Toowoomba, Queensland, Aust 30 January-2 February, 1996 3–6.

35.   Pang X, Li D, Peng A. 2002. Application of rare-earth elements in the agriculture of China and

its environmental behavior in soil. Environ Sci Pollut Res 9:143–148.

36.   Wahid PA, Valiathan MS, Kamalam N V., Eapen JT, Vijayalakshmi S, Prabhu RK, Mahalingam

TR. 2000. Effect of rare earth elements on growth and nutrition of coconut palm and root

competition for these elements between the palm and *calotropis gigantea*. J Plant Nutr

23:329–338.

37.   Xie ZB, Zhu JG, Chu HY, Zhang YL, Zeng Q, Ma HL, Cao ZH. 2002. Effect of lanthanum on rice

production, nutrient uptake, and distribution. J Plant Nutr 25:2315–2331.

38.   Wehrmann M, Berthelot C, Billard P, Klebensberger J. 2018. The PedS2/PedR2 Two-

Component System Is Crucial for the Rare Earth Element Switch in *Pseudomonas putida*

KT2440. mSphere 3:1–12.

39.   Farhan Ul Haque M, Kalidass B, Bandow N, Turpin EA, DiSpirito AA, Semrau JD. 2015. Cerium

Regulates Expression of Alternative Methanol Dehydrogenases in *Methylosinus trichosporium*

OB3b. Appl Environ Microbiol 81:7546–7552.

40.   Chu F, Beck DAC, Lidstrom ME. 2016. MxaY regulates the lanthanide-mediated methanol

dehydrogenase switch in *Methylomicrobium buryatense*. PeerJ 4:e2435.

41.   Ochsner AM, Hemmerle L, Vonderach T, Nüssli R, Bortfeld-Miller M, Hattendorf B, Vorholt JA.

2019. Use of rare-earth elements in the phyllosphere colonizer *Methylobacterium extorquens*

PA1. Mol Microbiol 0–2.

42.   Chu F, Lidstrom ME. 2016. XoxF Acts as the Predominant Methanol Dehydrogenase in the

Type I Methanotroph *Methylomicrobium buryatense*. J Bacteriol 198:1317–1325.

43.   Maniatis T, Fritsch E, Sambrook J, Laboratory CSH. 1982. Molecular Cloning⊡: A Laboratory

Manual. Cold Spring Harbor, N.Y. Cold Spring Harbor Laboratory.

44.   Kahm M, Kschischo M, Ludwig J, Lichtenberg-Fraté H, Hasenbrink G. 2015. grofit⊡: Fitting

Biological Growth Curves with R . J Stat Softw 33.

45.   Gibson DG. 2011. Enzymatic Assembly of Overlapping DNA Fragments, p. 349–361. *In*

Methods in Enzymology.

46.   Graf N, Altenbuchner J. 2011. Development of a method for markerless gene deletion in

Pseudomonas putida. Appl Environ Microbiol 77:5549–5552.

47.   Wessel D, Flügge UI. 1984. A method for the quantitative recovery of protein in dilute solution

in the presence of detergents and lipids. Anal Biochem 138:141–143.

48.   Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram

quantities of protein utilizing the principle of protein-dye binding. Anal Biochem 72:248–254.

49.   Rappsilber J, Ishihama Y, Mann M. 2003. Stop and Go Extraction Tips for Matrix-Assisted Laser

Desorption/Ionization, Nanoelectrospray, and LC/MS Sample Pretreatment in Proteomics.

Anal Chem 75:663–670.

50.   de Godoy LMF, Mortensen P, Makarov A, Li G, Macek B, Horning S, Pesch R, Mann M, Olsen J

V., Lange O. 2005. Parts per Million Mass Accuracy on an Orbitrap Mass Spectrometer via Lock

Mass Injection into a C-trap. Mol Cell Proteomics 4:2010–2021.

51. Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized

p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol

26:1367–1372.

52. Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen J V., Mann M. 2011. Andromeda: A

Peptide Search Engine Integrated into the MaxQuant Environment. J Proteome Res 10:1794–

1805.

53. Bateman A, Martin MJ, O'Donovan C, Magrane M, Alpi E, Antunes R, Bely B, Bingley M, Bonilla

C, Britto R, Bursteinas B, Bye-AJee H, Cowley A, Da Silva A, De Giorgi M, Dogan T, Fazzini F,

Castro LG, Figueira L, Garmiri P, Georghiou G, Gonzalez D, Hatton-Ellis E, Li W, Liu W, Lopez R,

Luo J, Lussi Y, MacDougall A, Nightingale A, Palka B, Pichler K, Poggioli D, Pundir S, Pureza L, Qi

G, Rosanoff S, Saidi R, Sawford T, Shypitsyna A, Speretta E, Turner E, Tyagi N, Volynkin V,

Wardell T, Warner K, Watkins X, Zaru R, Zellner H, Xenarios I, Bougueleret L, Bridge A, Poux S,

Redaschi N, Aimo L, ArgoudPuy G, Auchincloss A, Axelsen K, Bansal P, Baratin D, Blatter MC,

Boeckmann B, Bolleman J, Boutet E, Breuza L, Casal-Casas C, De Castro E, Coudert E, Cuche B,

Doche M, Dornevil D, Duvaud S, Estreicher A, Famiglietti L, Feuermann M, Gasteiger E, Gehant

S, Gerritsen V, Gos A, Gruaz-Gumowski N, Hinz U, Hulo C, Jungo F, Keller G, Lara V, Lemercier

P, Lieberherr D, Lombardot T, Martin X, Masson P, Morgat A, Neto T, Nouspikel N, Paesano S,

Pedruzzi I, Pilbout S, Pozzato M, Pruess M, Rivoire C, Roechert B, Schneider M, Sigrist C,

Sonesson K, Staehli S, Stutz A, Sundaram S, Tognolli M, Verbregue L, Veuthey AL, Wu CH,

Arighi CN, Arminski L, Chen C, Chen Y, Garavelli JS, Huang H, Laiho K, McGarvey P, Natale DA,

Ross K, Vinayaka CR, Wang Q, Wang Y, Yeh LS, Zhang J. 2017. UniProt: the universal protein

knowledgebase. Nucleic Acids Res 45:D158–D169.

54. Tyanova S, Temu T, Sinitcyn P, Carlson A, Hein MY, Geiger T, Mann M, Cox J. 2016. The

Perseus computational platform for comprehensive analysis of (prote)omics data. Nat

Methods 13:731–740.

55. Vizcaíno JA, Csordas A, Del-Toro N, Dianes JA, Griss J, Lavidas I, Mayer G, Perez-Riverol Y,

Reisinger F, Ternent T, Xu Q-W, Wang R, Hermjakob H. 2016. 2016 update of the PRIDE

database and its related tools. Nucleic Acids Res 44:D447–D456.

56. Galperin MY, Makarova KS, Wolf YI, Koonin E V. 2015. Expanded microbial genome coverage

and improved protein family annotation in the COG database. Nucleic Acids Res 43:D261–

D269.

57. Nikel PI, Romero-Campero FJ, Zeidman J a, Goñi-Moreno Á, de Lorenzo V. 2015. The Glycerol-

Dependent Metabolic Persistence of *Pseudomonas putida* KT2440 Reflects the Regulatory

Logic of the GlpR Repressor. MBio 6:1–13.

58. Nikel PI, Kim J, de Lorenzo V. 2014. Metabolic and regulatory rearrangements underlying

glycerol metabolism in *Pseudomonas putida* KT2440. Environ Microbiol 16:239–254.

59. Walsh M, Casey M, Kenny S, Narancic T, Blank L, Wierckx N, Ballerstedt H, O'Connor KE. 2019.

Insights into an alternative pathway for glycerol metabolism in a glycerol kinase deficient

Pseudomonas putida KT2440. Submitted to BioRxiv.

60. Tessier FJ, Monnier VM, Sayre LM, Kornfield JA. 2003. Triosidines: novel Maillard reaction

products and cross-links from the reaction of triose sugars with lysine and arginine residues.

Biochem J 369:705–719.

61. Benov L, Beema AF. 2003. Superoxide-dependence of the short chain sugars-induced

mutagenesis. Free Radic Biol Med 34:429–433.

62. Li XZ, Zhang L, Poole K. 1998. Role of the multidrug efflux systems of *Pseudomonas aeruginosa*

in organic solvent tolerance. J Bacteriol 180:2987–91.

63. Hishinuma S, Yuki M, Fujimura M, Fukumori F. 2006. OxyR regulated the expression of two

major catalases, KatA and KatB, along with peroxiredoxin, AhpC in *Pseudomonas putida*.

Environ Microbiol 8:2115–2124.

64. Sigrist CJA, de Castro E, Cerutti L, Cuche BA, Hulo N, Bridge A, Bougueleret L, Xenarios I. 2012.

New and continuing developments at PROSITE. Nucleic Acids Res 41:D344–D347.

65. Agarwal N, Kalra VK. 1983. Interaction of lanthanide cations and uranyl ion with the

631   calcium/proton antiport system in *Mycobacterium phlei*. Biochim Biophys Acta - Biomembr

632   727:285–292.

633   66.   Brown PH, Rathjen AH, Graham RD, Tribe DE. 1990. Chapter 92 Rare earth elements in

634   biological systems, p. 423–452. *In* Handbook on the Physics and Chemistry of Rare Earths.

635   67.   Peng L, Yi L, Zhexue L, Juncheng Z, Jiaxin D, Daiwen P, Ping S, Songsheng Q. 2004. Study on

636   biological effect of La^{3+} on *Escherichia coli* by atomic force microscopy. J Inorg Biochem 98:68–

637   72.

638   68.   Mattocks JA, Ho J V., Cotruvo JA. 2019. A Selective, Protein-Based Fluorescent Sensor with

639   Picomolar Affinity for Rare Earth Elements. J Am Chem Soc jacs.8b12155.

640
641

Tables

Table 1: Specific enzyme activities of purified PedE and PedH with the four tested growth substrates at 10 mM measured with 2,6-dichlorophenolindophenol (DCPIP) dependent colorimetric assay. Data represent the average of biological triplicates with according standard deviation. Activities below detection limit are indicated (n. d.).

Substrate	Mean Specific Activity (U mg^{-1}) ± SD	
	PedE 1 mM Ca^{2+}	PedH 1 µM La^{3+}
Citrate	n.d.	n.d.
Glucose	n.d.	n.d.
Glycerol	0.3 ± 0.1	0.9 ± 0.1
2-Phenylethanol	8.0 ± 0.4	6.3 ± 0.3

Table 2: Lag-times (λ), maximal OD_{600} during stationary phase (OD_{600}^{max}), and maximal growth rates
(μ_{max}) of different *P. putida* strains during growth with M9 medium supplemented with 20 mM
glycerol and 0 µM or 10 µM La^{3+} incubated in microtiter plates at 30°C and 250 rpm (see also Fig. 2
and Fig.5). Maximal growth rates and lag-times were determined by fitting growth curves to the
Richards model using "grofit" package in R (44). Cultures were incubated at 250 rpm and 30°C in
microplate reader with constant OD_{600} measurement. Data points represent average of biological
triplicates with corresponding error (λ, μ_{max}) or standard deviation (OD_{600}^{max}). No growth parameters
could be determined for cultures that did not reach stationary phase during incubation time (n. d.).

Strain	λ [h]		OD_{600}^{max}		μ_{max} [h^{-1}]	
	0 µM La^{3+}	10 µM La^{3+}	0 µM La^{3+}	10 µM La^{3+}	0 µM La^{3+}	10 µM La^{3+}
KT2440*	17.3 ± 0.2	10.2 ± 0.2	0.884 ± 0.004	0.680 ± 0.014	0.341 ± 0.010	0.201 ± 0.004
Δ*pedE* Δ*pedH*	17.5 ± 0.3	17.2 ± 0.4	0.869 ± 0.002	0.802 ± 0.013	0.276 ± 0.008	0.291 ± 0.012
Δ*garK*	17.1 ± 0.3	n.d.	0.805 ± 0.005	n.d.	0.321 ± 0.011	n.d.
Δ*calA*	16.0 ± 0.3	11.1 ± 0.3	0.890 ± 0.004	0.753 ± 0.006	0.336 ± 0.011	0.271 ± 0.008
Δ*glcDEF*	17.0 ± 0.2	10.1 ± 0.2	0.862 ± 0.016	0.678 ± 0.004	0.315 ± 0.009	0.207 ± 0.004

660 **Table 3:** List of upregulated proteins in presence of 10 µM La^{3+} compared to the absence of La^{3+}
661 when grown with glycerol as sole C-source.
662

Locus Tag	Protein name	Fold change induction ($\log_2$)	$-\log_{10}(p\text{-value})$
PP_2426	CalA	6.28	4.03
PP_2679	PedH	4.75	3.80
PP_3426	MexF	3.93	3.24
PP_3425	MexE	3.74	2.18
PP_4921		3.65	2.46
PP_2440	AhpF	3.13	3.08
PP_3745	GlcD	3.12	3.32
PP_3747	GlcF	2.67	3.50
PP_3746	GlcE	2.64	2.98
PP_4922	ThiC	2.21	3.84
PP_3748	GlcG	2.06	3.71
PP_3622		1.97	2.82
PP_3178	GarK	1.77	2.85
PP_3621	IorA-II	1.60	2.34
PP_0554	AcoB	1.54	2.27
PP_3623	AdhB	1.54	3.67
PP_2484		1.52	2.37
PP_0734	HemK	1.51	2.67
PP_2439	AhpC	1.39	2.02
PP_0556		1.35	2.34
PP_1125		1.30	3.01
PP_0555	AcoA	1.20	2.52
PP_1548		1.19	2.03
PP_1351	PanE	-1.45	2.06
PP_2258		-1.84	2.40
PP_5658		-1.99	2.93
PP_3557		-2.41	2.33
PP_3603		-2.46	2.73
PP_4313		-2.55	2.50
PP_0588		-2.75	2.62
PP_2674	PedE	-4.25	3.78
PP_2673		-5.37	3.71
PP_3732		-5.78	3.07

664 **Table S1:** Strains and plasmids used in the study

Strains	Relevant features	Source or reference
KT2440*	KT2440 with a markerless deletion of *upp* | Parent strain for deletion mutants	(46)
Δ*pedE*	KT2440* with a markerless deletion of *pedE*	(11)
Δ*pedH*	KT2440* with a markerless deletion of *pedH*	(11)
Δ*calA*	KT2440* with a markerless deletion of *calA*	this study
Δ*garK*	KT2440* with a markerless deletion of gene *garK*	this study
Δ*glcDEF*	KT2440* with a markerless deletion of gene cluster *glcDEF*	this study
Δ*pedE* Δ*pedH*	KT2440* with a markerless deletion of *pedE and pedH*	(11)
Δ*pedE* Δ*pedH* Δ*glpFKRD*	Δ*pedE*Δ*pedH* with a markerless deletion of gene cluster Δ*glpFKRD*	this study
E. coli TOP10	F- mcrA Δ(mrr-hsdRMS-mcrBC) φ80lacZΔM15 ΔlacX74 nupG recA1 araD139 Δ(ara-leu)7697 galE15 galK16 rpsL(Str^R) endA1 λ$^-$	Invitrogen
Plasmids		
pJOE6261.2	Suicide vector for gene deletions	(46)
pJOE-calA	pJOE6261.2 based deletion vector for gene *calA* (PP_2426)	this study
pJOE-garK	pJOE6261.2 based deletion vector for gene *garK* (PP_3178)	this study
pJOE-glp	pJOE6261.2 based deletion vector for gene cluster *glpFKRD* (PP_1076 to PP_1073)	this study
pMW08	pJOE6261.2 based deletion vector for gene cluster *glcDEF* (PP_3745 to PP_3747)	this study

667 **Table S2:** Primers used in the study

Primer		
Name	Sequence 5' → 3'	Annealing temperature
PcalA1	CGATGGCCGCTTTGGTCCCGAGCCGTTCCACACTTTCGC	67°C
PcalA2	AACCGATCAAGCAGGGCCTCGCAGTGAA	67°C
PcalA3	GAGGCCCTGCTTGATCGGTTGCCTGTACG	65°C
PcalA4	CCTGCAGGTCGACTCTAGAGCGTGGGTGAGCAAGGCAG	65°C
PgarK1	CGATGGCCGCTTTGGTCCCGAGTCGTTGCTGTGGTGCC	62°C
PgarK2	TACAGGGTGTGGGGTTTCTCCTGTCCTG	62°C
PgarK3	GAGAAACCCCACACCCTGTAGAAATGGCCTTATTG	67°C
PgarK4	CCTGCAGGTCGACTCTAGAGCAGCGGCAAACTGACCATC	67°C
Pglp1	CGATGGCCGCTTTGGTCCCGGTCATCAATAAAGGTCCG	55°C
Pglp2	ACCGTAGGTAGTATGACCTCGTTTTTTTTG	55°C
Pglp3	GAGGTCATACTACCTACGGTGAAGCCCTC	62°C
Pglp4	CCTGCAGGTCGACTCTAGAGGTTGTGAAGACCGCCTGC	62°C
MWH03	AGGCACGATGGCCGCTTTGGTCCCGGCCTGCTCGGGCAGTTGTTCC	63°C
MWH04	GCTAAGCATGGGCCATCGGCTCACTCGCAAC	63°C
MWH05	AGTGAGCCGATGGCCCATGCTTAGCAAGTTCGTTATCG	63°C
MWH06	GCATGCCTGCAGGTCGACTCTAGAGCCAGGGCAATGCGTATCAC	63°C

Table S3: List of upregulated proteins in presence of 10 µM La^{3+} compared to the absence of La^{3+}
when grown with 2-phenylethanol as sole C-source.

Locus Tag	Protein name	Fold change induction (log$_2$)	- log$_{10}$(p-value)
PP_2679	PedH	3.41	2.48
PP_3357	Vdh	2.65	2.92
PP_5125	MutM	2.24	3.15
PP_4905	MotA	1.84	2.01
PP_0091		1.71	2.26
PP_5157		1.59	2.20
PP_0342	WaaC	1.43	2.46
PP_3722	Alr	1.04	2.40
PP_2674	PedE	-1.49	5.90
PP_2673		-2.41	3.91

674   **Table S4:** List of upregulated proteins in presence of 10 μM La^{3+} compared to the absence of La^{3+}
675   when grown with glucose as sole C-source.
676

Locus Tag	Protein name	Fold change induction (log$_2$)	- log$_{10}$(p-value)
PP_2679	PedH	4.35	3.17
PP_4508		1.74	2.97
PP_3713	CatA	1.44	2.43
PP_5221		1.39	2.21
PP_0367		1.27	2.47
PP_4796	HolA	-1.30	2.15
PP_4374	FliT	-1.95	2.45
PP_2666		-2.01	2.51
PP_3951	PcaI	-2.13	2.68
PP_2680	AldB-II	-2.79	2.41
PP_4632	FolM	-3.54	2.53
PP_2662		-3.61	3.62
PP_1757	BolA	-3.64	3.99
PP_2674	PedE	-6.97	4.13

678 **Table S5:** List of upregulated proteins in presence of 10 µM La^{3+} compared to the absence of La^{3+}
679 when grown with citrate as sole C-source.
680

Locus Tag	Protein name	Fold change induction (log$_2$)	- log$_{10}$(p-value)
PP_2491		2.27	4.32
PP_0365	BioC	1.12	2.06
PP_4157	KdpE	-1.41	2.72
PP_2674	PedE	-2.56	2.80
PP_0634	PilA	-3.08	3.88

Figures

Figure 1: Venn-diagram (middle) of differentially produced proteins in response to 10 µM La^{3+} during
growth with glycerol, glucose, 2-phenylethanol and citrate. Proteins that showed up under several
growth conditions are stated under the diagram with colour code for classification (yellow dot =
glycerol; green dot = glucose; blue dot = 2-phenylethanol; red dot = citrate). Classifications of
differentially expressed proteins according to the Cluster of Orthologous Groups database are
depicted for each substrate.

692
693
694 **Figure 2:** Growth of strains (**A**) KT2440* and (**B**) Δ*pedE* Δ*pedH* in M9 minimal medium supplemented
695 with 20 mM glycerol in the absence (blue dots) or presence of 10 μM La^{3+} (orange dots) in 96-well
696 microtiter plates at 30°C and 250 rpm. Data represent average of biological triplicates with
697 corresponding standard deviation.
698

699
700

Figure 3: Proteins of the upstream central carbon metabolism of *P. putida* KT2440 including the
proposed glycerol degradation pathway via glycerate and the additional hypothetical route through
phosphorylation of glyceraldehyde (dotted line). Anticipated metabolic flux by the proteins that were
identified as differentially abundant in response to 10 µM La^{3+} during growth on glycerol are colour-
coded (green = increased, red = decreased, black = not affected). The figure is inspired by a scheme
originally published by Nikel *et al.* (58) and was adapted to include the novel metabolic route(s)
identified in this study.
708

709
710
Figure 4: (**A**) Growth of strains Δ*pedE* Δ*pedH* (blue dots), Δ*glpFKRD* (orange dots), and Δ*pedE* Δ*pedH*
Δ*glpFKRD* (grey dots) in M9 minimal medium supplemented with 20 mM glycerol in 96-well
microtiter plates. (**B**) Growth of strains Δ*pedE* Δ*pedH* Δ*glpFKRD* (grey dots) and Δ*garK* (yellow dots),
in M9 minimal medium supplemented with 20 mM DL-glycerate incubated in 96-well microtiter
plates at 28°C and 220 rpm. Data represent average of biological triplicates with corresponding
standard deviation.
717

718
719
720 **Figure 5:** Growth of strains (**A**) Δ*garK,* (**B**) Δ*calA*, and (**C**) Δ*glcDEF* in M9 minimal medium
721 supplemented with 20 mM glycerol in absence (blue dots) or presence 10 µM La^{3+} (orange dots)
722 incubated in 96-well microtiter plates at 30°C and 250 rpm. Data represent average of biological
723 triplicates with corresponding standard deviation.
724

725
726 **Figure S1**: **PCA comparing the four different carbon sources with and without La^{3+}.** Different carbon
727 sources are indicated by squares (2-phenylethanol), circles (citrate), diamonds (glucose) and stars
728 (glycerol). Samples with 10 µM La^{3+} or without La^{3+} in the medium are shown in blue and red,
729 respectively. Biological replicates are indicated in the same colour. Samples can be separated
730 according to different carbon sources while treatment with La^{3+} only showed minor effects.