Clustering co-abundant genes identifies components of the gut microbiome that are reproducibly associated with colorectal cancer and inflammatory bowel disease

Samuel S. Minot, Ph.D.*
Microbiome Research Initiative
Fred Hutchinson Cancer Research Center
Seattle, Washington, USA
sminot@fredhutch.org
ORCID: 0000-0003-1639-3905

Amy D. Willis, Ph.D.
Department of Biostatistics
University of Washington
Seattle, Washington, USA
ORCID: 0000-0002-2802-4317

* Corresponding author

Abstract

Background: Whole-genome "shotgun" (WGS) sequencing is an increasingly widely used tool for analyzing the metagenomic content of microbiome samples. While WGS data contains gene-level information, it can be challenging to analyze the millions of microbial genes which are typically found in microbiome experiments. To mitigate the ultrahigh dimensionality challenge of gene-level metagenomics, it has been proposed to cluster genes by co-abundance to form Co-Abundant Gene groups (CAGs). However, exhaustive co-abundance clustering of millions of microbial genes across thousands of biological samples has previously been intractable purely due to the computational challenge of performing trillions of pairwise comparisons.

Results: Here we present a novel computational approach to the analysis of WGS datasets in which microbial gene groups are the fundamental unit of analysis. We use the Approximate Nearest Neighbor heuristic for near-exhaustive average linkage clustering to group millions of genes by co-abundance. This results in thousands of high-quality CAGs representing complete and partial microbial genomes. We applied this method to publicly available WGS microbiome surveys and found that the resulting microbial CAGs associated with inflammatory bowel disease (IBD) and colorectal cancer (CRC) were highly reproducible and could be validated independently using multiple independent cohorts.

Conclusions: This powerful approach to gene-level metagenomics provides a powerful path forward for identifying the biological links between the microbiome and human health. By proposing a new computational approach for handling high dimensional metagenomics data, we identified specific microbial gene groups that are associated with disease that can be used to identify strains of interest for further preclinical and mechanistic experimentation.

Background

Metagenomic analysis of the microbiome typically falls into the categories of taxonomic classification, metabolic pathway reconstruction, or genome reconstruction. While each has been used to good effect, each also has its own limitations. Taxonomic analysis is constrained by the size and quality of reference databases, which have started to provide decreasing taxonomic precision as the number of sequenced genomes grows [1]. Metabolic analysis is limited by our ability to annotate biochemical function from primary sequence, with only a minority of genes receiving any sort of annotation. Genome reconstruction (or "genome-resolved metagenomics") has made immense contributions to our understanding of microbial diversity and evolution, but is challenging to exhaustively characterize environments like the human gut, which contain hundreds or thousands of strains. In contrast, we took the approach of quantifying each individual gene *de novo* from a given metagenome. While this approach presented considerable computational challenges, it is unconstrained by the limitations of reference databases or annotation systems and therefore presents the possibility of discovering novel biological patterns in the human microbiome.

The approach of gene-level metagenomics is not new to this study and has been proposed previously as an alternative to taxonomic or metabolic pathway analysis [2]. Indeed even the popular HUMAnN2 tool [3] includes gene-family abundance estimation using the UniRef database of proteins [4]. We took the previously-described approach of grouping together genes that are consistently found at a similar level of abundance across multiple samples [2]. Such co-abundant genes are likely to be found on the same chromosome or piece of DNA across multiple samples, such as in the core genome for a bacterial species or consortium, on a plasmid that may move between strains, or as part of an operon in the accessory genome of a species that is only found

in a subset of strains. Biologically speaking, co-abundant genes are not independent entities, and can be grouped together for purposes of inferring their relationship with human health and disease. In addition, grouping genes by co-abundance finds low-dimensional structure in high-dimensional gene-level data, mitigating challenges with the statistical analysis of high-dimensional metagenomics data.

Results and Discussion

The primary analytical challenge that we encountered in this project was that of efficiently clustering microbial genes based on co-abundance. This general approach has been proposed and implemented previously [2, 5], but existing implementations do not perform exhaustive searches for co-abundant genes because performing all pairwise comparisons of millions of genes in large microbiome datasets [6] is computationally intractable. To overcome this obstacle we took advantage of the Approximate Nearest Neighbor (ANN) heuristic, which is able to robustly identify candidate subsets of co-abundant genes without having to perform all pairwise comparisons [7, 8]. We implemented a Python package to perform exhaustive average linkage clustering using the cosine distance metric on any dataset containing gene abundance data across a set of samples. While this method is relatively computationally intensive, we were able to execute it in a reasonable amount of time using commodity "cloud" computational resources. In the ideal case this approach improves the precision of estimating gene-level abundance by combining data from multiple correlated observations, as well as reducing the number of hypotheses to test in an association study, while maintaining the interpretation advantages of distinct genetic elements (core genome, plasmid, virus, etc.).

We applied this novel approach to gene-level metagenomics to test for an association of the gut microbiome across two distinct human diseases: IBD and CRC. We selected these diseases because each has been studied by multiple groups who have collected stool samples and performed metagenomic WGS sequencing (Table S1) [9–15]. Because each of these previous studies used slightly different protocols for selecting patients, collecting samples, and performing sequencing, an integrated analysis of these datasets should serve to identify those signals in the microbiome which are most robust to the methodological and experimental confounders.

The CAGs identified in this project contained 2-23,856 genes, with the majority of genes found in CAGs ranging between 10 and 2,000 genes in size and containing the range of metabolic functions expected from complete and partial microbial genomes (Fig. S1). Visual inspection of the genes making up these CAGs also demonstrated the highly consistent patterns of abundance displayed by the genes which were ultimately grouped into these CAGs (Fig. 1). We also analyzed a published single-cell sequencing dataset from the stool microbiome [16] and found that genes from the same CAG were found in the same physical cell at 3-9X the rate expected by chance (Fig. S2). The size, functional content, and clear pattern of co-abundance displayed by the genes in this analysis suggest that the CAGs used for statistical analysis represent biological units that are meaningful reflections of the composition of the microbiome across multiple independent datasets.

144
Figure 1. *Patterns of gene-level co-abundance across all microbiome samples from a subset of*
CAGs. Each row represents a single microbial gene, each column represents a single biological
sample, and pixel color reflects the gene's relative abundance (sequencing depth) in the sample.
A subset of CAGs and genes was randomly selected for display from the CRC datasets (A) and
the IBD datasets (B). Unsupervised hierarchical clustering was used to group the rows and
columns, and the left-hand color bar indicates the CAG assignment for each gene.

151
Our approach to the bioinformatic and statistical analysis was to select a single study for each
disease as the "discovery" cohort, and to use that dataset to build a *de novo* catalog of microbial
genes and identify CAGs. Our statistical model was relatively straightforward and used random
effects modeling to estimate the difference in the centered-log ratio of the relative abundance of
each CAG in the samples from people with and without the disease state (accounting for multiple
sampling of some individuals with random effects models). We chose to group together all
participants with any form of the disease state, as the criteria for disease classification was not
consistent across studies. In this discovery-validation approach, those CAGs which had a q-value
of < 0.2 in the discovery cohort were subsequently tested in an additional "validation" cohort, and
those CAGs which also had a q-value < 0.2 in that second step and the same direction of effect
were considered to be associated with disease.

163
We found with this approach that the estimated coefficient of disease status in the set of CAGs
associated with disease in the discovery cohort was significantly associated with the estimated
coefficient in the validation cohort (Fig. 2A-B; CRC r=0.38 p=0; IBD r=0.32 p=0). We further
demonstrated the extent of this association by displaying the abundance of the most strongly
associated CAGs across a total of 3 (IBD) or 4 (CRC) cohorts (Fig 2E-F), suggesting that this
association is not limited to the cohorts selected for discovery and validation. When performing the
same analysis with unclustered gene-level abundances (a single gene randomly selected from
each CAG), we found a much lower correlation between the estimated coefficient of disease
status (Fig 2C-D), supporting the proposed utility of CAGs for detecting reproducible biological
associations of the microbiome with host disease. Over and above the claim that the microbiome
is associated with disease in both cohorts, we believe that these results indicate that a substantial

number of elements of the microbiome which are associated with disease in the discovery cohorts
are also associated with disease in the respective validation cohorts.
177

178
Figure 2. *Reproducible association of CAG abundance with disease status for CRC (A, E) and*
IBD (B, F). The estimated coefficient plotted in A-D represents the log10 change in relative
abundance associated with health (positive values) or disease (negative values), for each disease
state. The estimated coefficient for the discovery dataset is on the horizontal axis, and the
estimated coefficient for the validation dataset is on the vertical axis. The results from CAG-based

analysis are shown in A-B, while the results calculated from unclustered gene-level abundances
are shown in C-D. The abundance of four representative CAGs are shown in E-F across all
available datasets, with colors indicating the health status associated with each sample.
187
The pattern of association for the IBD datasets was dominated by the vast majority of CAGs which
had a positive coefficient, indicating that they were more abundant in participants without IBD (Fig
2B). We therefore investigated the gene-level richness, finding a lower level of gene richness
observed in IBD samples compared to healthy controls (Fig S3) [17], corroborating previous
observations of lower alpha diversity in IBD [18, 19]. Without our use of the centered log-ratio to
adjust for the compositional nature of these datasets the decreased abundance of a large fraction
of the microbiome may have resulted in a spurious finding that the remainder had increased in
abundance [20], but in fact we found that very few CAGs were increased in abundance in IBD
relative to the geometric mean of each sample. However, there was a subset of CAGs which were
consistently found to be more abundant in IBD, which may represent those bacteria which are
able to thrive in the environment of the inflamed gut. Indeed, the taxonomic annotation of the
genes in these CAGs is enriched for organisms which have been implicated in some previous
studies of IBD and gut pathogens, including Enterobacteriaceae such as Escherichia, Shigella,
and Salmonella [13, 18, 19]. There was also a set of KEGG annotations that are weakly but
consistently enriched in this set of IBD-associated genes related to colonization and pathogenesis,
such as fimA (K07345), fimD (K07347), iron transport (K02010), and putrescine transport
(K02052; K11072; K11076).
205
The pattern of association for the CRC datasets was generally balanced between CAGs that were
more or less abundant in healthy participants (Fig 2A). Of the largest CAGs that were reproducibly
associated with disease, those which were more abundant in healthy participants tended to be
classified as Clostridia (via alignment to NCBI RefSeq), while those which were more abundant in
participants with CRC were more taxonomically diverse (Fig 3A-B). Moreover, we found the
functional annotations of the genes in those CAGs to be particularly interesting. There were four
KEGG annotations that were significantly enriched in the set of CAGs found to be more abundant
in CRC samples (Fisher's exact test, Holm-Sidak alpha=0.01): 1) grdA (K10670) is involved in
metabolism of glycine/sarcosine/betaine, and higher levels of glycine is a recognized hallmark of
cancer cells [21, 22]; 2) oxyR (K04761) is a transcriptional regulator which regulates genes
protecting from the biochemical damage induced by reactive oxygen species, of which markedly
higher levels are associated with progressive tumors [23, 24]; 3) abgT (K12942) is a transporter
responsible for uptake of p-aminobenzoyl-glutamate, and may also import other dipeptides [25];
and 4) afuA/fbpA (K02012) are transporters responsible for importing iron [26], which is likely to be
more abundant in the gastrointestinal lumen of individuals with CRC due to bleeding. Three of
these four annotated functions have clear links to the altered environment of the gut microbiome
expected during CRC, and likely promote the growth of these organisms in that setting. It remains
to be seen whether those organisms which are able to thrive in the CRC gut microbiome also
contribute to progression of disease.
225

Figure 3. Association of individual microbes with CRC (A & C) and IBD (B & D). A & B show the estimated coefficient of abundance for individual CAGs with disease status (log10 mean and 90% confidence intervals, left panel), the taxonomic assignment (middle panel) and functional assignment (right panel) of genes within each of those CAGs. C & D show the number of genes from disease-associated CAGs that are found within bacterial genomes from NCBI RefSeq, showing both the total number of genes for each genome, as well as a heatmap showing which disease-associated genes are found in which genomes.

One advantage of a gene-based approach to metagenomic analysis is that any CAG of interest can be directly compared with the genomes of bacterial isolates in order to identify strains containing each gene. Of the set of genes that we identified as consistently associated with CRC

238 and IBD, we found a number of strains containing large fractions of these genes (Fig 3C-D). We
239 furthermore propose that this approach of aligning disease-associated genes to whole microbial
240 genomes may be used to identify the members of any culture collection which are likely to have
241 the largest effect in an experimental model of these human diseases.
242
243
244 **Conclusions**
245
By proposing an approach to the analysis of metagenomic data that produces consistent results
across multiple heterogeneous datasets, we are addressing one of the most important challenges
in metagenomics, namely, reproducibility. Our findings suggest that indeed co-abundant gene
groups are a reproducible and biologically meaningful unit of analysis. In addition, microbial genes
are a meaningful and useful unit of analysis because they can be linked to individual microbial
genomes, taxonomic annotations, and predicted metabolic functionality. Using this approach, we
identify a list of gene groups that are associated with human diseases in multiple cohorts, and we
identify specific microbial isolates that contain these genes. The development of diagnostics or
therapeutics based on this list of genes and genomes is left to future work.
255
256
257 **Methods**
258
259 Datasets
260

Group	Used For	Name	NCBI BioProject
IBD	Discovery	Schirmer, et al. [15]	PRJNA389280
IBD	Validation	Lewis, et al. [13]	SRP057027
IBD	Validation	Hall, et al. [12]	PRJNA385949
CRC	Discovery	Zeller, et al. [9]	PRJEB6070
CRC	Validation	Feng, et al. [14]	PRJEB7774
CRC	Validation	Yu, et al. [11]	PRJEB10878
CRC	Validation	Vogtmann, et al. [10]	PRJEB12449

261 **Table S1**. Published datasets analyzed in this study.
262
263 Gene-level metagenomic analysis pipeline
264
265 All microbiome WGS data was analyzed using a Docker-based workflow, with each individual step
266 executed inside a Docker image. The workflow outlined below was executed independently for the
267 set of samples from Schirmer, et al., as well as for the set of samples from Zeller, et al.
268
269 The sequence of analyses is as follows:
270
271   1. Each sample was individually downloaded from NCBI SRA with Entrez Direct
272   • Docker image: quay.io/fhcrc-microbiome/get_sra:v0.4

- Code: https://github.com/FredHutch/docker-sra
- Wrapper script: get_sra.py
- Software version(s):
  - sratoolkit.2.8.2-ubuntu64
  - CMake3.11
  - fastq-pair 4ae91b0d9074410753d376e5adfb2ddd090f7d85

2. Each sample was individually assembled with metaSPAdes
   - Docker image: quay.io/fhcrc-microbiome/metaspades:v3.11.1--10
   - Code: https://github.com/FredHutch/docker-metaspades
   - Wrapper script: run_metaspades.py
   - Software version(s): SPAdes-3.11.1-Linux

3. Each sample's metagenomic assembly was annotated using Prokka
   - Docker image: quay.io/fhcrc-microbiome/metaspades:v3.11.1--8
   - Code: https://github.com/FredHutch/docker-metaspades
   - Software version(s): Prokka v1.12; barrnap v0.9
   - Wrapper script: run_prokka.py

4. The protein-coding sequences from all of the metagenomic assemblies for a given dataset were clustered at 90% amino acid identity using mmSeqs2 to create a set of non-redundant protein sequences
   - Docker image: quay.io/fhcrc-microbiome/integrate-metagenomic-assemblies:v0.4
   - Code: https://github.com/FredHutch/integrate-metagenomic-assemblies
   - Software version(s): biopython==1.70; MMseqs2 v2-23394
   - Wrapper script: integrate_assemblies.py

5. Each sample was aligned against the non-redundant protein sequences using DIAMOND, with post-alignment filtering using FAMLI. The Docker image associated with this step includes both the DIAMOND aligner and the FAMLI filtering code.
   - Docker image: quay.io/fhcrc-microbiome/famli:v1.1
   - Code: https://github.com/FredHutch/famli
   - Software version(s): DIAMOND v0.9.10; famli==1.0
   - Wrapper script: famli
   - Parameters:
     - min_qual = 30
     - min_score = 20
     - query_gencode = 11

6. The non-redundant protein sequences were functionally annotated via eggNOG-mapper
   - Docker image: quay.io/fhcrc-microbiome/eggnog-mapper:v0.1
   - Code: https://github.com/FredHutch/docker-eggnog-mapper
   - Software version(s): eggNOG-mapper = 1.0.3--py27_0
   - Wrapper script: run_eggnog_mapper.py

7. The non-redundant protein sequences were analyzed via the taxonomic assignment functionality of DIAMOND (using NCBI's RefSeq as the reference database)
   - Docker image: quay.io/fhcrc-microbiome/famli:v1.3

321   • <u>Code</u>: https://github.com/FredHutch/famli
322   • <u>Software version(s)</u>: DIAMOND v0.9.22
323   • <u>Wrapper script</u>: diamond-tax.py
324   • <u>Parameters</u>: top_pct = 1
325
8. The non-redundant protein sequences were grouped into CAGs based on their abundance
profile across the dataset.
• <u>Docker image</u>: quay.io/fhcrc-microbiome/find-cags:v0.11.1
• <u>Code</u>: https://github.com/FredHutch/find-cags
• <u>Software version(s)</u>: nmslib = 1.7.3.5
• <u>Wrapper script</u>: find-cags.py
• <u>Parameters</u>:
• min_samples = 10
• max_dist = 0.3
• normalization = sum
336
337   9. Group the outputs of all previous steps into a single HDF file
338   • <u>Docker image</u>: quay.io/fhcrc-microbiome/experiment-collection:latest
339   • <u>Code</u>: https://github.com/FredHutch/minot-experiment-collection
340   • <u>Wrapper script</u>: make-experiment-collection.py
341
342 The validation datasets were analyzed by aligning the raw WGS reads against the non-redundant
343 protein sequences generated from the relevant discovery dataset as described in Step 5
344 described above. The final HDF file creation step (9) includes the results of that quantification step
345 for the validation datasets as well as the discovery datasets.
346
Given the difficulty of providing a workflow execution system that can be used effectively by a
broad range of users, we have elected to provide all of the individual tools needed to run a
complete analytical workflow, with public Docker images making up each individual step, instead
of providing a complete workflow system that each user would need to customize for their own
execution engine (Slurm, PBS, Kubernetes, AWS, GCP, Azure, etc.). This approach enables
execution of the exact code that we used in this analysis in a platform-independent manner using
the highest standard of reproducibility (Docker containers).
354
Our implementation of the analytical workflow described above relied upon the Amazon Web
Service and its Batch API, which allows users to submit individual jobs for analysis using utilities
from the boto3 library in Python. While this implementation does not represent a complete
workflow management system, the code used for this execution is available at
https://github.com/FredHutch/aws-batch-helpers/ in the batch_helpers/batch_task_manager.py
module.
361
362
363 <u>Grouping genes by co-abundance</u>
364
We did not find any public tools for grouping genes by co-abundance that were appropriate to the
scale of our datasets. To implement our own approach for finding CAGs, we utilized the Non-
Metric Space Library (`nmslib`, https://pypi.org/project/nmslib/) which implements the Approximate
Nearest Neighbor (ANN) algorithm [7, 8] and obviates the need for calculating the all-by-all
distance matrix typically used by clustering algorithms. The abundance matrix used for clustering

was created by calculating the depth of sequencing for each individual gene within each sample
and normalizing for total sequencing depth. The distance metric used to quantify the dissimilarity
of individual genes was the cosine distance. Gene clusters were identified iteratively by average
linkage clustering and a fixed cophenetic distance threshold. The ANN algorithm was used to
identify subsets of genes which were likely to be highly co-abundant, and which could be clustered
independently of the whole. The code executed for this analysis, as well as a Docker image
containing all required dependencies, can be found in the summary of the complete analysis
workflow (Step 8).
378
379
380 Correlating CAGs with health status
381
CAG discovery: For every CAG in the validation dataset, we tested the null hypothesis that the
mean difference in CLR abundance between patients with and without disease was zero using the
general linear model framework. Datasets with repeated measurements on subjects were
modelled using a linear mixed effects model with subject as a random effect. Using the `qvalue` R
package (v2.8.0), we calculate the q-values for each CAG. Our set of "discovered CAGs" for
validation is the set of CAGs with calculated q-value of 0.2 or less. These are the CAGs that would
be considered statistically significant while controlling the FDR at 20%.
CAG validation: *For only the discovered CAGs*, we tested the null hypothesis that the mean
difference in CLR abundance between patients with and without disease was zero in the validation
datasets. Our "validated CAGs" are the CAGs in this set with calculated q-values of 0.2 or less,
and that have an estimated difference in abundance between disease status groups of the same
sign as the estimated difference in the discovery dataset.
The probability of validating discovered CAGs: To calculate the probability of validating C_2 or
more out of C_1 CAGs under the global null hypothesis of no association between disease status
and any CAG's abundance, we bounded the p-value for validating discovered CAGs in the
following way. Let X be the number of CAGs with q-values less than 0.2 for the validation data and
with an estimated difference in CLR abundance across disease groups of the same sign in the
validation and discovery datasets, and Y be the number of CAGs with an estimated difference in
CLR abundance across disease groups of the same sign in the validation and discovery datasets.
Since under the null the test statistics are approximately Normal(0,1)-distributed,

$$Pr_{H_0}(X \geq C_2) \leq Pr_{H_0}(Y \geq C_2) = Pr(Binomial(C_1, 0.5) \geq C_2) \approx Pr\left(Normal(0,1) \geq \frac{C_2 - C_1/2}{\sqrt{C_1}/2}\right)$$

403 giving us a conservative p-value for the global null of no association.
404
405
406 Aligning protein-coding genes against RefSeq genomes
407
The alignment of individual protein-coding genes against the RefSeq collection of genomes in
NCBI was executed using the Docker image hosted at quay.io/fhcrc-microbiome/docker-
diamond:v0.9.23—0 and built using the Dockerfile hosted at https://github.com/FredHutch/docker-
diamond, running DIAMOND v0.9.23. The complete list of Prokaryotic RefSeq genomes was
downloaded from https://www.ncbi.nlm.nih.gov/genome/browse#!/prokaryotes/ and the query
proteins were aligned via DIAMOND against the annotated protein-coding sequences from each
genome individually. We implemented this analysis on the Amazon Web Service using the Batch
API for execution and resource management.
416
417
418 Quantification of co-abundant genes in uncultured single cells

Datasets from published single-cell sequencing microbiome experiments [16] were downloaded and split by 10X barcode (each corresponding to a single cell). The WGS data for each single cell was aligned against each reference gene catalog (for the CRC and IBD datasets) and filtered with FAMLI as described in workflow step 5, above. The result of this analysis was a count of the number of genes that were found in the same cell as another gene that is also part of the same CAG. As a comparison, we calculated the number of such genes that would be found with a randomly permuted set of CAG assignments.

Declarations

Ethics approval and consent to participate
Not applicable

Consent for publication
Not applicable

Availability of data and material
The data produced in this analysis is available on the Synapse platform at
https://www.synapse.org/#!Synapse:syn15623121 (doi:10.7303/syn15623121). The Synapse project will be made fully public upon acceptance for publication, and is currently available to reviewers. The repository includes documentation describing the organization and formatting of relevant data files and includes all of the outputs from the bioinformatic pipeline used for gene-level metagenomic analysis, as well as the Jupyter notebooks used to analyze those datasets and produce the figures and tables presented here.

Competing interests
AW: None to declare.
SM holds financial interest in Reference Genomics, Inc. (One Codex) and consults for the American Type Culture Collection (ATCC).

Funding
Not applicable

Authors' contributions
SM developed the novel CAG identification method and performed all primary data analysis; AW developed the statistical analysis and discovery-validation framework; both authors contributed to figure generation and writing the manuscript.

Acknowledgements
The authors are grateful for support from the Fred Hutch Microbiome Research Initiative (SM) and start-up funds from the Department of Biostatistics at the University of Washington (AW).

References

1. Nasko DJ, Koren S, Phillippy AM, Treangen TJ. RefSeq database growth influences the accuracy of k-mer-based lowest common ancestor species identification. Genome Biol.

2018;19:165. doi:10.1186/s13059-018-1554-6.

2. Nielsen HB, Almeida M, Juncker AS, Rasmussen S, Li J, Sunagawa S, et al. Identification and
assembly of genomes and genetic elements in complex metagenomic samples without using
reference genomes. Nat Biotechnol. 2014;32:822–8. doi:10.1038/nbt.2939.

3. Franzosa EA, McIver LJ, Rahnavard G, Thompson LR, Schirmer M, Weingart G, et al. Species-
level functional profiling of metagenomes and metatranscriptomes. Nat Methods. 2018;15:962–8.
doi:10.1038/s41592-018-0176-y.

4. Suzek BE, Wang Y, Huang H, McGarvey PB, Wu CH. UniRef clusters: a comprehensive and
scalable alternative for improving sequence similarity searches. Bioinformatics. 2015;31:926–32.
doi:10.1093/bioinformatics/btu739.

5. Plaza Oñate F, Le Chatelier E, Almeida M, Cervino ACL, Gauthier F, Magoulès F, et al.
MSPminer: abundance-based reconstitution of microbial pan-genomes from shotgun
metagenomic data. Bioinformatics. 2018;:bty830. doi:10.1093/bioinformatics/bty830.

6. Qin J, Li R, Raes J, Arumugam M, Burgdorf KS, Manichanh C, et al. A human gut microbial
gene catalogue established by metagenomic sequencing. Nature. 2010;464:59–65.
doi:10.1038/nature08821.

7. Har-Peled S, Indyk P, Motwani R. Approximate Nearest Neighbor: Towards Removing the
Curse of Dimensionality. Theory Comput. 2012;8:321–50. doi:10.4086/toc.2012.v008a014.

8. Kushilevitz E, Ostrovsky R, Rabani Y. Efficient Search for Approximate Nearest Neighbor in
High Dimensional Spaces. SIAM J Comput. 2000;30:457–74. doi:10.1137/S0097539798347177.

9. Zeller G, Tap J, Voigt AY, Sunagawa S, Kultima JR, Costea PI, et al. Potential of fecal
microbiota for early-stage detection of colorectal cancer. Mol Syst Biol. 2014;10:766–766.
doi:10.15252/msb.20145645.

10. Vogtmann E, Hua X, Zeller G, Sunagawa S, Voigt AY, Hercog R, et al. Colorectal cancer and
the human gut microbiome: Reproducibility with whole-genome shotgun sequencing. PLoS One.
2016;11:1–13. doi:10.1371/journal.pone.0155362.

11. Yu J, Feng Q, Wong SH, Zhang D, Liang Q yi, Qin Y, et al. Metagenomic analysis of faecal
microbiome as a tool towards targeted non-invasive biomarkers for colorectal cancer. Gut.
2015;66:70–8. doi:10.1136/gutjnl-2015-309800.

12. Hall AB, Yassour M, Sauk J, Garner A, Jiang X, Arthur T, et al. A novel Ruminococcus gnavus
clade enriched in inflammatory bowel disease patients. Genome Med. 2017;9:103.
doi:10.1186/s13073-017-0490-5.

13. Lewis JD, Chen EZ, Baldassano RN, Otley AR, Griffiths AM, Lee D, et al. Inflammation,
Antibiotics, and Diet as Environmental Stressors of the Gut Microbiome in Pediatric Crohn's
Disease. Cell Host Microbe. 2015;18:489–500. doi:10.1016/j.chom.2015.09.008.

14. Feng Q, Liang S, Jia H, Stadlmayr A, Tang L, Lan Z, et al. Gut microbiome development along
the colorectal adenoma–carcinoma sequence. Nat Commun. 2015;6:6528.
doi:10.1038/ncomms7528.

15. Schirmer M, Franzosa EA, Lloyd-Price J, McIver LJ, Schwager R, Poon TW, et al. Dynamics
of metatranscription in the inflammatory bowel disease gut microbiome. Nat Microbiol. 2018;1.
doi:10.1038/s41564-017-0089-z.

16. Bishara A, Moss EL, Kolmogorov M, Parada AE, Weng Z, Sidow A, et al. High-quality genome
sequences of uncultured microbes by assembly of read clouds. Nat Biotechnol. 2018.
doi:10.1038/nbt.4266.

17. Willis A, Bunge J. Estimating diversity via frequency ratios. Biometrics. 2015;71:1042–9.
doi:10.1111/biom.12332.

18. Gevers D, Kugathasan S, Denson LA, Vázquez-Baeza Y, Van Treuren W, Ren B, et al. The
Treatment-Naive Microbiome in New-Onset Crohn's Disease. Cell Host Microbe. 2014;15:382–92.
doi:10.1016/j.chom.2014.02.005.

19. Morgan XC, Tickle TL, Sokol H, Gevers D, Devaney KL, Ward D V, et al. Dysfunction of the

intestinal microbiome in inflammatory bowel disease and treatment. Genome Biol. 2012;13:R79.
doi:10.1186/gb-2012-13-9-r79.
20. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome datasets are
compositional: And this is not optional. Front Microbiol. 2017;8:2224.
doi:10.3389/fmicb.2017.02224.
21. Jain M, Nilsson R, Sharma S, Madhusudhan N, Kitami T, Souza AL, et al. Metabolite Profiling
Identifies a Key Role for Glycine in Rapid Cancer Cell Proliferation. Science (80-).
2012;336:1040–4. doi:10.1126/science.1218595.
22. Locasale JW. Serine, glycine and one-carbon units: cancer metabolism in full circle. Nat Rev
Cancer. 2013;13:572–83. doi:10.1038/nrc3557.
23. Kumari S, Badana AK, G MM, G S, Malla R. Reactive Oxygen Species: A Key Constituent in
Cancer Survival. Biomark Insights. 2018;13:1177271918755391.
doi:10.1177/1177271918755391.
24. Liou G-Y, Storz P. Reactive oxygen species in cancer. Free Radic Res. 2010;44:479–96.
doi:10.3109/10715761003667554.
25. Green JM, Hollandsworth R, Pitstick L, Carter EL. Purification and Characterization of the
Folate Catabolic Enzyme p-Aminobenzoyl-Glutamate Hydrolase from Escherichia coli. J Bacteriol.
2010;192:2407–13. doi:10.1128/JB.01362-09.
26. Willemsen PT, Vulto I, Boxem M, de Graaff J. Characterization of a periplasmic protein
involved in iron utilization of Actinobacillus actinomycetemcomitans. J Bacteriol. 1997;179:4949–
52.

539
540
541

Supplementary Figures

Supplementary Figure S1*. The distribution of CAG size (genes per CAG; A & B) and the functional annotation of genes in CAGs is shown by CAG size (C & D). Each gene can be annotated with a range of biological functions, and the proportion of CAGs of a given size containing at least one functional annotation is shown (C & D). The CAGs generated from the CRC datasets are shown in A & C, while the CAGs generated from the IBD datasets are shown in B & D.*

Supplementary Figure S2*. Single-cell microbiome datasets were analyzed using the gene catalogs and CAG groupings from the CRC and IBD datasets. Co-occurrence was measured as the number of genes that were found in the same cell with another gene from the same CAG. Simulations were performed by random permutation, with 1,000 replicates. Orange bars show mean and standard deviation.*

Supplementary Figure S3. Alpha diversity by diagnosis across cohorts. The number of total genes in each sample was estimated with breakaway for both the CRC (A) and IBD (B) cohorts.

Supplementary Tables

Supplementary Table S2. Description of genes associated with CRC, including the CAG grouping, correlation coefficient, taxonomic annotation, and functional annotation. Public repository URL: *https://www.synapse.org/#!Synapse:syn17104367*

Supplementary Table S3. Description of genes associated with IBD, including the CAG grouping, correlation coefficient, taxonomic annotation, and functional annotation. Public repository URL: *https://www.synapse.org/#!Synapse:syn17104250*