Mapping the human subcortical auditory system using histology, post mortem MRI and in vivo MRI at 7T

Kevin Richard Sitek[1,2†], **Omer Faruk Gulban**[3†], **Evan Calabrese**[4§], **G. Allan Johnson**[4], **Satrajit S. Ghosh**[1,2‡], **Federico De Martino**[3,5‡]

[1]MIT; [2]Harvard University; [3]Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht, The Netherlands; [4]Duke University; [5]Center for Magnetic Resonance Research, University of Minnesota, Minneapolis, USA

*For correspondence:
orcid.org/0000-0001-7761-3727 (OFG);
orcid.org/0000-0002-2172-5786 (KRS)

†These authors contributed equally to this work
‡These authors also contributed equally to this work

Present address: §Department of Radiology, University of California, San Francisco, USA

Abstract Studying the human subcortical auditory system non-invasively is challenging due to its small structures deep within the brain. Additionally, the elaborate three-dimensional (3-D) structure of the system can be difficult to understand based on currently available 2-D schematics and animal models. To address these issues, we measured functional magnetic resonance imaging (fMRI) responses from the human subcortical auditory system at 7 Tesla and validated the results by creating an atlas based on state of the art human histology (BigBrain) and post mortem MRI. Furthermore, using diffusion MRI tractography, we revealed structural connectivity maps of the human subcortical auditory pathway both in vivo (1050 μm isotropic resolution) and post mortem (200 μm isotropic resolution). This work contributes novel tools for researching the human auditory system to understand its structural organization as well as facilitates dissemination of this knowledge by making the created atlases openly available.

Introduction

Understanding the structure of the human subcortical auditory pathway is a necessary step to research its role in hearing, speech communication and music. However, due to methodological issues in human research, most of our understanding of the subcortical (thalamic, midbrain and brainstem) auditory pathway arises from research conducted in animal models. This might be problematic because, while the organization of the auditory pathway is largely conserved across mammalian species (*Malmierca and Hackett, 2010*; *Schofield, 2010*), the form and function of each structure may not be analogous (*Moore, 1987*). In this paper we show the convergence of three human imaging modalities — histology, post mortem magnetic resonance imaging (MRI), and in vivo MRI at ultra high-field (7 Tesla) — in the representation of the subcortical auditory pathway at high spatial resolution (between 50 and 1100 μm).

Although MRI has become increasingly powerful at imaging deep brain structures, anatomical investigation of the human subcortical auditory pathway has been primarily conducted in post mortem tissue dissection and staining. *Moore* (*1987*) stained both myelin and the cell bodies of subcortical auditory structures in four post mortem human brainstem samples and compared them to the analogous structures in cats (a common model for auditory investigations at the time). Later investigations from the same group (*Moore et al., 1995*) used myelin and Nissl cell body staining to investigate the timeline of myelination in human auditory brainstem development. More recently, *Kulesza* (*2007*) stained six human brainstems for Nissl substance, focusing on the superior olivary

complex, finding evidence of a substructure (the medial nucleus of the trapezoid body) whose existence in the human auditory system has been debated for decades.

To study the subcortical auditory system in living humans, MRI is the best available tool due to its high spatial resolution. Anatomical in vivo MRI investigations of the human subcortical auditory pathway so far have focused on thalamic nuclei (*Devlin et al., 2006*; *Moerel et al., 2015*), and the identification of the acoustic radiations between the auditory cortex and medial geniculate nucleus of the thalamus with diffusion-weighted MRI tractography (*Devlin et al., 2006*; *Behrens et al., 2007*; *Javad et al., 2014*; *Maffei et al., 2018*). Due to their small size and deep locations, identification of more caudal subcortical structures — the superior olivary complex and cochlear nucleus — remain challenging with in vivo anatomical MRI.

Although lower spatial resolution than anatomical MRI, functional MRI (fMRI) has been used to investigate the relevance of subcortical processing of auditory information in humans, but has been limited by the small size of the structures involved and the relatively low resolution attainable at conventional field strengths (3 Tesla and below) (*Guimaraes et al., 1998*; *Harms and Melcher, 2002*; *Griffiths et al., 2001*; *Hawley et al., 2005*). These acquisitions required trade-offs, such as low through-plane resolution (7 mm) in exchange for moderate in-plane resolution (1.6 mm), and in some cases researchers synchronized image collection to the cardiac cycle in order to overcame the physiological noise associated with blood pulsation in the brainstem (*Guimaraes et al., 1998*; *Sigalovsky and Melcher, 2006*).

More recent advances in MRI, especially the increased signal to noise ratio (SNR) available at ultra-high magnetic fields (7 Tesla and above), have enabled higher resolution functional imaging of subcortical structures and more advanced localization of human auditory subcortical structures as well as their functional characterization. Using a MRI at 7 Tesla (7T), *De Martino et al.* (*2013*) and *Moerel et al.* (*2015*) collected relatively high resolution (1.1-1.5 mm isotropic) fMRI with an auditory paradigm to identify tonotopic gradients in the inferior colliculus and medial geniculate nucleus. In these studies, high isotropic resolution and SNR provided an opportunity to investigate auditory responses throughout the subcortical auditory system.

Despite the methodological advances in investigating the human subcortical auditory system, a systematic comparison of their capabilities has not yet been undertaken. Here we use publicly available histological data (*Amunts et al., 2013*) to segment the main nuclei along the subcortical auditory pathway. We evaluate the ability of identifying the same structures through post mortem anatomical MRI, through functional MRI using natural sounds, and through estimating the connectivity between subcortical auditory structures with post mortem and in vivo diffusion MRI tractography. To compare the histological, post mortem and in vivo data, we project all images to MNI common reference space (*Fonov et al., 2009*, *2011*). To facilitate dissemination of our results, we make the resulting atlases publicly available.

Results

Definition of a subcortical auditory atlas from histology

To obtain a spatially accurate reference for all the subcortical auditory structures, we manually segmented publicly available histological data (100 μm version of the BigBrain 3-D Volume Data Release 2015 in MNI space from https://bigbrain.loris.ca (*Amunts et al., 2013*)).

Upon inspecting this dataset, we noticed that the area around the inferior colliculus was incorrectly transformed into MNI space. This was causing the colliculi to be larger and more caudal than in the MNI reference brain (Fig 7, second and third panels). Thus, our first step was to correctly register the area around the colliculi (Fig 7, fourth panel) — see Methods for details on the correction procedure.

The results of our BigBrain subcortical auditory segmentation in corrected MNI space are reported in Fig 1 together with schematics redrawn from *Moore* (*1987*) (for the cochlear nucleus, superior olivary complex and inferior colliculus) and the Allen Human Brain Atlas (*Hawrylycz et al.,*

90 *2012*; *Ding et al., 2016*) (for the medial geniculate body). These schematics were used as reference
91 during the segmentation. The 3-D rendering of the segmented structures highlighting the complex
92 shape of the cochlear nucleus and superior olivary complex is also presented in Fig 1. The rendering
93 is presented from a posterior lateral view in order to compare it with the Gray's Anatomy, Plate 719
94 (*Gray and Lewis, 1918*).

Post mortem MRI

Post mortem MRI atlas of the human subcortical auditory system

Magnetic resonance histology — i.e., the study of tissue at microscopic resolution using MRI —
provides several unique advantages over conventional histology: 1) it is non-destructive; 2) it suffers
minimal distortion from physical sectioning and dehydration; 3) it yields unique contrast based on
water in the tissue and how it is bound (e.g., diffusion); and 4) produces 3-D data. These advantages
make it an ideal medium for visualizing the 3-D organization of the deep brain structures (*Johnson
et al., 1993*). To delineate the subcortical auditory structures with MR histology, we acquired 50 μm
isotropic voxel size 3-D gradient echo (GRE) MRI on a human post mortem brainstem and thalamus
(described previously in (*Calabrese et al., 2015*); see Methods for additional details). These data are
presented in Figure 2 (second column) after transformation to MNI space and resampling to 100
μm isotropic resolution (see Methods section for details). The post mortem MRI data are presented
together with the histological data for comparison (first column).

Based on our segmentations of the subcortical auditory structures in the post mortem MRI
data, the resulting 3-D model is presented in Fig 2. A volumetric quantification of the identified
structures (in the BigBrain and post mortem MRI) is reported in Table 3 and the overlap between
the segmentations computed after projection in MNI space are reported in Table 2 (as inset in Fig 2).

3-D connectivity map of the human subcortical auditory system from post mortem
diffusion MRI

Identifying the connectivity between subcortical auditory nuclei is crucial for understanding the
structure of the pathway. However, methods for tracing neuronal pathways that are available in
other animal models are generally not available in human studies, even post mortem. Diffusion-
weighted MRI (dMRI) can be used to measure the orientation and magnitude of molecular motion
and infer patterns of white matter in brain tissue (post mortem and in vivo). Using 200 μm diffusion-
weighted MRI data acquired on the same post mortem sample (see Methods for details), we
modeled diffusion orientations and estimated likely connectivity pathways (or streamlines) using
tractography. Constraining the streamlines to only those that pass through auditory structures
(as identified from the anatomical MRI data), we visualized the connectivity map of the subcortical
auditory pathway in Fig 4, left panel.

Connectivity closely resembles the expected pattern of the human subcortical auditory wiring.
In particular, streamlines predominantly pass through the lateral lemniscus, the primary subcortical
auditory tract. Additional streamlines run through the brachium of the inferior colliculus, connecting
the inferior colliculus with the medial geniculate of the thalamus. Many streamlines then course
rostrally toward the auditory cortex (not present in this specimen).

At the caudal extent of the lateral lemniscus, streamlines pass through the superior olivary
complex. Streamlines also run through the root of CNVIII. In total, each expected step along the
subcortical auditory pathway is represented in this connectivity map.

Fig 4 (top right panel) shows the percentage of total streamlines connecting each of the sub-
cortical auditory structures as estimated from this post mortem diffusion MRI sample. Overall,
connections between structures is sparse in this representation, with the vast majority of stream-
lines only passing through one region (shown along the diagonal). However, some between-region
connections appear in this analysis, particularly ipsilaterally between left inferior colliculus and the
other auditory structures.

Figure 1. Literature diagrams (left columns) redrawn from *Moore* (*1987*) for the cochlear nucleus (CN), superior olivary complex (SOC), inferior colliculus (IC) and from the Allen Human Brain Atlas (*Hawrylycz et al., 2012*) for the medial geniculate body (MGB) compared to similar cuts from histology (BigBrain) in MNI (central column) and 3-D reconstructions of the segmented structures from the histology (bottom right column). The auditory structures are highlighted in gray in the left column, by a dotted line in the central column and in red on the modified Gray's anatomy Plate 719 (*Gray and Lewis, 1918*) and rendered as solid red surface meshes within the surface point cloud render of BigBrain MNI brainstem (right column).

BigBrain (100 μm)
MNI ICBM 152 (corrected)

7T Ex-vivo (100 μm)
T2*w

Post-mortem MRI
3D reconstruction

Table 2. Segmentation similarity comparison between BigBrain, post-mortem and in-vivo auditory nuclei

		DICE Coeff.		Avg. Hausdorff Dist.	
		Left	**Right**	**Left**	**Right**
BigBrain across segmenters	**MGN**	0.72	0.75	1.43	1.34
	IC	0.83	0.77	0.48	0.85
	SOC	0.61	0.51	4.30	5.80
	CN	0.80	0.74	0.77	2.63
BigBrain vs post-mortem	**MGN**	0.5	0.5	3.3	5.1
	IC	0.3	0.4	6.4	5.8
	SOC	0.2	0.01	5.8	7.8
	CN	0.2	0.2	7.1	6.6
BigBrain vs in-vivo	**MGN**	0.4	0.5	3.9	4.7
	IC	0.4	0.3	6.6	7.3
	SOC	0.1	0.03	8.9	11.5
	CN	0.04	0.1	14.6	11.6

Figure 2. BigBrain - 7T post mortem MRI image comparisons. Histological data (BigBrain) (left column) and T2*-weighted post mortem MRI data (100 μm - central column) in MNI space. Panels from bottom to top are chosen to highlight subcortical auditory structures (CN [bottom] to MGB [top]). Arrows (white with red outline) indicate the location of the subcortical auditory nuclei. The 3-D structures resulting from the segmentation of the post mortem data is presented on the top right panel. Table 2 quantifies (using DICE coefficient and average Hausdorff distance) the agreement (in MNI space) for all subcortical structures between: a) segmentations performed on the BigBrain dataset by the two raters (KS and OFG) [top]; 2) segmentations obtained from the BigBrain dataset and from the post mortem MRI data [middle]; 3) segmentations obtained from the BigBrain dataset and from in vivo functional MRI data [bottom])

Table 1. Volume comparisons (mm3) across different segmentations of the auditory brainstem regions of interest.

	Literature	BigBrain MNI	Post-mortem	In-vivo
CN	46	32	11	76
SOC	7	6	4	124
IC	65	63	73	200
MGN	58	75	134	285

Figure 3. Comparisons between the volume of auditory subcortical structures reported in the literature (*Glendenning and Masterton, 1998*) and the volume obtained in our BigBrain segmentation (in MNI space), post mortem MRI data segmentation and in vivo functional clusters.

Figure 4. Post mortem diffusion MRI tractography. Left: streamlines passing through subcortical auditory structures, defined from 50 μm post mortem MRI in the same specimen, warped to 200μm isotropic diffusion image space. Top right: connectivity heatmap of subcortical auditory structures. Bottom right: Diffusion orientation distribution functions (ODFs) for each voxel; axial slice at the level of the rostral inferior colliculus (IC), including the commissure of the IC (bottom center arrow) and brachium of the IC (top left arrow).

To investigate the sparsity of streamlines connecting auditory regions at 200 μm using 50
μm anatomical MRI-defined regions, we conducted two additional analyses. First, we dilated the
anatomical ROIs by 500 μm (2.5 voxels at 200 μm resolution), thereby including nearby white matter
tracts as well as adjacent subcortical structures. Using these dilated ROIs as tractography waypoints,
we found higher connectivity between subcortical ROIs, particularly on the right between inferior
colliculus (IC) and medial geniculate body (MGB) (Supplementary Fig 1).
Next, we resampled the diffusion MRI images to an in vivo-like resolution (1.05 mm isotropic). We
again estimated fiber ODFs using CSD and estimated white matter connections with deterministic
tractography. Using the (undilated) anatomically defined ROIs as tractography waypoints, we can
visualize streamline estimates connecting subcortical auditory structures (Supp. Fig 2). Again,
we see greater ipsilateral connectivity estimates between structures, particularly between left
structures.

Vasculature representations from post mortem MRI

Because T2*-weighted GRE imaging is sensitive to blood vessels, we processed our anatomical MR
image to highlight brainstem vasculature (Fig 5, right column, base image). These 3-D vasculature
images bear striking resemblance to post mortem data acquired with a stereoscopic microscope
after full clearing method (see *Duvernoy* (*2013*) for detailed diagrams of human brainstem vascula-
ture). These vasculature images in the MNI space can be helpful to understand the nature of the in
vivo functional signals (see next section).

In vivo MRI

We next sought to identify the structures and connections of the human subcortical auditory
system in living participants. By leveraging the increase signal and contrast to noise available at
ultra-high magnetic fields (7 Tesla) (*Vaughan et al., 2001*; *Ugurbil et al., 2003*; *Ugurbil, 2016*), we
collected high resolution anatomical (0.7 mm isotropic), diffusion-weighted (1.05 mm isotropic; 198
diffusion gradient directions across 3 gradient strengths) and functional (1.1 mm isotropic) MRI in
ten participants (see Methods for details).

Anatomical MRI

Visual inspection and comparison to the MNI dataset (Supp. Fig 5) showed that the MGB and IC
could be identified on the basis of the anatomical contrast, especially in the short inversion time
T1-weighted data (*Tourdias et al., 2014*; *Moerel et al., 2015*). However, while the superior olivary
complex (SOC) could be identified in the MNI dataset (Supp. Fig 5), it could not be identified in
average anatomical image from our 7T data. This is possibly due to the limited number of subjects
leading to the lower signal to noise in the average image. We have also explored the combination
of image contrasts within each individual using the compositional method proposed in (*Gulban
et al., 2018b*), but the results were inconclusive.

Functional MRI

The difficulty in delineating the CN and SOC from anatomical in vivo MRI data oriented our inves-
tigation towards the possibility of identifying the subcortical auditory pathway — in vivo and in
single individuals — on the basis of the functional responses to sounds. Functional responses to
168 natural sounds were collected at 7T using a sparse acquisition scheme and a fast even related
design.
In all our participants, we could identify clusters of significant activation in response to sounds
in the MGB, IC, SOC, and CN. On top of this, we created a probabilistic functional atlas based on the
overlap of statistically significant maps across individuals (after registration to a reference subject).
The left column of Figure 5 shows the probabilistic functional map (i.e., representing the number of
subjects in which each voxel was identified as significantly responding to sounds) overlaid on the in
vivo average anatomical MRI image (short inversion time T1-weighted image (*Tourdias et al., 2014*);
see Methods for details).

Projecting these data to the reference MNI space allowed evaluating the correspondence between in vivo functionally defined regions and histological data (Big Brain - Figure 5, center column).

At the level of the CN, the functionally active clusters correspond mostly to the ventral part of CN. The dorsal subdivision of the CN is not recovered on the basis of functional responses probably due to partial voluming with the nearby cerebrospinal fluid in combination with thinness (thickness around 0.5mm) of the dorsal CN as it wraps around inferior cerebellar peduncle (see Fig 1). Nearby, the location of the activation clusters identifying the SOC overlaps with the SOC as identified in the BigBrain data.

As the next step, we qualitatively investigated if the orientation of the vasculature at the level of the SOC may have an effect on size (and location) of the functionally defined regions. As a visual aid in this evaluation, we overlaid the functionally defined regions with the vasculature image obtained from the post mortem data (Fig 5, right column). In all subcortical regions the vasculature appears to have a specific orientation, and, at the level of the SOC, vessels drain blood from the center in a ventral direction (i.e., the direction of draining is towards the surface of the brainstem in the top of the image reported in the transverse view, bottom in Fig 5). This specific vasculature architecture may result in the displacement or enlargement of the functionally defined clusters towards the ventral surface of the brainstem (as highlighted in the correspondence with histological data in Fig 5).

The probability of corresponding voxels across subjects to be significantly modulated by sound presentation increased at the level of the IC and MGB, where the histologically defined regions corresponded (for the large part) to all subjects exhibiting significant responses to sounds. At the threshold of three subjects in the probabilistic maps, the IC seems to extend towards the superior direction, bordering and sometimes including parts of superior colliculus. On the other hand, similarly to what may happen in the SOC, the general directions of the vasculature penetrating the IC and draining blood towards the dorsal surface of brainstem angled in a superior direction (Fig 5 right panel) may also impact the functional definition of the IC.

The functional responses in the MGB cover an area that is in agreement with histological data. Interestingly, compared to the IC or SOC, there is no major direction of extension of functional responses as well as no clear direction (in comparison to SOC and IC) of vascular draining.

A quantification of the volume of functionally defined structures (significant activation in at least three of the ten subjects) is reported in Table 3, and the overlap between functional regions and the BigBrain segmentations after projection in MNI space is reported in Table 2 (as inset in Fig 2).

Diffusion MRI

With the successful identification of the subcortical auditory structures with functional MRI, we next sought to estimate the likely connections between these structures in vivo. We analyzed the high spatial and angular resolution diffusion data to estimate streamlines of white matter connectivity following a similar process as the post mortem MRI (see Methods for further details).

Fig 6 shows diffusion tractography streamlines that pass through at least one subcortical auditory structure (as defined by group-level probabilistic functional activation [significant response in at least three out of ten subjects]; see section above). The high spatial and angular resolution of these data allow for vastly improved estimation of white matter connections between these deep, small structures.

While not a measure of actual physical brain connections — and therefore requiring caution in interpretation — connectivity patterns resemble what we would expect to see based on animal model tracer investigations. Overall, the connectivity network appears to be dominated by laterality, in that left hemisphere structures are generally more connected with other left hemisphere structures.

However, there are a few notable exceptions to this pattern: the cochlear nuclei and superior olivary complexes are strongly connected bilaterally, which fits with animal research suggesting

Figure 5. In vivo functional MRI responses to auditory stimuli, combined across ten participants. Left column: Conjunction of participants plotted on top of one participant's short inversion T1-weighted anatomical MRI. Center column: Conjunction of participants' fMRI responses warped to MNI space and plotted on top of BigBrain MNI (corrected) image. Right column: Conjunction of fMRI responses plotted on top of post mortem MRI vasculature images (1.1 mm minimum intensity projection).

Figure 6. In vivo tractography of the subcortical auditory system from 7T diffusion-weighted MRI. Left: 3-D images from one participant. Fiber orientation distribution functions were estimated from diffusion-weighted MRI images of the brainstem and were used for deterministic tractography. Streamlines that passed through functionally defined auditory ROIs (dark grey) are shown here (excluding streamlines through the medulla). Colors represent the local orientation of the streamline at each point. Top right: connectivity between subcortical auditory ROIs as a percentage of total brainstem streamlines, averaged over 10 participants. Bottom right: schematic of auditory brainstem connectivity from Gray's Anatomy of the Human Body.

236 one-half to two-thirds of ascending auditory connections cross the midline at these early stages.
237 Additionally, there are a small number of connections between left and right inferior colliculi, likely
238 along the anatomical commissure of the inferior colliculus.

Discussion

The auditory pathway includes a number of subcortical structures and connections, but identifying
these components in humans has been challenging with existing in vivo imaging methods. To
facilitate research on the anatomy and function of the human subcortical auditory system, we
created and validated 3-D atlases of the human auditory pathway based on gold standard histology,
50 μm isotropic resolution post mortem anatomical MRI, and in vivo functional MRI at 7T. In addition,
we created 3-D connectivity maps of the human subcortical auditory pathway using diffusion MRI
tractography in a post mortem MRI sample and in living participants.
These atlases and connectivity maps are the first fully 3-D representations of the human
subcortical auditory pathway and are publicly available to make the localization of subcortical
auditory nuclei easier. In particular, the atlases are available in a common reference space (MNI152)
to make registration to other MRI data as straightforward as possible. As part of this registration
process, we have improved the registration of the brainstem of BigBrain histological data to the
MNI space, where the original MNI version presented a significant misregistration of the colliculi (as
noticeable in Fig 7). The result of our new registration allows to more correctly localize the colliculi

of BigBrain data in MNI without compromising the registration of other brainstem and thalamic nuclei.

In creating the atlas with three distinct modalities, we were able to compare the sensitivity of each of the methods in identifying the human subcortical auditory pathway as well as validate the ability of post mortem and in vivo MRI to delineate these structures in comparison with histological definitions. Each modality provided useful information to the segmentation of the auditory nuclei. All regions could be identified in the BigBrain histological data, that also allowed us to identify small auditory sub-nuclei such as the medial superior olive and lateral superior olive. High-resolution post mortem MRI also clearly delineated the medial geniculate and inferior colliculus (with less contrast for the superior olive and cochlear nucleus), while the overall image contrast facilitated registration with in vivo MRI. High-resolution in vivo functional MRI exhibited greater sensitivity to auditory structures than in vivo anatomical MRI that was even higher resolution. Together with strong overlap between the functional atlas and post mortem atlases, we showed that functional MRI is useful to localize structures throughout the auditory pathway despite their small size. In conclusion, we found that each of the MRI methods described here provides novel information about the structure of the human subcortical auditory pathway, suggesting that MRI is a capable tool for investigating this system.

While not representing specific cells, MRI holds a number of advantages over the gold standard method, histology (*Johnson et al., 1993*). First, MRI allows for visualization and analysis of an entire 3-D structure at once, with minimal geometric warping from (virtual) slice to slice (which can occur in slice-based histology if individual slices contract on a slide or are damaged during the physical slicing). Second, MRI can be used in vivo in human participants, opening up the possibility to address research questions on the functional and anatomical properties of human subcortical structures, their correspondence, and their involvement in human behavior.

Probing the connectivity of the human subcortical auditory pathway has been extremely limited, since gold standard (but invasive) tracer studies are largely unavailable for human specimens. In this study, we show that diffusion MRI tractography is sensitive to connections within the human subcortical auditory system, both post mortem and in vivo. In addition to streamlines corresponding to the lateral lemniscus — the major ascending auditory white matter tract — we can see streamlines crossing the midline at the level of the superior olivary complex and the inferior colliculus.

Interestingly, with the highest resolution data (200 μm post mortem diffusion-weighted MRI), we were able to estimate streamlines visually resembling the expected auditory pathway, but missing putative key connections between subcortical auditory structures themselves. In contrast, the relatively lower resolution in vivo diffusion-weighted MRI produced estimates of connectivity more like what we expected from the literature. We had two hypotheses as to why these results appeared. First, the higher resolution anatomical definition of the nuclei not including the immediately surrounding white matter could miss streamlines that terminate at the immediate proximity of the structures' borders (similar to issues in cortex (*Reveley et al., 2015*). Second, partial volume effects in the lower resolution data — combining white matter and grey matter in the same voxels — could actually *increase* streamlines terminating within the anatomical ROIs. Our follow-up analyses on the post mortem diffusion-weighted MRI data — dilating the ROIs and downsampling the data to the in vivo resolution — resulted in greater streamline connectivity between subcortical auditory structures, suggesting that our hypotheses were likely. Thus, while high spatial resolution diffusion-weighted MRI allowed for much finer, higher quality streamline estimates, it also places constraints on tractography analyses that must be accounted for and investigated further.

More generally, the density of brainstem and midbrain nuclei and frequent crossings between perpendicular white matter bundles pose a challenge to diffusion tractography estimations of white matter connectivity, so it was not clear beforehand if this methodology would be sufficient for visualizing these connections. Additionally, because a gold-standard connectivity method is not available in humans, we could not directly validate our tractography findings. However, our results suggest that, with continually improving diffusion-weighted MRI acquisition and analysis techniques,

focused investigations on the human subcortical auditory pathway can—and should—become more prominent in the near future.

In addition to high resolution anatomical post mortem MRI and diffusion MRI tractography, we were also able to identify the subcortical auditory system in vivo with functional MRI. Previous studies have identified these structures with functional MRI, but they typically required constrained acquisition parameters — for instance, incorporating single voxel slices at low through-plane resolution in order to support high in-plane resolution (*Guimaraes et al., 1998*; *Harms and Melcher, 2002*; *Griffiths et al., 2001*; *Hawley et al., 2005*; *Sigalovsky and Melcher, 2006*). In this study, by taking advantage of the increased signal of high-field (7-Tesla) MRI, we were able to image the brainstem using isotropic voxels at high resolution across a wider field-of-view that covers the human auditory pathway in coronal oblique slices. The use of slice acceleration (*Moeller et al., 2010*; *Setsompop et al., 2012*) allowed us to acquire enough slices to cover the whole brainstem, thalamus and cortical regions around Heschl's gyrus with the exclusion of anterior portions of the superior temporal gyrus and sulcus. Using isotropic voxels allowed us to better evaluate the 3-D volume of significantly activated regions, limiting partial volume effects that are inevitable when using thick anisotropic slices.

Similar to previous research at lower magnetic fields (*Hawley et al., 2005*; *Sigalovsky and Melcher, 2006*), the 7T MR images did not allow for an anatomical definition of the CN and SOC (although IC and MGB were clearly visible). A possible reason for this is the reduced signal- and contrast-to-noise ratio in these regions. It should be noted that we could identify the SOC in the MNI ICBM 152 dataset that results from the average of a much larger cohort. Therefore, future investigations should be tailored for optimizing anatomical image contrasts to auditory brainstem regions in single subjects. The (post mortem) atlases we provide here will prove a useful tool for these investigations by providing a reference for the expected location (and size) of these regions.

In contrast to in vivo anatomical localization, our data — in agreement with previous reports (*Hawley et al., 2005*; *Sigalovsky and Melcher, 2006*) — show that *functional* mapping of the subcortical auditory pathway is an effective method for localizing these structures. While histologically defined CN and SOC regions have been previously used to sample functional responses from in vivo fMRI data (*Hawley et al., 2005*; *Sigalovsky and Melcher, 2006*), the overlap between functionally and histologically defined subcortical auditory structures has not been reported before. Here we investigated the ability of BOLD fMRI (as an indirect measure of neuronal activity) to localize subcortical auditory regions. We show that functional definitions are possible, as distinct clusters of activation were detected in all subjects across the subcortical auditory pathway. We could identify the subcortical auditory nuclei despite not using cardiac gating, a method that previous studies showed to increase the signal-to-noise ratio in subcortical regions (*Guimaraes et al., 1998*; *Harms and Melcher, 2002*; *Griffiths et al., 2001*; *Hawley et al., 2005*; *Sigalovsky and Melcher, 2006*). We instead increased statistical power by presenting a large number of natural sounds with multiple repetitions. Using smaller voxels also reduced partial volume effects between cerebrospinal fluid (which is heavily affected by physiological noise) and the brain tissue (*Triantafyllou et al., 2016*). In addition, the correspondence of functionally defined regions across ten participants after anatomical alignment allowed us to build a functional probabilistic atlas.

Despite these positive outcomes, functionally defined regions exhibited overall larger volumes compared to the histological ones (see Table 3). Although relatively high resolution (1.1 mm isotropic), our functional voxel size and the spatial smoothing applied during preprocessing might be the source of this observation. Another source can be the residual anatomical misalignment across subjects that may have contributed to the lower degree of overlap at CN and SOC. In this case, the individual anatomical images not showing enough contrast might be the cause. Another factor might be the partial volume and draining effects due to the vascular architecture. Further, because we used only the overall response to sounds as functional definition, the regions we defined may include sub-regions not specific to the system under investigation (e.g., the inclusion of multisensory deep layers of the superior colliculus at the border with the IC) (*Sparks and Hartwich-Young, 1989*;

Jiang et al., 1997). This effect could be reduced by using different stimuli and statistical contrasts.
For instance, one could contrast unisensory and multisensory stimuli to identify – within the current
functional definition – the IC voxels that respond to visual stimulation and thus may represent
multisensory superior colliculus.

We also investigated the possibility of defining anatomical connections between subcortical
auditory nuclei using diffusion-weighted MRI. While affected by similar confounds as functional
MRI (e.g., partial voluming effects, physiological noise, and relative signal weighting), this technique
faces additional complications introduced by the number of orientations required, the gradient
strength (b-value) selected, the modeling of diffusion or fiber orientations within each voxel, and
the estimation of streamlines across brain regions. The post mortem and in vivo diffusion MRI
datasets in this study each implemented state-of-the-art acquisition techniques to optimize the MRI
signal-to-noise ratio and minimize MRI modeling errors. For example, as the fixation process likely
changes the diffusion characteristics of the tissue (*Pfefferbaum et al., 2004*; *Miller et al., 2011*), we
compensated for this effect by increasing the diffusion gradient strength (b-value). The constrained
spherical deconvolution modeling method takes advantage of the high angular resolution of each
dataset to provide fine-grained estimations of fiber orientation distributions. Additionally, the Euler
Delta Crossings (EuDX) deterministic tractography method is effective at generating streamlines
through voxels with multiple fiber orientation peaks, such as where white matter bundles cross.
However, as diffusion MRI and tractography are not measuring true neuronal connections, there is
still room for error in diffusion orientation and streamline estimation (*Schilling et al., 2019a*,b).

Our BigBrain histological segmentations are very similar in volume to those reported previously
in the literature (*Moore, 1987*; *Glendenning and Masterton, 1998*), with slightly smaller cochlear
nuclei and slightly larger medial geniculate bodies, but similar SOC and IC volumes. It has to be
noted that the physical slicing process potentially introduces deformations in the tissue, and while
the publicly available BigBrain dataset is of extremely high quality (with good registration from
slice to slice), subtle deformations may have affected the shape or volume of the structures we
identified.

Post mortem MRI segmentations differed more greatly, with smaller CN and SOC definitions but
larger MGB definitions compared to both the literature and BigBrain histological segmentations.
These differences could possibly be caused by the reduced contrast-to-noise ratio, compared to the
histological data, available in the post mortem MRI data (despite their high spatial resolution). This
reduced contrast-to-noise ratio may be caused by both reduced differences in magnetic properties
between the regions and their surrounding tissues as well as from residual partial volume effects
(especially for the very small sections of the dorsal CN, for example) that may have blurred the
borders of the auditory nuclei in the post mortem MRI data. Contrast-to-noise ratio may be
ameliorated by different acquisition/reconstruction techniques (*Wang et al., 2018*), and optimizing
parameters may improve the definition of auditory nuclei on the basis of post mortem MRI data.
Finally, slight misregistration between specimens (e.g. the histological data and the post mortem
MRI data) may still affect our comparisons, as registration between images remains a challenge. For
instance, Fig. 2 shows slightly different shapes and locations for the inferior colliculus between the
two datasets, despite non-linear registration to a common template. Although non-linear methods
significantly improve gross registration between specimens, large misregistrations are still possible
(as shown for the colliculi in the original BigBrain MNI registration). These issues can be addressed
manually using additional image registration techniques, as we did here with the BigBrain MNI
registration (see our "corrected" version above), but such hands-on, time-intensive edits are not
always possible. Further, vastly different image contrasts (like histology and MRI) result in different
regions or subregions being emphasized in the signal, creating an additional challenges in the
registration procedure.

More generally, post mortem imaging—whether MRI or histology—is prone to modest de-
formation of the specimen. Additionally, both post mortem specimens in this paper were from
sixty-five-year-old male donors, and age may have additionally affected the volume of the brain

structures we investigated.

Despite these limitations, the strong overlap across the varied datasets and methodologies in
this study suggest that each method is effective for localizing the subcortical auditory pathway. The
atlases derived from each localization method is publicly available (see "Data and code availability"
in Methods) to facilitate further investigations into the structure, function, and connectivity of the
human subcortical auditory system in vivo. Lastly, we hope that the three-dimensional representa-
tions found in this paper and in the available data will be beneficial to others in understanding the
immensely complex, but identifiable, structure of the human subcortical auditory pathway.

Methods

416 See Supplementary Figure 6 for a summary of data sources, data processing steps, and software
417 used in these analyses.

MRI acquisition parameters

In vivo MRI

The experimental procedures were approved by the ethics committee of the Faculty for Psychology
and Neuroscience at Maastricht University, and were performed in accordance with the approved
guidelines and the Declaration of Helsinki. Written informed consent was obtained for every
participant before conducting the experiments. All participants reported to have normal hearing,
had no history of hearing disorder/impairments or neurological disease.

Images were acquired on a 7T Siemens MAGNETOM scanner (Siemens Medical Solutions,
Erlangen, Germany), with 70 mT/m gradients and a head RF coil (Nova Medical, Wilmington, MA,
USA; single transmit, 32 receive channels) at Maastricht University, Maastricht, Netherlands.

We collected data for n=10 participants (age range 25 to 30, 6 females), in three separate sessions.
In the first session, we acquired the in vivo anatomical data set consisting of: 1) a T1-weighted (T1w)
image acquired using a 3-D MPRAGE sequence (repetition time [TR] = 3100 ms; time to inversion
[TI] = 1500 ms [adiabatic non-selective inversion pulse]; echo time [TE] = 2.42 ms; flip angle = 5°;
generalized auto-calibrating partially parallel acquisitions [GRAPPA] = 3 (*Griswold et al., 2002*); field
of view [FOV] = 224 × 224 mm^2; matrix size = 320 × 320; 256 slices; 0.7 mm isotropic voxels; pixel
bandwidth = 182 Hz/pixel; first phase encode direction anterior to posterior; second phase encode
direction superior to inferior); 2) a Proton Density weighted (PDw) image (0.7 mm iso.) with the
same 3-D MPRAGE as for the T1w image but without the inversion pulse (TR = 1380 ms; TE = 2.42
ms; flip angle = 5°; GRAPPA = 3; FOV = 224 × 224 mm^2; matrix size = 320 × 320; 256 slices; 0.7
mm iso. voxels; pixel bandwidth = 182 Hz/pixel; first phase encode direction anterior to posterior;
second phase encode direction superior to inferior); 3) a T2*-weighted (T2w) anatomical image
acquired using a modified 3-D MPRAGE sequence (*De Martino et al., 2015*) that allows freely setting
the TE (TR = 4910 ms; TE = 16 ms; flip angle = 5°; GRAPPA = 3; FOV = 224 × 224 mm^2; matrix size
= 320 × 320; 256 slices; 0.7 mm iso. voxels; pixel bandwidth = 473 Hz/pixel; first phase encode
direction anterior to posterior; second phase encode superior to inferior) and 4) a T1-weighted
images acquired with a short inversion time (SI-T1w) using a 3-D MPRAGE (*Tourdias et al., 2014*)
(TR = 4500 ms; TI = 670 ms [adiabatic non-selective inversion pulse]; TE = 3.37 ms; flip angle = 4°;
GRAPPA = 3; FOV = 224 × 224 mm^2; matrix size = 320 × 320; 256 slices; 0.7 mm isotropic voxels; pixel
bandwidth = 178 Hz/pixel; first phase encode direction anterior to posterior; second phase encode
direction superior to inferior). To improve transmit efficiency in temporal areas when acquiring
these anatomical images we used dielectric pads (*Teeuwisse et al., 2012*).

In the same session we acquired, for each participant, a diffusion-weighted MRI data set using a
protocol originating from the 7T Human Connectome Project (1.05 mm isotropic acquisition and
b-values = 1000 and 2000 s/mm^2) (*Vu et al., 2015*), extended in order to collect one additional shell
at b-value at b = 3000 s/mm^2 (*Gulban et al., 2018a*). Other relevant imaging parameters were (FOV
= 200 × 200 mm^2 with partial Fourier 6/8, 132 slices, nominal voxel size = 1.05 mm isotropic, TR/TE =

7080/75.6 ms, MB = 2, phase encoding acceleration (GRAPPA) = 3, 66 directions and 11 additional b = 0 volumes for every b-value). A total of 462 volumes were obtained (231 in each phase encoding direction anterior-posterior and posterior-anterior) for a total acquisition time of 60 minutes.

The other two sessions were used to collect functional data in order to identify sound responsive regions in the human thalamus and brainstem. Participants listened to 168 natural sounds (1 second long) coming from seven categories (speech, voice, nature, tools, music, animals and monkey calls) presented in silent gaps in between the acquisition of functional volumes and were asked to press a button every time the same sound was repeated. The experimental paradigm followed a rapid-event-related design in which sounds were presented with a mean inter stimulus interval of four volumes (minimum three maximum five volumes). The two sessions were identical and each session consisted of twelve functional runs and across the twelve runs each sound was presented three times (i.e. each sounds was presented six times across the two sessions). The 168 sounds were divided in four sets of 42 sounds, each set was presented in three (non consecutive) runs. As a result, the twelve functional runs of each session formed four cross validation sets each one consisting of nine training runs and three testing runs (i.e. 126 training and 42 testing sounds). Note that the testing runs were non overlapping across the cross validations. Catch trials (i.e. sound repetitions) were added to each run, and were excluded from all analyses.

Functional MRI data were acquired with a 2-D Multi-Band Echo Planar Imaging (2D-MBEPI) sequence (*Moeller et al., 2010*; *Setsompop et al., 2012*) with slices prescribed in a coronal oblique orientation in order to cover the entire brainstem and thalamus and covering primary and secondary cortical regions (TR = 2600 ms; Gap = 1400 ms ; TE = 20 ms; flip angle = 80°; GRAPPA = 3; Multi-Band factor = 2; FOV = 206 × 206 mm^2; matrix size = 188 × 188; 46 slices; 1.1 mm isotropic voxels; phase encode direction inferior to superior). Reverses phase encode polarity acquisitions were used for distortion correction. Respiration and cardiac information were collected during acquisition using a respiration belt and pulse oximeter respectively.

Post mortem MRI

A human brainstem and thalamus specimen were dissected at autopsy from a 65-year-old anonymous male. The specimen was flushed with saline and immersed for two weeks in 10% solution of neutral buffered formalin. Following this, the specimen was re-hydrated for one week in 0.1 M solution of phosphate buffered saline doped with 1% (5 mM) gadoteridol. Before the MRI acquisition, the specimen was placed in custom MRI-compatible tube immersed in liquid fluorocarbon.

Magnetic resonance imaging was conducted in a 210 mm bore Magnex/Agilent MRI at the Duke University Center for In Vivo Microscopy. Three-dimensional gradient echo images were collected at 50 μm^3 spatial resolution over a period of fourteen hours, with FOV = 80 × 55 × 45 mm, repetition time (TR) = 50 ms, echo time (TE) = 10 ms, flip angle = 60°, and bandwidth = 78 Hz/pixel.

Diffusion-weighted spin echo images were collected at 200 μm^3 spatial resolution with 120 diffusion gradient directions at strength b=4000 s/m^2 and 11 b=0 s/m^2 volumes over 208 hours. The FOV was 90 × 55 × 45 mm with TR = 100 ms, TE = 33.6 ms, and bandwidth = 278 Hz/pixel.

Anatomical image registration

SI-T1w, T1w, T2*w and PDw images (700 μm iso.) were transformed to Talairach space (500 μm iso.) using BrainvoyagerQX (*Goebel, 2012*) with default parameters in version 2.8.4. Intensity inhomogeneity correction as implemented in SPM12 unified segmentation (*Ashburner and Friston, 2005*) was used for all images. A smaller volume containing brainstem and thalamus in each image was extracted (in the Talairach space) using FSL version 5.0.9 (*Jenkinson et al., 2012*) and histogram matched using percentile clipping (1% and 99%).

Individual masks for each 10 brainstems were created semi-automatically using ITK-SNAP version 3.6.0 active contour segmentation mode followed by manual edits. These masks included regions starting from 2 cm below the inferior part of pons to 0.5 cm above the medial geniculate nucleus (MGN), with a lateral extend reaching until the lateral geniculate nucleus (LGN) and 3 cm

anterior from MGN, not including cerebellum or large arteries that lie on the surface of brainstem. These brainstem masks were then used with FSL-FNIRT (*Andersson et al., 2007*) to warp nine of the ten brainstems to the reference brainstem (subject 1) using SI-T1w images. We used the SI-T1w images to drive the non linear registration due to the enhanced anatomical contrast across structures within the thalamus and brainstem present in these images (*Tourdias et al., 2014*; *Moerel et al., 2015*). The FNIRT parameters were subsamp $= 2, 2, 1, 1$, miter $= 100, 100, 50, 50$, infwhm $= 2, 2, 1, 1$, reffwhm $= 2, 2, 0, 0$, lambda $= 100, 50, 20, 5$, estint $= 0, 0, 0, 0$, warpres $= 2, 2, 2$ with spline interpolation (parameters not mentioned here were the defaults as set in FSL 5.0.9).

To compare in vivo with post mortem MRI and histology data, we projected the averaged SI-T1w, T1w, T2*w and PDw images to the MNI reference space (ICBM 152 2009b non-linear symmetric, 500 μm iso.) (*Fonov et al., 2009*, *2011*) [1]. The ICBM 152 reference includes T1w, T2w and PDw data and projecting in vivo and post mortem MRI as well as histology data to this space allowed us also to evaluate the contrast that these commonly used template images have in subcortical auditory areas. To register our in vivo MRI data set to MNI, we used FSL-FNIRT but this time driven by the T1w images (abvailable both in our data set and in the MNI ICBM 152 2009b data).

The post mortem diffusion b0 image was transformed to the post mortem anatomical image space with an affine transformation in ANTs. Anatomical-space images (including the manually segmented atlas) could then be transformed into diffusion space using the 'antsApplyTransforms' command, with the affine transform matrix, a super-sampled diffusion image (from 200 μm to 50 μm to match the anatomical image resolution) as the reference image, and denoting the warp as an inverse transform.

In vivo and post mortem images were registered non-linearly using ANTs. The in vivo SI-T1w image was warped to the post mortem diffusion b0 image following a rigid, then affine, then non-linear SyN algorithm. This produced an in vivo brainstem image in post mortem diffusion space.

The ANTs non-linear registration also created warp and inverse warp transforms that could then be used to transform atlases from one space to another. To preserve the higher resolution of the post mortem MRI when inverse warping post mortem images to in vivo space, we supersampled the in vivo SI-T1w image to 200 μm (matching the post mortem diffusion image) or 50 μm (matching the post mortem anatomical image).

Finally, to transform the post mortem anatomical image (50 μm) to MNI space, we applied the inverse transform from post mortem anatomical to diffusion space (resampled to 50 μm), then the inverse transform from diffusion space to in vivo space (similarly supersampled to 50 μm), and finally from in vivo space to MNI space using the FSL-FNIRT inverse transform (described above).

BigBrain histology segmentation

In what follows we describe the main anatomical observations related to the auditory structures as segmented in the 100 μm histological data. Images were segmented independently by two raters (KRS, OFG). Overlap between the two raters was high (see Table 2 [top row - Big Brain across segmenters] in Fig 2); in the figures we show the regions that were consistently segmented by both raters.

Vestibulocochlear nerve

The vestibulocochlear nerve (the eighth cranial nerve, or CNVIII) enters the brainstem where the medulla and the pons meet (the pontomedullary junction). The cochlear component of the vestibulocochlear nerve is composed of spiral ganglion neurons, whose cell bodies are within the cochlea and which carry frequency-specific information to the brainstem.

In the BigBrain histology, CNVIII extends primarily laterally (but also anteriorly and inferiorly) from the pontomedullary junction, bound posteriorly by the cerebellum. Parts of the nerve root are

---

[1] http://www.bic.mni.mcgill.ca/ServicesAtlases/ICBM152NLin2009

551 still visible in the images although being cut. It is therefore not labeled in our histological atlas (but
552 see the post mortem MRI atlas below).

Cochlear nucleus

Once reaching the brainstem, the auditory nerves split into two main routes—one to the anterior
ventral cochlear nucleus (AVCN), and one to the posterior ventral cochlear nucleus (PVCN) and then
on to the dorsal cochlear nucleus (DCN) (*Webster, 1992*). Within each subnucleus, the neurons
maintain the tonotopic frequency representation they receive from the cochlea via the cochlear
nerve (*De No, 1933b*,a; *Rose et al., 1960*; *Sando, 1965*; *Evans, 1975*; *Ryugo and May, 1993*; *Ryugo*
and Parks, 2003) (see bottom panels of the two left most columns in Fig 2).
In the BigBrain data, the AVCN is situated anterior and medial to the root of CNVIII, while the
PVCN continues from the root of CNVIII and extends posteriorly towards the DCN. The DCN is clearly
visible as a dark band wrapping around the cerebellar peduncle posteriorly, becoming exposed on
the dorsal surface of the pons.

Superior olivary complex

The next structure along the auditory pathway is the superior olivary complex (SOC), which in
humans is located in the inferior pons. The SOC receives the majority of its ascending inputs
from the contralateral cochlear nucleus, although it also receives ipsilateral inputs as well. The
contralateral dominance is maintained throughout the remaining ascending pathway. The SOC is
comprised of the lateral superior olive (LSO), medial superior olive (MSO), and the medial nucleus
of the trapezoid body (MNTB). The size of each of these nuclei varies between species, and it
is debated whether the trapezoid body exists in the human SOC (*Moore, 1987*; *Strominger and*
Hurwitz, 1976) (but see *Kulesza and Grothe* (*2015*) review of recent findings affirming the existence
of the human MNTB).
Although the individual substructures within the SOC have unique anatomy that can be identified
from histology (*Moore, 1987*; *Kulesza, 2007*), here we outline the structure of the SOC as a whole in
order to include all identifiable substructures (namely the MSO and LSO - see second panel from
the bottom of the two left most columns in Fig 1). The MSO is the largest SOC nucleus in humans,
unlike in other animals. The MSO receives inputs from both the left and right AVCN and sends
outputs to the ipsilateral lateral lemniscus. The LSO receives inputs from the ipsilateral AVCN and
from the ipsilateral MNTB. Outputs are sent to both ipsilateral and contralateral lateral lemnisci.
The MNTB receives inputs from the contralateral AVCN, and its axons terminate in the ipsilateral
LSO.
The MSO and LSO are visible in the BigBrain images, despite their small size. The MSO is a
thin pencil-like collection of nuclei whose caudalmost point begins around the same axial plane
as the rostralmost extent of the AVCN, about 4 mm medial (and slightly anterior) to the AVCN. It
then extends about 1 cm rostrally (angled slightly laterally), where it eventually meets the lateral
lemniscal tract. The LSO neighbors the MSO near its caudalmost portion, forming a "V" shape
when viewed axially. In our histological atlas, these two structures are combined into a single SOC
segmentation. Cells of the MNTB are not clear to us in this sample, so we do not segment it in our
atlas.

Inferior colliculus

The inferior colliculus (IC) is a large, spherical structure in the dorsal midbrain and receives ascending
inputs from the auditory brainstem via the lateral lemniscus (see second panel from the top of the
two left most columns in Fig 1). The central nucleus of the inferior colliculus receives most of these
connections, with external nuclei primarily receiving descending connections (*Webster, 1992*). The
inferior colliculus sends axons to the medial geniculate body of the thalamus via the brachium of
the inferior colliculus.
In the BigBrain data, the inferior colliculus is clearly identifiable as the lower two of the four

bumps along the dorsal portion of the midbrain (or tectum). The darkest staining within these structures corresponds to the central nucleus of the inferior colliculus. An intensity gradient outside of the central nucleus likely corresponds to the external and dorsal nuclei, which were included in our segmentation of the IC. Bounding the IC superiorly is the superior colliculus; medially, the commissure of the IC connecting the two inferior colliculi, as well as the aqueduct and periaqueductal grey; and anteriorly, other midbrain nuclei such as the cuneiform nucleus (lateral and inferior to the IC are the borders of the midbrain).

Medial geniculate of the thalamus

The medial geniculate body (MGB) of the thalamus is the final subcortical auditory structure that sends auditory signals to the auditory cortex via the acoustic radiations (*Winer, 1984*) (see top panel of the two left most columns in Fig 1). The MGB contains two or three major subdivisions: the ventral MGB receives the majority of IC inputs, while the dorsal and medial subdivisions (at times grouped together, at times separately) receive more varied inputs from auditory and non-auditory subcortical structures.

In the BigBrain sample, the MGB is visible as a dark patch medial to the lateral geniculate nucleus (which can be easily identified by its striations) in a coronal view. Axially, the MGB takes an ovoid shape with a clear dorsolateral boundary next to the brachium of the superior colliculus, which appears light due to lack of cell nuclei being stained. Ventromedially, the MGB is bordered by a light band corresponding to the medial lemniscus. Rostrally, we marked the edge of the MGB where cell staining decreases, at the border with the pulvinar nucleus and ventral posterolateral nucleus of the thalamus.

Post mortem MRI segmentation

In what follows we describe the anatomical contrast that can be leveraged from these post mortem MRI data in order to identify structures in the auditory brainstem. We then used these segmentations to create an MRI-based atlas of the subcortical auditory system, separate from the BigBrain histology-based atlas.

Vestibulocochlear nerve

The CNVIII is visible in the post mortem MRI near the pontomedullary junction, extending laterally and anteriorly from the brainstem (see the lower panels in Fig 2).

Cochlear nucleus

The cochlear nuclei are challenging to identify in the post mortem MRI data, although the presence of the CNVIII root provides a landmark for localizing the other structures. Due to low signal contrast around the ventral cochlear nucleus area in the T2*-weighted GRE MRI, we segmented the VCN according to the literature: bound by the cochlear nerve root and wall of the pons laterally, and by cerebellar white matter tracks medially. We were able to segment the dorsal cochlear nucleus based on the T2*-weighted image, where it appears brighter and can be identified as running posteriorly from the VCN and dorsally along the surface of the pons, distal to the inferior cerebellar peduncle.

Superior olivary complex

As with the cochlear nuclei, the SOC are more difficult to identify in the post mortem MRI than in the histology, likely since the individual subnuclei like the MSO and LSO approach the size of a voxel in at least one direction and are therefore prone to partial voluming effects. However, the pencil-like MSO can still be identified in the coronal plane as a dark, elongated structure in the T2*-weighted image, starting around the level of the ventral cochlear nucleus. In the axial plane, the SOC (but not its individual subnuclei) can be seen as a dark spot in the T2*-weighted image between the facial nucleus and the trapezoid body (see the second row from the bottom in Fig 2).

Inferior colliculus

As in the BigBrain data, the inferior colliculus is relatively easy to identify based on its gross anatomical structure on the dorsal aspect of the midbrain. Additionally, the MR contrast provides relatively clear boundaries between the colliculi and surrounding structures. Indeed, it may even be possible to segment the inferior colliculus into its subnuclei—the central, external, and dorsal nuclei—based on T2*-weighted MR signal intensities (see the second row from the top in Fig 2). The external nucleus of the IC appears dark in the T2*-weighted image, on the lateral aspect of the IC. Medial to the external nucleus is the central nucleus, which has higher T2*-weighted intensity (appears brighter) in our MR images, and has clear boundaries on its ventral, medial, and dorsolateral sides. The dorsal nucleus is along the dorsal aspect of the IC and is the brightest subcomponent within the IC in terms of T2*-weighted MR signal.

Medial geniculate

Although the borders of the MGB are less clear in the post mortem MRI than in the BigBrain images, the structure itself is again relatively easy to identify by its gross anatomical location as well as MR signal intensity. In the coronal plane, the medial geniculate is medial to the lateral geniculate at the junction of the midbrain and thalamus. Axially, the medial geniculate has circular or ovoid shape, again medial to the lateral geniculate. In the axial plane, the medial geniculate is largely bordered dorsolaterally by the brachium of the superior colliculus, which appears as a thick, dark band of fibers in the T2*-weighted image. Medially, the medial geniculate is bound by the brachium of the inferior colliculus (also appearing as a dark fiber band), at least through the caudal half of the structure. We have included the portions of this fiber bundle in the segmentation of the medial geniculate, as the auditory fibers connecting the IC and the MGB are quite relevant to MRI connectivity investigations (including our own; post mortem tractography results below).

As with the inferior colliculus, it may be possible to identify separate divisions within the medial geniculate. Within the overall structure, there are two identifiable substructures based on T2*-weighted MR image intensity. Dorsomedially (and somewhat caudally), about half of the medial geniculate has high T2*-weighted contrast and appears bright; the ventrolateral (and slightly rostral) half appears darker in the T2*-weighted image. These segmentations largely (but not perfectly) align with the ventral and dorsal/medial nuclei of the medial geniculate in the Allen Human Brain Atlas (*Hawrylycz et al., 2012*), as well as with those of *Paxinos et al.* (*2019*). However, they vary somewhat from the the axial slice segmentation from *Merker* (*1983*) shown in *Amunts et al.* (*2012*), which show a largely horizontal delineation between the substructures.

Functional MRI analysis

Functional images were preprocessed using BrainvoyagerQX version 2.8.4 (*Goebel, 2012*). Slice-scan-time correction, motion correction, temporal high-pass filtering (GLM-Fourier, 6 sines/cosines) and temporal smoothing (Gaussian, width of kernel 5.2 s). The defaults in BrainvoyagerQX v2.8.4 were used for these steps aside from the explicitly stated values. The functional images were then distortion corrected using the opposite phase encoding images using FSL-TOPUP (*Andersson et al., 2003*). Conversion between Brainvoyager file types to NIfTI which was required to perform distortion correction was done using Neuroelf version 1.1 (release candidate 2) [2] in Matlab version 2016a. Spatial smoothing (Gaussian, FWHM 3 voxels) was also applied after distortion correction. The resulting functional images were then transformed to Talairach space using Brainvoyager at a resolution of 0.5 mm isotropic. We have previously used this procedure in order to reveal tonotopic maps in both the inferior colliculus and medial geniculate nucleus (*De Martino et al., 2013*; *Moerel et al., 2015*) and have shown that the upsampling has no consequence on the spatial distribution of the responses.

GLM-denoise (*Kay et al., 2013*) was used to estimate noise regressors. In brief, for each cross validation a noise pool of non responsive voxels (i.e. voxels with a response to sound representation

---

[2] http://neuroelf.net/

determined by an F-statistic below a given threshold) was determined on the training data set (16 runs across the two sessions) and used to obtain noise regressors defined as the principal components of the noise pool time course matrix that added to a GLM analysis (*Friston et al., 1994*) of the training data would result in an increased activation. The number of noise regressors was optimized using cross validation within the training set. The selected noise regressor spatial maps were projected on the test data to obtain the aggressors for the test data.

Similarly, the hemodynamic response function (HRF) best characterizing the response of each voxel in the brainstem was obtained using a deconvolution GLM (with 9 stick predictors) on the training data. Note that this procedure, while possibly overfitting information in the training data, produces noise regressors and an HRF for each test run (e.g. the noise regressors for runs 4, 6 and 9 of session one comes from an analysis performed on all other runs in the same session) that are not overfitted.

The resulting HRF and noise regressors were used in a GLM analysis of the test runs. We combined all test runs (for each individual voxel) within a fixed effect analysis.

Statistical maps of responses to sounds vs silence were corrected for multiple comparisons at the individual level using False Discovery Rate (FDR; q-FDR = 0.05) and warped to the reference brainstem (subject 1) by applying the warping field obtained on the anatomical data. After projection to the common space, single subject statistical maps were binarized and converted to a probabilistic map by summing maps across subjects at each single voxel (i.e. a value of 10 indicates that all 10 subjects exhibited a statistically significant response to sound presentation corrected for multiple comparisons). Maps are displayed with minimum threshold of at least three out of ten subjects exhibiting significant responses at each voxel and after the application of a cluster size threshold of 27 voxels. To compare the functional activation maps with histology data and post mortem MRI data, the probabilistic maps were projected to the MNI space using the warping field obtained from the anatomical dataset.

BigBrain data

Histology data were obtained by downloading the 100 μm version of the BigBrain (*Amunts et al., 2013*) 3-D Volume Data Release 2015 (from https://bigbrain.loris.ca). We downloaded both the original images and the dataset already aligned to MNI ICBM 152. The nuclei along the auditory pathway (cochlear nucleus, superior olive, inferior colliculus and medial geniculate nucleus) were manually segmented in the histology space image using ITK-SNAP (*Yushkevich et al., 2006*) largely following the definitions in *Moore* (*1987*) when possible.

Correction of the alignment of the inferior colliculi to MNI

Upon visual inspection of the BigBrain image in the MNI ICM 152 space, we detected a major registration error around the inferior colliculi (see Fig 7 - second panel from the left). The registration quality to MNI ICMBM 152 space in the rest of the brainstem was deemed satisfactory, but the the region of the inferior colliculus required correction in order to perform a valid comparison with the MRI data (in vivo and post mortem). Interestingly, the region of the colliculi of the BigBrain in the original histology space appeared to be closer in location to the position of the inferior colliculus in the MNI dataset (compare panel 1 and 3 in Fig 7) indicating that the highlighted misalignment in the original BigBrain MNI dataset originated during the registration procedure.

To perform a new registration to MNI of the brainstem and thalamus of the BigBrain data that observed the already correctly registered boundaries (e.g. the Pons) but corrected the region around the inferior colliculus bilaterally, we followed N steps. First, we defined a region of interest around the inferior colliculus using common anatomical landmarks that were visible in the BigBrain MNI and MNI (2009b) T1, PD, T2 images and where aligned satisfactorily. Second, this region was cut out from the BigBrain MNI and replaced by the same region (i.e. defined by the same anatomical landmarks) in the BigBrain histology space data (before projection to MNI). The convex hulls of the region of interest in the BigBrain histology and in the MNI space were matched using 3-D optimal

BigBrain (100 µm) BigBrain (100 µm) MNI ICBM 152 (500 µm) BigBrain (100 µm)
Histology space MNI ICBM 152 2009b T1w MNI ICBM 152 (corrected)

Figure 7. The registration error around the inferior colliculus is visible bilaterally when comparing Panel 2 and Panel 3. The dashed lines indicate the correct shape (and location) of the colliculi in MNI space. The arrows point to the inferior colliculus (IC). The last panel shows the corrected BigBrain MNI dataset.

transport as implemented in Geogram version 1.6.7 (*Lévy, 2015*; *Lévy and Schwindt, 2018*). Third, the convex hull matched region of the the BigBrain histology space was used to replace the incorrect region which was cut out at step 2. As a result of these three steps we obtained a version of the BigBrain in MNI (BigBrain MNI - implanted) that had the inferior colliculus in the right position but where the transitions between outside to inside of the region of interest that was corrected were visible and not respecting of the topology. To correct for these residual errors, we performed a new FSL-FNIRT alignment between the original BigBrain in histology space and the BigBrain MNI - implanted image. The resulting image (BigBrain MNI - corrected) preserved the actual topology inside the brainstem and at the same time resulted in a correct alignment of the regions around the inferior colliculus bilaterally (see Fig 7 - right panel).

Post mortem MRI vasculature analysis

Gradient echo (GRE) MRI is sensitive to vasculature within the imaged tissue. To highlight vasculature in the post mortem brainstem specimen, we computed the minimum intensity projection in coronal sagittal and axial direction from the 50 µm isotropic voxel GRE MRI data over slabs of 1.1 mm in thickness using Nibabel (*Brett et al., 2017*) and Numpy (*Van Der Walt et al., 2011*)). This image can be seen in Fig 5 right column.

Diffusion MRI analysis

Post mortem diffusion

Before analysis, post mortem diffusion volumes were each registered to the first b0 volume using an affine transformation in ANTs version 2.1.0 (*Avants et al., 2011*). To estimate white matter fiber orientations, we used the constrained spherical deconvolution (CSD) model as implemented in DIPY 0.14 (*Gorgolewski et al., 2011*; *Garyfallidis et al., 2014*; *Tournier et al., 2007*) as a Nipype pipeline (*Gorgolewski et al., 2011*). CSD posits that the observed diffusion signal is a convolution of the true fiber orientation distribution (FOD) with a response function, which can be estimated from the data. DIPY's 'auto-response' function takes a fractional anisotropy (FA threshold); as FA values are much lower in ex vivo tissue compared to in vivo, we used an FA threshold of 0.1 (compared to the default of 0.7).

White matter fiber streamlines were estimated deterministically with DIPY's EudX method (*Mori et al., 1999*; *Garyfallidis, 2013*) with 1,000,000 seeds per voxel and a 45° streamline angle threshold.

To define regions of interest (ROIs) for the fiber display, the auditory structures manually delineated in the post mortem T2*-weighted MR images were transformed to diffusion space using ANTs, and global streamlines were filtered by considering only the voxels in each one of the

ROIs as a seed and further constrained by using all auditory ROIs as tractography waypoints. This resulted in a high-resolution, high-quality auditory-specific subcortical tractogram, which were then visualized in TrackVis 0.6.1 (*Wang et al., 2007*).

In vivo diffusion

7T in vivo dMRI data was corrected for distortions with the HCP pipeline *Glasser et al.* (*2016*); *Sotiropoulos et al.* (*2013*). Specifically, geometric and eddy-current distortions, as well as head motion, were corrected by modeling and combining data acquired with opposite phase encoding directions *Andersson et al.* (*2003*); *Andersson and Sotiropoulos* (*2015*, 2016). The data were then masked to include just the brainstem and thalamus, matching the post mortem specimen.

Similar to the post mortem analysis, we estimated diffusion FODs with a CSD model implemented in DIPY. We used the same parameters as we did for post mortem analysis except for an FA threshold of 0.5 (vs. 0.1 for post mortem). White matter connectivity was estimated with deterministic tractography throughout the brainstem and thalamus, again using DIPY's EudX algorithm (*Mori et al., 1999*; *Garyfallidis, 2013*) with 1,000,000 seeds per voxel and a 45° streamline angle threshold.

For the tractography in the in vivo data we used subcortical auditory ROIs as defined by the analysis of the functional data (i.e. regions that exhibited significant [corrected for multiple comparisons] response to sound presentation in at least three out of ten subjects). The functional ROIs were transformed to individual diffusion space and used as tractography seeds, with all other auditory ROIs as waypoints, producing a subcortical auditory tractogram for each in vivo subject.

Data and code availability

Atlas segmentations and tractography streamlines will be made available through the Open Science Framework (https://osf.io/hxekn/?view_only=be9ec398304344e8bb694a0658d77ed6). Processing and analysis resources, including links to all data and software used in this paper, will be available at https://github.com/sitek/subcortical-auditory-atlas. See Supplementary Figure 6 for an overview of currently available data and code (full resolution version available at the Github repository).

Acknowledgements

We would like to thank to Christophe Lenglet for helping with in vivo dMRI processing and analysis; Agustin Lage-Castellanos for helping with fMRI analysis; and Michelle Moerel, Marta Bianciardi, Jean Augustinack and George Paxinos for giving anatomical advice at various stages. F.D.M. and O.F.G. were supported by NWO VIDI grant 864-13-012. S.S.G. and K.R.S. were supported by NIH grant 5R01EB020740. K.R.S. was additionally funded by NIH NRSA fellowship 5F31DC015695, the Amelia Peabody Scholarship from the Eaton Peabody Laboratory at Mass. Eye and Ear (Boston, MA, USA), and a travel grant from the Harvard Brain Science Institute. MR Histology was acquired at the Duke Center for In Vivo Microscopy, an NIH/NIBIB National Resource (P41EB015897 to G.A.J.), NIH 1S10OD010683-01 (to G.A.J.).

810 ## Glossary

811 ### Anatomical abbreviations

AVCN	Anteroventral cochlear nucleus.
CN	Cochlear nucleus.
CNVIII	8th nerve, vestibulocochlear nerve.
DCN	Dorsal cochlear nucleus.
IC	Inferior colliculus.
LGN	Lateral geniculate nucleus.
LSO	Lateral superior olive.
MGB/MGN	Medial geniculate body/nucleus.
MNTB	Medial nucleus of the trapezoid body.
MSO	Medial superior olive.
PVCN	Posteroventral cochlear nucleus.
SOC	Superior olivary complex.

813 ### MRI acquisition abbreviations

7T	7 Tesla.
dMRI	diffusion magnetic resonance imaging.
FOV	Field of view.
fMRI	functional magnetic resonance imaging.
GRAPPA	Generalized auto-calibrating partially parallel acquisitions.
MB	Multi-band.
MPRAGE	Magnetization prepared rapid acquisition gradient echo.
MRI	Magnetic resonance imaging.
PDw	Proton density weighted.
SI-T1w	Short inversion time T1-weighted.
T1w	T1-weighted.
T2*w	T2*-weighted.
TE	Echo time.
TR	Repetition time.

815 ### Data analysis abbreviations

CSD	Constrained spherical deconvolution.
FA	Fractional anisotropy.
FDR	False discovery rate.
FOD	Fiber orientation distribution.
GLM	General linear model.
HCP	Human connectome project.
HRF	Hemodynamic response function.
ICBM	Internation Consortium for Brain Mapping.
M0	T2 signal with no diffusion weighting.
MD	Mean diffusivity.
MNI	Montreal Neurological Institude.
MSMT	Multi-shell multi-tissue
ODFs	Orientation distribution functions.
ROI	Region of interest.

References

Amunts K, Lepage C, Borgeat L, Mohlberg H, Dickscheid T, Rousseau MÉ, Bludau S, Bazin PL, Lewis LB, Oros-Peusquens AM, Shah NJ, Lippert T, Zilles K, Evans AC. BigBrain: An ultrahigh-resolution 3D human brain model. Science. 2013; doi: 10.1126/science.1235381.

Amunts K, Morosan P, Hilbig H, Zilles K. Auditory system. In: *The Human Nervous System* Elsevier; 2012.p. 1270–1300.

Andersson JLR, Jenkinson M, Smith S. Non-linear registration, aka spatial normalisation. FMRIB Technial Report TR07JA2.; 2007.

Andersson JLR, Skare S, Ashburner J. How to correct susceptibility distortions in spin-echo echo-planar images: application to diffusion tensor imaging. NeuroImage. 2003 oct; 20(2):870–88. doi: 10.1016/S1053-8119(03)00336-7.

Andersson JLR, Sotiropoulos SN. Non-parametric representation and prediction of single- and multi-shell diffusion-weighted MRI data using Gaussian processes. NeuroImage. 2015 nov; 122:166–76. doi: 10.1016/j.neuroimage.2015.07.067.

Andersson JLR, Sotiropoulos SN. An integrated approach to correction for off-resonance effects and subject movement in diffusion MR imaging. NeuroImage. 2016 jan; 125:1063–1078. doi: 10.1016/j.neuroimage.2015.10.019.

Ashburner J, Friston KJ. Unified segmentation. NeuroImage. 2005 jul; 26(3):839–851. doi: 10.1016/j.neuroimage.2005.02.018.

Avants BB, Tustison NJ, Song G, Cook PA, Klein A, Gee JC. A reproducible evaluation of ANTs similarity metric performance in brain image registration. NeuroImage. 2011 feb; 54(3):2033–44. http://www.ncbi.nlm.nih.gov/pubmed/20851191http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3065962, doi: 10.1016/j.neuroimage.2010.09.025.

Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. Probabilistic diffusion tractography with multiple fibre orientations: What can we gain? NeuroImage. 2007 jan; 34(1):144–55. http://www.ncbi.nlm.nih.gov/pubmed/17070705, doi: 10.1016/j.neuroimage.2006.09.018.

Brett M, Hanke M, Côté MA, Markiewicz C, Ghosh S, Wassermann D, Gerhard S, Larson E, Lee GR, Halchenko Y, Kastman E, M C, Morency FC, Maloney B, Rokem A, Cottaar M, Millman J, jaeilepp, Gramfort A, Vincent RD, et al., nipy/nibabel: 2.2.0; 2017. https://doi.org/10.5281/zenodo.1011207, doi: 10.5281/zenodo.1011207.

Calabrese E, Hickey P, Hulette C, Zhang J, Parente B, Lad SP, Johnson GA. Postmortem diffusion MRI of the human brainstem and thalamus for deep brain stimulator electrode localization. Human brain mapping. 2015 aug; 36(8):3167–78. http://www.ncbi.nlm.nih.gov/pubmed/26043869http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4652933, doi: 10.1002/hbm.22836.

De Martino F, Moerel M, van de Moortele PF, Ugurbil K, Goebel R, Yacoub E, Formisano E. Spatial organization of frequency preference and selectivity in the human inferior colliculus. Nature communications. 2013; 4:1386. http://www.ncbi.nlm.nih.gov/pubmed/23340426http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3556928, doi: 10.1038/ncomms2379.

De Martino F, Moerel M, Xu J, van de Moortele PF, Ugurbil K, Goebel R, Yacoub E, Formisano E. High-Resolution Mapping of Myeloarchitecture In Vivo: Localization of Auditory Areas in the Human Brain. Cerebral cortex (New York, NY : 1991). 2015 oct; 25(10):3394–405. http://www.ncbi.nlm.nih.gov/pubmed/24994817http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4585494, doi: 10.1093/cercor/bhu150.

De No RL. III. General Plan of Structure of the Primary Cochlear Nuclei. The Laryngoscope. 1933 apr; 43(4):327–349. http://doi.wiley.com/10.1288/00005537-193304000-00014, doi: 10.1288/00005537-193304000-00014.

De No RL. The central projection of the nerve endings of the internal ear. The Laryngoscope. 1933 jan; 43(1):1–38. http://doi.wiley.com/10.1288/00005537-193301000-00001, doi: 10.1288/00005537-193301000-00001.

Devlin JT, Sillery EL, Hall DA, Hobden P, Behrens TEJ, Nunes RG, Clare S, Matthews PM, Moore DR, Johansen-Berg H. Reliable identification of the auditory thalamus using multi-modal structural analyses. NeuroImage. 2006 may; 30(4):1112–20. http://www.ncbi.nlm.nih.gov/pubmed/16473021http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1458525, doi: 10.1016/j.neuroimage.2005.11.025.

Dhollander T, Raffelt D, Connelly A. Unsupervised 3-tissue response function estimation from single-shell or multi-shell diffusion MR data without a co-registered T1 image. In: *ISMRM Workshop on Breaking the Barriers of Diffusion MRI*; 2016. p. 5.

Dhollander T, Raffelt D, Connelly A. Unsupervised 3-tissue response function estimation from single-shell or multi-shell diffusion MR data without a co-registered T1 image. In: *Proceedings of the 26th annual meeting of the International Society of Magnetic Resonance in Medicine*; 2018. p. 1569.

Ding SL, Royall JJ, Sunkin SM, Ng L, Facer BAC, Lesnar P, Guillozet-Bongaarts A, McMurray B, Szafer A, Dolbeare TA, Stevens A, Tirrell L, Benner T, Caldejon S, Dalley RA, Dee N, Lau C, Nyhus J, Reding M, Riley ZL, et al. Comprehensive cellular-resolution atlas of the adult human brain. . 2016; doi: 10.1002/cne.24097.

Duvernoy HM. Human Brain Stem Vessels: Including the Pineal Gland and Information on Brain Stem Infarction, vol. 11. Springer Science & Business Media; 2013. https://books.google.com/books?id=K87sCAAAQBAJ{&}pgis=1.

Evans EF. Cochlear Nerve and Cochlear Nucleus. In: *Handbook of Sensory Physiology* Springer Berlin Heidelberg; 1975.p. 1–108. http://www.springerlink.com/index/10.1007/978-3-642-65995-9{_}1, doi: 10.1007/978-3-642-65995-9_1.

Fonov V, Evans AC, Botteron K, Almli CR, McKinstry RC, Collins DL. Unbiased average age-appropriate atlases for pediatric studies. NeuroImage. 2011; 54(1):313–327. http://dx.doi.org/10.1016/j.neuroimage.2010.07.033, doi: 10.1016/j.neuroimage.2010.07.033.

Fonov V, Evans A, McKinstry R, Almli C, Collins D. Unbiased nonlinear average age-appropriate brain templates from birth to adulthood. NeuroImage. 2009; 47:S102. http://linkinghub.elsevier.com/retrieve/pii/S1053811909708845, doi: 10.1016/S1053-8119(09)70884-5.

Friston KJ, Holmes AP, Worsley KJ, Poline JP, Frith CD, Frackowiak RSJ. Statistical parametric maps in functional imaging: A general linear approach. Human Brain Mapping. 1994; doi: 10.1002/hbm.460020402.

Garyfallidis E. Towards an accurate brain tractography. PhD thesis, University of Cambridge; 2013.

Garyfallidis E, Brett M, Amirbekian B, Rokem A, van der Walt S, Descoteaux M, Nimmo-Smith I, Dipy Contributors. Dipy, a library for the analysis of diffusion MRI data. Frontiers in neuroinformatics. 2014; 8:8. doi: 10.3389/fninf.2014.00008.

Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, Ugurbil K, Andersson J, Beckmann CF, Jenkinson M, Smith SM, Van Essen DC. A multi-modal parcellation of human cerebral cortex. Nature. 2016; doi: 10.1038/nature18933.

Glendenning KK, Masterton RB. Comparative Morphometry of Mammalian Central Auditory Systems: Variation in Nuclei and Form of the Ascending System. Brain Behav Evol. 1998; 51:59–89. http://www.karger.chhttp://biomednet.com/karger.

Goebel R. BrainVoyager — Past, present, future. NeuroImage. 2012 aug; 62(2):748–756. doi: 10.1016/j.neuroimage.2012.01.083.

Gorgolewski K, Burns CD, Madison C, Clark D, Halchenko YO, Waskom ML, Ghosh SS. Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. Front Neuroinform. 2011 08; 5. https://doi.org/10.3389/fninf.2011.00013, doi: 10.3389/fninf.2011.00013.

Gray H, Lewis W. Anatomy of the human body. 20 ed. Philadelphia: Lea \& Febiger; 1918.

Griffiths TD, Uppenkamp S, Johnsrude I, Josephs O, Patterson RD. Encoding of the temporal regularity of sound in the human brainstem. Nature Neuroscience. 2001; doi: 10.1038/88459.

Griswold MA, Jakob PM, Heidemann RM, Nittka M, Jellus V, Wang J, Kiefer B, Haase A. Generalized Autocalibrating Partially Parallel Acquisitions (GRAPPA). Magnetic Resonance in Medicine. 2002; 47(6):1202–1210. doi: 10.1002/mrm.10171.

Guimaraes AR, Melcher JR, Talavage TM, Baker JR, Ledden P, Rosen BR, Kiang NYS, Fullerton BC, Weisskoff RM. Imaging subcortical auditory activity in humans. Human Brain Mapping. 1998; 6(1):33–41. http://doi.wiley.com/10.1002/{%}28SICI{%}291097-0193{%}281998{%}296{%}3A1{%}3C33{%}3A{%}3AAID-HBM3{%}3E3.0.CO{%}3B2-M, doi: 10.1002/(SICI)1097-0193(1998)6:1<33::AID-HBM3>3.0.CO;2-M.

Gulban OF, De Martino F, Vu AT, Yacoub E, Uğurbil K, Lenglet C. Cortical fibers orientation mapping using in-vivo whole brain 7 T diffusion MRI. NeuroImage. 2018 sep; 178(December 2017):104–118. http://linkinghub.elsevier. com/retrieve/pii/S1053811918304087https://linkinghub.elsevier.com/retrieve/pii/S1053811918304087, doi: 10.1016/j.neuroimage.2018.05.010.

Gulban OF, Schneider M, Marquardt I, Haast RAM, De Martino F. A scalable method to improve gray matter segmentation at ultra high field MRI. PLOS ONE. 2018 jun; 13(6):e0198335. http://dx.plos.org/10.1371/journal. pone.0198335, doi: 10.1371/journal.pone.0198335.

Harms MP, Melcher JR. Sound repetition rate in the human auditory pathway: representations in the waveshape and amplitude of fMRI activation. Journal of neurophysiology. 2002 sep; 88(3):1433–50. http://www.ncbi.nlm. nih.gov/pubmed/12205164, doi: 10.1152/jn.2002.88.3.1433.

Hawley ML, Melcher JR, Fullerton BC. Effects of sound bandwidth on fMRI activation in human auditory brainstem nuclei. Hearing research. 2005 jun; 204(1-2):101–10. http://www.ncbi.nlm. nih.gov/pubmed/15925195http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1855158, doi: 10.1016/j.heares.2005.01.005.

Hawrylycz MJ, Lein ES, Guillozet-Bongaarts AL, Shen EH, Ng L, Miller JA, van de Lagemaat LN, Smith KA, Ebbert A, Riley ZL, Abajian C, Beckmann CF, Bernard A, Bertagnolli D, Boe AF, Cartagena PM, Chakravarty MM, Chapin M, Chong J, Dalley RA, et al. An anatomically comprehensive atlas of the adult human brain transcriptome. Nature. 2012 sep; 489(7416):391–399. http://www.ncbi.nlm.nih.gov/pubmed/22996553http: //www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4243026, doi: 10.1038/nature11405.

Javad F, Warren JD, Micallef C, Thornton JS, Golay X, Yousry T, Mancini L. Auditory tracts identified with combined fMRI and diffusion tractography. NeuroImage. 2014 jan; 84:562–74. http://www.ncbi.nlm. nih.gov/pubmed/24051357http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3898984, doi: 10.1016/j.neuroimage.2013.09.007.

Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. NeuroImage. 2012 aug; 62(2):782–790. doi: 10.1016/j.neuroimage.2011.09.015.

Jeurissen B, Tournier JD, Dhollander T, Connelly A, Sijbers J. Multi-tissue constrained spherical deconvolution for improved analysis of multi-shell diffusion MRI data. NeuroImage. 2014 dec; 103:411–426. http://www. ncbi.nlm.nih.gov/pubmed/25109526, doi: 10.1016/j.neuroimage.2014.07.061.

Jiang ZD, Moore DR, J King A. Sources of subcortical projections to the superior colliculus in the ferret. Brain Research. 1997 may; 755(2):279–292. http://linkinghub.elsevier.com/retrieve/pii/S0006899397001054, doi: 10.1016/S0006-8993(97)00105-4.

Johnson GA, Benveniste H, Black RD, Hedlund LW, Maronpot RR, Smith BR. Histology by magnetic resonance microscopy. Magnetic resonance quarterly. 1993 mar; 9(1):1–30. http://www.ncbi.nlm.nih.gov/pubmed/ 8512830, doi: 10.1139/m63-008.

Kay KN, Rokem A, Winawer J, Dougherty RF, Wandell BA. GLMdenoise: A fast, automated technique for denoising task-based fMRI data. Frontiers in Neuroscience. 2013; doi: 10.3389/fnins.2013.00247.

Kulesza RJ. Cytoarchitecture of the human superior olivary complex: medial and lateral superior olive. Hearing research. 2007 mar; 225(1-2):80–90. http://www.ncbi.nlm.nih.gov/pubmed/17250984, doi: 10.1016/j.heares.2006.12.006.

Kulesza RJ, Grothe B. Yes, there is a medial nucleus of the trapezoid body in humans. Frontiers in neuroanatomy. 2015 may; 9(5):35. http://www.ncbi.nlm.nih.gov/pubmed/4379933http://www.ncbi.nlm. nih.gov/pubmed/25873865http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4379933, doi: 10.3389/fnana.2015.00035.

Lévy B. A Numerical Algorithm for L 2 Semi-Discrete Optimal Transport in 3D. ESAIM: Mathematical Modelling and Numerical Analysis. 2015 nov; 49(6):1693–1715. http://www.esaim-m2an.org/10.1051/m2an/2015055, doi: 10.1051/m2an/2015055.

Lévy B, Schwindt EL. Notions of optimal transport theory and how to implement them on a computer. Computers and Graphics (Pergamon). 2018; doi: 10.1016/j.cag.2018.01.009.

Maffei C, Jovicich J, De Benedictis A, Corsini F, Barbareschi M, Chioffi F, Sarubbo S. Topography of the human acoustic radiation as revealed by ex vivo fibers micro-dissection and in vivo diffusion-based tractography. Brain structure & function. 2018 jan; 223(1):449–459. http://www.ncbi.nlm.nih.gov/pubmed/28866840, doi: 10.1007/s00429-017-1471-6.

Malmierca MS, Hackett TA. Structural organization of the ascending auditory pathway. The Auditory Brain. 2010; p. 9–41.

Merker B. Silver staining of cell bodies by means of physical development. Journal of Neuroscience Methods. 1983 nov; 9(3):235–241. http://linkinghub.elsevier.com/retrieve/pii/0165027083900869, doi: 10.1016/0165-0270(83)90086-9.

Miller KL, Stagg CJ, Douaud G, Jbabdi S, Smith SM, Behrens TEJ, Jenkinson M, Chance SA, Esiri MM, Voets NL, Jenkinson N, Aziz TZ, Turner MR, Johansen-Berg H, McNab JA. Diffusion imaging of whole, post-mortem human brains on a clinical MRI scanner. NeuroImage. 2011 jul; 57(1):167–181. http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3115068, doi: 10.1016/j.neuroimage.2011.03.070.

Moeller S, Yacoub E, Olman CA, Auerbach E, Strupp J, Harel N, Uğurbil K. Multiband multislice GE-EPI at 7 tesla, with 16-fold acceleration using partial parallel imaging with application to high spatial and temporal whole-brain fMRI. Magnetic resonance in medicine. 2010 may; 63(5):1144–53. http://www.ncbi.nlm.nih.gov/pubmed/20432285http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC2906244, doi: 10.1002/mrm.22361.

Moerel M, De Martino F, Uğurbil K, Yacoub E, Formisano E. Processing of frequency and location in human subcortical auditory structures. Scientific reports. 2015 nov; 5:17048. doi: 10.1038/srep17048.

Moore JK. The human auditory brain stem: a comparative view. Hearing research. 1987; 29:1–32. doi: 10.1016/0378-5955(87)90202-4.

Moore JK, Perazzo LM, Braun A. Time course of axonal myelination in the human brainstem auditory pathway. Hearing research. 1995 jul; 87(1-2):21–31. http://www.ncbi.nlm.nih.gov/pubmed/8567438, doi: 10.1016/0378-5955(95)00218-9.

Mori S, Crain BJ, Chacko VP, van Zijl PC. Three-dimensional tracking of axonal projections in the brain by magnetic resonance imaging. Annals of neurology. 1999 feb; 45(2):265–9. http://www.ncbi.nlm.nih.gov/pubmed/9989633, doi: 10.1002/1531-8249(199902)45:2<265::AID-ANA21>3.0.CO;2-3.

Paxinos G, T F, C W. Human Brainstem: Cytoarchitecture, Chemoarchitecture, Myeloarchitecture. 1 ed. Acedemic Press; 2019.

Pfefferbaum A, Sullivan EV, Adalsteinsson E, Garrick T, Harper C. Postmortem MR imaging of formalin-fixed human brain. NeuroImage. 2004 apr; 21(4):1585–95. http://www.ncbi.nlm.nih.gov/pubmed/15050582, doi: 10.1016/j.neuroimage.2003.11.024.

Reveley C, Seth AK, Pierpaoli C, Silva AC, Yu D, Saunders RC, Leopold DA, Ye FQ. Superficial white matter fiber systems impede detection of long-range cortical connections in diffusion MR tractography. Proceedings of the National Academy of Sciences of the United States of America. 2015 may; 112(21):E2820–8. http://www.ncbi.nlm.nih.gov/pubmed/25964365http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4450402, doi: 10.1073/pnas.1418198112.

Rose J, Galambos R, Hughes J. Organization of frequency sensitive neurons in the cochlear nuclear complex of the cat. In: *Neural Mechanisms of the Auditory and Vestibular Systems* Thomas Springfield, IL; 1960.p. 116–136. http://scholar.google.com/scholar?hl=en{&}btnG=Search{&}q=intitle:Organization+of+frequency+sensitive+neurons+in+the+cochlear+nuclear+complex+of+the+cat{#}0.

Ryugo DK, May SK. The projections of intracellularly labeled auditory nerve fibers to the dorsal cochlear nucleus of cats. The Journal of comparative neurology. 1993 mar; 329(1):20–35. http://www.ncbi.nlm.nih.gov/pubmed/8454724, doi: 10.1002/cne.903290103.

Ryugo DK, Parks TN. Primary innervation of the avian and mammalian cochlear nucleus. Brain Research Bulletin. 2003; 60(5-6):435–456. doi: 10.1016/S0361-9230(03)00049-2.

Sando I. The Anatomical Interrelationships of the Cochlear Nerve Fibers. Acta Oto-Laryngologica. 1965 jan; 59(2-6):417–436. http://www.tandfonline.com/doi/full/10.3109/00016486509124577, doi: 10.3109/00016486509124577.

Schilling KG, Daducci A, Maier-Hein K, Poupon C, Houde JC, Nath V, Anderson AW, Landman BA, Descoteaux M. Challenges in diffusion MRI tractography - Lessons learned from international benchmark competitions. Magnetic resonance imaging. 2019 apr; 57:194–209. http://www.ncbi.nlm.nih.gov/pubmed/30503948http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6331218, doi: 10.1016/j.mri.2018.11.014.

Schilling KG, Nath V, Hansen C, Parvathaneni P, Blaber J, Gao Y, Neher P, Aydogan DB, Shi Y, Ocampo-Pineda M, Schiavi S, Daducci A, Girard G, Barakovic M, Rafael-Patino J, Romascano D, Rensonnet G, Pizzolato M, Bates A, Fischi E, et al. Limits to anatomical accuracy of diffusion tractography using modern approaches. NeuroImage. 2019; doi: 10.1016/j.neuroimage.2018.10.029.

Schofield BR. Structural organization of the descending auditory pathway. The Auditory Brain. 2010; p. 43–64.

Setsompop K, Gagoski BA, Polimeni JR, Witzel T, Wedeen VJ, Wald LL. Blipped-controlled aliasing in par-allel imaging for simultaneous multislice echo planar imaging with reduced g-factor penalty. Magnetic resonance in medicine. 2012 may; 67(5):1210–24. http://www.ncbi.nlm.nih.gov/pubmed/21858868http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3323676, doi: 10.1002/mrm.23097.

Sigalovsky IS, Melcher JR. Effects of sound level on fMRI activation in human brainstem, thalamic and cortical centers. Hearing research. 2006 may; 215(1-2):67–76. http://www.ncbi.nlm.nih.gov/pubmed/16644153http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC1794213, doi: 10.1016/j.heares.2006.03.002.

Sotiropoulos SN, Jbabdi S, Xu J, Andersson JL, Moeller S, Auerbach EJ, Glasser MF, Hernandez M, Sapiro G, Jenkinson M, Feinberg DA, Yacoub E, Lenglet C, Van Essen DC, Ugurbil K, Behrens TEJ. Advances in diffusion MRI acquisition and processing in the Human Connectome Project. NeuroImage. 2013; doi: 10.1016/j.neuroimage.2013.05.057.

Sparks DL, Hartwich-Young R. The deep layers of the superior colliculus. Reviews of oculomotor research. 1989; 3:213–55. http://www.ncbi.nlm.nih.gov/pubmed/2486324.

Strominger NL, Hurwitz JL. Anatomical aspects of the superior olivary complex. The Journal of comparative neu-rology. 1976 dec; 170(4):485–97. http://www.ncbi.nlm.nih.gov/pubmed/826550, doi: 10.1002/cne.901700407.

Teeuwisse WM, Brink WM, Webb AG. Quantitative assessment of the effects of high-permittivity pads in 7 tesla MRI of the brain. Magnetic resonance in medicine. 2012; 67(5):1285–1293.

Tourdias T, Saranathan M, Levesque IR, Su J, Rutt BK. Visualization of intra-thalamic nuclei with opti-mized white-matter-nulled MPRAGE at 7T. NeuroImage. 2014 jan; 84:534–45. http://www.ncbi.nlm.nih.gov/pubmed/24018302http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3927795, doi: 10.1016/j.neuroimage.2013.08.069.

Tournier JD, Calamante F, Connelly A. Robust determination of the fibre orientation distribution in diffu-sion MRI: Non-negativity constrained super-resolved spherical deconvolution. NeuroImage. 2007; doi: 10.1016/j.neuroimage.2007.02.016.

Tournier JD, Calamante F, Gadian DG, Connelly A. Direct estimation of the fiber orientation density function from diffusion-weighted MRI data using spherical deconvolution. NeuroImage. 2004; doi: 10.1016/j.neuroimage.2004.07.037.

Triantafyllou C, Polimeni JR, Keil B, Wald LL. Coil-to-coil physiological noise correlations and their impact on functional MRI time-series signal-to-noise ratio. Magnetic Resonance in Medicine. 2016 dec; 76(6):1708–1719. http://doi.wiley.com/10.1002/mrm.26041, doi: 10.1002/mrm.26041.

Ugurbil K. What is feasible with imaging human brain function and connectivity using functional mag-netic resonance imaging. Philosophical Transactions of the Royal Society B: Biological Sciences. 2016 oct; 371(1705):20150361. http://rstb.royalsocietypublishing.org/lookup/doi/10.1098/rstb.2015.0361, doi: 10.1098/rstb.2015.0361.

Ugurbil K, Adriany G, Andersen P, Chen W, Garwood M, Gruetter R, Henry PG, Kim SG, Lieu H, Tkac I, Vaughan T, Van De Moortele PF, Yacoub E, Zhu XH. Ultrahigh field magnetic resonance imaging and spectroscopy. Magnetic Resonance Imaging. 2003 dec; 21(10):1263–1281. https://linkinghub.elsevier.com/retrieve/pii/S0730725X03003382, doi: 10.1016/j.mri.2003.08.027.

Van de Moortele PF, Auerbach EJ, Olman C, Yacoub E, Uğurbil K, Moeller S. T1 weighted brain images at 7 Tesla unbiased for Proton Density, T2* contrast and RF coil receive B1 sensitivity with simultaneous vessel visualization. NeuroImage. 2009 jun; 46(2):432–46. doi: 10.1016/j.neuroimage.2009.02.009.

Van Der Walt S, Colbert SC, Varoquaux G. The NumPy array: a structure for efficient numerical computation. Computing in Science & Engineering. 2011; 13(2):22–30.

Vaughan JT, Garwood M, Collins CM, Liu W, DelaBarre L, Adriany G, Andersen P, Merkle H, Goebel R, Smith MB, Ugurbil K. 7T vs. 4T: RF power, homogeneity, and signal-to-noise comparison in head images. Magnetic resonance in medicine. 2001 jul; 46(1):24–30. http://www.ncbi.nlm.nih.gov/pubmed/11443707, doi: 10.1002/mrm.1156.

Vu AT, Auerbach E, Lenglet C, Moeller S, Sotiropoulos SN, Jbabdi S, Andersson J, Yacoub E, Ugurbil K. High resolution whole brain diffusion imaging at 7T for the Human Connectome Project. NeuroImage. 2015; doi: 10.1016/j.neuroimage.2015.08.004.

Wang N, Anderson RJ, Badea A, Cofer G, Dibb R, Qi Y, Johnson GA. Whole mouse brain structural connectomics using magnetic resonance histology. Brain structure & function. 2018 dec; 223(9):4323–4335. http://www.ncbi.nlm.nih.gov/pubmed/30225830, doi: 10.1007/s00429-018-1750-x.

Wang R, Benner T, Sorensen AG, Wedeen VJ. Diffusion toolkit: a software package for diffusion imaging data processing and tractography. In: *Proc Intl Soc Mag Reson Med* Berlin; 2007. p. 3720.

Webster DB. An Overview of Mammalian Auditory Pathways with an Emphasis on Humans. In: *Handbook of Auditory Research* Springer; 1992.p. 1–22. http://link.springer.com/10.1007/978-1-4612-4416-5{_}1, doi: 10.1007/978-1-4612-4416-5_1.

Winer Ja. The human medial geniculate body. Hearing Research. 1984; 15(3):225–247. doi: 10.1016/0378-5955(84)90031-5.

Yushkevich PA, Piven J, Hazlett HC, Smith RG, Ho S, Gee JC, Gerig G. User-guided 3D active contour segmentation of anatomical structures: significantly improved efficiency and reliability. NeuroImage. 2006 jul; 31(3):1116–28. doi: 10.1016/j.neuroimage.2006.01.015.

1083 **Appendix 1**

1084

1085 **Appendix 1 Figure 1.** Post mortem human diffusion-weighted MRI tractography (from 200 μm
1086 isotropic voxels) with anatomically defined subcortical auditory seeds, dilated 500 μm.

Appendix 1 Figure 2. Post mortem human diffusion-weighted MRI tractography with anatomically defined subcortical auditory seeds. MRI data were downsampled from 200 μm to 1050 μm to match in vivo data acquisition and then processed in the same manner as other diffusion tractography analyses

Appendix 1 Figure 3. Diffusion-weighted MRI tractography streamlines passing through each subcortical auditory region of interest for the ten in vivo participants.

Appendix 1 Figure 4. In vivo anatomical group average images.

Appendix 1 Figure 5. Anatomical images from MNI ICBM 152 dataset compared to BigBrain histology in MNI152 space (left column).

Appendix 1 Figure 6. Summary of data processing steps, including availability of data and code. This figure will be updated as resources as shared online.