Pacing behaviour in laboratory macaques is an unreliable indicator of acute stress.

Colline Poirier[1*], Caitlin J. Oliver[1], Paul Flecknell[2] and Melissa Bateson[1]

1 Institute of Neuroscience and Centre for Behaviour and Evolution, Newcastle University

2 Comparative Biology Centre, Newcastle University

*Corresponding author: Colline Poirier

Institute of Neuroscience

Newcastle University

Framlington Place,

Newcastle upon Tyne

NE2 4HH

UK

Tel: +44 (0)191 222 3445

Fax: +44 (0)191 208 5731

colline.poirier@ncl.ac.uk

Pacing behaviour, the most frequent stereotypic behaviour displayed by laboratory rhesus macaques

(*Macaca mulatta*) is often used as an indicator of acute stress. In this study, we investigated how

reliable this welfare indicator is by testing the reaction of macaques to the stressful event of being

exposed to a fight between conspecifics housed in the same room but in a different cage. Pacing,

agitated locomotion, and stress-related displacement behaviours were quantified before, during and

after fight exposure, based on video recordings of 13 socially-housed macaques in their home cage.

Displacement behaviours increased after fight exposure, confirming that the events were experienced

as stressful by the focal individuals. The occurrence of pacing did not increase during or after the

fights. Instead, agitated locomotion increased during the fights. These results suggest either, that

pacing as an indicator of acute stress is prone to false negative results, increasing in some stressful

situations but not others, or that agitated locomotion has been mistaken for pacing in previous studies

and that pacing is in fact unrelated to current acute stress. Both interpretations lead to the conclusion

that pacing is unreliable as an indicator of acute stress in laboratory rhesus macaques.

Of the laboratory animal species commonly used in biomedical research, non-human primates (NHPs)

are phylogenetically closest to humans. This proximity makes NHPs a crucial animal model in

biomedical science, but also makes their welfare of particular concern to the general public. Current

regulations reflect this public concern[1,2], and require that researchers ensure the best possible welfare

for the animals that they use. To achieve this goal, reliable welfare indicators are necessary.

Stereotypies, which are repetitive, unvarying and apparently functionless behaviours, are often used as

indicators of acute or chronic stress[3]. In rhesus macaques, one of the main NHP models used in

research, pacing is the most common stereotypic behaviour observed in the laboratory[4]. In this study

we sought to establish specifically whether pacing is a reliable indicator of acute stress in this species.

Pacing in rhesus macaques (hereafter macaques) is defined as repetitive walking of an

individual in the exact same pattern[4,5]. Previous studies have reported that macaques increase their

pacing during or immediately after short periods of acute stress (e.g. placing in a transport box for few

seconds, exposure to a 10-minute cue of impending social isolation)[6,7]. Injecting macaques with a high

dose of the anxiogenic drug FG7142 also increases pacing[8]. An increase in pacing frequency is thus

usually interpreted as an increase in acute stress[9,10], where the term 'stress' is used to refer to an

unpleasant, arousing affective state.

However, in a recent study where macaques were submitted to the human intruder test, a

classical test to induce acute stress, pacing frequency was found to decrease during the stressful

event[11]. This result suggests that pacing can increase or decrease with acute stress, depending on the

stressful situation. This finding is important from a welfare perspective, because if the frequency of a

behaviour can increase in some stressful situations and decrease in others, variation in the behaviour

in a new situation (whose stressful nature is unknown) cannot be interpreted unequivocally; in other

words, the behaviour is unreliable as an indicator of acute stress.

In the current study we sought to determine whether the behavioural response to the human

intruder test is an exceptional case, or whether other types of stressors also induce a decrease in

pacing. To do this, we needed to identify an alternative source of acute stress that occurs relatively

frequently in laboratory macaques and hence had the potential to explain the prevalence of pacing in

primate research facilities. We chose to investigate the effect of passive exposure to fights between

conspecifics on the assumption that this is stressful. Agonistic interactions in macaques are used to

challenge or reassert the social position of protagonists and can induce severe injuries. In free-ranging

macaque colonies, agonistic support (where one or several individuals help one of the protagonists) is

frequent[12]. Hence, witnessing a fight implies potential imminent involvement in a stressful situation,

which is likely to be stressful in itself. Exposure to fights has ecological relevance, since it occurs in

research colonies where individuals are housed in pairs or small social groups, which is now the norm

in European primate research facilities and an emerging trend in North America. Guided by ethical

considerations, we made use of fights that occurred spontaneously in our research colony. We

measured the occurrence of pacing before and during passive exposure to fights. In the recent intruder

test study[11], the authors did not include behavioural observation of macaques after exposure to the

acute stressor had ceased. It is thus possible that an increase of pacing frequency happened

subsequently and was not recorded. To control for the possibility that the expected increase in pacing

could be delayed, we also measured the occurrence of pacing after exposure to the fights.

Results

Fights

Video footage of home-cage behaviour of 13 focal individuals previously recorded for another

purpose was used for this project. Fights were defined as any period lasting at least 10 s during which

threatening/alarm vocalisations associated with loud object banging noises were heard. This

identification was made based on the soundtrack of the video recordings with the video images

concealed in order to avoid any selection bias caused by the behaviour of the focal individuals.

Fourteen fights were identified. Analyses of video images confirmed that in all but one case the focal

individuals were not involved in these fights. Data related to that fight were discarded. Video

recordings of each focal individual were not always available for each fight. The number of fights for

which video existed varied between 4 and 12 fights per individual, resulting in a total of 108

behavioural responses to fights in which the focal individual was not involved.

Pacing

Behaviour of focal individuals was scored in three time intervals relative to the fight: during each

fight (the time interval designated as [0]), during the 15 min preceding the fight (designated [-15 – 0])

and during the 15 min following the end of the fight (designated [0 – 15]). The three time intervals

were divided into 10-s time bins and the presence (1) or absence (0) of pacing within each time bin

was recorded (for the operational definition of pacing, see table 1). Pacing behaviour was displayed

by 7 out of 13 individuals. Among pacing individuals, occurrence of pacing varied from 0.7 to 46.7 %

(percentage of 10-s time bins in which pacing occurred).

102   Generalised linear mixed models of pacing data revealed a significant effect of time interval

103   relative to fight (Fig. 1; Table 1, model 1 vs. model 0, χ^2 (2) = 11.2, P = 0.004), driven by a decrease

104   in occurrence of pacing after the fights, compared to before (Table 2, model 1: β (s.e.) = -0.24 (0.07),

105   Z = -3.4, P < 0.001). There was no difference in occurrence of pacing before and during the fights

106   (Table 2, model 1: β (s.e.) < 0.01 (0.34), Z = 0.005, P > 0.99).

107   Exposure to a stressful event induces an immediate increase in catecholamines and a slower

increase in cortisol, associated with behavioural responses occurring at different time scales[13–15]. Our

previous analyses focused on testing the hypothesis that the fast released cathecholamines would

induce an increase of pacing. However, if an increase of pacing is triggered by cortisol, one would

expect it to take place 20 to 30 minutes after the stressful event, when cortisol is at its peak. To take

into account this possibility, we subsequently analysed additional videos of pacing individuals

corresponding to the time interval [15 – 40] min post-fight. The analysis of the dataset including the

four time intervals ([-15 – 0], [0], [0 – 15] and [15 – 40]) revealed a significant effect of time interval

(Table 2, model 2 vs. model 0, χ^2 (3) = 93.9, P < 0.001) due to a decrease in occurrence of pacing

during the two time intervals after the fights, compared to before (Table 2, model 2: [0 – 15] vs. [-15 –

0]: β (s.e.) = -0.23 (0.07), Z = -3.3, P < 0.001; [15 – 40] vs. [-15 – 0]: β (s.e.) = -0.61 (0.07), Z = -9.5,

P < 0.001).

119   Finally, we tested the possibility that the decrease in pacing might be due to an effect of

absolute time of the day, because we have unpublished data suggesting that occurrence of pacing

follows a circadian rhythm. Despite the fact that our analyses were restricted to a narrow time window

of 55 min, circadian changes might still potentially confound the effect of time intervals relative to

fight. Including time of day as an additional fixed effect significantly improved model fit (Table 2,

model 3 vs. model 2, χ^2 (1) = 5.9, P = 0.02). Occurrence of pacing was found to decrease with time of

day (Table 2, model 3: β (s.e.) = -14.84 (6.04), Z = -2.46, P = 0.01) and the effect of time interval

relative to fight in the new model was no longer significant (Table 2, model 4 vs. model 3: χ^2 (3) =

2.9, P = 0.41).

Agitated locomotion

In addition to pacing behaviour, we also assessed the frequency of agitated locomotion, a behaviour

defined as 'moving rapidly between locations, often via jumps, with a stiff un-relaxed gait' (Table 1).

In our macaque facility where individuals are housed in large and enriched cages, this behaviour

differs from pacing not only in the gait of the subject (which is stiff in agitated locomotion and elastic

in pacing) but also in the flexibility of the path used within and between occurrences, making the two

behaviours easy to distinguish (the inter-observer reliability measured with Kappa scores were of 1

for pacing and 0.97 for agitated locomotion in the present study). However, it is possible that when

macaques are housed in small, barren cages, the two behaviours could be conflated. We thus

hypothesised *a priori* that agitated locomotion might be the behaviour increasing during or after

exposure to an acute stressor.

Presence (1) or absence (0) of agitated locomotion within each time bin was recorded and was

displayed by 11 out of 13 individuals, including all pacers (N = 7) and 4 non-pacers. The occurrence

of agitated locomotion among individuals that displayed the behaviour at least once varied from 0.03

to 1.1 %. Occurrence of agitated locomotion displayed by each individual and occurrence of pacing

were not correlated (r = 0.002, t (11) < 0.01, P > 0.99).

Generalised linear mixed models of the occurrence of agitated locomotion revealed a significant

effect of time interval relative to fight (Table 2, model 6 vs. model 5: χ^2 (3) = 149, P < 0.001), driven

by an increase in agitated locomotion during fights, compared to before (Table 2, model 6: [0] vs. [-15

− 0]: β (s.e.) = 4.1 (0.28), Z = 14.4, P < 0.001). There was no difference in occurrence of agitated

locomotion before and after the fights ([0 − 15] vs. [-15 − 0]: β (s.e.) = -0.48 (0.28), Z = -1.69, P =

0.09; [15 − 40] vs. [-15 − 0]: β (s.e.) = 0.12 (0.24), Z = 0.48, P = 0.63). Generalised linear mixed

models applied separately to pacers and non-pacers revealed similar effects of time interval relative to

fight in both groups (Fig. 2, pacers: χ^2 (3) = 120, P < 0.001, [0] vs. [-15 – 0]: β (s.e.) = 4.28 (0.344), Z

= 12.7, P < 0.001; non-pacers: χ^2 (3) = 34, P < 0.001, [0] vs. [-15 – 0]: β (s.e.) = 3.57 (0.60), Z = 5.9,

P < 0.001). Adding the absolute time of day as an additional fixed effect did not improve model fit

(Table 2, model 7 vs. model 6: χ^2 (1) = 0.5, P = 0.49).

Displacement behaviours

To verify that fight exposure was experienced as stressful by the macaques, we recorded and analysed

the occurrence of displacement behaviours[16] displayed by macaques, namely scratching, body shaking

and self-grooming (Table 1). These behaviours were chosen based on their pharmacological

validation as indicators of stress or anxiety by Schino and colleagues[17]. Following these authors

approach, scratching, body shaking and self-grooming were combined to create a single category of

'displacement behaviours'. Presence of at least one of the behaviours (1) or absence of all of them (0)

within each time bin was recorded.

Displacement behaviours were displayed by all individuals, with occurrence varying between 5.1

and 26.9 %. Generalised linear mixed models of occurrence of displacement behaviours revealed an

effect of time interval relative to fight (Table 2, model 9 vs. model 8: χ^2 (3) = 43.0, P < 0.001), driven

by an increase in displacement behaviour frequency in the second time interval after the fight,

compared to before ([15 – 40] vs. [-15 – 0]: β (s.e.) = 0.34 (0.06), Z = 6.0, P < 0.001). Adding the

absolute time of the day as an additional fixed effect did not improve model fit (Table 2, model 10 vs.

model 9: χ^2 (1) = 1.5, P = 0.2).

Discussion

This study investigated the behavioural reaction of laboratory rhesus macaques to fights occurring

between conspecifics housed in separate, but nearby cages within the same room. Displacement

behaviours, which are pharmacologically validated indicators of stress or anxiety, increased after the

fights, confirming that fights were perceived as stressful by the macaques. Our study reveals that

stereotypic pacing did not increase during or after the fights. Peterson and colleagues previously

showed that pacing did not increase during an intruder test, a paradigm established to induce stress in

macaques[11]. Here we extend this result by testing a different stressful situation (passive exposure to a

fight between conspecifics) and by providing a more comprehensive assessment of pacing by testing

its frequency not only during the stressful event but also during a sustained period of time (up to 40

min) after the stressful event. These data contrast with previous studies where pacing was found to

increase with acute stress[6,7]. These conflicting results can be interpreted in two different ways that we

describe in the next paragraphs.

The first possible explanation for the discrepancy in results from different studies is that

pacing frequency increases in some stressful situations but not in others. If this is the case, it means

that pacing when used as an indicator of acute stress is prone to false negative results, where

macaques experience acute stress but do not express it by an increase in pacing (Fig. 3). From a

welfare perspective, this possibility is problematic for two reasons. First, using pacing as an indicator

of acute stress might lead stressful husbandry or experimental procedures to fail to be identified as

such, preventing researchers, veterinarians and/or technical staff from refining them and thus

improving animal welfare. Second, false-negative results also undermine the use of an absence of

pacing frequency increase during or after a procedure as a reliable indicator that the procedure is not

stressful. This is problematic because determining which husbandry and scientific procedures are not

experienced as stressful is as important as identifying those which are stressful. It should also be noted

that whether pacing increases after positively-valenced (i.e. pleasant) arousing events has never been

investigated. The potential of pacing to generate false positive results (Fig. 3) when used as an

indicator of acute stress is thus currently unknown. As a consequence, when pacing frequency is

found to be increased by an event or procedure, one cannot conclude with confidence that this event

or procedure induced acute stress rather than a pleasant arousing emotion. Because optimising the

welfare of animals means reducing negative emotions but also improving positive ones, attempts to

reduce pacing frequency could be counter-productive if increased pacing has been caused by

positively-valenced arousal.

Alternatively, the fact that pacing seems to increase during some stressful situations[6,7], but not

others (ref 11, our study), might be due to the possible conflation between pacing and agitated

locomotion in previous studies. Indeed, our study revealed an increase in agitated locomotion during

fights. Differences between pacing and agitated locomotion are summarised in Table 1. One notable

difference is that agitated locomotion is not stereotypic, the path used by an individual varying from

one occurrence to another. In our research facility where macaques are housed in big cages (15 m^3)

with shelves and ropes, the two behaviours are easy to distinguish. However, it is possible that in

small cages agitated locomotion appears stereotypic due to the lack of options in the paths individuals

can walk, making it visually indistinguishable from true pacing. Interestingly, studies reporting an

increase in pacing frequency induced by acute stress, including the anxiogenic drug FG7142[6–8], all

come from macaques housed in very small cages (below 1.3 m^3). It is thus possible that existing

literature linking acute stress and pacing should be reinterpreted as showing a link between acute

stress and agitated locomotion. If this is the case, pacing would never be related to present acute

stress.

In some individuals, occurrence of pacing was found to be high before, during and after the fights

(fig. 1B). Source(s) of acute stress during the whole period could not be identified. This result

indicates that pacing is likely to be caused by factor(s) other than current acute stress. We have

recently reviewed these potential factors[18]. Our review suggests that pacing might be the consequence

of brain abnormalities induced by a past stressful event, and be unrelated to the current state of the

individual. Alternatively, pacing might be caused by chronic stress, boredom or a specific need to

walk. These different hypotheses have different welfare consequences with the welfare of pacing

macaques being either better, worse or equivalent to that of non-pacing individuals[18]. As long as the

cause(s) of pacing remain unresolved, we argue that pacing should not be used as an indicator of acute

or chronic stress, especially in the absence of other indicators of poor welfare.

Our results raise the possibility that agitated locomotion could be a useful indicator of acute

stress. In sharp contrast with pacing behaviour, agitated locomotion increased during the fights but

was virtually non-existent outside periods of fights, when no source of acute stress could be detected.

These results however need to be replicated with other stressful situations. If agitated locomotion is

systematically observed during different stressful situations (i.e. if no/few false negative results are

observed), this behaviour might be a convenient welfare marker since it is easy to identify (at least in

large, enriched cages) and its baseline frequency near to zero makes any increase easy to detect.

Whether agitated locomotion also increases during positively-valenced arousing events (susceptibility

to false positive results) will also need to be determined.

To our knowledge, the precise time scale of behavioural indicators of acute stress in macaques has

never been systematically investigated. From an applied perspective, this information might help in

designing better behavioural observation protocols. At the physiological level, acute stress induces a

well-established cascade of events, including the release of different hormones over different time

scales. In the present study, we used this knowledge to guide our observation periods. Agitated

locomotion increased during the very brief period of fights, suggesting that this behaviour might be

triggered by the fast-released catecholamines. By contrast, displacement behaviours increase was

observed during the 15 - 40 minute interval after the stressor, suggesting that these behaviours might

be induced by the slowly-released cortisol. Displacement behaviours, however, correspond to a

combination of different behaviours (scratching, self-grooming and body shaking). Further studies

will need to determine whether the exact timing of these three behaviours is similar or not.

To conclude, by showing that pacing does not increase following exposure to conspecific

fights this study shows that pacing behaviour is not a reliable indicator of acute stress. Our data also

suggest that agitated locomotion might potentially be a useful indicator of acute stress, or at least

arousal, but more studies are needed to confirm this. Welfare research on captive animals is often

done using experimental paradigms that involve manipulating the affective state of the animals by

imposing a stressor[19,20]. While this approach is undoubtedly the most powerful, the current study

illustrates how new knowledge can also be gained by using observations of animals without inducing

additional stress.

Materials and methods

Subjects and ethical statement

Subjects were 13 male rhesus macaques (*Macaca mulatta*) aged between 4 and 10 years (weight

range: 5 to 18 kg). Individuals had been raised in a British breeding centre, first with their mother for

at least 6 months and then with other juveniles in large rooms. When adolescent, subjects were moved

to the Newcastle University research facility, which complies with the NC3Rs Guidelines for

'Primate Accommodation, care and use' (www.nc3rs.org.uk/primatesguidelines). There, they were

housed in small social groups (2 to 3 individuals) in large cages (2.1 x 3.0 x 2.4 m) exceeding the

minimal space requirement under the UK legislation (1.8 m^3 per animal). The height of the cages and

the presence of high perches and shelves allowed macaques, if alarmed, to flee upwards. Besides the

presence of at least one cage mate, enrichment was provided by daily foraging opportunity (the food

was scattered in the litter on the floor), wooden shelves, swings, ropes, objects to manipulate (changed

on a regular basis) and natural light. Cages were located in a large room housing more than 40

individuals. Focal individuals were thus in visual and auditory contact with many other conspecifics

in addition to their cage mates. Some of the individuals were routinely involved in neuroscience

experiments while others had not yet been subjected to any experimental procedures. This study

consisted of behavioural observation of macaque home cage behaviour recorded via remotely-

controlled cameras, inducing no interference with the animals. Females were not included in this

study for the sole reason that video recordings of these individuals were not available in sufficient

quantity. The stressor consisted of passive exposure to fights occurring naturally between other

animals; the frequency of fights was not manipulated for this study. No licence was required for this

study.

Behavioural observation

Videos were recorded with remotely-controlled cameras (Cube HD 1080, Y-cam Solutions Limited,

Twickenham, UK). Recordings from between 1600 and 1800 on Saturdays and Sundays between

January and December 2015 were analysed. This period of the week was chosen as it was the only

period when macaques were in their home cage, awake and no people were present in the facility.

This strategy avoided the necessity of having to control for the potentially stressful effects of human

presence and cage mate absence.

Behavioural scoring and analysis

Statistical analyses were performed in R, version 3.3.2[21] using the *lme4* package[22]. Data were

analysed using generalised linear mixed models (*glmer* function) with binomial error distribution and

logistic link function. The *time interval relative to fight* was declared as a fixed effect (categorical

effect with 3 or 4 levels), while *subjects* and *fights* were declared as random effects, with *fights* being

nested within *subjects*. Maximum-likelihood estimation was employed throughout. Note that these

models cope with highly unbalanced designs such as this one (the amount of data in the [0] interval

being much less than in the other time intervals) and non-independence due to repeated measures of

subjects and fights. Significance testing was carried out by the likelihood ratio test, which compares

the change in deviance when a term is excluded from the model with the χ^2 distribution.

Data availability

The datasets generated during and/or analysed during the current study are available from the

corresponding author on reasonable request.

Acknowledgements

This work was supported by the Association for the Study of Animal Behaviour, the NC3Rs

(NC/K000802/1) and internal funding from Newcastle University. The funders had no role in the

writing of the report nor in the decision to submit the article for publication.

Author contributions statement

Funding acquisition: C.P., P.F., M.B; Conceptualisation and Methodology: C.P and M.B.; Resources:

P.F.; Investigation: C.P. and C.O.J.; Formal analysis: C.P.; Supervision: C.P. and M.B.; Visualisation:

C.P.; Writing – original draft: C. P.; Writing – review and editing: C.P., C.O.J, P.F., M.B.

Competing interests

The authors declare no competing interests.

References

1.   United States Department of Agriculture. Animal Welfare Act and Animal Welfare

320   Regulations. Chapter 45 (2013). doi:10.2307/4443437

321   2.   European Union. *Directive 2010/63/EU of the european parliament and of the council of 22*

322   *September 2010 on the protection of animals used for scientific purposes.* (2010).

323   3.   Mason, G. J. Stereotypies : a critical review. *Anim. Behav.* **41,** 1015–1037 (1991).

324   4.   Lutz, C., Well, A. & Novak, M. Stereotypic and self-injurious behavior in rhesus macaques: A

325   survey and retrospective analysis of environment and early experience. *Am. J. Primatol.* **60,** 1–

326   15 (2003).

327   5.   Gottlieb, D. H., Capitanio, J. P. & Mccowan, B. Risk Factors for Stereotypic Behavior and

328   Self   Biting in Rhesus Macaques (Macaca mulatta): Animal ' s History , Current

329   Environment , and Personality. *Am. J. Primatol.* **75,** 995–1008 (2013).

330   6.   Mitchell, G. & Gomber, J. Moving laboratory rhesus monkeys (Macaca mulatta) to unfamiliar

331   home cages. *Primates* **17,** 543–546 (1976).

332   7.   Willott, J. F. & McDaniel, J. Changes in the behavior of laboratory-reared rhesus monkeys

333   following the threat of separation. *Primates* **15,** 321–326 (1974).

334   8.   Major, C. A. *et al.* The anxiogenic drug FG7142 increases self-injurious behavior in male

335   rhesus monkeys (Macaca mulatta). *Life Sci.* **85,** 753–758 (2009).

336   9.   Waitt, C. & Buchanan-Smith, H. M. What time is feeding?: How delays and anticipation of

337   feeding schedules affect stump-tailed macaque behavior. *Appl. Anim. Behav. Sci.* **75,** 75–85

338   (2001).

339   10.   Gottlieb, D. H., Coleman, K. & McCowan, B. The Effects of Predictability in Daily

340   Husbandry Routines on Captive Rhesus Macaques (Macaca mulatta). *Appl. Anim. Behav. Sci.*

341   **143,** 117–127 (2013).

342   11.   Peterson, E. J. *et al.* Rhesus macaques (Macaca mulatta) with self-injurious behavior show

343   less behavioral anxiety during the human intruder test. *Am. J. Primatol.* **79,** e22569 (2017).

344   12.   Maestripieri, D. & Hoffman, C. L. in *Bones, Genetics, and Behavior of Rhesus Macaques* (ed.

345   Q, W.) 247–262 (Springer, 2012). doi:10.1007/978-1-4614-1046-1

346   13.   Joëls, M., Fernandez, G. & Roozendaal, B. Stress and emotional memory: A matter of timing.

347   *Trends Cogn. Sci.* **15,** 280–288 (2011).

348   14.   Koelsch, S. *et al.* The impact of acute stress on hormones and cytokines, and how their

349   recovery is affected by music-evoked positive mood. *Sci. Rep.* **6,** 23008 (2016).

350   15.   Hermans, E. J., Henckens, M. J. A. G., Joels, M. & Fernandez, G. Dynamic adaptation of

351   large-scale brain networks in response to acute stressors. *Trends Neurosci.* **37,** 304–314

352   (2014).

353   16.   Maestripieri, D., Schino, G., Aureli, F. & Troisi, A. A modest proposal: displacement activities

354   as an indicator of emotions in primates. *Anim. Behav.* **44,** 967–979 (1992).

355   17.   Schino, G., Perretta, G., Taglioni, A. M., Monaco, V. & Troisi, A. Primate Displacement

356   Activities As an Ethopharmacological Model of Anxiety. *Anim. Behav.* **44,** 186–191 (1996).

357   18.   Poirier, C. & Bateson, M. Pacing stereotypies in laboratory rhesus macaques: implications for

358   animal welfare and the validity of neuroscientific findings. *Neurosci. Biobehav. Rev.* **83,** 508–

359   515 (2017).

360   19.   Edgar, J. *et al.* Social buffering in a bird. *Anim. Behav.* **105,** 11–19 (2015).

361   20.   White, L. J., Thomson, J. S., Pounder, K. C., Coleman, R. C. & Sneddon, L. U. The impact of

362   social context on behaviour and the recovery from welfare challenges in zebrafish, Danio

363   rerio. *Anim. Behav.* **132,** 189–199 (2017).

364   21.   R Core Team. A language and environment for statistical computing. (2017).

365   22.   Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting Linear Mixed-Effects Models Using

366   lme4. *J. Stat. Softw.* **67,** 1–48 (2015).

Figure captions

Figure 1: Occurrence of pacing as a function of the time interval relative to fight. Occurrence of

pacing corresponds to the proportion of 10-s time bins when the behaviour was displayed (with 1

corresponding to 100 %). The dotted line indicates the 'during fight' interval. a: Data averaged over

fights and subjects. b: Data averaged over fights. c: Data per fight and per subject. Note that points

appear darker when data from several fights are superimposed. The n value indicated in the corner of

each individual plot corresponds to the number of fights. For the legend of the x axis, see plots a and

b.

Figure 2: Occurrence of agitated locomotion as a function of the time interval relative to fight in

pacers (a-c) and non-pacers (d-f). Occurrence of agitated locomotion corresponds to the proportion of

time bins when the behaviour was displayed (with 1 corresponding to 100 %). The dotted line

indicates the 'during fight' interval. a/d: Data averaged over fights and subjects. b/e: Data averaged

over fights. c/f: Data per fight and per subject. Note that points appear darker when data from several

fights are superimposed. The n value indicated on each plot corresponds to the number of fights. For

the legend of the x axis, see plots a/d and b/e.

Figure 3: Different possible relationships between pacing frequency and acute stress.

Table 1: Ethogram

Behaviour	Description
Pacing	Repetitive walking of the same path (at least twice)
Agitated Locomotion	Moving fast between locations with a stiff un-relaxed gait
Self-grooming	Stroking, picking, or otherwise manipulating own body surface
Body shaking	Dog-like shake of whole body
Self-scratching	Scratching the skin vigorously with nails

392 Table 2: Generalised linear mixed models

Dependent variable	Model	Fixed predictors	Degrees of freedom
Pacing	0	none	3
	1	Time interval relative to fights (3 levels)	5
	2	Time interval relative to fights (4 levels)	6
	3	Time interval relative to fights (4 levels); Time of day	7

Agitated locomotion	4	Time of day	4
	5	none	3
	6	Time interval relative to fights (4 levels)	6
	7	Time interval relative to fights (4 levels); Time of day	7
Displacement behaviours	8	none	3
	9	Time interval relative to fights (4 levels)	6
	10	Time interval relative to fights (4 levels); Time of day	7

393 All models include random effects for focal subject and fight to control for non-independence arising from
394 repeated measures on subjects and fights.

397 Table 3: Differences between Pacing and Agitated Locomotion

	Pacing	Agitated locomotion
Form	Elastic gait	Inelastic gait
Flexibility	Invariant path	Variable path
Prevalence[1]	About half of individuals (7/13)	Almost all individuals (11/13)
Frequency[1]	Frequent (up to 46.7%)	Rare (up to 1.1 %)
Duration[1]	Long (up to 10 min)	Short (≤ 20 s)
Context[1]	Before, during and after fights	Almost exclusively during fights

398 [1] in this study

		Induction of acute stress	
		Yes	No
Increase in pacing frequency	Yes	True positives	False positives
	No	False negatives	True negatives