Hydrology-informed metapopulation modeling of liver fluke transmission in the Lawa Lake complex of northeast Thailand

Tomás M. León[1,3]*, Vichian Plermkamon[2], Kittiwet Kuntiyawichai[2], Banchob Sripa[3],
Robert C. Spear[1]

[1]School of Public Health, University of California, Berkeley, USA

[2]Faculty of Engineering, Khon Kaen University, Khon Kaen, Thailand

[3]Tropical Disease Research Center, Faculty of Medicine, Khon Kaen University, Khon
Kaen, Thailand

* Corresponding author

Email: tomas.leon@berkeley.edu (TML)

Present address: 2121 Berkeley Way #5302, UC-Berkeley School of Public Health,
Berkeley, CA, 94720-7360, USA

Abstract:

While hydrologic processes are intuitively understood to influence transmission dynamics of water-related diseases, limited research exists that explicitly links hydrologic and infectious disease data. In the case of the life cycle of liver flukes, hydrology influences several transmission processes that mediate infection risk for multiple hosts. Northeast Thailand is a hotspot for liver fluke transmission and has strong seasonal flooding patterns. A metapopulation model linking local hydrologic processes with transmission of the liver fluke *Opisthorchis viverrini* in a lake system in northeast Thailand was developed and parameterized using infection data from 2008-2016. A rainfall-runoff model and other hydrologic data were used to assess level of connectivity between villages and the influence of upstream communities on parasite distribution in the study area. Disease transmission was modeled with metapopulations representing six village clusters around the lake using known prevalence data from humans, cats and dogs, snails, and fish. The metapopulation model improved upon the single-village model in its match to historical data patterns for the six village clusters with the introduction of the new time-variable parameters. Results suggest there are three unique hydrologic-epidemiologic regimes within the Lawa Lake system in response to upstream watersheds and risk of overland flooding that contribute to risk for *O. viverrini* infection. While available data may be insufficient to specifically characterize exact transmission dynamics, the practical implications of such findings are the importance of addressing connectivity for any intermediate host-based intervention. Similar approaches using hydrologic data to assess the impacts of water on pathogen transmission dynamics and inform mechanistic disease transmission models could be applied across other water-related disease systems.

Keywords: liver fluke, *Opisthorchis viverrini*, mathematical modeling, food-borne

trematodiases, Thailand, hydrology

Introduction

Opisthorchiasis, infection with liver flukes of the genus *Opisthorchis*, is a disease

whose transmission and distribution are significantly influenced by hydrologic

processes. The parasites' egg and cercarial forms require sufficient water and hydraulic

transport to the next intermediate host for the transmission cycle to be sustained. Eggs

are excreted in the feces of infected final hosts (humans, and reservoir cats, dogs, and

other mammals to a limited extent); if not safely treated or contained, these eggs reach

the aqueous habitat of intermediate host snails, which consume the eggs and mediate

maturation to the cercarial stage. The cercariae are then released back into water, where

they swim and seek out the second intermediate host cyprinid fish. They encyst in the

fish, which if then consumed raw or undercooked by humans or other host mammals

can migrate to their bile ducts and mature into adult worms. Water is a sustaining force

for this parasitic life cycle, and its movement permits viable infection at each successive

host stage.

The major liver fluke of interest in Thailand is *Opisthorchis viverrin*i. Given its

known disease burden in Thailand, opisthorchiasis (infection with *O. viverrini)* has been

a public health priority there, where it is transmitted to humans via the consumption of

popular local raw and fermented fish dishes such as *koi pla* and *pla som*. The highest

prevalence of opisthorchiasis and cholangiocarcinoma (CCA), the fatal bile duct cancer

associated with *O. viverrini* infection, are found in northeast Thailand in the region

surrounding Khon Kaen (Sithithaworn et al. 1997). Historical hotspots of

opisthorchiasis and CCA include the villages around Lawa Lake. While much research

has been conducted into the pathology of opisthorchiasis and CCA, there is limited

literature addressing the ecologic and hydrologic aspects of parasite transmission in the

environment (Grundy-Warr et al. 2012, Wang et al. 2017).

Lawa Lake is an approximately 4000-acre body of water that is highly vegetated

and subject to significant hydrologic changes caused by seasonal variation in northeast

Thailand. A peak in liver fluke infections is seen with lag following the rainy season in

Thailand, as flooding facilitates the spread of fecal contamination and coincides with

the rapid increase in snail populations (Sithithaworn et al. 1997). Since several weeks

are required for the parasite to mature through its life stages, high infection rates in fish

are seen in the late rainy season and late summer (July-January), and there is lower

infection risk in the dry season and early summer (March-June). A primary industry on

the lake is fishing, which contributes to sustaining local liver fluke transmission

(Aunpromma et al. 2012). The hydrology of the Lawa Lake region is exceedingly

complex and disturbed, as significant development in recent years has led to

construction of new irrigation canals and ditches, new culverts and spillways that are

opened and closed in the flooding season, and fish ponds that have become increasingly

popular as a source of food and revenue.

Metapopulation modeling is commonly used to better understand the

connectivity and influence of discrete populations and environmental patches on each

other. This type of modeling is especially powerful for understanding pathogen

transmission in heterogeneous environments for complex life cycles. Connectedness

between these environmental and host patches can occur in multiple ways, including

movement of humans between villages and dispersion of a waterborne pathogen from a

section of river or lake adjacent to one population to another section adjacent to a

separate population. The second example demonstrates hydrologic influence on a

disease transmission system, as waterborne diseases as diverse as cholera and

schistosomiasis rely on advective transport to expose new susceptible individuals with

pathogens excreted or shed by infected individuals. Hydrologic patterns are time-

varying and markedly local in nature. Given this environmental complexity, hydrologic

fate and transport of pathogens are difficult to study. In addition, motile waterborne

parasitic forms, such as liver flukes and schistosome cercariae, have independent

mobility behaviors, making hydrologic flows not entirely representative of how these

parasites reach their next host, though the distances covered are much smaller

compared with the potential impact of flooding or river flow (Haas et al. 1990,

Krishnamurthy et al. 2017).

Research connecting hydrology with waterborne disease transmission is an

emerging field with recent work on cholera and schistosomiasis (Remais et al. 2008,

Rinaldo et al. 2012, Perez-Saez et al. 2016). Tracing the spread of pathogens in the

environment is challenging, and countervailing forces and feedback loops make it

difficult to ascribe an increase or decrease in human infection to trends in meteorology

and climate or consequent hydrology. Long-term studies of climatic changes in rainfall

patterns or the influence of dams allow more definitive statements about impacts on

disease transmission, though these are also complicated by shifts in host and vector

habitat and in seasonal patterns that may disrupt or exacerbate host and vector growth

and reproduction (Tompkins et al. 2013, Ziegler et al. 2013).

In this work, a metapopulation disease transmission model is developed and

parameterized to assess hydrologic connectivity and *O. viverrini* parasite movement

between six village clusters around Lawa Lake in Khon Kaen Province, Thailand, and

how that is reflected by opisthorchiasis prevalence in hosts. Understanding liver fluke

transmission in this seasonal, hydrologically connected environment with modeling can

help define the scale of transmission processes and thereby optimize environmental

control and treatment to have maximum impact on reducing disease transmission and

prevalence in this setting and others.

Materials and Methods

The model structure is an extension of the modeling framework presented in

León et al. (2018). The six village clusters studied (Figure 1) are now connected in a

mechanistic metapopulation framework to account for exchange of parasites and hosts

between village clusters and their associated environments. This enables the model to

include the influence of population-level factors, spatial heterogeneity, and degrees of

connectedness between patches. This metapopulation model leverages information

about hydrologic connectivity between village and host clusters to understand the

movement of the liver fluke parasite's various forms in the environment as mediated by

water. To consider local hydrologic impacts on the liver fluke transmission cycle, five

main factors are included: 1) egg inputs into the system from upstream watersheds; 2)

egg inputs into the system from overland flooding; 3) snail and fish mobility due to

hydrologic connectivity; 4) snail and fish available habitat; and 5) hotspots where

infectious snails come into contact with susceptible fish. These factors are modeled

monthly with seasonality to account for changing patterns throughout the year, i.e.,

they are modeled in a piecewise fashion and updated monthly.

The metapopulation model connects the six villages or geographically proximate

village clusters shown in Figure 1 around Lawa Lake in Khon Kaen Province of

northeast Thailand. The villages or village clusters described here are CCK, Lawa, BT

(cluster of 5 villages), NNK, KSR, and DPD (cluster of 2 villages), which were chosen

and clustered based on geographical location and how historical human infection

survey data was collected. The six clusters further sort into three groups based on

impact or lack thereof of flooding and upstream watersheds. The two upstream

watersheds (Figure 2) flow into Lawa Lake at locations adjacent to NNK and BT and

contribute to egg input there. Villages CCK and Lawa are in proximity to the Chi River

and are most susceptible to seasonal overland flooding. KSR and DPD are the villages

most "downstream" and are relatively isolated from major flooding or upstream

drainage impacts. Villages within the upstream watershed have not had as significant

treatment and control programs as the villages around Lawa Lake, and infection

surveys suggest that upstream villages still have high infection prevalence values over

50% (unpublished data).

Figure 1: Connectivity between six village clusters around Lawa Lake (Map data: Landsat, USGS).

Figure 2: Two upstream sub-basins flow into Lawa Lake near NNK and BT, respectively (QGIS).

These hydrologic impacts seem to mirror trends in epidemiological patterns.

Previous research highlighted the north-to-south gradient of nitrogen and salinity in the

water that predicted higher snail abundance towards the south side of the lake (Kim et

al. 2016). NNK, the southernmost village cluster, had the highest human infection

prevalence at the time points when it was studied (Table 1). KSR and DPD, the

northernmost villages and the farthest from the Chi River, had the lowest baseline

prevalence of *O. viverrini* infection before the control program started. In the single-

village model, varying transmission parameters by village did not fully capture the

different patterns that occurred in the six village clusters when connectivity was not

included. Therefore, the role of hydrologic processes and connectivity between villages

needs to be considered in the disease transmission model to better account for the

patterns observed.

Village Cluster	2008	2010	2011	2012	2014	2015
CCK	54.9% (593)		33.0%*	44.3% (139)		
Lawa	67.1% (501)	63.1% (108)	19.0%*	(16)		8.7%*
BT	61.9% (346)	37.2% (131)		35.0% (136)	9.0%*	14.2%*
NNK	74.1% (499)			50.0% (61)		
KSR	16.4% (101)					
DPD	22.1% (112)	36.5% (82)				14.6%*

Table 1: Infection prevalence (%) and mean intensity in positive individuals (EPG) for six village clusters around Lawa Lake. * indicates less sensitive diagnostic method (Kato-Katz or Kato thick smear).

To understand the effects of complex hydrologic factors on snail habitat, fish

access to these habitats, and the pathways of parasite transmission, a hydrologic model

of the Lawa Lake system was utilized to simulate flow patterns and changes in water

levels over time. This model is a rainfall/runoff model common in hydrology that

considers the transport of water through a system originating from upstream in the

catchment basin or from precipitation. It uses a Soil & Water Assessment Tool (SWAT)

model to generate runoff for the PCSWMM hydraulic model to determine hydraulic

parameters of Lawa Lake such as flood depth and extent, flow velocity, and travel time

(CHI 2018). The inputs for the SWAT model include meteorological data (rainfall,

temperature, relative humidity, and windspeed) from the Thai Meteorological

Department, soil type, land use, and a digital elevation model (DEM) generated from

satellite imagery. For the hydraulic model, hydrologic structures and key parts of Lawa

Lake were surveyed for elevation at 1m x 1m resolution using a drone, and a 2D model

integrating runoff, the improved DEM, and meteorological data from 2008 to 2016 was

developed using PCSWMM. Outputs include water level and flow vectors for the time

points modeled between 2008 and 2016. Calibration was conducted with available

precipitation and gauge data in the area from the Bureau of Water Management and

Hydrology, Royal Irrigation Department, in Thailand. Figure 3 demonstrates an

example of the variation in flows predicted by the model over the course of a calendar

year encompassing the rainy and dry seasons; arrow direction and thickness represent

the relative change in flow vectors. All of the villages are adjacent to Lawa Lake, and

inflows and outflows as well as the impacted snail and fish populations are dynamic

and heterogeneously distributed. The rainy season is characterized by high and active

flows that generally peak in October with flooding from the Chi River varying from

year to year. The dry season has relatively little hydrologic activity establishing

connectivity between village clusters.

Figure 3: Hydrologic flows change dynamically throughout the year (semiquantitative interpretation of hydrology model results for an annual cycle) (Map data: Landsat, USGS).

Because gauge data was not available for the sub-basins upstream of Lawa Lake,

a Soil and Water Assessment Tool – Calibration and Uncertainty Procedures (SWAT-

CUP) model was used to calibrate and validate runoff into the Lawa Lake system from

these sources generated by the SWAT model. Inputs for the upstream sub-basins

included a 30m x 30m DEM, land use data, and soil type data from the Land

Development Department of Thailand. To map the hydrologic features of Lawa Lake

and finer scale structures, a drone was used to chart these areas in greater detail and

determine elevations where water was flowing into or out of Lawa Lake. Sub-basin

calibration and validation graphs are shown in Figure 4; 2006-2010 data was used for

calibration, and 2011-2013 data was used for validation. The R^2 values range from 0.61

to 0.81; both the calibration and validation models miss late peaks in their runs. For the

calibration phase, fitting the less extreme peaks may have disadvantaged the model

from predicting the major discharge in 2010. For the validation phase, the 2011 peak

discharge was better modeled, but 2013 was missed by a large margin for reasons that

are not entirely clear but may be related to the different timing of precipitation-driven

flooding in 2013 compared with other years.

Based on these data and tools, the presence/absence of connections between

parcels of water associated with different village clusters were assessed. For example,

KSR would not contribute to transmission in Lawa or NNK year-round but the reverse

would be true. In March, when less rainfall and flooding occur, each village cluster is

relatively isolated, with the exception of the relatively weak influence of near neighbors

(Lawa to CCK or BT to DPD). The origin of flows is also subject to a differential dilution

effect; contaminated waste with parasite eggs from the upstream watershed (Figure 2)

would more strongly impact NNK than DPD or KSR, and the upstream villages (such

as NNK) would experience the effects closer to discharge time.

The mathematical model of disease transmission (which incorporates data from

the hydrologic model as variables and parameters) is an expansion of the single-village

model described in León et al. (2018) to connect the six village clusters and uses as its

state variables the infection prevalence in humans, reservoir hosts (cats and dogs), and

snails, and infection counts in fish (equations in Appendix). Initial values were set from

the baseline surveys in 2008, and base transmission parameters are carried over from the

single-village model found using Markov chain Monte Carlo (MCMC) maximum

likelihood methods to fit the model onto known infection prevalence data as described

below. The infection prevalence data includes surveys using two different methods:

formalin-ether concentration technique (FECT) and Kato-Katz (or Kato thick smear). In

Thailand, FECT has been found to be the significantly more sensitive method to detect *O.*

viverrini infection based on available data and because the protocol is intended to make

microscopic examination easier (Laoprom et al. 2016, Saijuntha et al. 2018). Within the

fish state variable is a fish demography model that captures the small window of time in

the first few months of a fish's life when it is susceptible to cercarial infection (before its

scales harden and it becomes more resistant). This model assumes a maximum fish

lifespan of 4 years before either being caught or natural death.

Figure 4: Calibration and validation curves for sub-basins 1 & 2. Calibration data
encompasses 2006-2010 and validation 2011-2013.

Figure 5 shows the general model flow diagram and how available data sources

and village-specific factors are used in the model. The fecal waste-related egg inputs

come from the Chi River and the two upstream watersheds and affect the snail state

variable, contributing to the force of infection in that linkage; these time-varying

parameters are derived on a monthly basis from the rainfall-runoff model. Other egg

inputs from open defecation and disposal of septic tank sludge are not modeled due to

lack of information about where and when they occur. The egg inputs from overland

flooding of the Chi River were assumed to affect CCK and Lawa villages equally and

were calculated by using flow measurements from the river and multiplying by a scalar

to relate the impact of that water source with the upstream sub-basins. The first and

second upstream sub-basins' outflow were modeled to contribute eggs to the systems in

BT and NNK exclusively and multiplied by their own scalars to translate those flows

into contributions to human and reservoir host infection.

Figure 5: Model flow diagram showing submodels and input data that impact population numbers and transmission rates.

Connectivity rates between patches were varied on a monthly basis between

2008 and 2016 based on the hydrology model results to account for snail and fish

mobility due to hydrologic connectivity. Figure 6 shows a model representation of how

this works in a metapopulation model. These connectivity parameters were derived by

assessing the fate and transport of parcels of water in a village cluster's area and what

proportions reached other village clusters in the Lawa Lake system. These r parameters

are unitless and vary from 0 to 1, describing the proportion of each village cluster's

force of infection for that host stage that affects each other village. Figure 7 shows

examples of how these connectivity parameters varied by village pairing, month, and

year (Figures 7-10 were produced in R using the ggplot2 package (Wickham 2016, R

Core Team 2020)).

Figure 6: Model flow diagram showing how the metapopulation model works, with r connectivity parameters affecting transmission between each state variable. Subscript i denotes the source village and subscript j the receptor village for the parasite load in that transmission process. H = Human, R = Reservoir, S = Snail, F = Fish.

The parameters carried over from the single-village model are listed in Table S1.

The β values are transmission parameters and are village cluster-specific (Table S2). The

r connectivity parameters are specific to each patch-to-patch relationship (Figure 1), γ is

the fish catch rate describing the fraction of the total fish population caught at each time

step, $\lambda(t)$ is a gating function to control fish birth, death, and aging processes, μ are

mortality rates, and α_{PZQ} are praziquantel (PZQ) treatment events.

Figure 7: Village connectivity for BT-DPD, CCK-Lawa, and NNK-BT in 2008, 2012, and 2012

While a daily time scale could be preferable for assessing hydrologic

impact, historical data only captured month-to-month variability. Because human

infection survey data only measures prevalence and not incidence, this time scale is

reasonable for this study. From the hydrology model, the extent of water surface area at

a suitable depth (under 0.3m) for the contact events between juvenile cyprinid fish and

the aquatic snail intermediate hosts ("hotspots" for infection transfer) were used and to

estimate snail populations $N_{S,i}$ and fish populations $N_{F,i}$. The transport time for a parcel

of water between each village cluster was employed to estimate the time scale of

movement between the locations, but these interactions happened on the order of days

and not months and therefore the model's time scale was not granular enough to

introduce a time lag component. There was insufficient data to track fish mobility

between patches, so fishermen's movement data from Kim et al. (2017) was used to

inform the exchange of fish in inter-village commerce as contributing to human

infection from other village clusters. Table 2 summarizes these linkages and use of

hydrology-related data in the disease transmission model. Figure 7 shows examples of

the variability between village connectivity across months and years for the

human/reservoir-to-snail and snail-to-fish transmission processes (the remaining

connectivity parameter graphs are shown in the Appendix). The general trends persist

from year to year, but the timing of peaks differ and affect village connections

differently. The year 2008 produced stronger connectivity for BT-DPD and NNK-BT,

while 2010 had stronger connectivity for CCK-Lawa.

Hydrology-related transmission impact	Quantification method	Retained in model?
Egg inputs from overland flooding	Rainfall-runoff model output from Chi River summarized on monthly basis	Yes
Egg inputs from upstream water basins	Sub-basin model output summarized on monthly basis	Yes
Snail and fish mobility	Snail: Patch connectivity from rainfall-runoff model; Fish: fishermen catch data	Yes
Snail and fish available habitat	Snail: Rainfall-runoff model output and GIS analysis; Fish: N/A	Yes
Hotspots for snail-to-fish contact	Rainfall-runoff model output and GIS analysis	No

Table 2: Description of linkages between hydrology model and disease transmission model

Results

Figure 8 shows the metapopulation model results for the six village clusters in

the base scenario with the metapopulation model establishing relative connectivity

between villages. Prevalence data points are included, distinguishing between the more

sensitive FECT surveys and the less sensitive Kato method surveys. The heterogeneity

of outcomes reflects the data: some villages saw reductions in infection prevalence to

less than 10% (KSR, DPD), yet a few villages continued to have predicted prevalence

values greater than 20% (BT and NNK). The steep drops in the graph were treatment

events, when a subgroup of villagers was tested for infection and given praziquantel if

they tested positive (the model assumes 100% drug efficacy). Model simulations were

run for eight years between 2008 and 2016.

Figure 8: Metapopulation model run for human infection prevalence in villages around
Lawa Lake

316   Figures 9 and 10 show infection prevalence values for intermediate snail and fish

317 hosts. In snails, the prevalence cycles seasonally with most patch populations

318 decreasing to 0.1% from initial values of 0.2% (with the exception of Lawa where

Figure 9: Snail prevalence values for metapopulation model

Figure 10: Fish prevalence values for metapopulation model

prevalence approaches 0%). NNK has the highest final snail prevalence value at 0.18%,

which is well within the range of what would be expected in this type of environment.

For fish prevalence, because the initial conditions are disparate and based on baseline

survey data, the model behavior is quite different. There is a seasonal aspect to their

dynamics though this is dampened for most villages except NNK, where it is readily

apparent. The end prevalence values range from 8-41%, with CCK, Lawa, BT, and DPD

having the lowest values and NNK having the highest.

Discussion

Compared with the single-village model, the metapopulation model no longer

over-predicts final infection prevalence for the villages in 2016. Instead, the model now

occasionally under-predicts prevalence for some data points, notably, CCK in 2012. This

specific issue can be partially accounted for by the number of treatment-naïve

individuals included in the 2012 infection surveys. Interpretation of the data leads to

some specuation about the meaning of the surveys and their different diagnostic

methods. BT stands out as the modeled village with the least improvement (though it

overpredicts the 2012 and 2014 data). According to control program managers, BT

villagers were the least participatory in the Lawa Project and may therefore have reaped

the least benefit from the control program. Given that this mathematical model is

primarily concerned with infectious indviduals actively shedding parasites into the

environment rather than asymptomatic cases, the Kato-based surveys from 2011, 2014,

and 2015 may actually capture the most infectious and relevant individuals in the

system and therefore be useful for thinking about infection prevalence patterns.

However, fitting all of the data points accurately would be nearly impossible since most

villages show non-monotonic patterns, infection burden builds up over time, and other

unmeasured factors not included in the model may play a a major role in mediating

these dynamics.

Two major questions are, given the discrepancy between model fits and the data,

whether the data accurately reflects the reality of the disease transmission situation and

whether the model should be tuned with yet more parameters to get a closer fit to the

data. While this dataset is relatively complete and informative by the standards of NTD

surveys, it still lacks enough time points, consistency in collection methods, and large

enough sample proportions of the populations to give a detailed picture of the *O.*

viverrini infection situation around Lawa Lake. The data (Table 1) show major swings

across time points and discrepancies between the survey methods. Therefore, while the

FECT data was used to fit via MCMC the transmission parameters in the single-village

model, no parameters were fit for the metapopulation model because of the increase in

model complexity and number of parameters (eighteen transmission parameters alone

compared with three per single-village model, plus seventy-two time-varying

connectivity parameters between the six village clusters). For this reason, the

connectivity parameters were derived exclusively from hydrology submodel results.

The metapopulation model is more believable than the single-village model in part

because of its inclusion of external hydrologic influences and connecitivity and in part

because the modeled behavior is more nuanced in the clustered patterns of village

prevalence change it exhibits. The lack of parameters fitted to this model structure and

the higher quality data that informed the hydrology model lend a realism to the

underlying mechanics of the metapopulation model that improves upon the relatively

straightforward transmission framework of the single-village model. The

metapopulation model is meant to capture patterns of transmission rather than exact

fits to the data. Nonetheless, the lack of information about differences in snail infection

and raw fish eating patterns between villages remains a weakness, as they could not be

incorporated into the model. Better data on these aspects of the transmission cycle

would further strengthen the modeling framework and bring it into greater alignment

with reality.

The patterns observed in these results support the sorting of the six village

clusters into three geographical clusters that exhibit different trends based on human

prevalence values. The first cluster, consisting of CCK and Lawa, is in close proximity

to the Chi River and is most susceptible to overland flooding experienced during the

rainy season. Its villages had high human prevalence values at the start of the control

program, which decreased sharply during the period of treatment and control activity.

These villages were the headquarters and major focal area of Lawa Project activities,

suggesting that they benefited the most from health education and health volunteer

engagement. The model is able to accurately account for the decrease in prevalence

without making any assumptions about reinfection. Additionally, these villages are in

close proximity to heavily fished waters in the lake, as supported by GPS evidence and

interviews described in Kim et al. (2017).

The second cluster consists of BT and NNK, the villages to the south and

southeast of the lake. These village also had high prevalence values at the start of the

treatment and control program but experienced much more modest decreases when

comparing data from later infection surveys. These villages are most impacted by

upstream watersheds draining into Lawa Lake (Figure 2.2), where some villages still

have over 50% *O. viverrini* infection prevalence (unpublished data). Consequently, if

open defecation or unsafe disposal of human or reservoir waste is occurring in these

watersheds, the runoff will disperse parasite eggs into the canals, ponds, and sections of

the lake in close proximity to the second cluster's villages. These villages are also

adjacent to the highest concentration of fish ponds in the system and were not a focal

area for Lawa Project activities.

The third cluster is KSR and DPD, which are located to the northeast of Lawa

Lake. At the start of the control program, these villages had significantly lower

prevalence values, which may be attributable to a lower degree of baseline

environmental contamination. These villages were significantly affected by neither the

Chi River nor the upstream watersheds, so they experienced fewer external inputs of

infectious individuals or waterborne forms of the parasite into their local systems.

These villages were not adjacent to high concentrations of fish ponds or fishing activity

in their local waters and received less emphasis by the Lawa Project compared to the

first cluster.

Because of the low prevalence of snail infection, the lack of field survey data, and

the difficulty measuring snail prevalence precisely, strong claims cannot be made about

the model results for the snail intermediate host. However, most field surveys indicate

that snail prevalence in this region ranges between 0 and 0.2%, agreeing with the model

results (Kim et al. 2016). With current diagnostic methods, differentiating between the

clusters would require surveys of tens of thousands of snails at least. In recent years,

the highest snail prevalence values found are still less than 10% (Kiatsopit et al. 2012).

Much like other snail-borne diseases such as schistosomiasis though, only a few snails

are required to maintain transmission in an area because of the high number of

cercariae they shed into water bodies. Further understanding of where snails are most

likely to be infected will help with environmental measurement and control.

Bottlenecks of water flow, such as certain ditches and culverts, would concentrate fecal

waste and parasite eggs and be zones of likely contact with susceptible snail hosts.

Eliminating or protecting these areas could be an effective method of environmental

control subject to proper coverage in the environment of interest and patch/cluster

connectivity.

Fish infection prevalence is much higher than for snails and therefore it is easier

to discern differences between clusters. Because transmission is foodborne, we are

interested in the supply chain of food to consumers, which is not necessarily related to

proximity between where fish lived and where they are eaten. Fishermen, middlemen,

and merchants are all mobile and may choose to sell and distribute fish to other villages

to expand their market. In the model simulation, the relative ordering of low to high

fish prevalence values in fish hosts differ considerably from the results in humans, with

NNK and KSR having the highest infection prevalence. NNK's is driven largely by the

initial value, but KSR's is driven by dynamics, as its outcome is quite different from

Lawa's, which started with a similar prevalence level. Fish are infected by having

infectious snails releasing cercariae into areas with juvenile fish, so KSR is the cluster

with the greatest magnitude of this process taking place. Paying attention to fish

prevalence results and how they interact with patterns of fish commerce can help

identify where to target health education interventions related to cooking.

The model and the data that inform the model have limitations due to their

fragmentary nature. Other model assumptions include ignoring the impact of different

parasite burden levels in hosts and the age structure of human populations. The

hydrology model was calibrated and validated against available data, leading to a

plausible interpretation of the observed hydrologic behavior of the system. However, it

could not account for very local effects that eluded its time and spatial scale and may

have relevance for the points of contact between stages of the life cycle. The infection

survey data may not be representative of the host populations because of sampling bias,

but its overall spatial and temporal patterns align with local understanding and

experience.

Conclusions

This work highlights a major driver of persistent *O. viverrini* infection in

northeast Thailand: a disturbed and dynamic hydrologic environment that mediates

parasite transmission between connected village clusters and environments. This was

accomplished by discussing and modeling five different means of hydrologic processes

influencing parasite transmission and showing that its effects are significant and

complex, acting heterogeneously across the Lawa Lake system. While local infection

and contamination may be the main factor driving transmission at higher prevalence

levels, as prevalence declines and villages move toward elimination connectivity will

play a bigger role in maintaining the transmission cycle and preventing local

elimination of the parasite.

The connectivity between water bodies and villages ensures that elimination of

local infection is not possible without addressing upstream and adjacent environments.

If infectious fecal waste from human and reservoir hosts is continually flushed

downstream and the source is not treated, transmission will be restarted even if

previously disrupted. This argument supports an approach that first targets villages

and populations near headwaters and then proceeds further downstream while still

accounting for human mobility and migration that could reintroduce infection into a

previously cleared environment. Flood pulses and overland flooding also risk moving

parasites into isolated and disconnected environmental patches on an annual basis,

which requires constant treatment and attention to infection status of individuals in

those patches. Snail and fish mobility remain little understood but have major relevance

for *O. viverrini*'s life cycle, specifically how infection propagates in the environment.

Targeting the locations where snails and juvenile fish come into close proximity with

each other could be a promising environmental control technique but requires greater

knowledge about the conditions that enable this transmission process.

A linked disease transmission-hydrologic modeling approach was employed

here that uses hydrology model outputs as time-varying inputs in the disease

transmission model to account for seasonal effects of flooding and water movement

relevant to the intermediate hosts and waterborne forms of *O. viverrini*. Based on model

results, village clusters were grouped into three disease prevalence curve patterns based

on presence/absence of upstream and flooding impacts and history of control program

intensiveness. Considering these findings, we argue for the use of this modeling

approach and its results to inform environmental control of *O. viverrini* and for the need

for environmental surveillance. While the specifics of the hydrology, population

structure, and pathogen transmission cycle are local and specific in nature, this

approach can be replicated across a variety of disease systems that are impacted by

seasonality and a dynamic hydrologic regime.

Acknowledgments: The authors would like to thank Barry Croke for helpful comments

on the hydrology model and the Tropical Disease Research Laboratory staff at Khon

Kaen University for their technical and logistical support for collecting the presented

data.

Funding: This work was supported by a US National Institutes of Health (NIH)

(www.nih.gov) grant [R21AI104513] (TML, VP, KK, RCS) and by the National Science

Foundation (www.nsf.gov) Graduate Research Fellowship under Grant No. [DGE

1106400] (TML). The funders had no role in study design, data collection and analysis,

decision to publish, or preparation of the manuscript.

The data used for this study are included in the main text and supplement. Analysis

scripts are available from the corresponding author on reasonable request.

References

Aunpromma S, Tangkawattana P, Papirom P, Kanjampa P, Tesana S, Sripa B,

Tangkawattana S, 2012. High prevalence of Opisthorchis viverrini infection in reservoir

hosts in four districts of Khon Kaen Province, an opisthorchiasis endemic area of

Thailand. Parasitology international. 61(1):60-4.

Computational Hydraulics International [CHI] (2018). PCSWMM Modeling Software.

Available at: https://www.pcswmm.com

Grundy-Warr C, Andrews RH, Sithithaworn P, Petney TN, Sripa B, Laithavewat L,

Ziegler AD, 2012. Raw attitudes, wetland cultures, life-cycles: socio-cultural dynamics

relating to Opisthorchis viverrini in the Mekong Basin. Parasitology International.

61(1):65-70.

Haas W, Granzer M, Brockelman CR, 1990. Opisthorchis viverrini: finding and

recognition of the fish host by the cercariae. Experimental Parasitology. 71(4):422-31.

Kiatsopit N, Sithithaworn P, Saijuntha W, Boonmars T, Tesana S, Sithithaworn J,

Petney TN, Andrews RH, 2012. Exceptionally high prevalence of infection of Bithynia

siamensis goniomphalos with Opisthorchis viverrini cercariae in different wetlands in

Thailand and Lao PDR. The American Journal of Tropical Medicine and Hygiene.

86(3):464-9.

Kim CS, Echaubard P, Suwannatrai A, Kaewkes S, Wilcox BA, Sripa B, 2016. Seasonal

and spatial environmental influence on Opisthorchis viverrini intermediate hosts,

abundance, and distribution: insights on transmission dynamics and sustainable

control. PLoS NTDs. 10(11):e0005121.

Kim CS, Smith JF, Suwannatrai A, Echaubard P, Wilcox B, Kaewkes S, Sithithaworn P,

Sripa B, 2017. Role of socio-cultural and economic factors in cyprinid fish distribution

networks and consumption in Lawa Lake region, Northeast Thailand: Novel

perspectives on Opisthorchis viverrini transmission dynamics. Acta Tropica. 170:85-94.

Krishnamurthy D, Katsikis G, Bhargava A, Prakash M, 2017. Schistosoma mansoni

cercariae swim efficiently by exploiting an elastohydrodynamic coupling. Nature

Physics. 13(3):266.

Laoprom N, Laithavewat L, Kopolrat K, Kiatsopit N, Kaewkes S, Chantalux S,

Mongkolsin C, Chanmaha B, Andrews RH, Petney TN, Sithithaworn P, 2016. Evaluation

of a commercial stool concentrator kit compared to direct smear and formalin-ethyl

acetate concentration methods for diagnosis of parasitic infection with special reference

to *Opisthorchis viverrini sensu lato* in Thailand. The Southeast Asian Journal of

Tropical Medicine and Public Health. 47(5):890-900.

León TM, Porco TC, Kim CS, Kaewkes S, Kaewkes W, Sripa B, Spear RC, 2018.

Modeling liver fluke transmission in northeast Thailand: Impacts of development,

hydrology, and control. Acta Tropica. 188:101-7.

Perez-Saez J, Mande T, Ceperley N, Bertuzzo E, Mari L, Gatto M, Rinaldo A, 2016.

Hydrology and density feedbacks control the ecology of intermediate hosts of

schistosomiasis across habitats in seasonal climates. PNAS. 201602251.

R Core Team, 2020. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Remais J, Liang S, Spear RC, 2008. Coupling hydrologic and infectious disease models

to explain regional differences in schistosomiasis transmission in southwestern China.

Environmental Science & Technology. 42(7):2643-9.

Rinaldo A, Bertuzzo E, Mari L, Righetto L, Blokesch M, Gatto M, Casagrandi R, Murray

M, Vesenbeckh SM, Rodriguez-Iturbe I, 2012. Reassessment of the 2010–2011 Haiti

cholera outbreak and rainfall-driven multiseason projections. PNAS. 109(17):6602-7.

Saijuntha W, Duenngai K, Tangkawattana S, Petney TN, Andrews RH, Sithithaworn P,

2018. Recent Advances in the Diagnosis and Detection of Opisthorchis viverrini Sensu

Lato in Human and Intermediate Hosts for Use in Control and Elimination Programs.

Advances in Parasitology. 101:177-214.

Sithithaworn P, Pipitgool V, Srisawangwong T, Elkins DB, Haswell-Elkins MR, 1997.

Seasonal variation of Opisthorchis viverrini infection in cyprinoid fish in north-east

Thailand: implications for parasite control and food safety. Bulletin of the World Health

Organization. 75(2):125.

Tompkins AM, Ermert V, 2013. A regional-scale, high resolution dynamical malaria

model that accounts for population density, climate and surface hydrology. Malaria

Journal. 12(1):65.

Wang YC, Yuen R, Feng CC, Sithithaworn P, Kim IH, 2017. Assessing the role of

landscape connectivity on Opisthorchis viverrini transmission dynamics. Parasitology

international. 66(4):402-12.

Wickham H, 2016. ggplot2: Elegant Graphics for Data Analysis. Springer.

Ziegler AD, Petney TN, Grundy-Warr C, Andrews RH, Baird IG, Wasson RJ,

Sithithaworn P, 2013. Dams and disease triggers on the lower Mekong river. PLoS

NTDs. 7(6):e2166.

 Supporting information

Table S1: Parameter values for single-village model

Parameter	Value	Units	Source	Symbol
Natural mortality of snails	1.37E-03	per day	Kruatrachue et al. 1982	μ_S
Parasite dependent mortality of snails	1.37E-03	per day	Chanawong & Waikagul 1991	α_S
Mortality of fish	6.85E-04	per day	Suvarnaraksha et al. 2011	μ_F
Parasite dependent mortality of fish	0	per day	Assumption (unstudied)	α_F
Natural mortality of humans	3.69E-05	per day	CIA 2015 (Factbook)	μ_H
Human infection clearance by praziquantel	Variable	Episodic	Treatment data from clinics	α_{PZQ}
Mortality of reservoir host	2.74E-04	per day	Local interview data	μ_R
Reservoir infection clearance by praziquantel	Variable	Episodic	Cat/dog treatment data from veterinarians	$\alpha_{PZQ,R}$
Transmission parameters (fish-to-human, fish-to-reservoir, human-to-snail, snail-to-fish)	See Table 3.5.1	per day per infectious host/worm	Equilibrium conditions and MCMC	$\beta_{FH}\beta_{FR},$ β_{HS}, β_{SF}
Fish population	3000	fish	Estimate	N_F
Snail population	30000	snails	Estimate	N_S
Human population	Variable	humans	Village censuses	N_H
Cat and dog population	100	reservoir hosts	Estimate from village censuses	N_R
Mean human worm count	Variable	worms	Infection survey data	W_H
Mean reservoir worm count	Variable	worms	Infection survey data	W_R

Table S2: Beta transmission parameters for single village model

Equilibrium	CCK	Lawa	BT	NNK	KSR	DPD
Fish to Human	2.50E-10	7.24E-11	2.53E-10	1.81E-11	3.32E-07	2.68E-07
Human to Snail	8.90E-09	1.01E-08	4.92E-09	8.92E-09	6.56E-12	1.18E-11
Snail to Fish	7.34E-06	7.10E-06	3.40E-06	8.96E-06	1.67E-05	1.35E-05
MCMC	CCK	Lawa	BT	NNK	KSR	DPD
Fish to Human	2.95E-07	3.45E-07	7.26E-07	2.28E-07	3.30E-08	7.75E-08
Human to Snail	1.25E-08	2.53E-09	3.45E-09	2.26E-08	1.28E-08	1.48E-08
Snail to Fish	4.48E-06	7.03E-06	2.23E-06	2.28E-05	7.08E-06	3.89E-06

 Model Equations

$$\frac{dI_{H,i}}{dt} = \beta_{FH,i}\left(1 - I_{H,i}(t)\right)\sum_{j=1}^{n} r_{FH_{i,j}}(t) * \gamma \sum_{k=1}^{3} I_{F_{j,k}}(t) - (\mu_H + \alpha_{PZQ}(t))I_{H,i}(t) \tag{1}$$

$$\frac{dI_{R,i}}{dt} = \beta_{FR,i}\left(1 - I_{R,i}(t)\right)\sum_{j=1}^{n} r_{FR_{i,j}}(t) * \gamma \sum_{k=1}^{3} I_{F_{j,k}}(t) - (\mu_R + \alpha_{PZQ,R}(t))I_{R,i}(t) \tag{2}$$

$$\frac{dI_{S,i}}{dt} = \beta_{HS,i}\left(1 - I_{S,i}(t)\right)\left(\sum_{j=1}^{n} r_{HS_{i,j}}I_{H,j}(t)N_{H,j}(t) + \sum_{j=1}^{n} r_{RS_{i,j}}I_{R,j}(t)N_{R,j}(t)\right) - \mu_S I_{S,i}(t) \tag{3}$$

$$\frac{dI_{F,i}^0}{dt} = \beta_{SF,i}\left(N_{F,i}^0(t) - I_{F,i}^0(t)\right)\lambda(t)\sum_{j=1}^{n} r_{SF_{i,j}}I_{S,j}(t)N_{S,j}(t) - \mu_F I_{F,i}^0(t) \tag{4}$$

$$\frac{dI_{F,i}^{1-3}}{dt} = -(\gamma + \mu_F)I_{F,i}^{1-3}(t) \tag{5}$$