Title: Extreme midsummer rainfall event drives early onset cyanobacterial bloom

Running head: Extreme rainfall drives early bloom

Megan L. Larsen[1]*, Helen M. Baulch[2], Sherry L. Schiff [3], Dana F. Simon[4], Sébastien Sauvé [4],

and Jason J. Venkiteswaran[1]

[1]Department of Geography and Environmental Studies, Wilfrid Laurier University, 75 University

West, Waterloo, Ontario Canada N2L 3C5

[2]School of Environment and Sustainability, Global Institute for Water Security, University of

Saskatchewan, 323 Kirk Hall, 117 Science Place, Saskatoon Saskatchewan Canada, S7N 5C8

[3]Department of Earth and Environmental Sciences, University of Waterloo, 200 University

Avenue West, Waterloo, Ontario Canada N2L 3G1

[4] Department of Chemistry, Université de Montréal, 2900 Édouard-Montpetit, Montréal, Québec

Canada H3C 3J7

* corresponding author: mlarsen@wlu.ca

Number of words: 4099

Number of manuscript pages: 34

Number of tables: 2

Number of figures: 4

Abstract

The prevalence and increasing global distribution of cyanobacteria-dominated harmful algal blooms is strongly associated with changing climatic patterns and local biogeochemical and hydrological processes. Changes to precipitation frequency and intensity, as predicted by current climate models, are likely to alter bloom development and composition due changes in nutrient fluxes and water column mixing on short time scales during the open water season. However, few studies have directly documented the effects of precipitation events on cyanobacterial composition, biomass, and toxin production. In this study, we describe an early-initiated cyanobacterial bloom in Conestogo Lake, a eutrophic flood control reservoir located in southwestern Ontario, following heavy rainfall and subsequent flooding within the catchment. An increase in bioavailable phosphorus by more than 25-fold in surface waters resulted in biomass increases of *Aphanizomenon flos-aquae* throughout the reservoir approximately 2 weeks post-flooding. Anabaenopeptin-A and three microcystin congeners (microcystin-LR, -YR, and -RR) were detected at varying levels across sites during the bloom period, which lasted between 3 to 5 weeks. Together, these findings indicate that water column mixing and elevated phosphorus concentrations induced by increased water flow were the key drivers for the early cyanobacterial bloom in Conestogo Lake. Mitigation strategies for bloom-related water quality impairment must be both responsive and adaptive to the complexity of drivers affecting reservoir mixing dynamics, nutrient loads, and blooms. However, bloom mitigation efforts can be severely constrained by the large watersheds and broad suite of operational goals of many reservoirs. that are most often focussed on water quantity (flood water control and flow augmentation) rather than water quality or cyanobacterial bloom management.

47   **Key words**: cyanobacterial bloom, phosphorus, climate, precipitation, toxin, harmful algal

48   bloom, Ontario, reservoir, extreme rainfall

Introduction

Cyanobacteria are critical to the structure and function of aquatic communities (Reynolds 1984,
Wetzel 2001). However, prolific cyanobacterial growth often negatively impacts ecosystems by
reducing water quality and driving disruptions to aquatic food chains (Pearl 1988). Several
bloom-forming species also synthesize an array of bioactive compounds that pose chronic and
acute health risks to humans and animals through dermal contact, inhalation, and/or ingestion of
contaminated waters (Chorus et al. 2000, Codd 2000, Carmichael 2001). As a result, blooms and
their associated bioactive metabolites have economic consequences through impacts on tourism
and recreation, and via costs for drinking water treatment (Dodds et al. 2009, Bullerjahn et al.
2016).

The increased prevalence of cyanobacterial blooms has been historically
attributed to increased anthropogenic eutrophication, with emphasis on the relative
abundances, contributions, and impacts of nitrogen and phosphorus (Winter et al. 2012,
Paerl and Otten 2013). Changing climatic conditions are also now strongly considered as
significant drivers in bloom intensity and distribution (Paerl and Huisman 2009, O'Neil
et al. 2012, Sukenik et al. 2015, Paerl et al. 2016). Current climate models predict
increasing regional temperatures likely resulting in warmer surface waters, increased
thermal stratification, and water column stability that promote the growth of certain
bloom-forming species (Jöhnk et al. 2008, Paerl and Huisman 2009). Precipitation
frequency and intensity are also predicted to change, but less attention has been given to
how these factors may influence bloom development and cyanobacterial biomass
(Reichwaldt and Ghadouani 2012).

Increasing precipitation variability may impact external nutrient and sediment delivery to

waterbodies, alter residence time and flushing, and reduce vertical stratification, which, in turn,

may affect bloom development (Jacobsen and Simonsen 1993, Mitrovic et al. 2003, Wood et al.

2017). For example, Wood *et al.* (2017) reported decreased cyanobacterial biomass following an

extreme rainfall event that lead to water column cooling and destratification in a shallow,

eutrophic New Zealand lake. Intense precipitation, preceded or followed by extensive drought

may create episodic or pulsed nutrient loads potentially further favoring cyanobacterial

development (Bouvy et al. 2003, Reichwaldt and Ghadouani 2012). In addition to climatic

factors, bloom development is also dependent on a suite of site-specific characteristics including

hydrology, lake geomorphology, catchment size, and nutrient loading from internal and external

sources. Thus, the sensitivity of each lake to these drivers, including precipitation, will vary.

Cyanobacterial blooms are of particular concern for lake and reservoir managers due to

the potential for toxin production and impacts on water quality that may affect ecosystem

processes and the recreational uses of the water body (Chorus and Bartram 1999, Chorus et al.

2000). Cyanobacterial metabolites are both chemically variable and bio-actively diverse

(Carmichael 1997, Welker and Von Döhren 2006) with various modes of cellular action that may

cause superficial skin irritation at low exposures and sickness or even death if ingested at high

enough concentrations (Pouria et al. 1998, Chorus et al. 2000, Carmichael et al. 2001,

Carmichael 2001). Often, the recommended course of action for recreational waterbodies is to

limit activity in and exposure to bloom-affected waters due to the potential for accidental

ingestion and/or inhalation during recreational activities (Backer et al. 2010). Here, we explore

the effects of an extreme precipitation event on cyanobacterial blooms within Conestogo Lake

(Ontario, Canada; Figure 1), a eutrophic flood-control and river augmentation reservoir, which

has experienced recurring, late-fall cyanobacterial blooms dominated by the nitrogen-fixing

species *Aphanizomenon flos-aquae.* On 23 June 2017, the Upper Conestogo watershed received a

substantial amount of rainfall (daily avg. 78 mm measured at Conestogo Dam) that resulted in

severe flooding. According to the GRCA, nearly 80% of the volume of Conestogo Lake was

flushed downstream during this 2-day event (Grand River Conservation Authority 2018). In this

study, we were specifically interested in how significant rainfall from the catchment altered

water quality and nutrient concentrations and if so, how these changes impacted cyanobacterial

bloom development and toxin production.

Materials and methods

Study Site

Conestogo Lake is located in southwestern Ontario, Canada. It is a bottom-draw reservoir (7.35

km^2) operated by the Grand River Conservation Authority (GRCA; Mapleton Township, ON).

Built in 1958, it primarily serves as a flood-control and down-stream flow augmentation system

while providing ancillary recreational benefits for fishing, boating, and ~ 400 summer cottages.

Higher flows in spring from the Upper Conestogo River basin (566 km^2), a predominately

agricultural catchment (> 80 %, Figure 1, *left*), are captured in the reservoir through the end of

April and used during the summer drawdown (*i.e.* augmentation) period from 01 May to 30

September to provide consistent downstream flow of the Conestogo and Grand Rivers. As a

result of the reservoir drawdown, mean lake depth varies throughout the season with the deepest

point always near the dam. Reservoir storage volume during the summer augmentation period

has historically ranged between 12.7 and 56.3 Mm^3 (59.3 Mm^3 max cap.) with an average drop

between 5 and 7 m in stage elevation.

Sample collection and analysis

Sampling at Conestogo Lake began on 21 Jun 2017 with physical and biological samples

collected from the center (CLC) site closest to the dam. Two additional sites in the east (CLE)

and west (CLW) arms of the reservoir were added on 05 Jul and 11 Jul, respectively, following

the flooding event to assess the potential differences between catchment inputs into the reservoir

(Figure 1). From each location we collected a suite of physical, chemical, and biological data

throughout the water column. Physical water column profiles were collected at 0.5 m increments

using an EXO2 sonde (YSI, Yellow Springs, OH) equipped with chlorophyll-a, pH, temperature,

and dissolved oxygen sensors. The Secchi disk transparency at each site was used to collect

samples for cyanobacterial toxin analysis described in further detail below. In addition, we

collected discrete samples from 2 m for water chemistry and phytoplankton taxonomy and

enumeration approximately once per week from 21 June 2017 to 17 August 2017.

Whole water samples were collected for total phosphorus (TP) and phytoplankton

enumeration. Subsamples were field-filtered using a 0.45 μm syringe filter (Whatman) for

soluble reactive phosphorus (SRP), total dissolved phosphorus (TDP), ammonia (NH_4-N), nitrate

(NO_3^-), total dissolved nitrogen (TDN), anions, and cations. All samples were transported on ice,

stored at -20 °C, and analyzed at the Environmental Geochemistry Laboratory at the University

of Waterloo (Waterloo, ON) as per AHPA (1998; see Table S1 for instrument listing and

detection limits).

Phytoplankton identification and enumeration were completed by D. Findlay (Plankton-

R-Us, Winnipeg, MB, http://www.plankton-r-us.ca) as per Findlay & Kling (2003). Briefly,

phytoplankton samples were collected in a dark bottle, preserved with Lugol's solution, and

stored at 4 °C until analysis. Ten-mL aliquots of Lugol's preserved sample were gravity

settled for 24 h and counted using a modified Ütermohl technique (Nauwerk 1963) on

an inverted microscope with phase contrast illumination. Cell counts were converted to

wet weight biomass by approximating cell volume, which were obtained by

measurements of up to 50 cells of an individual species and applying the geometric

formula best fitted to the shape of the cell (Vollenweider 1968, Rott 1981). A specific

gravity of 1 was assumed for cellular mass. All biomass estimates are expressed as

mg/m^3 (D. Findlay, pers. communication).

Total (*i.e.*, intracellular and extracellular) cyanobacterial metabolite

concentrations were determined from a 115 mL whole water sample calculated from the

site-specific Secchi depth (2× Secchi depth) at each site. Each sample was collected in

an amber NalgeneTM polyethylene terephthalate glycol (PETG) bottle to limit

adsorption and overflow during freezing (Fisher reference: 322021-0125). All samples

were stored at -20 °C until analysis at the University of Montreal (Montréal, QC).

Samples were prepared and screened for each of seventeen cyanobacterial compounds

(Table S2) as per Fayad *et al.* (2015) via on-line solid phase extraction ultra-high

performance liquid chromatography high resolution mass spectrometry (SPE−UHPLC–

HRMS) using standards purchased from Enzo Life Science, Abraxis, or Cyano Biotech

GmbH. The limit of detection (LOD) and limit of quantification (LOQ) were calculated

for every batch of samples. Only results that exceeded the LOQ were included in our

analysis. Therefore, some samples may have had detectable, but unquantifiable toxin

concentrations.

Environmental and dam-related data

Meteorological and other reservoir-related data for this study were obtained from the GRCA's

online data portal (https://data.grandriver.ca/). Because GRCA data are provided under a

provisional status, all GRCA data were passed through quality assurance and quality checking

metrics for outliers to ensure sound data structure before proceeding with analysis. Spatial data

were obtained from open source portals at the GRCA (bathymetry), the United States Geological

Society, Statistics Canada, and the Ontario Ministry of Natural Resources (land-use).

Data analysis and statistics

All data analysis was completed in R version 3.5.1 (R Core Team 2018). Water column profile

data derived from sonde measurements as well as the discrete chemical profiles were constructed

using a multilevel B-spline interpolation from the *MBA* package (Finley et al. 2017) and were

adjusted to reflect the reservoir stage elevation at the time of sampling. One sampling in the east

arm contained a high chlorophyll-a concentration that skewed the profiles for all other sites.

Therefore, we log transformed (base 10) the data to allow for better visual comparison.

Comparisons of chemical and physical data across sites were completed using principal

coordinates analysis (PCoA) with a Euclidean matrix on log-normalized data to account for

scaling differences in the dataset and tested for difference through time and by site using

permutational analysis of variance (PERMANOVA) with *adonis* in the *vegan* package

(Okasanen et al. 2016). Similarly, we tested for differences in the phytoplankton community

composition by biomass using PCoA with a Bray-Curtis dissimilarity matrix on log-transformed

data to account for rare species biomass in the community and tested for difference through time

and by site using PERMANOVA. Biomass and toxin trends were analyzed using repeated

measures ANOVA (RM-ANOVA) with a linear mixed effects model (*lme*) from the *nlme*

package (Pinheiro et al. 2018) where time was a fixed effect, sampling location as a random

effect, and the correlation matrix selected based on Akaike Information Criterion (AIC).

Relationships between cyanobacterial biomass and environmental drivers including surface

water temperature and phosphorus concentrations were also completed using RM-ANOVA. In

addition, we tested for the correlation of biomass with toxin concentrations using Kendall's rho.

All code used for this project are available on github at

https://github.com/biogeochem/formbloom_conestogo_2017HAB.

Results

Flood event

The Upper Conestogo watershed received an intense rainfall event on 23-24 June 2017. This

two-day event increased mean daily flow from 4.56 m^3/s to 219 m^3/s (max 521 m^3/s), increased

the total reservoir volume by 3.5 Mm^3 (million cubic meters), and is estimated to have replaced

~80% of the reservoir volume (Grand River Conservation Authority 2018). Within several days,

the reservoir storage and drawdown returned to engineered levels (Figure 1*b-e*). The sizable

influx of water from the catchment during the flood event increased measurable phosphorus

between 1.9 to 33× within the water column at the center site by the next sampling on 05 July

2017 but did not affect any measured nitrogen species (Figures 2 and 3).

Hydrology and Descriptive Limnological Characteristics

All measured forms of phosphorus increased in the water column following the severe rainfall

event (Figures 2*a* and 3*a*, Tables 1 and S1). Epilimnetic concentrations declined throughout the

sampling period while hypolimnetic concentrations, notably TDP and TP, increased between 12

July (DOY 193) and 26 July (207) across all three sites. All nitrogen species (Figure 3b and S1,

Tables 1 and S1) were well mixed throughout the water column and gradually decreased

throughout the sampling campaign.

Conestogo Lake was initially and only briefly thermally stratified at the center site on 21

June 2017. By 05 July (DOY 186), the lake was isothermal across the lake and remained

isothermal throughout the rest of the summer augmentation period (Figure S2a). Mean surface

water temperatures (0 - 2 m) ranged from 21.2 to 25.5 °C. Subsequent flow from the catchment

following the 23 July rainfall reduced the surface water temperatures in the east by

approximately 2 °C from the previous sampling but did not affect the surface temperatures in the

west arm or center sites. Dissolved oxygen (Figure S2b, Tables 1 and S1) at the surface was

highest during peak biomass. Hypoxia developed in the hypolimnion at all three sampling

locations just as bloom biomass started to increase. Epilimnetic (0 - 2 m) chemical (TDN, NO_3^-,

DOC, SRP, TDP, TP, SO_4^{2-}, Mn, Ca^{2+}, K^+, Na^+, Mg^{2+}, F,$^-$, Cl^-) and physical (surface

temperature, dissolved oxygen, residence time) properties were significantly different between

sites (PERMANOVA, $P = 0.013$) and changed over the course of the sampling period

(PERMANOVA, $P = 0.001$; Figure S2).

Flooding induced cyanobacterial bloom

GRCA managers have visibly recorded cyanobacterial blooms in Conestogo Lake in late August

through September (S. Cooke, pers. communication), similar to other eutrophic lake systems in

this region (Winter et al. 2012). Within two weeks of the flooding event measurable increases in

biomass of *Aphanizomenon flos-aquae*, a filamentous nitrogen-fixer, were detected in each of the

sampling locations (Figure 4). Epilimnetic phytoplankton composition significantly changed

over the course of the sampling campaign with assemblages in the west and center more similar

than those in the east (Figure S3, PERMANOVA, site: $F = 2.22$, $P = 0.001$, time: $F = 2.69$, $P =$

0.003). While other phytoplankton groups were represented (Figure S4), cyanobacteria exceeded

50% of the phytoplankton community in the east and west arms near 12 July and approximately

a week later in the center. Total cyanobacterial biomass varied across sites and was significantly

affected by eplimnetic SRP, mean daily flow rates, and surface water temperatures

(RMANOVA, $F_{1,462} = 778.79$, $P < 0.0001$). The east arm contained the highest measured

biomass on 19 July (DOY 200, 10,900 mg/m^3, 4,500 cells/mL), which declined following

another, smaller, rainfall event on 23 Jul 2017. The center and west arm biomass peaked near 25

Jul (DOY 206; center: 4,710 mg/m^3, west: 5,230 mg/m^3). Bloom duration, measured as the

period during which cyanobacterial biomass composed > 50% of the fractional biomass,

persisted between 26 and 34 days and occurred 4 to 6 weeks earlier than has been previously

documented by the GRCA (S. Cooke, pers. communication). The cyanobacterial fraction across

all sampling sites also included *Woronichinia compacta*, *Anabaena (Dolichospermum) flos-*

aquae, *Microcystis aeruginosa*, *Anabaena* (*Dolichospermum) crassa*, *Planktolyngbya limnetica*,

and *Pseudoanabaena sp.* (Figure 4*a*). *W. compacta* exhibited similar biomass trends to *A. flos-*

aquae, though at much smaller biomass levels; by 17 Aug, its biomass increased to 8% (193.8

mg/m^3), 13% (108.9 mg/m^3), and 18% (585.9 mg/m^3) of the cyanobacterial biomass in the west,

center, and east, respectively.

Cyanotoxins were detectable at low concentrations

Detectable anabaenopeptin-A (AP-A), microcystin-LR (MC-LR), -YR, and -RR were present in

90 % of the samples ($n = 19$) but were only quantifiable in 81 % of samples ($n = 17$; Table 2).

Total microcystin concentration did not exceed 1.0 µg/L throughout the sampling season and

were comparable to previously measured concentrations during the summer months

(Yakobowski 2008). Metabolite type and concentration varied through time, but not by site

(RMANOVA, toxin × time, $F_{3,60} = 9.90$, $P < 0.0001$; site $P > 0.05$). Quantifiable concentrations

of all three microcystins were present in the east arm, MC-LR and MC-YR in the center site, and

only MC-LR in the west. MC-LR was the dominant microcystin variant across all sites with the

highest quantified values in the east arm at peak *A. flos-aquae*, *M. aeruginosa*, and *P. limnetica*

biomass (Figure 4*b*, Table 2). Further, MC-YR at the center did not correspond with the MC-LR

peak suggesting its possible origin in the east arm. AP-A was present at all sampling locations

throughout the bloom period, was the dominant metabolite at all sites by 25 July (DOY 206) and

had replaced all microcystin variants by the end of the sampling campaign. Toxin concentrations

were not significantly correlated with the biomass of any particular cyanobacterial species

(Kendall rho, $P > 0.05$).

Discussion

Conestogo Lake has experienced annual cyanobacterial blooms over the last decade, some of

which have resulted in the issuance of health advisories for recreational users and cottagers (S.

Cooke, GRCA, pers. communication). Consistent with other eutrophic lake and reservoir

systems in the region (Winter et al. 2012), cyanobacterial blooms typically develop in mid- to

late-August and persist through September or early October (S. Cooke, GRCA, pers.

Communication, Larsen *et al.* in prep). However, the marked increase in soluble P as a result of

rainfall combined with rapid physical change from disruption of water column structure during

flooding to reestablishment of a stable water column made 2017 an atypical season with an

unusually early bloom. This highlights the potential for changing windows of bloom-related risk

and associated management considerations.

Global increases in extreme rainfall events are predicted to outpace changes in

total precipitation under various climate models (Allen and Ingram 2002, IPCC 2007)

suggesting that heavy rain and flooding events will likely become more common in the

near future. The impact of changes to precipitation intensity and frequency on

cyanobacterial bloom development and dynamics is inherently complex and may lead to

either promotion or disruption of bloom dynamics (Reichwaldt and Ghadouani 2012,

Wood et al. 2017). Interactions between physical, chemical, and biological lake

parameters are influenced by other factors such as catchment size, land use, soil type,

etc. and ultimately, the overall effect of the rainfall event is regulated by the rainfall

intensity, water inflow, and seasonal timing (reviewed in Reichwaldt and Ghadouani

2012). Limited studies investigating such events on cyanobacterial bloom development

have identified changes to flushing rates, water column mixing, and nutrient inputs from

rainfall events as the main factors affecting cyanobacteria and phytoplankton

communities (Bouvy et al. 2003, Reichwaldt and Ghadouani 2012) – all factors that

appeared to impact the bloom dynamics in Conestogo Lake. Regional climate models

suggest we will continue to see impacts of extreme events that can affect bloom timing,

and risk (Bouvy et al. 2003, IPCC 2007, McDermid et al. 2015).

Sizable influxes of dissolved and particulate nutrient pools are common following

heavy rainfall and erosion, particularly if they are preceded by a warm, dry periods (Jones and

Poplawski 1998, Chiew et al 1995). For example, in Australian reservoirs, a 440 mm rainfall in a

3-day period resulted in an input equivalent of 80% and 400% of the average annual in-lake N

and P, respectively (Jones and Poplawski 1998). Here we observed a 27x increase in SRP levels

in surficial waters, which was likely key to the proliferation and dominance of cyanobacteria.

Emerging cyanotoxins

Many lake monitoring programs focus on the measurement of microcystins because of their vast

distribution (Loftin et al. 2016), ecotoxicity (Chorus and Bartram 1999), and relatively

inexpensive and rapid detection in whole water samples (*e.g.* ELISA test kits). However, several

studies using cyanobacterial extracts have reported harmful/toxic effects that could not be

explained solely by microcystin concentration or presence, suggesting the possibility of other

toxic compounds (Keil et al. 2002, Teneva et al. 2005, Baumann and Jüttner 2008, Smutná et al.

2014, Lenz et al. 2019). Improved analytical techniques have identified numerous bioactive

compounds such as cyanopeptolins and anabaenopeptins that are readily detectable in

freshwaters and often produced simultaneously with microcystin variants (Harada et al. 1995,

Welker and Von Döhren 2006, Gkelis et al. 2015, Beversdorf et al. 2017, 2018).

In some cases, and consistent with our results, anabaenopeptins are reported in equal or

higher concentrations than microcystins (Janssen 2019). For example, Beversdorf *et al.* (2017)

reported an average of 0.65 μg/L total microcystin in Lake Koshkonong, Wisconsin, while

anabaenopeptin-B and -F were measured at 6.56 μg/L combined. Though there are no case

studies of human toxicity caused by anabaenopeptins, the compound inhibits carboxypeptidase

A, and like microcystin, also inhibits protein phosphatases with slightly overlapping inhibitory

concentration ranges (Honkanen et al. 1990, Sano et al. 2001, Spoof et al. 2016). At present, the

concentration to which anabaenopeptins and other metabolites would have to reach to affect

human health is currently unknown resulting in a lack of regulations or advisories for

recreational or drinking water. However, ecological effects may be observed at relatively low

concentrations. A recent study by Lenz *et al.* (2019) reported induced toxicity by low

concentrations (10 μg/L) of anabaenopeptins, including AP-A, on the nematode *Caenorhabditis*

elegans resulting in reduced reproduction, reduced lifespan, and delayed hatching. Though

compounds like AP-A have been previously considered non-toxic, they may now represent a

new class of emerging toxins, whose potential impacts to human health and toxicity to aquatic

organisms require immediate attention and therefore, inclusion in risk assessment for lake

mitigation and monitoring programs

Recommendations for management

Watershed managers have various options to mitigate cyanobacterial blooms by disrupting

conditions that favor growth. These strategies, including increasing flushing rates, reducing

internal/external nutrient loads, destratisfying the water column through mixing, application of

chemicals, and/ or biological manipulations, each come with varying economic costs and have

been extensively reviewed elsewhere (Paerl 2018). Different short-term and long-term targets for

water management regimes and/or nutrient loads will likely be needed to mitigate cyanobacterial

bloom formation in Conestogo Lake. However, manipulating water flow to reduce residence

time is not likely a feasible option since the reservoir operates with base-flow augmentation as a

primary deliverable. Therefore, management goals will likely need to incorporate strategies to

mitigate internal and external nutrient loads. TDN nor NO_3 concentrations increased in response

to the flood event and steadily decreased throughout the campaign. This suggests that N is not

likely the regulating nutrient for biomass in this system though it may play a role in selecting for

cyanobacterial species, bloom timing, or contributing to other biogeochemical cycles.

Concurrent with other work on Conestogo Lake (Guildford 2006), our results suggest that P

management strategies should be the main given the increased SRP concentrations in the water

column following the flood and hypolimnetic P release from anoxic sediments. Further work will

be needed to determine whether, if in a typical year, internal P release is the main driver for the

late August or early September cyanobacterial blooms that have been previously documented

anecdotally.

Short-term strategies such as chemical binding (*e.g.* Phoslock or alum) will precipitate

suspended P to the sediment and could therefore reduce the size and/or severity of the bloom

(Robb et al. 2003, Lürling and Faassen 2012, Cooke et al. 2013). However, keeping P

immobilized for long periods is difficult in systems with strong anoxic bottom waters like

Conestogo Lake. Longer-term strategies could include reduced P catchment loads via changes to

the release of P-rich wastewater from upstream water treatment facilities, the incorporation of an

increased riparian zone and/or dredging (Lürling and Faassen 2012, Aguiar et al. 2015).

Increased riparian zones have been effective in reducing particulate P in various other systems

(Aguiar et al. 2015) and may be a possible strategy since the predominant Tavistill soil type in

the Upper Conestogo River catchment is highly susceptible to erosion (Macrae et al. 2007, Grand

River Conservation Authority 2018). At present, only a small strip of forested land surrounds the

main body of the reservoir, hence reduction of localized inputs may be achievable, even if

riverine inputs are more difficult to control.

This task of identifying solutions to mitigate cyanobacterial blooms is challenging – yet

urgent – given widespread reports of worsening bloom risk (Pick 2016). Development of

appropriate mitigation strategies must incorporate an understanding of watershed nutrient

sources, physiochemical lake structure, biological conditions, and climatic predictions into an

adaptive framework considering what is possible, what are the costs (financial costs, and trade-

offs to ecosystem services), and what are the goals of ecosystem and watershed management.

Within Conestogo Lake we demonstrate that extreme rainfall events can trigger early bloom

development. Management strategies and risk reduction activities, which currently target the

low flow period in late summer may need to be altered in light of predictions of more frequent

extreme rainfall events which may trigger an earlier bloom season in some ecosystems.

Acknowledgments

Funding for this work was provided by the Canada First Research Excellence Fund program.

This was a part of the Global Water Futures Initiative FORMBLOOM: FORecasting tools and

Mitigation options for diverse BLOOM-affected lakes as well as the ATRAPP project (Algal

Blooms, Treatment, Risk Assessment, Prediction and Prevention through Genomics) with from

Genome Canada and Genome Quebec. We thank S. Cooke, D. McFadden, and the Grand River

Conservation Authority for their cooperation and knowledgeable insight, R. Elgood, E. McQuay,

T. Cornell, B. Gruber, S. Sine, and M. Soares-Paquin for assistance with sample collection and

analysis, J. Atkins for her GIS assistance, and the members of the Venkiteswaran and Schiff lab

groups for their reviews of manuscript drafts and friendly discussion.

Data availability statement

Data and code that support the analysis and findings in this study are available in figshare at

http://doi/10.6084/m9.figshare.7811963.

Literature Cited

Aguiar, T. R., K. Rasera, L. M. Parron, A. G. Brito, and M. T. Ferreira. 2015. Nutrient removal

effectiveness by riparian buffer zones in rural temperate watersheds: The impact of no-till

crops practices. Agricultural Water Management 149:74–80.

Allen, M. R., and W. J. Ingram. 2002. Constraints on future changes in climate and the

hydrologic cycle. Nature 419:224–232.

Backer, L. C., S. V. McNeel, T. Barber, B. Kirkpatrick, C. Williams, M. Irvin, Y. Zhou, T. B.

Johnson, K. Nierenberg, M. Aubel, R. LePrell, A. Chapman, A. Foss, S. Corum, V. R. Hill,

S. M. Kieszak, and Y. S. Cheng. 2010. Recreational exposure to microcystins during algal

blooms in two California lakes. Toxicon 55:909–921.

Baumann, H. I., and F. Jüttner. 2008. Inter-annual stability of oligopeptide patterns of

Planktothrix rubescens blooms and mass mortality of Daphnia in Lake Hallwilersee.

Limnologica 38:350–359.

Beversdorf, L. J., K. Rude, C. A. Weirich, S. L. Bartlett, M. Seaman, C. Kozik, P. Biese, T.

Gosz, M. Suha, C. Stempa, C. Shaw, C. Hedman, J. J. Piatt, and T. R. Miller. 2018.

Analysis of cyanobacterial metabolites in surface and raw drinking waters reveals more than

microcystin. Water Research 140:280–290.

Beversdorf, L. J., C. A. Weirich, S. L. Bartlett, and T. R. Miller. 2017. Variable cyanobacterial

toxin and metabolite profiles across six eutrophic lakes of differing physiochemical

characteristics. Toxins 9.

Bouvy, M., S. M. Nascimento, R. J. R. Molica, A. Ferreira, V. Huszar, and S. M. F. O. Azevedo.

2003. Limnological features in Tapacurá reservoir (northeast Brazil) during a severe

drought. Hydrobiologia 493:115–130.

Bullerjahn, G. S., R. M. McKay, T. W. Davis, D. B. Baker, G. L. Boyer, L. V. D'Anglada, G. J.

Doucette, J. C. Ho, E. G. Irwin, C. L. Kling, R. M. Kudela, R. Kurmayer, A. M. Michalak,

J. D. Ortiz, T. G. Otten, H. W. Paerl, B. Qin, B. L. Sohngen, R. P. Stumpf, P. M. Visser, and

S. W. Wilhelm. 2016. Global solutions to regional problems: Collecting global expertise to

address the problem of harmful cyanobacterial blooms. A Lake Erie case study. Harmful

Algae 54:223–238.

Carmichael, W. W. 1997. The Cyanotoxins. Advances in Botanical Research 27:211–256.

Carmichael, W. W. 2001. Health Effects of Toxin-Producing Cyanobacteria: "The

CyanoHABs." Human and Ecological Risk Assessment: An International Journal 7:1393–

1407.

Carmichael, W. W., S. M. F. O. Azevedo, J. S. An, R. J. R. Molica, E. M. Jochimsen, S. Lau, K.

L. Rinehart, G. R. Shaw, and G. K. Eaglesham. 2001. Human fatalities form cyanobacteria:

Chemical and biological evidence for cyanotoxins. Page Environmental Health

Perspectives.

Chorus, I., and J. Bartram, editors. 1999. Toxic Cyanobacteria in Water: A guide to their public

health consequences, monitoring and management. E & FN Spon, London and New York.

Chorus, I., I. R. Falconer, H. J. Salas, and J. Bartram. 2000. Health Risks Caused By Freshwater

Cyanobacteria in Recreational Waters. Journal of Toxicology and Environmental Health,

Part B 3:323–347.

Codd, G. A. 2000. Cyanobacterial toxins, the perception of water quality, and the prioritisation

of eutrophication control. Ecological Engineering 16:51–60.

Cooke, G. D., E. B. Welch, S. A. Peterson, and S. A. Nichols. 2013. Restoration and

Management of Lakes and Reservoirs. Page Restoration and Management of Lakes and

Reservoirs. Third. Taylor & Francis, New York, NY, USA.

Dodds, W. K., W. W. Bouska, J. L. Eitzmann, T. J. Pilger, K. L. Pitts, A. J. Riley, J. T.

Schloesser, and D. J. Thornbrugh. 2009. Eutrophication of U. S. freshwaters: Analysis of

potential economic damages. Environmental Science and Technology 43:12–19.

Fayad, P. B., A. Roy-Lachapelle, S. V. Duy, M. Prévost, and S. Sauvé. 2015. On-line solid-phase

extraction coupled to liquid chromatography tandem mass spectrometry for the analysis of

cyanotoxins in algal blooms. Toxicon 108:167–175.

Findlay, D. . L., and H. J. Kling. 2003. Protocols for measuring biodiversity: phytoplankton in

freshwater. Winnipeg: Department of Fisheries and Oceans.

Finley, A., S. Banerjee, and Ø. Hjelle. 2017. MBA: Multilevel B-Spline Approximation.

Gkelis, S., T. Lanaras, K. Sivonen, and O. Taglialatela-Scafati. 2015. Cyanobacterial toxic and

bioactive peptides in freshwater bodies of Greece: Concentrations, occurrence patterns, and

implications for human health. Marine Drugs 13:6319–6335.

Grand River Conservation Authority. 2018. Belwood and Conestogo Water Management

Reservoirs : An Assessment of Surface and Groundwater Conditions DRAFT January 2018.

Guildford, S. J. 2006. Factors controlling cyanobacteria blooms in three Grand River Basin

reservoirs during 2005.

Harada, K., K. Fujii, T. Shimada, M. Suzuki, H. Sano, K. Adachi, and W. W. Carmichael. 1995.

Two cyclic peptides, anabaenopeptins, a third group of bioactive compounds from the

cyanobacterium *Anabaena flos-aquae* NRC 525-17. Tetrahedron Letters 36:1511–1514.

Honkanen, R. E., J. Zwiller, R. E. Mooren, S. L. Daily, B. S. Khatra, M. Dukelow, and A. L.

Boynton. 1990. Characterization of Microcystin- LR, a potent inhibitor of type 1 and type

2A protein phosphatases. Journal of Biological Chemistry 265:19401–19404.

IPCC. 2007. Climate Change 2007: The Scientific Basis. Contribution of Working Group I to the

Fourth Assessment report of the Intergovernmental Panel on Climate Change. New York,

NY, USA.

Jacobsen, B. A., and P. Simonsen. 1993. Disturbance events affecting phytoplankton biomass,

composition and species diversity in a shallow, eutrophic, temperate lake. Page

Hydrobiologia.

Janssen, E. M.-L. 2019. Cyanobacterial peptides beyond microcystins – A review on co-

occurrence, toxicity, and challenges for risk assessment. Water Research 151:488–499.

Jöhnk, K. D., J. Huisman, J. Sharples, B. Sommeijer, P. M. Visser, and J. M. Strooms. 2008.

Summer heatwaves promote blooms of harmful cyanobacteria. Global Change Biology

14:495–512.

Jones, G. J., and W. Poplawski. 1998. Understanding and management of cyanobacterial blooms

in sub-tropical reservoirs of Queensland, Australia. Water Science and Technology 37:161–

168.

Keil, C., A. Forchert, J. Fastner, U. Szewzyk, W. Rotard, I. Chorus, and R. Krätke. 2002.

Toxicity and microcystin content of extracts from a Planktothrix bloom and two laboratory

strains. Water Research 36:2133–2139.

Lenz, K. A., T. R. Miller, and H. Ma. 2019. Anabaenopeptins and cyanopeptolins induce

systemic toxicity effects in a model organism the nematode *Caenorhabditis elegans*.

Chemosphere 214:60–69.

Loftin, K. A., J. L. Graham, E. D. Hilborn, S. C. Lehmann, M. T. Meyer, J. E. Dietze, and C. B.

Griffith. 2016. Cyanotoxins in inland lakes of the United States: Occurrence and potential

recreational health risks in the EPA National Lakes Assessment 2007. Harmful Algae

56:77–90.

Lürling, M., and E. J. Faassen. 2012. Controlling toxic cyanobacteria: Effects of dredging and

phosphorus-binding clay on cyanobacteria and microcystins. Water Research 46:1447–

1459.

Macrae, M. L., M. C. English, S. L. Schiff, and M. Stone. 2007. Intra-annual variability in the

contribution of tile drains to basin discharge and phosphorus export in a first-order

agricultural catchment. Agricultural Water Management 92:171–182.

Mitrovic, S. M., R. L. Oliver, C. Rees, L. C. Bowling, and R. T. Buckney. 2003. Critical flow

velocities for the growth and dominance of *Anabaena circinalis* in some turbid freshwater

rivers. Freshwater Biology 48:164–174.

Nauwerk, A. 1963. Die Beziehungen zwischen Zooplankton und Phytoplankton im See Erken.

Symbolae Botanicae Uppsaliensis 8:5–162.

O'Neil, J. M., T. W. Davis, M. A. Burford, and C. J. Gobler. 2012. The rise of harmful

cyanobacteria blooms: The potential roles of eutrophication and climate change. Harmful

Algae 14:313–334.

Okasanen, J., F. G. Blanchet, R. Kindt, P. Legendre, P. R. Minchin, R. B. O'hara, G. L. Simpson,

P. Solymos, M. H. H. Stevens, and H. Wagner. 2016. vegan: Community Ecology Package.

Paerl, H. W. 2018. Mitigating toxic planktonic cyanobacterial blooms in aquatic ecosystems

facing increasing anthropogenic and climatic pressures. Toxins 10:1–16.

Paerl, H. W., W. S. Gardner, K. E. Havens, A. R. Joyner, M. J. McCarthy, S. E. Newell, B. Qin,

and J. T. Scott. 2016. Mitigating cyanobacterial harmful algal blooms in aquatic ecosystems

impacted by climate change and anthropogenic nutrients. Harmful Algae 54:213–222.

Paerl, H. W., and J. Huisman. 2009. Climate change: A catalyst for global expansion of harmful

cyanobacterial blooms. Environmental Microbiology Reports 1:27–37.

Paerl, H. W., and T. G. Otten. 2013. Harmful Cyanobacterial Blooms: Causes, Consequences,

and Controls. Microbial Ecology 65:995–1010.

Pearl, H. W. 1988. Nuisance phytoplankton blooms in coastal, estuarine, and inland waters. Page

Limnology and Oceanography.

Pick, F. R. 2016. Blooming algae: a Canadian perspective on the rise of toxic cyanobacteria.

Canadian Journal of Fisheries and Aquatic Sciences 73:1149–1158.

Pinheiro, J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team. 2018. {nlme}: Linear and

Nonlinear Mixed Effects Models.

Pouria, S., A. De Andrade, J. Barbosa, R. L. Cavalcanti, V. T. S. Barreto, C. J. Ward, W. Preiser,

G. K. Poon, G. H. Neild, and G. A. Codd. 1998. Fatal microcystin intoxication in

haemodialysis unit in Caruaru, Brazil. Lancet 352:21–26.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. Vienna,

Austria.

Reichwaldt, E. S., and A. Ghadouani. 2012. Effects of rainfall patterns on toxic cyanobacterial

blooms in a changing climate: Between simplistic scenarios and complex dynamics. Water

Research 46:1372–1393.

Reynolds, C. S. 1984. Ecology of Freshwater Phytoplankton. Cambridge University Press.

Robb, M., B. Greenop, Z. Goss, G. Douglas, and J. Adeney. 2003. Application of Phoslock™, an

innovative phosphorus binding clay, to two Western Australian waterways: Preliminary

findings. Page Hydrobiologia.

Rott, E. 1981. Some results from phytoplankton counting intercalibrations. Schweizerische

Zeitschrift für Hydrologie 43:34–62.

Sano, T., T. Usui, K. Ueda, H. Osada, and K. Kaya. 2001. Isolation of new protein phosphatase

inhibitors from two cyanobacteria species, *Planktothrix* spp. Journal of Natural Products

64:1052–1055.

Smutná, M., P. Babica, S. Jarque, K. Hilscherová, B. Maršálek, M. Haeba, and L. Bláha. 2014.

Acute, chronic and reproductive toxicity of complex cyanobacterial blooms in *Daphnia*

magna and the role of microcystins. Toxicon 79:11–18.

Spoof, L., A. Błaszczyk, J. Meriluoto, M. Cegłowska, and H. Mazur-Marzec. 2016. Structures

and activity of new anabaenopeptins produced by Baltic Sea cyanobacteria. Marine Drugs

14:8.

Sukenik, A., A. Quesada, and N. Salmaso. 2015. Global expansion of toxic and non-toxic

cyanobacteria: effect on ecosystem functioning. Biodiversity and Conservation 24:889–908.

Teneva, I., B. Dzhambazov, L. Koleva, R. Mladenov, and K. Schirmer. 2005. Toxic potential of

five freshwater *Phormidium* species (Cyanoprokaryota). Toxicon 45:711–725.

Vollenweider, R. A. 1968. Scientific fundamentals of the eutrophication of lakes and flowing

waters, with particular reference to nitrogen and phosphorus as factors in eutrophication.

Page Paris: Organisation for Economic Co-operation and Development. Technical Report

DAS/CS1/68.27.

Welker, M., and H. Von Döhren. 2006. Cyanobacterial peptides - Nature's own combinatorial

biosynthesis. FEMS Microbiology Reviews 30:530–563.

Wetzel, R. G. 2001. Limnology : lake and river ecosystems. Third. Academic Press, San Diego.

Winter, J. G., E. T. Howell, and L. K. Nakamoto. 2012. Trends in nutrients, phytoplankton, and

chloride in nearshore waters of Lake Ontario: Synchrony and relationships with physical

conditions. Journal of Great Lakes Research 38:124–132.

Wood, S. A., H. Borges, J. Puddick, L. Biessy, J. Atalah, I. Hawes, D. R. Dietrich, and D. P.

Hamilton. 2017. Contrasting cyanobacterial communities and microcystin concentrations in

summers with extreme weather events: insights into potential effects of climate change.

Hydrobiologia 785:71–89.

Yakobowski, S. J. 2008. Ecological factors controlling microcystin concentrations in the Bay of

Quinte, Maumee Bay, and three Grand River Reservoirs. University of Waterloo.

Table 1. Chemical water column summary with mean (SD) [minimum – maximum] at sampling

each site.

Site	NO$_3^-$ (mg-N/L)	TDN (mg-N/L)	SRP (µg-P/L)	TDP (µg-P/L)	TP (µg-P/L)
Center	4.82 (0.59) [3.99 - 5.65]	5.32 (0.64) [4.24 - 5.86]	13.66 (22.86) [1.52 - 65.2]	26.41 (26.32) [9.96 - 85.77]	39.84 (27.59) [14.99 - 103.48]
East	4.37 (0.79) [3.34 - 5.63]	4.74 (0.89) [3.39 - 5.85]	5.89 (8.45) [1.56 - 24.61]	18.53 (11.14) [7.79 - 41.59]	49.54 (11.08) [35.2 - 68.65]
West	4.69 (0.57) [3.89 - 5.43]	5.1 (0.83) [3.94 - 6.03]	4.09 (4.73) [1.23 - 13.48]	15.99 (7.7) [8.37 - 29.83]	39.35 (4.26) [34.69 - 47.04]

564 **Table 2**. Quantified cyanobacterial metabolites across sampling sites as measured by LC-HMRS

565 with values which exceeded the limit of detection (LOQ) presented as mean ng/L (standard

566 deviation ng/L), those in italics exceeded the limit of detection (LOD) but not the limit of

567 quantification (LOQ), and those with (-) < LOD. The metabolites MC-LA, -LY, -LW, -LF, -

568 HiIR, -HtyR, and CPA were not detected in samples.

Date	Site	cyanobacterial secondary metabolites						
		AP-A	MC-LR	MC-RR	MC-YR	ATX	HATX	CYN
21 Jun	Center	-	-	-	-	-	-	-
05 Jul		-	-	-	*33.26 (16.08)*	-	-	-
14 Jul		*23.72 (4.89)*	-	-	*20.6 (29.13)*	-	-	-
18 Jul		109.25 (78.62)	406.88 (18.06)	-	-	-	-	-
25 Jul		189.65 (35.58)	88.18 (6.22)	*35.47 (9.17)*	*19.95 (28.21)*	-	-	-
01 Aug		278.84 (27.42)	16.77 (10.62)	-	59.22 (16.5)	-	-	-
10 Aug		619.33 (140.95)	-	-	*19.41 (1.29)*	-	-	-
17 Aug		343.99 (50.32)	-	-	-	-	-	-
05 Jul	East	-	-	-	-	-	-	-
11 Jul		*18.59 (26.29)*	-	-	-	-	-	-
19 Jul		-	763.25 (154.17)	120.97 (1.56)	97.26 (36.16)	-	-	-
25 Jul		174.25 (34.91)	72.07 (23.34)	*27.59 (2.67)*	-	-	-	-
31 Jul		188.58 (108.52)	-	-	-	-	-	-
10 Aug		411 (30.56)	-	-	-	-	-	-
17 Aug		856.77 (37.15)	-	-	-	-	-	-
11 Jul	West	51 (24.31)	-	-	-	-	-	-
19 Jul		160.79 (14.53)	211.23 (23.56)	-	-	-	-	-
25 Jul		33.89 (25.24)	106.53 (28.81)	*30.03 (8.44)*	-	-	-	-
31 Jul		242.35 (29.74)	-	-	-	-	-	-
10 Aug		471.57 (75.91)	-	-	*26.43 (37.38)*	-	-	-
17 Aug		846.72 (0.53)	-	-	-	-	-	-

Abbreviations: anabaenopeptin A (AP-A), microcystin (MC-), anatoxin (ATX), homoanatoxin-a (HATX), cylindrospermopsin (CYN), cyanopeptolin-a (CPA)

Figure 1. Watershed and bathymetric (*a*) map of Conestogo Lake, Mapleton Township, Ontario

with sampling sites for the west arm (CLW), center (CLC), and east arm (CLE) of the reservoir.

Bathymetric data is scaled to the maximum dam stage elevation (meters at sea level; m.a.s.l.).

Total precipitation in the Upper Conestogo watershed (*b*, gauge in Arthur, ON) is directly related

to the mean daily flow rate (*c*) into the east arm (gauges in Drayton and Moorefield, ON) and the

calculated residence time (*d*) in the reservoir. Average flow rate (red line, *c*) is typically near

0.01 m/s. An intense rainfall event on 23-24 Jun (grey vertical bar) resulted in a notable increase

in the mean daily storage volume (1000 m^3) of the reservoir (*e*). Dark gray bars represent

standard deviation. This figure contains information made available under GRCA's Open Data

Licence v3.0 (panel a) and v2.0 (panels *b* - *e*).

Figure 2. Multiplicative changes in measured water chemistry between samplings before (21 Jun

2017) and after the flood event (05 Jul 2017) at 2, 7, and 16 m depths at the center site. Values

above 1 correspond to increased concentrations while those below 1 indicate reduced

concentrations. (*a*) Total phosphorus (TP), total dissolved phosphorus (TDP), and soluble

reactive phosphorus (SRP), (*b*) anions, (*c*) Nitrogen species including total dissolved nitrogen

(TDN), nitrate (NO$_3^-$), nitrite (NO$_2^-$), ammonium (NH$_4^+$), and dissolved organic carbon (DOC),

and (*d*) dissolved cations.

Figure 3. Nutrient profiles for phosphorus (*a***;** P, μg-P/L) and nitrogen species (*b*; N, mg-N/L)

collected at 2, 7, and 0.5m from the bottom at each of the sampling locations. The flood event

(white vertical bar) introduced increased levels of all measured P species but did not greatly

affect N species (see Figure S1). The dynamic area of the colored box reflects the changing

depths within each sample location. Profile depths at each site were standardized to the dam

stage elevation (m) at the time of sampling and illustrate both the differences in bottom depth

and reservoir drawdown during the sampling campaign.

Figure 4. Cyanobacterial biomass (*a*) and cyanotoxin concentration trends > LOQ for

anabaenopeptin-A (AP.A) and microcystin-LR (MC.LR), -YR (MC.YR), and -RR (MC.RR) (*b*)

in each of the three sampling locations (*l* to *r:* West, Center, and East) were detected

approximately 2 weeks following the flood event on 23-24 Jun (DOY 172; grey vertical bar).

Aphanizomenon flos-aquae dominated (> 50%) phytoplankton biomass in the west and east arms

of Conestogo Lake near 10 Jul 2017 (DOY 191) and in the center near 16 Jul 2017 (DOY 197).

The bloom period lasted between 26-34 days depending on the sampling site with a site-specific

peak bloom biomass occurring near 25 Jul 2017 (DOY 206). Of the seven cyanobacterial species

detected in Conestogo Lake, various strains of *A. flos-aquae*, *D. flos-aquae* (*Anabaena flos-*

aquae), *M. aeruginosa*, and *W. compacta* have been documented as potential toxin producers.

Figure 1
612

614   **Figure 2**

Figure 3

   Figure 4