A high-throughput delayed fluorescence method reveals underling differences in the control

of circadian rhythms in *Triticum aestivum* and *Brassica napus*

Hannah Rees[1,2], Susan Duncan[1], Peter Gould[2], Rachel Wells[3], Mark Greenwood[4,5], Thomas

Brabbs[1], Anthony Hall[1*]

[1]Earlham Institute, Norwich Research Park, Norwich, NR4 7UG, UK

[2]Institute of Integrative Biology, University of Liverpool, Crown Street, Liverpool, L69 7ZB,

UK

[3]John Innes Centre, Norwich Research Park, Norwich NR4 7UH, UK

[4]Sainsbury Laboratory, University of Cambridge, Cambridge, CB2 1LR, UK

[5]Department of Biochemistry, University of Cambridge, Cambridge, CB2 1QW, UK

*Email anthony.hall@earlham.ac.uk

Abstract

Background

A robust circadian clock has been implicated in plant resilience, resource-use efficiency, competitive growth and yield. A huge number of physiological processes are under circadian control in plants including: responses to biotic and abiotic stresses; flowering time; plant metabolism; and mineral uptake. Understanding how the clock functions in crops such as *Triticum aestivum* (bread wheat) and *Brassica napus* (oilseed rape) therefore has great agricultural potential. Delayed fluorescence (DF) imaging has been shown to be applicable to a wide range of plant species and requires no genetic transformation. Although DF has been used to measure period length of both mutants and wild ecotypes of *Arabidopsis*, this assay has never been systematically optimised for crop plants. The physical size of both *B. napus* and *T. aestivum* led us to develop a representative sampling strategy which enables high-throughput imaging of these crops.

Results

In this study, we describe the plant-specific optimisation of DF imaging to obtain reliable circadian phenotypes with the robustness and reproducibility to detect diverging periods between cultivars of the same species. We find that the age of plant material, light regime and temperature conditions all significantly effect DF rhythms and describe the optimal conditions for measuring robust rhythms in each species. We also show that sections of leaf can be used to obtain period estimates with improved throughput for larger sample size experiments.

Conclusions

We present an optimized protocol for high-throughput phenotyping of circadian period

specific to two economically valuable crop plants. Application of this method revealed

significant differences between the periods of several widely grown elite cultivars.

This method also identified intriguing differential responses of circadian rhythms in *T.*

aestivum compared to *B. napus*; specifically the dramatic change to rhythm robustness

when plants were imaged under constant light versus constant darkness. This points

towards diverging networks underling circadian control in these two species.

Keywords: Circadian period, delayed fluorescence, free-running conditions, Hexaploid

wheat, oilseed rape, aging, L:L, D:D, temperature, rhythm robustness

Background

A circadian clock is an endogenous oscillator entrained by external temporal cues. Circadian

control of gene expression is a ubiquitous feature which appears to have arisen

independently in bacteria, fungi, plants and animals(1). Since the discovery of the first

Arabidopsis circadian mutant in 1995(2), the significance of the circadian clock in plants has

become increasingly evident. Approximately 30% of genes in *Arabidopsis* are predicted to

be under circadian control, regulating photosynthetic, metabolic and developmental

pathways(3,4). Moreover, a selective advantage resulting from a clock which is matched to

the exogenous day-length has been demonstrated in mammals, insects, bacteria and

plants(5–9).

The most recent model for the molecular control of the *Arabidopsis* clock is comprised of a

series of interlocking negative transcriptional feedback loops regulated by key activators

which control the oscillation of clock gene expression(10). To ascertain the underlying

nature of circadian rhythms, a clock-controlled output representing the pace of the clock

must be measured in constant (free-running) conditions. Previously this research has been

conducted by studying leaf movement rhythms or by following luciferase gene expression

under the control of a circadian regulated promoter(11–13). Delayed fluorescence (DF)

imaging provides an alternative to these methods that does not require plant

transformation. It has previously been shown to work in a variety of plants for which leaf

movement assays are not feasible(14,15). Delayed fluorescence occurs when excited

electrons in photosystem II (PSII) undergo spin-conversion to a triplet excited state before

charge recombination allows them to return to their ground state releasing light energy(16).

Measurements of DF have been correlated with the photosynthetic state of PSII(17) and the

amount of DF production is regulated by the circadian clock. DF can be measured with a

low-light imaging system identical to that used for luciferase imaging and output rhythms

have been shown to oscillate with a comparable period to those estimated from luciferase

reporter experiments(14). The output from a DF experiment is a waveform which has

parameters that can be mathematically defined and therefore quantified. These parameters

include 'period' (the time taken to complete one cycle), 'phase' (the time of day at which

this peaks) and 'amplitude' (the distance between the peak and the baseline of the

oscillation). Important to circadian dynamics is also the idea of 'rhythm robustness' i.e.

whether these parameters change over time. In this paper, rhythm robustness was assessed

by: the percentage of samples classified as rhythmic; the relative amplitude error (RAE); the

period CV and the average period error threshold (all defined in Supplementary Materials).

Together, these parameters allow the effects of different imaging conditions to be

quantified.

As DF measurement is correlated with the oscillations in photosynthetic status of PSII, leaf

material is the logical choice for a representative sample. Rhythms have been shown to

persist in excised leaves in several species(18–22). However, previous research has

demonstrated that independent clocks run at different periods throughout the plant under

constant conditions, coordinated by a degree of intercellular coupling(23–27). The extent to

which the clock is affected by dissecting leaf material into small segments is investigated in

this paper.

Alongside these spatial differences, the clock has also been shown to be temporally dynamic

and is affected by the life history of the plant. Both the systemic age of the plant and the

'emergence age' of the individual leaves on a plant have been reported to effect the clock in

Arabidopsis, with increasing age associated with period reduction(28). Conversely, the

timing of leaf senescence has also been shown to be directly regulated by core circadian

genes(29).

In addition to this endogenous entrainment, the clock is also responsive to external stimuli;

the most well characterized of which are light and temperature cues. Increasing light

intensity causes a shortening of period in free-running conditions(30–32) and these rhythms

rapidly dampen in amplitude under continuous darkness(33). Circadian systems are

relatively buffered against temperature changes compared to other biochemical reactions

but are not completely independent of it(34). Period shortening of 1.8–4.2h have been

reported following temperature increases from 17°C to 27°C determined by both leaf-

movement assays and luciferase reporters under circadian regulation in

Arabidopsis(31,35,36). Seedlings grown at 17°C also have rhythms with lower period

variability and RAE values than plants grown at 27°C(35,37). The extent to which rhythms

are temperature compensated is described using the inverse of the temperature coefficient

Q10; the change in the rate of a process over a temperature change of 10°C(38).

Here we present an optimized protocol for high-throughput phenotyping of circadian period

using two crop plant models; *Triticum aestivum* (bread wheat) and *Brassica napus* (oilseed

rape). Both *T. aestivum* and *B. napus* are polyploid organisms that have been influenced by

human domestication, genome duplication events and geographical speciation as the use of

these crops became globalized. The specific and combined effects of these factors on the

control of the clock is yet to be investigated.

Within this paper we show that both the age of the plant and the developmental age of

leaves have significant effects on period with older material displaying shorter rhythms.

Brassica and wheat seedlings exhibited a shortening of period of approximately half an hour

to almost three hours per week after sowing. Period differences of up to 3.14h were

observed between leaves of plants of a single systemic age. To make our method high-

throughput whilst still providing reliable rhythms, tissues were segmented into various sizes

and compared to whole leaf samples. We identify regions of the plant leaves which are the

most robustly rhythmic and give the most consistent period estimates.

Both the light regime and temperature conditions also had large effects on period

estimation. Constant light (L:L) produced robust rhythms in *B. napus* and constant darkness

(D:D) produced more robust rhythms in *T. aestivum.* Warmer temperatures accelerated DF

rhythms in both *Brassicas* and wheat, indicating a curtailed temperature compensation

response. Intriguingly, we also demonstrate that rhythm robustness is positively correlated

with increasing temperature in wheat; opposite to what has previously been observed in

Arabidopsis (29).

Finally, we applied our optimized, high-throughput DF method to investigate differences

between elite cultivars in both *B. napus* and *T. aestivum* and demonstrate it to be a useful

tool for assaying circadian rhythms in these crop species.

147 Results

<u>Circadian variability due to leaf development and age of plant</u>

We tested the effect of both plant and leaf aging on period estimates from *Brassica* and

wheat seedlings. Previous studies in *Arabidopsis* have reported that the pace of the clock

increases as the plant ages and that earlier emerged leaves have a shorter period than those

which emerge later within the same individual(28). Our results mirror these findings for

young wheat and *Brassica* plants, however this association was lost for older material

(Figure 1A). For wheat we calculated period estimates from the second leaf of plants at 18,

25, 32 and 39 days after sowing and show that between 18-32 days period decreases

linearly at a rate of approximately half an hour per week while maintaining a near constant

relative amplitude error (RAE) (Figure 1A-C). However, in leaves from 39 day old plants

there was an increase in both average period and relative amplitude error, potentially due

to metabolic changes as a leaf changes from a source to a sink tissue or due to the onset of

senescence in these samples. A one-way analysis of variance yielded a significant effect of

wheat age on both period and RAE $(F(3,90)=12.13, p<0.001)$ and $(F(3,90)=7.018, p<0.001)$

respectively. Based on our investigation, we recommend using plants between 25 to 32 days

after sowing. At 25 days 100% of samples were classified as rhythmic and period CV was

1.52h. 32 day old samples were also robust, having the lowest RAE (0.15) and period error

(0.43) averages (see Supplementary Materials S1.)

In a separate experiment, we analyzed 4 leaves from 25 day old wheat plants as is shown in

Figure 1D, where leaves 1 and 3 were the oldest leaves and leaves 2 and 4 the second oldest

leaves from the main and secondary tiller, respectively. There was a statistically significant

difference between the mean periods at each leaf age (one-way ANOVA ($F_{(3,83)}=7.434$,

$p<0.001$). Within each tiller pair, the older leaf had a shorter period than the younger leaf

and had higher RAE averages. The mean period for leaf 4 (24.50h) was found to be

significantly longer than both leaf 1 (23.15h) and leaf 3 (23.32h) (Tukey HSD). We

recommend using leaf 2 as it had the best overall circadian robustness with regards to the %

samples returned (100%), RAE (0.18) and period error (0.50) (Supplementary Materials S2).

For *Brassica* seedlings, plants were grown to 4 different ages: 20, 25, 30 and 35 days after

sowing and leaf 1, 3 and 5 were sampled in the same experiment with leaf 1 being the

earliest emerged leaf and leaf 5 the most recently emerged leaf (Figure 2). We conducted a

nested ANOVA to test the effects of both plant age and within-plant leaf-age on period. We

found that variation in plant-age had a significant effect on period, with increasing age

causing a shortening of period ($F_{(3, 53)}=8.48$, $p<0.001$). The nested effect of leaf age within

each plant age-group was also found to be significant ($F_{(8, 53)}=5.45$, $p<0.001$). The largest

difference between leaves in each age-group was seen for 20 day old plants where a

difference of 3.14h was observed between leaf 1 and 5 ($p<0.001$, Tukey HSD) (Figure 2A).

Brassica plant-age was also found to have a significant effect on RAE averages with younger

plants having a lower mean RAE ($F_{(3,53)}=5.953$, $p<0.01$) (Figure 2B). Supplementary

Materials S3 shows robustness statistics for all plant ages and leaf-ages tested. We

recommend using leaf 1 from 20 day old plants as they had the lowest RAE (0.15) and

period error threshold (0.47).

To approximate the period shortening due to plant aging in *Brassica* we followed the

changes in average period in leaf 5 across plant ages from 20 days old to 30 days old as is

shown in Figure 2C. Our analysis revealed that period shortened by approximately 3 hours

per week from a mean of 26.50h(SD 1.17) to 22.38h (SD 0.62).

<u>Finding an optimal size of leaf sample</u>

We needed to identify representative leaf sections which allowed a sufficient number of

samples to be analyzed on one plate without compromising the robustness of rhythms for

period estimation. For wheat, we selected leaf 2 from 25 day old plants and analyzed the

periods and circadian robustness given by whole leaves compared to leaves cut into 10cm

sections and leaves cut into 4cm sections as shown in Figure 3A. By taking 4cm samples

from 2 regions on the same leaf (5 or 15cm down from the tip) we could investigate changes

in period across the length of the leaf. For *Brassica* seedlings we selected leaf 1 from 21 day

old plants and then kept them whole, took 3cm square samples from the centre or

quartered them (Figure 3B). This helped inform whether any changes in circadian

characteristics were a result of size reduction or from sub-sectioning regions of the leaf. Our

data showed that period and RAE averages were not significantly affected by cutting

samples in wheat (Figure 3C and 3D), (F(3,75)= 2.066, p>0.1, one-way ANOVA). However,

cutting *Brassica* leaves did significantly affect period estimates; quartered segments had a

slightly longer period than whole samples (F(2,113)=5.46, p<0.01, one-way ANOVA, Tukey

HSD (Whole-Quarter p<0.01) but RAE means were similar (Figure 3E and 3F) (Supplementary

Materials S4). From this data we recommend using 10cm segments for wheat and 3cm

square sections for *Brassica* imaging as these gave similar results to whole leaves and

increased throughput by 44% for *Brassica* and 100% for wheat.

We next wanted to investigate whether period estimates changed across the axis of the

leaf. We selected only the whole leaf images and digitally sectioned them into 10 or 5

regions of interest for wheat and *Brassica* leaves respectively (Figure 3G and 3H). Using this

approach we observed an average within-leaf variance of 0.45h in wheat and 0.42h in

Brassica leaves. This variation was larger than the leaf-to-leaf variation determined for

wheat (0.04h) and *Brassica* (0.3h) leaves. The mean period and RAE for each section across

these leaves was calculated and plotted (Figure 3G and 3H). No significant difference was

observed between the period of wheat or *Brassica* segments; however the RAE was

significantly different across wheat leaves ($F_{(9,139)}$=6.077, $p<0.001$, One-way ANOVA). The

middle segments (4, 5, 6, 7 and 8) had significantly lower RAE averages compared to the tip

(segment 1) suggesting that this middle region may give the most robust DF rhythms.

Constant free-running conditions: Dark versus Light

Two light regimes were tested which allowed free-running DF rhythms to be recorded. We

entrained plants for 4 days in 12:12 light:dark (L:D) cycles at 22°C before sampling and

imaging every hour under constant conditions, as described in Figure 4A. In both D:D and L:L

conditions the exposure time was kept at 1 minute. Figure 4B shows the striking differences

in period estimate accuracy obtained from wheat and *Brassica* under the two light regimes.

For wheat, periods from leaves under a D:D regime had much lower variance than those

under the L:L regime (D:D mean =23.29h, SD=0.53; L:L mean =23.54h, SD=3.19). For *Brassica*

the opposite was observed; rhythms were more accurate under L:L (D:D mean=24.89h,

SD=2.91; L:L mean=22.92h, SD=0.31). A shortening of period was observed for both *Brassica*

and wheat under L:L compared to D:D based on median values, however the increased

variance observed within wheat-L:L and *Brassica*-D:D resulted in these differences having

low significance (wheat t(23.16)=-0.38, p>0.5; *Brassica* t(10.14)=2.24, p=0.048 Welch's t-

test).

RAE ratios reflected the accuracy seen in period estimation between the regimes (Figure

4C). RAE averages were smaller in D:D for wheat (D:D mean=0.20, SD=0.08; L:L mean=0.38,

SD=0.18) and in L:L for *Brassica* (D:D mean=0.53, SD=0.13; L:L mean=0.13, SD=0.02). RAE

differences were significant between regimes for both species (Wheat t(29.23)=-4.38,

p<0.001; *Brassica* t(10.39)=9.92, p<0.001 Welch's t-test). Figure 4D and E show mean

oscillation traces which demonstrate how DF rhythms were sustained in wheat and *Brassica*

under the different light conditions. Interestingly, DF rhythms also had a dawn-phased peak

in wheat and a dusk-phased peak in *Brassica* which became shifted as different light

conditions were applied (Figure 4F).

Wheat samples under the D:D conditions returned 100% of samples from period estimation

(L:L=95.83%), an average RAE ratio of 0.20 (L:L=0.38), a period CV of 2.25% (L:L=13.53%) and

a period error threshold of 0.55 (L:L=1.17). For *Brassica*, rhythms under the L:L regime

returned 100% from period estimation (D:D=66.67%), a RAE average of 0.13 (D:D=0.53), a

period CV of 1.34% (D:D= 11.80) and a period error threshold of 0.41 (D:D=1.62). See

Supplementary Materials S5. We would therefore recommend running wheat DF

experiments under D:D conditions and *Brassica* DF experiments under L:L.

<u>Finding an optimum free-running temperature</u>

To investigate the effect of temperature on period and rhythm robustness we tested

Brassica and wheat seedlings at a range of constant temperatures. We used the optimal

conditions from the variables so far tested and entrained each batch of plants at the

imaging temperature for four days prior to imaging (see Methods). Both *Brassica* and wheat

experienced an acceleration of the clock at higher temperatures, with the rate increasing

most dramatically at lower temperatures (Figure 5A). Periods decreased from 26.40h

(SD=3.60) at 17°C to 22.48h (SD=0.31) at 32°C in wheat. Periods decreased from 26.28h

(SD=0.72) at 12°C to 23.16h (SD=0.52) at 22°C in *Brassica*. The temperature coefficient Q10

was calculated as an average across all temperatures (Supplementary Materials S7). Q10

was found to be 1.12 for wheat and 1.14 for *Brassica* indicating a degree of thermal

compensation, but to a lesser extent than has been previously reported in *Arabidopsis* (36).

We next looked at which temperatures gave the best rhythmicity in each crop. Rhythms

were most robust in wheat grown at 27°C: 100% of period estimates were returned, the

average RAE ratio was 0.15, period CV was 2.48% and period error threshold was 0.48

(Supplementary S6). There was a clear negative trend in period CV as the temperature

increased in wheat from 13.63% at 17°C to 1.38% at 32°C. Mean RAE at each temperature

can be seen in Figure 5B.

Across the temperatures tested *Brassica* rhythm robustness remained consistent; all

samples were returned from FFT-NLLS and RAE, period CV and Period error were similar

(Figure 5B, Supplementary Materials S6). We recommend 22°C for DF using *Brassica* as it

had the lowest period CV of 2.24%.

<u>An optimized DF method can be used in circadian analysis for crops</u>

To see whether the optimized method could be used to investigate circadian differences

between cultivars of the same species, we looked at circadian rhythms from seven *T.*

aestivum cultivars and three *B. napus* cultivars. For the *B. napus* lines, seeds were obtained

from 3 different harvest years to see whether period was constant between batches. The

optimized imaging parameters we used for these elite cultivars is outlined in Table 1.

Species	Plant age (days after sowing)	Leaf age (1=oldest leaf)	Cut sample	Light regime	Temperature	Throughput (N/imaging cabinet)	% return (from period estimation algorithms)
Brassica	20	1	3cm square	L:L	22°C	36	96.3
Wheat	25	2	10cm section	D:D	27°C	44-48	98.8

Table 1. Optimised DF method for circadian phenotyping of *Brassica* and wheat leaves

There was significant variation in the periods of the wheat lines tested as shown in Figure

6A ($F_{(6,152)}=9.81$, $p<0.001$, one-way ANOVA). A Tukey HSD test showed that Paragon

(mean=23.48h, SD=0.54) and Norin 61 (mean=23.50h, SD=1.30) both have longer periods

than Chinese Spring (mean=22.70h, SD=0.40), Claire (mean=22.48, SD=0.54) and Robigus

(mean=22.32h, SD=0.34) α=0.01).

Figure 6A shows the variation in period across three *Brassica* lines taken from three seed

batches. We conducted a two-way analysis of variance to compare cultivar ID and seed

batch effects as well as the interaction between the two factors. The cultivar ID was found

to have a significant effect on period ($F_{(2,60)}=25.47$, $p<0.001$) but batch year did not

significantly account for any variation in period either as a main effect ($F_{(2,60)}=1.73$, $p>0.1$)

or as an interaction with the cultivar ID ($F_{(4,60)}=2.27$, $p=0.72$). This suggests that the

observed differences in period are due to heritable genetic differences. The *Brassica* cultivar

Norin had the shortest overall period of 22.29h (SD 0.34); shorter than either Cabriolet

(23.32h, SD=0.57) or Chuanyou II (23.18h, SD=0.72) ($p<0.001$, Tukey HSD).

DF oscillations in both *Brassica* and wheat remained rhythmic throughout the experiment

allowing confident period estimation over 4 days (24-120h following T_0). The average DF

oscillations for the two most divergent wheat lines is shown in Figure 6C; the other lines

have been omitted for clarity. DF expression from all three years was averaged to make the

oscillation plots for the *Brassica* lines as shown in Figure 6D. The percentage of DF rhythms

returned from period estimation was high for both *Brassica* (96.3%) and wheat (98.8%)

proving that the method is both efficient and reliable.

Discussion

Manipulating the circadian clock has potential for influencing crop productivity, efficiency

and resilience; however research has been hindered by the lack of high-throughput

circadian protocols which can be reliably applied to crop plants. Transcriptional assays,

luciferase constructs and fluorescent markers have been used to investigate circadian

rhythms in tobacco[13], tomato[39], potato[40,41], *Brassica rapa*[42], rice[43–45],

barley[46] and wheat[47]. However, these approaches are either manually intensive,

technologically expensive or require genetic modification to systematically investigate each

component and so are low throughput. We have optimized a delayed fluorescence imaging

method for reliable circadian phenotyping of either *Brassica* or wheat seedlings. *Brassica*

napus (AACC) and *Triticum aestivum* (AABBDD) are both recent polyploids still undergoing

genomic rearrangements. The contribution of each genome to clock function remains to be

investigated. *B. napus* is a dicot recently diverged from *Arabidopsis*[48] and so is likely to

have clock homologs with similar functions. *T. aestivum* is a monocot with an incompletely

understood clock mechanism[49]. These species therefore provide interesting insights into

two genetically diverse families. Several differences between the function of these clocks

have been exposed through the factors examined in this paper. The opposing robustness of

clocks under D:D or L:L and the fact that DF rhythms peak with distinct phases under each

condition is indicative of diverging networks underlying circadian control of each species.

Lower temperatures (17°C) also seem to have a detrimental effect on the robustness of the

clock in *T. aestivum* but not *B. napus*, suggesting that temperature may be a stronger

zeitgeber for wheat than for *Brassica* within this temperature range. The DF rhythms in both

T. aestivum and *B. napus* have reduced temperature compensation compared to those

reported for leaf movement in *Arabidopsis*(36,37). However, it is important to recognize

that our rhythms were measured in dissected sections of leaves and may not be truly

analogous to rhythms from whole *Arabidopsis* individuals. The difference between intact

and excised leaves has been previously reported in Hall et al. 2001(22). Our analysis of the

homogeneity of periods across a single leaf also revealed variability of period robustness

across the axis of wheat leaves but relatively little variation across *Brassica* leaves.

In this study, we have shown that in both *Brassica* and wheat there is a strong interaction

between circadian period and age due to both systemic aging and leaf-specific

developmental aging. Previous research in *Arabidopsis* has asked whether the onset of

senescence is a result of a faster running clock or vice versa(28,50). Our results suggest that

the acceleration of the clock occurs in very young plants before senescence phase, raising

the possibility that the clock could be artificially manipulated to moderate senescence and

control timing of peak productivity in crops.

Natural variation of circadian phenotypes has been previously demonstrated in wild

Arabidopsis accessions (35,36,51) revealing a selection pressure for circadian traits specific

to different ecological settings. The extent to which circadian fitness has been selected-for

in modern crop plants has not yet been investigated. Application of our optimized protocol

in this study demonstrates that diverging rhythms are present within elite cultivars of the

same species. This variation in circadian period suggests that some level of circadian

diversity exists, but the question remains as to whether each cultivar is currently optimized

to enhance individual plant fitness. Crop plants with 'optimized circadian clocks' may have

the capacity to improve yield, efficiency and resilience potentially overlooked by traditional

plant breeding methods.

Conclusions

In this study, we investigated several important factors influencing circadian rhythms in

Brassica napus and *Triticum aestivum* and reveal intriguing differences between the two

crops. We provide an optimized DF methodology which can be reliably used for high-

throughput measurement of circadian rhythms. This research highlights the considerable

plasticity of the circadian clock under free-running conditions. It is our hope that these

results may inform future research by showing the extent to which controllable variables

can affect period estimation and how these may differ depending on the model species

being studied.

Method

<u>Plant material and growth conditions</u>

Brassica seedlings used were from the winter varieties Cabriolet and Norin and the semi-

winter variety Chuanyou II from the OREGIN *Brassica napus* Diversity Fixed Foundation Set

(BnDFFS)(52). Wheat seedlings used were all hexaploid elite cultivars ordered from the

Genome Resource Unit (John Innes Centre) (Supplementary Materials S8).

Brassica plants were grown in Levington's F2 mix in FP11 pots, spaced 5 plants to a pot.

They were grown in controlled greenhouse conditions, (16:8h L:D at 22:20°C). After 17 days,

plants were transferred to a plant growth chamber set at 12:12 L:D cycle at 22°C under

approximately 200μmol m^{-2} s^{-1} white light for 4 days entrainment (light spectra can be seen

in Supplementary Materials S9).

Wheat plants were imbibed at 4°C for 6 days before being planted in Petersfield cereal mix

in FP9 pots, spaced two to a pot. They were then grown in controlled greenhouse conditions

(16:8h L:D 17:12°C). After 21 days plants were transferred to a plant growth chamber set at

the cabinet conditions above. For temperature experiments, plants were entrained at the

temperatures in which they would be imaged.

<u>Image acquisition-standard conditions</u>

Leaves were removed just after entrainment dawn and placed face up onto 24cm square

petri dishes (Stratlab LTD, cat no. 163-PB-007) containing 0.5% water agar (Sigma-Aldrich,

SKU A1296). Unless otherwise stated, 3cm squares were cut from the second true leaf of 21

day old *B. napus* seedlings. A segment of 10cm was taken from the second leaf of the main

tiller of 25 day old *T. aestivum* seedlings, beginning 5cm down from the tip. Plates were

secured with masking tape around the periphery.

The imaging set-up is adapted from that described by Southern et al(53). A set-up schematic

can be seen in Supplementary Materials S10. We use Lumo Reteiga CCD cameras

(QImaging, Canada), which we have found to have comparable image quality to the Orca II

(Hamamatsu Photonics, Japan) without the need to run a water-cooling pump. Cameras

were fitted with a Xenon 0.95/25mm lens (Schneider-Kreuznach, Germany).

A custom built 25x25 red/blue LED rig (approx. 60 μmol m^{-2} s^{-1}) was controlled by μManager

software (v1.4.19, Open Imaging) through an Arduino Uno microcontroller board(54). LED

spectra for cabinets can be viewed in Supplementary Materials S9. μManager was used to

configure both the supplied camera driver software (PVCam v3.7.1.0) and program the

Arduino after installing the firmware source code available online (55)).

Both camera and LEDs were housed in a temperature controlled growth cabinet (Sanyo

MIR-553) in a dark room. The temperature was set to 22°C unless otherwise specified

(changed for the temperature experiments (Figure 5) and for the wheat cultivar experiment

(Figure 6)). Camera properties were kept the same in each experiment (Binning=4, Gain=1,

Readout-Rate=0.650195MHz 16 bit) and camera exposure was initiated 500ms after the

lights were turned off. A 'L:L' script refers to a regime of 59min of light followed by a 1

minute exposure in the dark. A 'D:D' script refers to 54min of darkness followed by 5min

light and then the 1min exposure. BeanShell scripts run by μManager have been adapted

from scripts used previously(56) and are available to view as Additional files 15 and 16.

Wheat imaging used the D:D script and *Brassica* imaging used the L:L script with the

exception of experiments in Figure 4.

Processing in FIJI and BioDare2 parameters

Image stacks were imported into FIJI(57) and regions of interest were selected.

Measurements for integrated density were taken for these regions across the stack using

the Multi-measure plugin. Each region was then labeled in Excel and an offset time series

added. The 'offset time' is the difference between the time of the first image (T1) and

entrainment dawn (ZT) in decimal hours. Data can then be uploaded to BioDare2 as

described online(58,59). BioDare2 is an open-access web tool for analyzing timeseries data

and predicting circadian parameters. For our data we found that Baseline and amplitude

(BAMP) de-trending was most appropriate but recommend visual inspection of the

detrending methods available to find the least intrusive method which removes any

baseline trends. Period estimation was done using the Fast Fourier Transform Non-Linear

Least Squares (FFT-NLLS) algorithm[60] on a data window of 24-120h with expected periods

set to 18-34h. Manual inspection of resulting periods ensured that all arrhythmic traces

were excluded from further analysis.

<u>Rhythm Robustness analysis</u>

We summarized rhythm robustness metrics based on several BioDare2 outputs. '%

returned' is the number of samples for which periods could be estimated out of the number

of samples originally imaged. The RAE (relative amplitude error) is the ratio of amplitude

error to amplitude and represents amplitude robustness. A RAE of 1 indicates the most

irregular waveform which can still be classified as rythmic whereas a RAE of 0 indicates a

perfect sine wave with no amplitude error. The period coefficient of variation (CV) is the

standard deviation of period estimates adjusted for the mean period and represents

between sample variation[58,61]. Period error is the extent to which the period estimate

could vary and still give a good fit to the model. Error scores close to 0 indicate a tight fit of

the model to the observed data and a high within sample period robustness. See

Supplementary data S1-6 for statistic tables and further descriptions.

<u>Normalization for experimental effects</u>

After circadian parameters were estimated in BioDare2, data was normalized to account for

the following random experimental effects. For the wheat plant age and leaf age

experiments (Figure 1), *Brassica* plant-leaf age experiments (Figure 2) and the *Brassica*

cultivar experiments (Figure 6B) samples were split between two imaging cabinets run in

parallel in a single experiment. The predicted parameters (e.g. period) for each sample from

the two cabinets were adjusted so that the cabinet means were then equivalent. This was

achieved by dividing the cabinet means by the overall mean to get an adjustment factor for

each cabinet and then dividing each individual value by that factor to get a cabinet-

normalized value. For the cutting data (Figure 3), the experiments were replicated in two

separate imaging weeks and then adjusted for the between-experiment effects. For the

wheat cultivar experiments (Figure 6A) data was obtained from two cabinets over two

separate experiments and was normalized for both effects in a similar way. The light regime

(Figure 4) and the temperature experiments (Figure 5) measured each variable in one

cabinet at a time and therefore did not require any normalization. The conclusions from

these experiments is consistent with preliminary experiments presented in Supplementary

Materials S11-14.

Statistical analysis was carried out in RStudio v1.1.423 using aov and t.test functions fit with

an appropriate linear model in the format specified in the Results.

<u>Declarations</u>

<u>Ethics approval and consent to participate</u>

Not applicable

<u>Consent for publication</u>

Not applicable

<u>Availability of data and materials</u>

The datasets generated during the current study are available as additional files and

supplementary materials in the online version of this article. Raw image files are available

from the corresponding author on reasonable request.

<u>Competing interests</u>

The authors declare that they have no competing interests.

Funding

This project was supported by the BBSRC via the Earlham institute CSP (BB/P016774/1, AH

SD, HR) and BBSRC Design Future Wheat (BB/P016855/1, AH)

Authors' contributions

This project was conceptualized by HR and AH. HR designed and conducted experiments,

carried out data processing and analysis and wrote the manuscript with contributions from

AH. All authors read and approved the final manuscript.

References

1.	Young MW, Kay SA. Time zones: a comparative genetics of circadian clocks. Nat Rev

Genet. 2001 Sep 1;2(9):702–15.

2.	Millar AJ, Straume M, Chory J, Chua NH, Kay SA. The regulation of circadian period by

phototransduction pathways in Arabidopsis. Science. 1995 Feb 24;267(5201):1163–6.

3.	Harmer SL, Hogenesch JB, Straume M, Chang HS, Han B, Zhu T, et al. Orchestrated

transcription of key pathways in Arabidopsis by the circadian clock. Science. 2000 Dec

15;290(5499):2110–3.

4.	Covington MF, Maloof JN, Straume M, Kay SA, Harmer SL. Global transcriptome

analysis reveals circadian regulation of key pathways in plant growth and

development. Genome Biol. 2008;9(8):R130.

5.	Dodd AN, Salathia N, Hall A, Kévei E, Tóth R, Nagy F, et al. Plant circadian clocks

increase photosynthesis, growth, survival, and competitive advantage. Science.

2005;309(5734):630–3.

6.   Ouyang Y, Andersson CR, Kondo T, Golden SS, Johnson CH. Resonating circadian

clocks enhance fitness in cyanobacteria. Proc Natl Acad Sci U S A. 1998 Jul

21;95(15):8660–4.

7.   DeCoursey PJ, Walker JK, Smith SA. A circadian pacemaker in free-living chipmunks:

essential for survival? J Comp Physiol A. 2000 Feb;186(2):169–80.

8.   von Saint Paul U, Aschoff J. Longevity among blowfliesPhormia terraenovae R.D. kept

in non-24-hour light-dark cycles. J Comp Physiol. 1978 Sep;127(3):191–5.

9.   Pittendrigh CS, Minis DH. Circadian systems: longevity as a function of circadian

resonance in Drosophila melanogaster. Proc Natl Acad Sci U S A. 1972

Jun;69(6):1537–9.

10.   Hernando CE, Romanowski A, Yanovsky MJ. Transcriptional and post-transcriptional

control of the plant circadian gene regulatory network. Vol. 1860, Biochimica et

Biophysica Acta - Gene Regulatory Mechanisms. 2017. p. 84–94.

11.   Edwards KD, Millar AJ. Analysis of Circadian Leaf Movement Rhythms in Arabidopsis

thaliana. In: Rosato E, editor. Circadian Rhythms: Methods and Protocols. Totowa, NJ:

Humana Press; 2007. p. 103–13.

12.   Welsh DK, Imaizumi T, Kay SA. Real-Time Reporting of Circadian-Regulated Gene

Expression by Luciferase Imaging in Plants and Mammalian Cells. In: Methods in

enzymology. 2005. p. 269–88.

13.   Millar AJ, Short SR, Chua N-H, Kay SA. A novel circadian phenotype based on firefly

luciferase expression in transgenic plants. Plant Cell. 1992;4(9):1075–87.

14.   Gould PD, Diaz P, Hogben C, Kusakina J, Salem R, Hartwell J, et al. Delayed

fluorescence as a universal tool for the measurement of circadian rhythms in higher

plants. Plant J. 2009 Jun;58(5):893–901.

15.   Gawroński P, Ariyadasa R, Himmelbach A, Poursarebani N, Kilian B, Stein N, et al. A

520   distorted circadian clock causes early flowering and temperature-dependent variation

521   in spike development in the Eps-3Am mutant of einkorn wheat. Genetics. 2014

522   Apr;196(4):1253–61.

523   16.   Goltsev V, Zaharieva I, Chernev P, Strasser RJ. Delayed fluorescence in

524   photosynthesis. Photosynth Res. 2009 Sep 23;101(2–3):217–32.

525   17.   Kurzbaum E, Eckert W, Yacobi YZ. Delayed fluorescence as a direct indicator of diurnal

526   variation in quantum and radiant energy utilization efficiencies of phytoplankton.

527   Photosynthetica. 2007 Dec;45(4):562–7.

528   18.   Gould PD, Diaz P, Hogben C, Kusakina J, Salem R, Hartwell J, et al. Delayed

529   fluorescence as a universal tool for the measurement of circadian rhythms in higher

530   plants. Plant J. 2009;58(5):893–901.

531   19.   Boxall SF, Dever L V, Kneřová J, Gould PD, Hartwell J. Phosphorylation of

532   Phosphoenolpyruvate Carboxylase Is Essential for Maximal and Sustained Dark CO_2

533   Fixation and Core Circadian Clock Operation in the Obligate Crassulacean Acid

534   Metabolism Species Kalanchoë fedtschenkoi. Plant Cell. 2017 Oct 1;29(10):2519–36.

535   20.   Thain SC, Hall A, Millar AJ. Functional independence of circadian clocks that regulate

536   plant gene expression. Curr Biol. 2000 Aug 14;10(16):951–6.

537   21.   Takahashi N, Hirata Y, Aihara K, Mas P. A Hierarchical Multi-oscillator Network

538   Orchestrates the Arabidopsis Circadian System. Cell. 2015 Sep 24;163(1):148–59.

539   22.   Hall A, Kozma-Bognár L, Tóth R, Nagy F, Millar AJ. Conditional circadian regulation of

540   PHYTOCHROME A gene expression. Plant Physiol. 2001 Dec;127(4):1808–18.

541   23.   Endo M, Shimizu H, Nohales MA, Araki T, Kay SA. Tissue-specific clocks in Arabidopsis

542   show asymmetric coupling. Nature. 2014 Nov 20;515(7527):419–22.

543   24.   Thain SC, Murtas G, Lynn JR, McGrath RB, Millar AJ. The circadian clock that controls

544   gene expression in Arabidopsis is tissue specific. Plant Physiol. 2002;130(1):102–10.

545   25.   James AB, Monreal JA, Nimmo GA, Kelly CL, Herzyk P, Jenkins GI, et al. The circadian

546   clock in Arabidopsis roots is a simplified slave version of the clock in shoots. Science.

547   2008 Dec 19;322(5909):1832–5.

548   26.   Gould PD, Domijan M, Greenwood M, Tokuda IT, Rees H, Kozma-Bognar L, et al.

549   Coordination of robust single cell rhythms in the Arabidopsis circadian clock via

550   spatial waves of gene expression. Elife. 2018 Apr 26;7:e31700.

551   27.   Wenden B, Toner DLK, Hodge SK, Grima R, Millar AJ. Spontaneous spatiotemporal

552   waves of gene expression from biological clocks in the leaf. Proc Natl Acad Sci U S A.

553   2012 Apr 24;109(17):6757–62.

554   28.   Kim H, Kim Y, Yeom M, Lim J, Nam HG. Age-associated circadian period changes in

555   Arabidopsis leaves. J Exp Bot. 2016;67(9):2665–73.

556   29.   Kim H, Kim HJ, Vu QT, Jung S, McClung CR, Hong S, et al. Circadian control of *ORE1* by

557   PRR9 positively regulates leaf senescence in *Arabidopsis*. Proc Natl Acad Sci.

558   2018;201722407.

559   30.   Harris PJC, Wilkins MB. Light-induced changes in the period of the circadian rhythm of

560   carbon dioxide output in Bryophyllum leaves. Planta. 1976;129(3):253–8.

561   31.   Somers DE, Webb AA, Pearson M, Kay SA. The short-period mutant, toc1-1, alters

562   circadian clock regulation of multiple outputs throughout development in Arabidopsis

563   thaliana. Development. 1998 Feb;125(3):485–94.

564   32.   Aschoff J. Circadian Rhythms: Influences of Internal and External Factors on the

565   Period Measured in Constant Conditions1. Z Tierpsychol. 2010 Apr 26;49(3):225–49.

33.   Salomé PA, Xie Q, McClung CR. Circadian timekeeping during early Arabidopsis

development. Plant Physiol. 2008 Jul;147(3):1110–25.

34.   Colin Pittendrigh BS, by N Harvey CE. On temperature independence in the clock-

system controlling emergence Time in Drosophila. Vol. 102, Proc. Soc. Exptl. Biol.

Med. 1954.

35.   Kusakina J, Gould PD, Hall A. A fast circadian clock at high temperatures is a

conserved feature across Arabidopsis accessions and likely to be important for

vegetative yield. Plant Cell Environ. 2014 Feb;37(2):327–40.

36.   Edwards KD, Lynn JR, Gyula P, Nagy F, Millar AJ. Natural allelic variation in the

temperature-compensation mechanisms of the Arabidopsis thaliana circadian clock.

Genetics. 2005;170(1):387–400.

37.   Gould PD, Locke JCW, Larue C, Southern MM, Davis SJ, Hanano S, et al. The Molecular

Basis of Temperature Compensation in the Arabidopsis Circadian Clock. Plant Cell.

2006 May 1;18(5):1177–87.

38.   Edwards KD, Lynn JR, Gyula P, Nagy F, Millar AJ. Natural allelic variation in the

temperature-compensation mechanisms of the Arabidopsis thaliana circadian clock.

Genetics. 2005 May;170(1):387–400.

39.   Facella P, Lopez L, Carbone F, Galbraith DW, Giuliano G, Perrotta G. Diurnal and

circadian rhythms in the tomato transcriptome and their modulation by

cryptochrome photoreceptors. PLoS One. 2008 Jul 30;3(7):e2798.

40.   Kloosterman B, Abelenda JA, Gomez M del MC, Oortwijn M, de Boer JM, Kowitwanich

K, et al. Naturally occurring allele diversity allows potato cultivation in northern

latitudes. Nature. 2013 Mar 6;495(7440):246–50.

41.   Navarro C, Abelenda JA, Cruz-Oró E, Cuéllar CA, Tamaki S, Silva J, et al. Control of

flowering and storage organ formation in potato by FLOWERING LOCUS T. Nature.

2011 Oct 25;478(7367):119–22.

42.   Xu X, Xie Q, McClung CR. Robust circadian rhythms of gene expression in Brassica

rapa tissue culture. Plant Physiol. 2010 Jun;153(2):841–50.

43.   Izawa T, Mihara M, Suzuki Y, Gupta M, Itoh H, Nagano AJ, et al. Os-GIGANTEA confers

robust diurnal rhythms on the global transcriptome of rice in the field. Plant Cell.

2011 May;23(5):1741–55.

44.   Sugiyama N, Izawa T, Oikawa T, Shimamoto K. Light regulation of circadian clock-

controlled gene expression in rice. Plant J. 2001 Dec 23;26(6):607–15.

45.   Nagano AJ, Sato Y, Mihara M, Antonio BA, Motoyama R, Itoh H, et al. Deciphering and

Prediction of Transcriptome Dynamics under Fluctuating Field Conditions. Cell. 2012

Dec 7;151(6):1358–69.

46.   Ford B, Deng W, Clausen J, Oliver S, Boden S, Hemming M, et al. Barley (Hordeum

vulgare) circadian clock genes can respond rapidly to temperature in an EARLY

FLOWERING 3-dependent manner. J Exp Bot. 2016;67(18):5517–28.

47.   Beales J, Turner A, Griffiths S, Snape JW, Laurie DA. A Pseudo-Response Regulator is

misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (Triticum

aestivum L.). Theor Appl Genet. 2007 Aug 24;115(5):721–33.

48.   Yang Y-W, Lai K-N, Tai P-Y, Li W-H. Rates of Nucleotide Substitution in Angiosperm

Mitochondrial DNA Sequences and Dates of Divergence Between Brassica and Other

Angiosperm Lineages. J Mol Evol. 1999 May;48(5):597–604.

49.   Campoli C, Shtaya M, Davis SJ, von Korff M. Expression conservation within the

circadian clock of a monocot: natural variation at barley Ppd-H1 affects circadian

expression of flowering time genes, but not clock orthologs. BMC Plant Biol. 2012 Jun

21;12:97.

50.   Kim H, Kim HJ, Vu T, Jung S, Robertson Mcclung C, Hong S, et al. Circadian control of

ORE1 by PRR9 positively regulates leaf senescence in Arabidopsis.

2018;115(33):8448–53.

51.   Michael TP, Salomé PA, Yu HJ, Spencer TR, Sharp EL, McPeek MA, et al. Enhanced

Fitness Conferred by Naturally Occurring Variation in the Circadian Clock. Science (80-

). 2003;302(5647):1049–53.

52.   Pink D, Bailey L, McClement S, Hand P, Mathas E, Buchanan-Wollaston V, et al.

Double haploids, markers and QTL analysis in vegetable brassicas. Euphytica. 2008

Nov 20;164(2):509–14.

53.   Southern MM, Brown PE, Hall A. Luciferases as Reporter Genes. In: Arabidopsis

Protocols. New Jersey: Humana Press; 2006. p. 293–306.

54.   Arduino [Internet]. [cited 2018 Nov 14]. Available from: https://www.arduino.cc/

55.   uManager-Arduino [Internet]. [cited 2018 Nov 15]. Available from: https://micro-

manager.org/wiki/Arduino#Firmware

56.   Litthauer S, Battle MW, Lawson T, Jones MA. Phototropins maintain robust circadian

oscillation of PSII operating efficiency under blue light. Plant J. 2015 Sep

1;83(6):1034–45.

57.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an

open-source platform for biological-image analysis. Nat Methods. 2012 Jul

1;9(7):676–82.

58.   Zielinski T, Moore AM, Troup E, Halliday KJ, Millar AJ. Strengths and Limitations of

Period Estimation Methods for Circadian Data. Yamazaki S, editor. PLoS One. 2014

May 8;9(5):e96462.

59.   BioDare2 - circadian period analysis [Internet]. [cited 2018 Nov 14]. Available from:

https://biodare2.ed.ac.uk/welcome

60.   Straume M, Frasier-Cadoret SG, Johnson ML. Least-Squares Analysis of Fluorescence

Data. In: Topics in Fluorescence Spectroscopy. Boston: Kluwer Academic Publishers;

1991. p. 177–240.

61.   Toth R, Kevei E, Hall A, Millar AJ, Nagy F, Kozma-Bognár L. Circadian Clock-Regulated

Expression of Phytochrome and Cryptochrome Genes in Arabidopsis. PLANT Physiol.

2001 Dec 1;127(4):1607–16.

Figure 1. DF rhythms in wheat change with the age of the plant and between leaves on the same plant. The wheat plant age experiment (A-C) used 'leaf 2' from plants grown for 18, 25, 32 or 39 days. Blue boxplots show differences in period (A) and RAE (B) at each plant age. Figure C shows the mean periods from which the increase in rate was approximated (error bars are standard deviations and the blue dashed line is a linear trend for first three ages). The wheat leaf variation experiment used 4 leaves sampled from 25 day old plants following the leaf numbering system described in D. Orange boxplots show differences in period (E) and RAE (F) at each leaf age. Color scales reflect an ageing gradient with lighter colors representing younger material. Data represents results from two imaging cabinets run in parallel as technical replicates and normalized for the between-cabinet effects. Period estimates were calculated using FFT-NLLS (BAMP de-trended data, 24-120h cut-off). N values reflect the number of samples for which period was estimated out of the total number of individuals sampled. Age 18 (N=26/26), age 25 (N=24/24), age 32 (N=25/26), age 39 (N=19/23). Leaf 1 (N=22/22), leaf 2 (N=22/22), leaf 3 (N=22/22), leaf 4 (N=21/22).

Figure 2. DF rhythms in *Brassica* change with the age of the plant and between different leaf ages. *Brassica* seedlings were grown to 4 different ages: 20, 25, 30 and 35 days after sowing Leaves 1, 3 and 5 were sampled from each plant in the experiment with leaf 1 being the earliest emerged leaf and leaf 5 the most recently emerged leaf. Boxplots show differences in period (A) and RAE (B) for each leaf age within each plant age. Figure C shows the mean periods from which the increase in rate was approximated using leaf 5 (error bars are standard deviations and the blue dashed line is a linear trend for first three ages). Color scales reflect an ageing gradient with lighter colors representing younger material. Periods and RAE estimates were calculated using FFT NLLS (BAMP dtr, 24-120h cut-off). Data represents results from two imaging cabinets run in parallel as technical replicates and normalized for the between-cabinet effects. Age 20: leaf 1 (N=6/6), leaf 3 (N=6/6), leaf 5 (N=6/6). Age 25: leaf 1 (N=4/6), leaf 3 (N=6/6), leaf 5 (N=6/6). Age 30: leaf 1 (N=2/6), leaf 3 (N=5/6), leaf 5 (N=6/6). Age 35: leaf 1 (N=6/6), leaf 3 (N=6/6), leaf 5 (N=6/6).

Figure 3. Cut sections of leaf material can be used to accurately make period estimates. The second leaf from 25 day old wheat seedlings was either left whole or sectioned into 10 cm or 4cm fragments cut either 5cm or 15cm from the tip (shown as dark grey sections in A). The first leaf from 21 day old *Brassica* plants was either left whole, sectioned into a 3cm square or quartered (B). Orange boxplots show differences in period (C) and RAE (D) for wheat sections. Purple boxplots show differences in period (E) and RAE (F) for brassica sections. Period and RAE were estimated using FFT NLLS, BAMP dtr, 24-120h cut-off. Whole leaves were digitally sectioned along the axis of the leaf post image-acquisition. Wheat period and RAE means for each section are shown in G. *Brassica* period and RAE means are plotted corresponding to the sectioning shown in H. Error bars show standard deviation. Data represents results from two experiments normalized for the between-experiment effects. Wheat: Whole (N=15/15), 10cm (N=23/23), 4cm top (N=20/21), 4cm bottom (N=22/22). *Brassica*: Whole (N=20/20), Square (N=21/21), Quarter (N=75/76).

Figure 4. Effect of either L:L or D:D free-running light conditions on DF rhythms. Wheat and *Brassica* seedlings were entrained for 4 days in L:D at 22°C before sections were cut, plated and imaged. A D:D free run consisted of a loop of 54 minutes of darkness followed by 5 minutes of light exposure and then image capture. A L:L free-run consisted of 59 minutes of light exposure before image capture (A). Boxplots of period (B) and RAE (C) are shown for *Brassica* and wheat in D:D and L:L conditions where *Brassica* data is displayed in purple and wheat data in orange. Period and RAE were estimated using FFT NLLS, BAMP dtr, 24-120h cut-off. Examples of oscillation traces are shown in D:D (black lines) and L:L (red lines) for wheat (D) and *Brassica* (E). Thick lines represent the mean trace of 6 mean-normalized individuals with error bars representing standard deviations. Estimated individual circadian phases are shown in the clock plot in F where the length of the line reflects the inverse circadian phase error (longer lines imply more confidence in the phase prediction). All phase estimates are relative to 0 where 0 represents entrainment dawn and 12 represents dusk.
D:D *Brassica* (N=12/18); D:D Wheat (N=24/24); L:L *Brassica* (N=18/18); L:L Wheat (N=23/24). Data is consistent with additional preliminary experiments which can be seen in Supplementary S11 and S12.

Figure 5. Increasing temperature causes a shortening of period and effects rhythm robustness. Wheat and *Brassica* seedlings were entrained for 4 days in L:D at the temperature being assessed before imaging. Each temperature point represents a separate imaging experiment. Period means decrease with increasing temperatures as is shown in A for *Brassica* (purple circles) or wheat (orange triangles). Error bars represent standard deviation. Box plots show the effect of temperature on RAE for wheat (orange) or *Brassica* (purple) (B). Period and RAE were estimated by FFT NLLS, BAMP dtr, 24-120h cut-off. Wheat: 17°C (N=17/23); 22°C (N=32/32); 27°C (N=11/11); 32°C (N=15/15). *Brassica*: 12°C (N=24/24); 17°C (N=35/35); 22°C (N=30/30). An additional preliminary experiment consistent with these observations can be seen in Supplementary S13.

Figure 6. DF can be used to measure period differences between elite cultivars in *Brassica* and wheat. 10cm sections from the second leaf of 25 day old wheat seedlings were imaged under D:D at 27°C. 3cm square sections from the first leaf of 21 day old *Brassica* seedlings were imaged under L:L at 22°C. Period boxplots based on the DF oscillations from different cultivars are shown for wheat (A) and *Brassica* (B). The three replicates in the *Brassica* data represent different seed batches. Period values were estimated using FFT NLLS, BAMP dtr, 24-120h cut-off window. BAMP de-trended DF data was normalized to the mean DF

intensity across all cultivars and plotted against time in hours after dawn (C and D). *Brassica* data represents results from two cabinets normalized for between-cabinet effects. Wheat data represents results from two identical experiments with two imaging cabinets run in parallel as technical replicates and normalized for the between-cabinet and experimental-run effects. Wheat: Robigus (N=25/25), Cadenza (N=12/12), Chinese Spring (N=24/25), Claire (N=27/27), Weebil (N=22/24), Paragon (N=24/4), Norin-61 (N=25/25). *Brassica*: for each seed batch of Cabriolet, Chuanyou II, Norin (N=8/8). *Brassica* data is consistent with previous experiments shown in supplementary materials S14.

A

12h 12h

4 days entrainment

'D:D' 54 minutes in dark   5 min

7 days imaging   image capture

'L:L' 59 minutes in light

B

Period (hours)

| DD | LL |

Brassica Wheat   Brassica Wheat

C

Relative Amplitude Error

| DD | LL |

Brassica Wheat   Brassica Wheat

D

DF intensity (normalised to mean)

Wheat

D:D
L:L

Time (hours since T0)

E

DF intensity (normalised to mean)

Brassica

D:D
L:L

Time (hours since T0)

F

Brassica DD
Wheat DD
Brassica LL
Wheat LL