Eco-evolutionary dynamics further weakens mutualistic interaction and coexistence under population decline

Avril Weinbach[1,2], Nicolas Loeuille[1], Rudolf P. Rohr[2]

1. Sorbonne Université, CNRS, Institut of Ecology and Environmental Sciences (iEES-Paris), place Jussieu, 75005, Paris, France;

2. Department of Biology, Ecology and Evolution, University of Fribourg, Switzerland.

Abstract

With current environmental changes, evolution can rescue declining populations, but what happens to their interacting species? Mutualistic interactions can help species sustain each other when their environment worsens. However, mutualism is often costly to maintain, and evolution might counter-select it when not profitable enough. We investigate how evolution of mutualism affects the coexistence of two mutualistic species, e.g. a plant-pollinator or plant-fungi system. Specifically, using eco-evolutionary dynamics, we study the evolution of the focal species investment in the mutualistic interaction of a focal species (e.g. plant attractiveness via flower or nectar production for pollinators or carbon exudate for mycorrhizal fungi), and how it is affected by the decline of the partner population with which it is interacting. We assume an allocation trade-off so that investment in the mutualistic interaction reduces the species intrinsic growth rate. First, we investigate how evolution changes species persistence, biomass production, and the intensity of the mutualistic interaction. We show that concave trade-offs allow evolutionary convergence to stable coexistence. We next assume an external disturbance that decreases the partner population by lowering its intrinsic growth rate. Such declines result in the evolution of lower investment of

the focal species in the mutualistic interaction, which eventually leads to the extinction of the partner species. With asymmetric mutualism favouring the partner, the evolutionary disappearance of the mutualistic interaction is delayed. Our results suggest that evolution may account for the current collapse of some mutualistic system like plant-pollinator ones, and that restoration attempts should be enforced early enough to prevent potential negative effects driven by evolution.

Keywords: adaptive dynamics, plant attractiveness, pollinators decline, evolutionary murder, asymmetrical interactions, alternative stable states

1 Introduction

Facing current global change, evolutionary mechanisms can help maintain biodiversity. Evolutionary rescue [1,2] corresponds to the selection of new traits in population collapsing with environmental changes, that allow for a demographic bounce. The signature for evolutionary rescue is the increase in frequency of the allele and corresponding phenotype robust to the new environment, correlatively to the population bounce. While this can be easily highlighted in lab experiments, it has so far been seldom observed in nature. In their review Carlson and collaborators [2] cite, for example, a previous study showing the adaptation of some *Chlorella* species, but not all, after strong acidification of many Canadian lakes with industrial pollution.

However, species are not isolated from one another and interaction might interfere with this evolutionary process. Mutualism is an interaction that has already been intensively studied, and proven to be fragile to Global Changes [3]. Echoing to that loss of interactions is that of biodiversity and system services such as pollination [4] and seed dispersal [3] and effective carbon and nutrient cycles [6]. While several reviews like that of Potts and collaborators [7]

point out the critical ecological crises we are undergoing, Toby Kiers and collaborators [3] add that mutualism, by binding species to a common fate, could create an evolutionary breakdown. With environmental changes, mutualism can become costly to maintain. Aside from co-extinction of the two interacting species, evolution can lead to mutualism loss, partner switch, or even a shift to antagonism. They thoroughly present how the different types of mutualisms are specifically sensitive or resistant to breakdown depending on the global change drivers. For example, plant-pollinator mutualism could be strongly affected by climate change and habitat fragmentation, while plant-rhizosphere mutualisms will be more affected by nutrient enrichment and the introduction of exotic species.

Mutualist systems, like all other systems of interacting species, will respond differently to global change [8]. One species can strongly decrease its density, following for example an environmental worsening, while the other might adapt faster to the new environment. This inertia decreases the interaction strength. We then observe a perverse positive feedback loop, where the decrease of one species is worsened by the evolution of the other, as illustrated in [9]. For mutualism, and especially obligatory ones, this might lead to species extinction, driven by the evolutionary disinvestment of its interactor. This effect is called an evolutionary murder (name suggested by [10]).

For instance, plants have been shown to evolve rapidly to changing pollinator populations [11–13]. A recent study from Gervasi and Schiestl [14] experimentally showed that changes in pollinator communities affect plant trait evolution after only eleven generations. Exposed to bumblebees, which are very efficient pollinators of *Brassica rapa,* the plants evolved toward more attractive traits to those pollinators (e.g. traits attracting pollinators such as volatile organic compounds, flower size, or plant height). Moreover, hoverflies, which are less efficient pollinators of *B. rapa*, caused a 15-fold increase in self-reproduction and a reduction in plant attractiveness. Given these experimental results, the current change and

reshaping of pollinator communities may affect the evolution of plant species, which in turn could influence coexistence with their interacting pollinators, i.e., an eco-evolutionary feedback loop.

Plant-mycorrhizal fungi interaction is another type of mutualism affected by global changes. The mycorrhizal fungi can for example fix inorganic nitrogen and provides this essential nutrient to the plant, who, in exchange, transfers via its roots carbon products to the fungi. Several experiments already showed that enriching the soil in nitrogen (often from anthropogenic sources in natural environments) disturbs this mutualistic exchange, by inducing a shift in the allocation to the mycorrhizal structures [15] and the composition of the mycorrhizal community [16]. This can in turn affects the plant community, and can even facilitate the invasion of alien plant species [17].

Theoretical studies have investigated the ecological [18–21] and evolutionary dynamics [3,22–25] of mutualistic communities such as plant-pollinators or plant-fungi. In particular, the evolution of plant selfing with changing pollinator communities has been studied in several papers [26–28]. Thomann and collaborators [29] even suggested that the decrease in pollinator richness and density could intensify pollen limitation. They propose that plants could in turn adapt either by increasing autonomous selfing or reinforcing the interaction with pollinators. Here we study the consequences of a declining population (e.g. pollinator collapse) on the eco-evolutionary dynamics of a two-species mutualistic system (e.g. plant-pollinator or plant-fungi). Specifically, we study the eco-evolution of the investment in mutualism of a focal species. To do so, we use the adaptive dynamics framework. This framework explicitly accounts for the eco-evolutionary feedback loop between the two species. We clarify when evolution leads to high or low investment in mutualism and determine the conditions under which evolution leads to the coexistence of the whole system. We then show that a declining partner population often results in a counter-selection of the

investment in mutualism of the focal species, which eventually enhances the population declines. For simplicity, in the following, we will use the example of a plant-pollinator

70   system. The adaptive trait is the plant investment, and the declining population is the pollinator. However, our approach remains general and can be applied to other mutualistic systems.

2 Plant-pollinator model and ecological dynamics

We consider a simple system with two interacting species; a plant with biomass density P,

75   and a pollinator with biomass density A. Note that this model is formulated as a general model of mutualism rather than very specifically tied to plant-pollinator interactions so that results may also concern other mutualistic systems. The community dynamics are given by a Lotka-Volterra type model:

$$\begin{cases} \dfrac{dA}{dt}=A\left(r_A-c_A A+\alpha\, \gamma_P P\right) \\ \dfrac{dP}{dt}=P\left(r_P-c_P P+\alpha\, \gamma_A A\right) \end{cases} \qquad (1)$$

A schematic view of the system is given in figure 1. The parameters r_A and r_P correspond to

80   the intrinsic growth rate of the pollinator and plant populations, respectively. We assume r_P to be strictly positive because of other reproduction means, e.g. vegetative reproduction or autogamy. The intrinsic growth rate of the pollinator (r_A) can be positive (e.g., interaction with other plants) or negative. Parameters c_A and c_P modulates intraspecific competition for the two species. Mutualistic interactions are given by $\alpha\gamma_A$ and $\alpha\gamma_P$, with γ_P the energetic

85   gain provided by the plant (via nectar, pollen and/or other plant exudates) to the pollinator, and γ_A the fertilisation provided by the pollinator to the plant. Because we consider mutualism as the net benefice obtained by both species, both γ parameter values are assumed positive in our model. We modulate the intensity of the interaction between the two

species with the parameter α. While the interaction depends on biological traits from both interactors (e.g. pollinator morphology or flight capacities, plant attractiveness), we have chosen to model it as a plant dependant trait and have therefore linked it to other plant traits via a trade-off function, Fig 2. We interpret it here as the attractiveness of the plant for the pollinator, and it corresponds to the trait that is under selection in the rest of the study. This plant attractiveness includes investment in various characters such as the number of flowers, their shape, their colour, volatile organic compound (VOCs) that attract insects with their odour, plant height, flowering duration or nectar quantity and quality (see part II in Willmer (2011) [28]).

*Figure 1: **Population variation rates of plant P and pollinator A.** Blue arrows indicate the density variations independent of the mutualistic interaction, green arrows the effects of the mutualistic interaction, and red arrows the effects of intraspecific competition. The parameters are described in the main text. B)*

Extrapolating from previous results [21], coexistence is stable provided:

$$\begin{cases} \dfrac{\alpha\gamma_P r_P + c_P r_A}{c_A c_P - \alpha^2 \gamma_A \gamma_P} > 0 \\[2ex] \dfrac{\alpha\gamma_A r_A + c_A r_P}{c_A c_P - \alpha^2 \gamma_A \gamma_P} > 0 \\[2ex] \alpha^2 \gamma_A \gamma_P < c_A c_P \end{cases} \tag{2}$$

The first two inequalities give the condition for the existence of an equilibrium point allowing positive densities (i.e. feasibility conditions). The last inequality ensures the dynamical

stability of the equilibrium. In the case of two interacting species, this local stability condition implies the global stability of the feasible equilibrium. The globally stable equilibrium is then:

$$
\begin{cases}
A^* = \dfrac{\alpha\, \gamma_P\, r_P + c_P\, r_A}{c_A\, c_P - \alpha^2\, \gamma_A\, \gamma_P} \\[3mm]
P^* = \dfrac{\alpha\, \gamma_A\, r_A + c_A\, r_P}{c_A\, c_P - \alpha^2\, \gamma_A\, \gamma_P}
\end{cases}
\tag{3}
$$

If the stability condition is not fulfilled, i.e., interspecific mutualism is stronger than intraspecific competition, the positive feedback loop resulting from interspecific mutualism may drive the system towards infinite growth. In such cases, other limiting factors (e.g. pathogen, predators, or new competitors) eventually regulate the populations. Since these factors are not taken into account in our model assumptions, we define a maximum plant attractiveness α_{cl} below which stability is warranted:

$$
\alpha_{cl} = \sqrt{\frac{c_A\, c_P}{\gamma_A\, \gamma_P}}.
\tag{4}
$$

We allow the evolution of α between zero (no investment in attractiveness) and this maximal level $\alpha_{max} < \alpha_{cl}$. We could also have controlled the infinite growth of our system by choosing a saturating function for the mutualistic interaction (Holling type II or other, e.g. [30]). Our choice of a linear functional response, however, allows explicit analytical computations and has the advantage to keep the model general and applicable to mutualistic interactions other than pollination.

3 Evolution of plant attractiveness

We study the evolution of plant attractiveness (α), assuming an allocation trade-off affecting the plant intrinsic growth rate r_P [31]. Its biomass can grow either via a reproduction process dependent on the interaction with its mutualist (e.g. pollination) whose intensity is controlled

120 by its attractiveness α, or via intrinsic growth (e.g. vegetative growth) and self-reproduction. The plant has a given quantity of energy that is divided between these two growth modes. [31,32], so that we assume r_P to be a decreasing function of the attractiveness α:

$$\left(\frac{r_P}{r_{Pmax}}\right)^s + \left(\frac{\alpha}{\alpha_{max}}\right)^s = 1.$$

(5)

The plant maximal intrinsic growth rate r_{Pmax} can be fixed to one without loss of generality,

125 by rescaling time unit:

$$r_P = \left(1 - \left(\frac{\alpha}{\alpha_{max}}\right)^s\right)^{1/s}.$$

(6)

The s exponent controls the trade-off shape. When $s=1$ there is a linear relationship between r_P and α. When $0 < s < 1$ the trade-off is convex. On the opposite, $s > 1$ produces a concave trade-off (as shown in figure 2).

Figure 2: **Variation of the attractiveness ratio** $\frac{\alpha}{\alpha_{max}}$ **with the plant intrinsic growth rate** r_P **depending on the trade-off strength.** Continuous lines show convex trade-offs, the dashed line a linear trade-off, and dashed-dotted lines concave trade-offs.

We follow the evolution of plant attractiveness using adaptive dynamics techniques [32,33].

130   Under adaptive dynamics hypotheses (see supplementary material for a full description of the

method and hypotheses) we can model the evolution of plan attractiveness and its

consequences on species density dynamics, and the feedback of species density on the

evolutionary process [33]. Evolution then proceeds by the successive invasions and

replacements of resident by mutant populations. Such dynamics are approximated, given rare

135   and small mutations, by the canonical equation [33]:

$$\frac{d\alpha}{dt} = \frac{1}{2}\mu\sigma^2 P^*(\alpha)\left.\frac{\partial\omega(\alpha_m,\alpha)}{\partial\alpha_m}\right|_{\alpha_m\to\alpha} \tag{7}$$

The term $\frac{1}{2}\mu\sigma^2 P^*(\alpha)$ encapsulates the phenotypic variability brought by the mutation

process on which selection can act. The last term, called the selective gradient, is based on the

variations of the relative fitness of mutants α_m given a resident α. It gives the direction of

evolution; a positive gradient selects larger attractiveness, while a negative gradient selects

140 smaller trait values. The relative fitness of the mutant is computed as the *per capita* growth

rate of a rare mutant population in a resident population at ecological equilibrium (3):

$$\omega(\alpha_m, \alpha) = \frac{1}{P_m} \frac{dP_m}{dt}\bigg|_{P_m \to 0} = r_P(\alpha_m) - c_P P^*(\alpha) + \alpha_m \gamma_A A^*(\alpha), \tag{8}$$

Eco-evolutionary equilibrium (called a singular strategy) occurs when the phenotypic trait

stops varying, i.e. equation 7 equals 0. Since its first part is always positive, it corresponds to

$\hat{\alpha}$ values for which the selective gradient is null:

$$\frac{\partial \omega(\alpha_m, \alpha)}{\partial \alpha_m}\bigg|_{\alpha_m, \alpha \to \hat{\alpha}} = \frac{dr_P(\hat{\alpha})}{d\hat{\alpha}} + \gamma_A A^*(\hat{\alpha}) = 0. \tag{9}$$

145 At singularities, costs in terms of energy dedicated to alternative means of reproduction (

$dr_P(\hat{\alpha})/d\hat{\alpha}$) therefore match pollination benefits ($\gamma_A A^*(\hat{\alpha})$). The existence of a singular

strategy is not enough to guarantee that evolutionary dynamics locally lead to it (convergence

condition) or that it persists (non-invasibility condition, i.e. resistance to invasion by nearby

mutants). A singular strategy that is both convergent and non-invasible is called a

150 continuously stable strategy (CSS) [35]. Evolution toward a CSS guarantees the coexistence

of the two species. This and other singularity types are presented in figure 3. Calculation of

the second and cross-derivative of the fitness function determines criteria for convergence

and invasibility [36]. The mathematical computation for the existence of singular strategies

and their convergence and invasibility properties are detailed in the supplementary material.

155 Equation (9) can be solved analytically for particular sets of parameters (see supplementary

material). For other cases, we graphically determine convergence and invasibility using

pairwise invasibility plots (figure 3). It is possible to show (supplementary material), as

illustrated in figure 2, that among the particular trade-offs that we study (eq. 6), only concave

allocation trade-offs lead to non-invasible strategies. Therefore, CSSs are only obtained with

160 a concave trade-off function. Convergence depends on the pollinator's intrinsic growth rate

(figure 3c and 3d). Mathematical analyses show that linear trade-offs lead to singular strategies that are not convergent. For non-linear trade-offs, convergence criteria cannot be solved and we rely on numerical investigations and PIPs. For positive pollinator intrinsic growth rate, given concave trade-offs, we obtain only one convergent stable singular strategies (CSS) at which ecological coexistence is granted. For negative pollinator intrinsic growth rate, the system exhibits a second singular strategy that is a Garden of Eden (non-convergent and non-invasible), i.e. a stable strategy that can never be reached by nearby mutants. For convex trade-offs (figure 3a and 3b), we always observe repellors (non-convergent and invasible). Starting above the repellor, attractiveness increases to reach the maximum value ($\alpha = \alpha_{max}$) and the plant growth relies only on the mutualistic interaction. Starting below the repellor, attractiveness evolves to zero, so that the two species no longer interact at the end of the evolutionary dynamics (e.g. complete selfing or clonal reproduction). In the following, we only study concave trade-off functions (i.e. $s > 1$).

Figure 3: ***Pairwise invasibility plots (PIPs) representing the invasibility potential of a rare mutant*** α_m ***within a resident plant population*** α ***at ecological equilibrium.*** *Grey areas indicate that the mutant relative fitness* $\omega(\alpha_m, \alpha)$ *is positive, so that it invades and replaces the resident population. In panels a and c, arrows show the direction of evolutionary trajectories. The system exhibits several singular strategies depending on the parameter values. Circles represent convergent strategies, whereas squares are non-convergent. Filled symbols represent invasible strategy, while not filled symbols are non-invasible. In panels a and b, the singular strategy is non-convergent and invasible (repellor). In panel c, the singular strategy is convergent and non-invasible (CSS). Panel d displays two strategies, one CSS and one which is non-convergent and non-invasible (Garden of Eden). Parameter values are:* $c_A = c_P = \gamma_A = \gamma_P = 1$ *, and* $\alpha_{max} = 0.8 * \alpha_{cl}$ *.*

4 Consequences of pollinator population decline

175 Now that we have characterised the eco-evolutionary dynamics of the plant-pollinator system, we study how pollinator decline may affect its outcome. We simulate less favourable environmental conditions for pollinators (e.g. habitat fragmentation, pesticides, diseases) by decreasing their intrinsic growth rate (r_A) . We illustrate the effects of this disturbance through Ecology-Evolution-Environment (E^3) diagrams [1,37]. These diagrams, presented

180 in figures 4 and 5, show the outcome of eco-evolutionary dynamics as a function of the environmental parameter, here the pollinator intrinsic growth rate r_A . Figure 4 exhibits two types of singular strategies: a convergent and stable singular strategy (CSS, continuous line), and Garden of Eden (GOE, dashed line). The Garden of Eden singular strategy is present only for negative pollinator intrinsic growth rates, i.e. in bad environments for the

185 pollinator. In this case, the system exhibits evolutionary alternative stable states. When plant attractiveness is above the Garden of Eden value, the evolution converges toward the CSS, while when below the GOE value, selection leads to ever-decreasing attractiveness, weakening the mutualistic interaction and eventually leading to the extinction of the pollinator (arrow (7) in figure 3a).

190 For positive pollinator intrinsic growth rates ($r_A > 0$), i.e. in "good pollinator environments", the system converges toward intermediate attractiveness α (arrows (1) and (2) in figure 4a). Now consider an environmental degradation (red arrow (3) in figure 4a). In the absence of evolution, both plant and pollinator populations have positive biomass densities. However, considering evolution, plant attractiveness is counter selected as pollinators are too rare to

195 compensate for the intrinsic costs of attractiveness. Eventually, evolution drives pollinator populations to extinction; an evolutionary murder depicted by arrow (4) in figure 4a. Faced with the crash of pollinator populations, restoration attempts may be undertaken (i.e. an increase in r_A value, e.g. by suppressing pesticides or adding other plant resources for

pollinators). Early intervention, depicted by arrow (5), can restore a stable mutualistic interaction. Delayed restoration attempts (white arrow (6)), do not allow such a rescue, as evolutionary trajectories will counteract their effects and lead to the extinction of the pollinator (arrow (7)). Note that here we separate timescales for simplicity, and consider that deterioration and restoration are fast compared to the evolutionary dynamics, hence horizontal arrows for these environmental changes.

*Figure 4: **Ecology-evolution-environment (E^3) diagram representing the impact of pollinator environmental deterioration on the evolution of plant attractiveness and on pollinator (panel a) and plant (panel b) equilibrium biomass densities .** White areas show parameters for which extinction occurs for either plants or pollinators. The blue intensity correlates with population densities of pollinators (panel a) or plants (panel b). Black lines show the position of singular strategies; continuous lines show convergent and non-invasible singular strategies (CSS), and dashed lines show Garden of Edens (non-invasible, divergent). Vertical black arrows (1, 2, 4, 7) display the direction of evolution. Environmental disturbance is represented by a red arrow (3). White arrows (5, 6) represent restoration attempts at different times along the evolutionary trajectory. On panel b) the red point and dotted lines represent the lowest r_A and $\frac{\alpha}{\alpha_{max}}$ values for allowing a CSS, therefore the maintenance of the mutualistic interaction. This point is what we call an eco-evolutionary tipping point. Parameters values are $s=2.5$, $c_A=c_P=\gamma_P=1$, $\gamma_A=0.2$, and $\alpha_{max}=0.8*\alpha_{cl}$. Similar E³ diagrams can be found in [1,37]*

Pollinator density at **ecological** equilibrium

Figure 5: **Influence of trade-off shape and mutualistic gains on** E^3 **diagrams.** *Columns differ in trade-off concavity. Lines differ in the asymmetry of mutualistic gains: in the top line (panels a,b, and c) pollinators benefit more than plants; the middle line (panels d,e, and f) shows equal gains while in the bottom line plant gains are larger. Red points and dotted lines represent the lowest* r_A *and* $\frac{\alpha}{\alpha_{max}}$ *values for maintaining a CSS, allowing the maintenance of the mutualistic interaction. Colours and lines are the same as in figure 4. The parameter values are* $c_A = c_P = 1$ *and* $\alpha_{max} = 0.8 * \alpha_{cl}$.

205   Finally, we study how trade-off shapes and asymmetry of mutualistic gains affect the eco-

evolutionary outcome (figure 5). More concave trade-offs allow a larger coexistence domain.

For less concave trade-offs, s < 2, only a positive pollinator intrinsic growth rate r_A allows

coexistence (figure 5a, d, g). Negative pollinator intrinsic growth rates lead to small benefits

for the plant, so that attractiveness is counter selected, eventually leading to the pollinator extinction . For stronger concave trade-offs, s > 2, (figure 5b, c, e, f, h, i) we observe qualitatively the same dynamics as in figure 4. For those trade-offs, asymmetric mutualistic gains favouring pollinators allow a larger range of disturbance, including negative intrinsic growth rates r_A , before attractiveness is counterselected and extinction occurs. Therefore, an increased mutualistic gain of the pollinator relative to that of the plant facilitates the long-term coexistence of the plant-pollinator system. This produces a more robust system that eases a potential restoration process. Note, however, that favouring pollinators gain over plants leads to lower selected levels of attractiveness.(compare figure 5 a, b, c vs g, h, i).

5 Discussion

While from a one species perspective, evolution can help to avoid extinction by fostering adaptation and restoring positive growth rates (evolutionary rescue), we here show an example in which accounting for mutualistic interactions largely modifies this optimistic view. Here, the evolution of one species in response to disturbances acting on its mutualistic interactor selects further decreases in the interactor population, eventually leading to the demise of the mutualistic interaction. This shows that evolution within mutualistic systems can actually be detrimental to the system's persistence and could undermine restoration attempts. Because we have used a general model of mutualism, this mechanism may concern various systems. This clearly suggests that when investigating the impact of global changes, we need to account for eco-evolutionary dynamics of the species and their interactors.

The model we use is voluntarily simple to allow a more complete mathematical study of eco-evolutionary dynamics and to highlight the role of key parameters (e.g. trade-off shapes or mutualistic gains). However, it may be linked to other models that study various types of mutualism. For instance considering pollination systems and plant reproduction, in line with

the presentation of the results, our model recalls previous theoretical works on plant evolution, that detail furthers the reproductive implications (e.g. [26–28]). For instance,

235 Lepers *et al.* [27] explicitly modelled the evolution of a plant reproduction system by taking prior selfing and inbreeding depression into account. In particular, they showed that evolution toward high prior selfing (for us of lesser attractiveness) leads first to pollinator extinction (our evolutionary murder). Because they also model the cost resulting from the inbreeding depression, they show that this evolutionary murder may further lead to the extinction of the

240 plant population. However, the model we propose may also be adapted to consider other mutualistic systems. For instance, in plant-mycorrhizae interactions, a resource exchange takes place, where plants provide carbon-based resources (e.g., sugars) to mycorrhizae while they get nutrients from the interaction. Such a situation does fit our hypotheses. The trait α would then be the quantity of resource provided by the plant (ie, its investment in the

245 mutualistic interaction), and this production diverts resources from growth and reproduction, therefore fitting our trade-off hypotheses. As such, our model recalls the results of a study by de Mazancourt & Schwartz [23]. They show that mutualism can arise and be evolutionarily selected from a two-species competing model by including trading. Each species can trade the resource it extracts in excess with the other. In our system, this trading would correspond

250 to the benefit $\alpha \gamma$ provided by the mutualistic interaction. Depending on the resource availability (the intrinsic growths in our model) the plant can either perform better on its own (possibly at the detriment of the fungi, as in our model) or can benefit from the mutualistic association with a mycorrhizal fungus. The mutualistic trading interaction can extend the coexistence boundaries, i.e. the resource space the two species can live in.

255 Our results also highlight that mutualistic interactions could be more or less vulnerable to environmental changes and population declines. For instance, here, only concave allocation trade-offs between plant intrinsic growth rate and investment in mutualism lead to the maintenance of the mutualistic interaction. These trade-offs favour intermediate investments

in the mutualistic interaction, while in the case of convex trade-offs, either complete

260 investment or no investment is eventually selected, depending on initial conditions. We kept

our study general because trade-off shapes are extremely difficult to measure *in vivo*, and can

vary deeply depending on the environment or the species types [39].

Bistability and critical transitions have been highlighted in a variety of ecological situations

(e.g. [40,41] in mutualistic system), and result from a strong positive feedback loop. Here we

265 have a similar phenomenon but on an eco-evolutionary scale. If the evolved investment in

mutualism before environment deterioration is above a certain threshold, evolution reinforces

the interaction, by increasing the attractiveness values, eventually leading to a stable,

coexisting system. On an ecological scale, this interaction reinforcement increases the

abundance of both species, which in turn favours the evolution of the focal species

270 investment toward higher value. Below a critical level of evolved investment, the population

of the mutualistic partner species is low. Evolution then further decreases investment in

mutualism, eventually leading to complete disinvestment in the mutualistic interaction. This

runaway selection for decreased investment leads in our case to the evolutionary murder of

the partner population by the evolving species [42]. Note that the trade-off shape modulates

275 the strength of the positive feedback loop. More concave trade-offs decrease the threshold

value above which interaction is maintained, thereby facilitating the persistence of the

system. Such dynamics have important implications. For instance, consider pollination as the

mutualistic interaction. Current data suggest large decreases in pollinator abundances [43].

Such pollinator declines are often considered to be directly linked to environmental changes

280 (eg, habitat change, pesticides [7]). However, our results suggest that evolutionary

components may also be present. If these declines favour plant strategies that offer less

resource, plant evolution may enhance the observed declines. In line with these predictions,

empirical observations suggest a decline of flower resources parallel to the pollinator decline

[43].

285 On a management side, alternative stable states and critical transitions have large implications, as systems may then shift abruptly, and large restorations are needed to recover previous states [41]. The eco-evolutionary alternative stable states we describe here have similar implications. Restoration can either be a reduction of the mortality causes of the declining species (banning pesticide, ploughing controlling pests and predators) or the

290 increase in their alternative resource source (plant sowing or nutrient addition). Here we consider that restorations are faster than evolutionary timescales. Evolution can however act fast [14], while restoration timescales largely vary from a few months (e.g., sowing high reward plants) to much longer timescales. Changes in pesticide regulations and applying these regulations may require national or international consensus. Similarly, while a change

295 in the agricultural mode does occur (eg, from intensive to agroecology), its dynamics happen over decades, while the evolution of plant reward may happen in just a few generations [14]. Note that, were we to consider longer restoration attempts we would still observe eco-evolutionary tipping points in our system. Such tipping points also make restoration attempts more difficult from two different points of view. First, the timing of the attempt becomes

300 important. Restoration is only successful when achieved before the threshold attractiveness is evolved. Second, if the system becomes degraded, a small restoration attempt may not be sufficient to recover large populations, but large efforts will have to be undertaken.

While in the face of current changes in the environment, evolution can play a key role in restoring populations and maintaining diversity, our results suggest that in the case of

305 mutualistic interactions, evolution may also favour strategies that eventually further threatens species coexistence. As such, our model echoes recent analyses that highlight the evolutionary fragility of mutualisms, given current changes [3,9]. Because our model is voluntarily simple, restricted in its number of interaction types and species, we expect evolutionary effects in complex ecological networks to be more complex and context-

310 dependent. However, we expect that accounting for these covariations of evolutionary

dynamics and changes in ecological interactions will be important, and that the effects of

evolution will then not systematically be positive.

6 Bibliography

1. Ferriere R, Legendre S. 2013 Eco-evolutionary feedbacks, adaptive dynamics and evolutionary rescue theory. *Phil Trans R Soc B* **368**, 20120081. (doi:10.1098/rstb.2012.0081)

2. Carlson SM, Cunningham CJ, Westley PAH. 2014 Evolutionary rescue in a changing world. *Trends Ecol. Evol.* **29**, 521–530. (doi:10.1016/j.tree.2014.06.005)

3. Toby Kiers E, Palmer TM, Ives AR, Bruno JF, Bronstein JL. 2010 Mutualisms in a changing world: an evolutionary perspective. *Ecol. Lett.* **13**, 1459–1474. (doi:10.1111/j.1461-0248.2010.01538.x)

4. Willmer P. 2011 *Pollination and Floral Ecology*. Princeton University Press.

5. Jordano P, Forget P-M, Lambert JE, Böhning-Gaese K, Traveset A, Wright SJ. 2010 Frugivores and seed dispersal: mechanisms and consequences for biodiversity of a key ecological interaction. *Biol. Lett.* , rsbl20100986. (doi:10.1098/rsbl.2010.0986)

6. Wilson GWT, Rice CW, Rillig MC, Springer A, Hartnett DC. 2009 Soil aggregation and carbon sequestration are tightly correlated with the abundance of arbuscular mycorrhizal fungi: results from long-term field experiments. *Ecol. Lett.* **12**, 452–461. (doi:10.1111/j.1461-0248.2009.01303.x)

7. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353. (doi:10.1016/j.tree.2010.01.007)

8. Tylianakis JM, Didham RK, Bascompte J, Wardle DA. 2008 Global change and species interactions in terrestrial ecosystems. *Ecol. Lett.* **11**, 1351–1363. (doi:10.1111/j.1461-0248.2008.01250.x)

9. Loeuille N. 2019 Eco-evolutionary dynamics in a disturbed world: implications for the maintenance of ecological networks. *F1000Research* **8**. (doi:10.12688/f1000research.15629.1)

10. Parvinen K. 2005 Evolutionary suicide. *Acta Biotheor.* **53**, 241–264. (doi:10.1007/s10441-005-2531-5)

11. Darwin C. 1877 *On the various contrivances by which British and foreign orchids are fertilised by insects*.

12. Parmesan C. 2006 Ecological and Evolutionary Responses to Recent Climate Change. *Annu. Rev. Ecol. Evol. Syst.* **37**, 637–669. (doi:10.1146/annurev.ecolsys.37.091305.110100)

13. Hopkins R, Rausher MD. 2012 Pollinator-Mediated Selection on Flower Color Allele

Drives Reinforcement. *Science* **335**, 1090–1092. (doi:10.1126/science.1215198)

14. Gervasi DDL, Schiestl FP. 2017 Real-time divergent evolution in plants driven by pollinators. *Nat. Commun.* **8**, 14691. (doi:10.1038/ncomms14691)

15. Johnson NC, Rowland DL, Corkidi L, Egerton-Warburton LM, Allen EB. 2003 Nitrogen Enrichment Alters Mycorrhizal Allocation at Five Mesic to Semiarid Grasslands. *Ecology* **84**, 1895–1908. (doi:10.1890/0012-9658(2003)084[1895:NEAMAA]2.0.CO;2)

16. Egerton-Warburton LM, Johnson NC, Allen EB. 2007 Mycorrhizal Community Dynamics Following Nitrogen Fertilization: A Cross-Site Test in Five Grasslands. *Ecol. Monogr.* **77**, 527–544. (doi:10.1890/06-1772.1)

17. Nijjer S, Rogers WE, Lee C-TA, Siemann E. 2008 The effects of soil biota and fertilization on the success of Sapium sebiferum. *Appl. Soil Ecol.* **38**, 1–11. (doi:10.1016/j.apsoil.2007.08.002)

18. Thébault E, Fontaine C. 2010 Stability of Ecological Communities and the Architecture of Mutualistic and Trophic Networks. *Science* **329**, 853–856. (doi:10.1126/science.1188321)

19. Rohr RP, Saavedra S, Bascompte J. 2014 On the structural stability of mutualistic systems. *Science* **345**, 1253497. (doi:10.1126/science.1253497)

20. Saavedra S, Rohr RP, Olesen JM, Bascompte J. 2016 Nested species interactions promote feasibility over stability during the assembly of a pollinator community. *Ecol. Evol.* **6**, 997–1007. (doi:10.1002/ece3.1930)

21. Goh BS. 1979 Stability in Models of Mutualism. *Am. Nat.* **113**, 261–275.

22. Ferriere R, Bronstein JL, Rinaldi S, Law R, Gauduchon M. 2002 Cheating and the evolutionary stability of mutualisms. *Proc. R. Soc. Lond. B Biol. Sci.* **269**, 773–780. (doi:10.1098/rspb.2001.1900)

23. De Mazancourt C, Schwartz MW. 2010 A resource ratio theory of cooperation. *Ecol. Lett.* **13**, 349–359. (doi:10.1111/j.1461-0248.2009.01431.x)

24. Georgelin E, Loeuille N. 2016 Evolutionary response of plant interaction traits to nutrient enrichment modifies the assembly and structure of antagonistic-mutualistic communities. *J. Ecol.* **104**, 193–205. (doi:10.1111/1365-2745.12485)

25. Valdovinos FS, Brosi BJ, Briggs HM, Moisset de Espanés P, Ramos-Jiliberto R, Martinez ND. 2016 Niche partitioning due to adaptive foraging reverses effects of nestedness and connectance on pollination network stability. *Ecol. Lett.* **19**, 1277–1286. (doi:10.1111/ele.12664)

26. Cheptou P, Massol F. 2009 Pollination Fluctuations Drive Evolutionary Syndromes Linking Dispersal and Mating System. *Am. Nat.* **174**, 46–55. (doi:10.1086/599303)

27. Lepers C, Dufay M, Billiard S. 2014 How does pollination mutualism affect the evolution of prior self-fertilization? A model. *Evolution* **68**, 3581–3598. (doi:10.1111/evo.12533)

28. Astegiano J, Massol F, Vidal MM, Cheptou P-O, Jr PRG. 2015 The Robustness of Plant-

Pollinator Assemblages: Linking Plant Interaction Patterns and Sensitivity to Pollinator Loss. *PLOS ONE* **10**, e0117243. (doi:10.1371/journal.pone.0117243)

29. Thomann M, Imbert E, Devaux C, Cheptou P-O. 2013 Flowering plants under global pollinator decline. *Trends Plant Sci.* **18**, 353–359. (doi:10.1016/j.tplants.2013.04.002)

30. Holland JN, DeAngelis DL. 2010 A consumer–resource approach to the density-dependent population dynamics of mutualism. *Ecology* **91**, 1286–1295. (doi:10.1890/09-1163.1)

31. Willmer P. 2011 Rewards and costs: the environmental economics of pollination. In *Pollination and Floral Ecology*, pp. 234–257. Princeton University Press.

32. Obeso JR. 2002 The costs of reproduction in plants. *New Phytol.* **155**, 321–348. (doi:10.1046/j.1469-8137.2002.00477.x)

33. Dieckmann U, Law R. 1996 The dynamical theory of coevolution: a derivation from stochastic ecological processes. *J. Math. Biol.* **34**, 579–612. (doi:10.1007/BF02409751)

34. Geritz S a. H, Kisdi E, Mesze NA G, Metz J a. J. 1998 Evolutionarily singular strategies and the adaptive growth and branching of the evolutionary tree. *Evol. Ecol.* **12**, 35–57. (doi:10.1023/A:1006554906681)

35. Christiansen FB. 1991 On Conditions for Evolutionary Stability for a Continuously Varying Character. *Am. Nat.* **138**, 37–50.

36. Marrow P, Dieckmann U, Law R. 1996 Evolutionary dynamics of predator-prey systems: an ecological perspective. *J. Math. Biol.* **34**, 556–578. (doi:10.1007/BF02409750)

37. Dieckmann U, Ferrière R. 2004 Adaptive dynamics and evolving biodiversity. In *Evolutionary conservation biology*, pp. 188–224. Cambridge: Cambridge University Press.

38. Bond WJ. 1994 Do Mutualisms Matter? Assessing the Impact of Pollinator and Disperser Disruption on Plant Extinction. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **344**, 83–90. (doi:10.1098/rstb.1994.0055)

39. Reekie E, Bazzaz FA. 2011 *Reproductive Allocation in Plants*. Elsevier.

40. Dercole F, Ferriere R, Rinaldi S. 2002 Ecological Bistability and Evolutionary Reversals Under Asymmetrical Competition. *Evolution* **56**, 1081–1090. (doi:10.1111/j.0014-3820.2002.tb01422.x)

41. Scheffer M, Carpenter SR. 2003 Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends Ecol. Evol.* **18**, 648–656. (doi:10.1016/j.tree.2003.09.002)

42. Dieckmann U, Marrow P, Law R. 1995 Evolutionary cycling in predator-prey interactions: population dynamics and the red queen. *J. Theor. Biol.* **176**, 91–102. (doi:10.1006/jtbi.1995.0179)

43. Biesmeijer JC *et al.* 2006 Parallel Declines in Pollinators and Insect-Pollinated Plants in Britain and the Netherlands. *Science* **313**, 351–354. (doi:10.1126/science.1127863)