1

3   **A chronology of multicellularity evolution in cyanobacteria**

Katrin Hammerschmidt[1*], Giddy Landan[1*], Fernando Domingues Kümmel Tria[1,†], Tal

Dagan[1]

[1] Genomic Microbiology Group, Institute of General Microbiology, Kiel University, Kiel,

Germany

[†] Present address: Institute for Molecular Evolution, Heinrich Heine University

Düsseldorf, Düsseldorf, Germany

[*] These authors contributed equally to this work.

Corresponding author: khammerschmidt@ifam.uni-kiel.de

Abstract

The transition from unicellular to multicellular organisms is one of the most significant events in the history of life. Key to this process is the emergence of Darwinian individuality at the higher level: groups must become single entities capable of reproduction for selection to shape their evolution[1-4]. Evolutionary transitions in individuality are characterized by cooperation between the lower level entities[5-7] and by division of labour[8,9]. Theory suggests that division of labour may drive the transition to multicellularity by eliminating the trade-off between two incompatible processes that cannot be performed simultaneously in one cell[1,9,10]. Here we examine the evolution of the most ancient multicellular transition known today, that of cyanobacteria[11,12]. We developed a novel approach for the precedence polarization of phenotypic traits that employs gene phylogenies and does not require a species tree. Applying our procedure to cyanobacterial genomes we reconstruct the chronology of ecological and phenotypic trait evolution in cyanobacteria. Our results show that the prime driver of multicellularity in cyanobacteria was the expansion in metabolic capacity offered by nitrogen fixation, which was accompanied by the emergence of the filamentous morphology and a reproductive life cycle. This was followed by a range of niche expansions and interactions with other species, and the progression of multicellularity into higher complexity in the form of differentiated cells and patterned multicellularity.

Introduction

Multicellularity is considered a characteristic trait of eukaryotes, but has evolved independently several times in diverse prokaryote taxa, including actinobacteria, myxobacteria, and cyanobacteria[5]. Bacterial multicellularity ranges from transient associations, such as colonies, biofilms and cellular aggregations, to permanent multicellular forms[13]. Instances of multicellular bacterial species present the major traits of eukaryotic multicellularity, including cell-to-cell adhesion, peri- or cytoplasmic continuity, intercellular communication, patterning, programmed cell death (PCD), and division of labour, often visible as cell differentiation[14]. Aggregative forms of multicellularity are common among bacterial species, for example, those that form a biofilm under specific external conditions[15]. *Bacillus subtilis*, for example, forms biofilms upon nutrient deprivation[16] in which cells differentiate into motile, matrix producing, or spore cells depending on the environmental cues[17]. Notably, cell differentiation in aggregates is adaptive at the level of the individual cell as it directly confers a fitness benefit to that particular cell. In contrast, under true division of labour, cells are interdependent upon each other and specialize in performing complementary tasks. These tasks (e.g., somatic functions or PCD) cannot be regarded beneficial on the level of the individual cell, but are considered advantageous for the colony; thus, they are emergent properties on a higher level of organization[18].

Such true division of labour in bacteria is best described in actinobacteria and cyanobacteria[18]. In cyanobacteria, the most complex of the filamentous species can differentiate up to five different cell types: vegetative (photosynthetic) cells, akinetes (spore-like cells), hormogonia (reproductive, motile filaments), necridia (dead cells resulting from PCD for hormogonia release), and heterocysts[14,19]. Heterocysts differentiate under nitrogen deprivation and are specialized in nitrogen (N_2) fixation by

the enzyme nitrogenase[20]. As this enzyme is sensitive to oxygen (O_2), these cells are

characterized by the absence of photosynthesis and by a thick cell wall, which maintains

an anaerobic environment. Heterocyst and vegetative cells in the filament are

metabolically interdependent with the heterocysts providing combined nitrogen to the

other cells within the filament and receiving fixed carbon compounds in return. With

cyanobacteria possessing the hallmark traits reminiscent of complex eukaryotic

multicellularity, except for tissues and organs, the trajectory of the evolution of

multicellularity in cyanobacteria is of general interest and has implications beyond

prokaryotic multicellularity.

Results & Discussion

To reconstruct the order of trait emergence in the evolution of cyanobacterial

multicellularity, we evaluated 25 phenotypic traits variably present in 199

cyanobacterial species and determined the order of their first appearance (Table 1;

Supplementary Tables 1 and 2). The ability to perform photosynthesis is not included in

the analysis as it is common to all cyanobacteria. The chronology of trait emergence is

derived from the ancestor-decedent relations of pairs of traits, hereby termed *trait-pair*

polarity.

The problem of trait-pair polarity poses a general challenge in microbial

phylogenetic reconstruction due to two confounding aspects: the absence of reliable

species phylogenies and the frequent absence of clear outgroups for the resolution of

ancestor-descendent relations. We applied a novel phylogenomics approach for trait-

pair polarity inference (see Fig. 1 and legend; Methods). The phylogenomic procedure

does not rely on a particular species tree or an outgroup, but extracts the total evidence

for trait precedence from a genomic sample of gene trees for 1671 protein coding gene

families. The statistical inference from large samples is effective in overcoming

conflicting signals originating from individual gene phylogenies.

The reconstructed trait-pairs polarity matrix (Fig. 2a) reveals several groups of

traits that are inferred to appear simultaneously, that we term temporal phases. These

include: (phase i) the cyanobacteria ancestor (traits 1-7), (phase ii) the transition to

multicellular individuality (traits 8-16), (phase iii) the expansion of niche occupation

and species interactions (traits 17-20), and (phase iv) the evolution of higher complexity

(traits 21-25). Apart from providing an overall ordering of traits during the evolution of

multicellularity, the pairwise tests are instrumental in resolving several debates about

trait precedencies in cyanobacterial evolution (Table 2).

Considering the set of traits in its entirety, the polarity matrix leads to the

following scenario for the evolution of multicellularity in cyanobacteria (Fig. 2b). The

prime driver of multicellularity in cyanobacteria was the expansion in metabolic

capacity offered by N FIXATION (phase ii). This emergence was accompanied by two other

cardinal innovations: the emergence of the FILAMENTOUS morphology and the emergence

of HORMOGONIA, and thus a reproductive life cycle. Together, these traits form the

essential elements of true multicellularity. Secondary traits that played a role in

stabilizing the nascent multicellular organism emerged in close succession, and include

GAS VESICLES, SHEATH, and MUCILAGE. There followed a cascade of niche expansions and

species interactions, and the culmination of complex multicellularity in the form of

differentiated cells and patterned multicellularity.

To elaborate, N_2 fixation – the reduction of molecular dinitrogen (N_2) to

ammonium (NH_3) – is catalysed by the enzyme nitrogenase. Whereas today's

cyanobacteria, other microorganisms, and most plants are able to take up nitrogen in

various combined forms, such as nitrate, ammonium, organic nitrogen, or urea, these

combined forms of nitrogen are scarce in most environments (e.g., open oceans or

terrestrial habitats[21]). Combined nitrogen, which is critical for the biosynthesis of amino

and nucleic acids, was likely a limiting resource in the early Earth environment[22]. Hence,

the capability of N FIXATION was key for cyanobacterial radiation into new habitats and

subsequent diversification (phases ii and iii).

The realization of the full metabolic potential of N_2 fixation, however, faced the

challenge of the incompatibility of nitrogenase with intracellular oxygen[23]. When the

cyanobacterial ancestor first acquired the capacity of N_2 fixation, it must have imposed a

strong selection pressure on the individual cells. The trade-off between photosynthesis

and nitrogen fixation led to the evolution of multiple solutions, which are still present in

today's cyanobacteria: the circadian rhythm of N_2 fixation in unicellular cyanobacteria[24]

and the differentiation of the highly specialized heterocyst[25].

Theory predicts that within a population of genetically identical unicellular

nitrogen fixing cyanobacteria, cell differentiation and phenotypic heterogeneity would

have been adaptive if this increased the fitness of the organisms in groups[10]. In the case

of unicellular cyanobacteria this means that cells evolved adhesion and exchanged fixed

nitrogen and carbon products within early cell groups such as filaments. Indeed, our

results indicate that the FILAMENTOUS morphology emerged simultaneously with N

FIXATION (phase ii). In filamentous cyanobacteria, where dividing cells remain linked in a

chain, surface-associated growth, SHEATH, and MUCILAGE lead to a localization of cells in

close spatial proximity, facilitating metabolite exchange between the individual cells.

When compared to the more transient associations in spatially structured communities,

such as in extracellular polymeric substance (EPS) imbedded biofilms, the development

of filaments opens possibilities for a more direct exchange of molecules with high

specificity. Metabolic exchange could have evolved as described for the evolution of

metabolic cross-feeding[26], as the exchange of carbon and nitrogen against other

products is generally common in photosynthetic or nitrogen-fixing organisms[27].

The trait HORMOGONIA is inferred to occur simultaneously with the appearance of

the FILAMENTOUS morphology. Its co-occurrence with other traits, such as NOT FREE-LIVING

BENTHIC/SESSILE, EPILITHIC/ ENDOLYTHIC, SHEATH, and MUCILAGE underline the transition from

a planktonic to a benthic lifestyle (phase ii). The differentiation of HORMOGONIA can be

induced by environmental stimuli, such as nitrogen deprivation[28]. After breaking off

from the mother filament at the necridia, HORMOGONIA disperse via gliding motility or

float thanks to GAS VESICLES, ensuring the reproduction of benthic species[29,30]. The

differentiation into HORMOGONIA is reversible – they change back to the sessile lifestyle,

where they grow into a new vegetative filament[28]. Here we observe the emergence of a

two-phases life cycle, which is important for the transition to multicellularity[31,32].

Traits that are indicative of higher complexity emerged late in the evolution of

cyanobacteria and dominate (phase iv) of the polarity matrix. We observe the

occurrence of AKINETES and the irreversibly differentiated HETEROCYSTS. HETEROCYSTS

represent not only a morphological adaptation to the obstacle of N_2 fixation under oxic

conditions but also an elaborate and highly specialized communication and metabolite

exchange system. In *Anabaena* sp., for example, where several hundred cells

communicate within a filament, a regular heterocyst formation pattern along the

filament must be achieved to guarantee that every cell is adequately supplied with fixed

nitrogen compounds[19]. For this, the inhibitory signalling peptide PatS needs to be

distributed along the filament with heterocyst formation occurring only in cells with low

PatS concentration[33]. Whether the exchange of metabolites and regulators happens via

the continuous periplasm[34,35] or through septal junctions[36,37] is still not fully resolved.

Another central innovation that occurs in this phase of the polarity matrix is the

ability to FISSION IN MULTIPLE PLANES. This trait co-occurred with the ability to produce

BAEOCYTES, differentiated cells, which are the reproductive stages in the order

Pleurocapsales[38]. Notably, baeocyte-forming cyanobacteria, that have been traditionally

grouped together with unicellular cyanobacteria[29], appear to immediately predate the

evolution of AKINETES and HETEROCYSTS and thus emerge much later than filamentous

forms. The ability to FISSION IN MULTIPLE PLANES is known to underlie the TRUE-BRANCHING

morphology, where cells in a filament divide in more than one plane, and which is the

last trait to emerge in our analysis. Members of the Haphalosiphon/Stigonematales

clade having true-branching and multiseriate filament morphology are considered as the

latest evolutionary innovations[29] and this is further observed in cyanobacterial

phylogenies[39,40].

Common features of evolutionary transitions in individuality comprise

cooperation between the lower level units[5-7] and the division of labour[8,9]. The latter

might be of particular advantage, and serve as the driver of the transition to

multicellularity when there is a strong trade-off between processes that cannot be

performed in a single cell at one time[9,10]. Our current findings support this theory and

point to nitrogen fixation, and its incompatibility with photosynthesis, as the trigger for

the evolution of multicellularity in cyanobacteria. One open question concerns how the

underlying genetics of novel traits, such as the division of labour, arise within a newly

emerging multicellular individual. In the case of cyanobacteria multicellularity, we

suggest that no new genes were required and that higher complexity was achieved by

regulatory changes in gene expression patterns. Basic communication and metabolite

exchange was pre-existing as single-celled bacteria frequently engage in cell-cell

communication and cross-feeding of metabolites via the external environment[26].

Division of labour between photosynthesis and nitrogen fixation was likely first

established by the regulatory mechanism of temporal switching. Once simple forms of

division of labour and metabolic exchange existed, the transition into spatial separation

in differentiated cells could have evolved mainly by regulatory modifications.

Differentiated cells are one of the hallmarks of complex multicellularity. It is

therefore significant that we observe six distinct cell types in cyanobacteria:

photosynthetic, hormogonia, necridia, akinetes, baeocytes, and heterocysts. Such a

plurality indicates that the underlying regulatory mechanisms are well developed and

that their plasticity and adaptability are a matter of course. It is also significant that

three of the differentiated cell types, hormogonia, akinetes, and baeocytes, offer novel

reproductive potential and the establishment of a multicellular life cycle. Moreover,

signs of a nascent developmental plan can be observed in both the distribution of

heterocysts along filaments and in the patterning of true branching cyanobacteria. These

elements have no fitness value for the individual cell, but are selectable adaptations on

the higher level, the filament. The chronology of the evolution of multicellularity in

cyanobacteria shows that, once established, multicellular individuality opens new vistas

of opportunities.

METHODS

Data

The data underlying this study consists of the genomic sequences and phenotypic traits

of 199 cyanobacterial species. These were selected from the available genomes so that

the number of represented taxa will be as large as possible and genus-level redundancy

will be reduced (See Supplementary Table 1 for the complete list of species).

Phenotypic traits

Phenotypic traits were chosen for their potential relevance to the evolution of multicellularity in cyanobacteria, such as environmental factors that might facilitate multicellularity and markers that are indicative for the transition to multicellularity (Table 1). Information on presence and absence of traits was obtained from the published literature and from the Pasteur Culture Collection of cyanobacteria, extending the work by Uyeda et al[44], and coded as binary trait states. Traits included MORPHOLOGY, NITROGEN FIXATION, FRESHWATER, MARINE, BAEOCYTES, HORMOGONIA, THERMOPHILIC, AKINETES, HETEROCYSTS, TRUE BRANCHING, EPI/ ENDOLITHIC, EPIPHYTIC, PERIPHYTIC, MATS, FREE-LIVING, HABITAT, SHEATH, MUCILAGE, GAS VESICLES, MOTILITY, and FISSION IN MULTIPLE PLANES (Supplementary Table 1).

Protein families and gene trees

The cyanobacteria protein families were constructed from completely sequenced genomes available in RefSeq database[51] (ver. May 2016). For the construction of protein families, at the first stage, all protein sequences annotated in the genomes were blasted all-against-all using stand-alone BLAST[52] ver. 2.2.26. Protein sequence pairs that were found as reciprocal best BLAST hits (rBBHs)[53] with a threshold of E-value $\leq 1 \times 10^{-5}$ were further compared by global alignment using needle[54]. Sequence pairs having $\geq 30\%$ identical amino acids were clustered into protein families using the Markov clustering algorithm (MCL)[55] ver. 12-135 with the default parameters. By requiring a gene to be present in the genomes of the SynProCya clade and in at least one other major cyanobacterial clade, we identified 1671 gene families that are present on both sides of the cyanobacterial root, i.e., ancient proteins families. Protein sequences of these families were aligned using MAFFT version 7.027b employing the L-INS-i strategy[56].

239 Maximum likelihood trees were reconstructed with PhyML version 20120412[57] with

240 parameters -b -4 -v e -m LG -c 4 -s SPR.

Trait ordering

We consider four possible polarizations of a pair of traits *A* and *B*: *A* originates before *B*;

B originates before *A*; *A* and *B* emerge simultaneously; and *A* and *B* are not nested but

originate in independent lineages. We infer the pairwise emergence order of the traits

by collecting evidence from a set of phylogenetic trees of independent protein families,

and conducting a formal statistical test for the best supported polarization. Finally, the

set of all pairwise polarities is used to derive the order of emergence of all traits (Fig. 1).

Polarization evidence from a single phylogenetic tree

Given an unrooted phylogenetic tree, we first consider all possible rooted versions of the

tree. In an *n*-OTU (*Operative Taxonomic Unit*) unrooted tree, the root can be placed on

any of the 2*n-3* branches of the tree. For each of the 2*n-3* rooted trees, we label the OTUs

by the presence of the two traits *A* and *B*, and infer the most parsimonious *Last Common*

Ancestor (LCA) of each trait, assuming a single trait origin and possible trait losses. We

then record the polarity induced by the putative root position according to whether the

LCAs of *A* and *B* coincide, are descended one from the other, or are located on

independent lineages.

Next we conduct a MAD analysis[48], yielding an *Ancestor Deviation* (AD) statistic

for every branch of the tree. The AD measure quantifies the amount of lineage rate

heterogeneity that is induced by postulating a branch as harbouring the root of tree. We

have previously shown that the AD measure provides robust evidence for the inference

of the root of the tree. In the current study we do not infer a single root, but use the AD

measure to assess the relative strength of alternative rootings of the same tree.

Thus, for each possible root position we obtain a polarity state for *A* and *B* and a

corresponding AD value. For each of the four polarities, we take the minimal AD value as

the tree support of that polarity. Note that for certain trees and trait pairs, some

polarities may not occur. We differentiate between two such possibilities. First, a certain

tree is *Uninformative* regarding a specific trait-pair polarity if the species composition of

the tree renders the observation of the polarity impossible in any possible topology. For

example, the polarity **'A precedes B'** is impossible to observe in trees where the *A* OTUs

are a subset of the *B* OTUs, regardless of the specific topology or root position. In

contrast, a certain polarity may be unobserved in any of the rootings of a specific

unrooted tree, while still being a possible observation for a different tree topology. An

example of such *Informative absent-observation* is when *A* and *B* label disjoint sets of

OTUs yet the **'Not nested'** polarity is not observed in the actual tree topology. In the

latter case, the absence of an observation is evidence *against* that specific polarity, and

the polarity is assigned the maximal observed AD as its score. When the tree is

uninformative regarding a polarity, on the other hand, it is assigned a *'missing'* value,

and is excluded from subsequent analyses.

Statistical inference of pairwise polarity

Repeating the preceding procedure for trees derived from all different single-copy

proteins families yields a phylogenomic sample of four variables - the AD scores of each

of the four polarities for the pair of traits. The four distributions are paired, as each tree

brings in a 4-tuple of values, and a significant difference in support values can be tested

using the non-parametric Mann-Whitney U test[58]. In all, we conduct twelve one-sided

tests of contrasting polarities, while employing an FDR[59] correction for multiple

comparisons. A trait-pair is considered polarized if there exists one polarity where all

three tests against the other polarities recover significantly lower AD values at the 1%

FDR level. In exceptional situations we again encounter polarity contrasts that cannot be

tested. For some polarity contrasts the sample size may not meet our threshold (10

paired observations), and the test is conservatively considered valid but not significant

with a p-value of 1.0. A second exception occurs when one trait (*A*) is present only in a

strict subset of the species that possess the other trait (*B*). In this case it is impossible to

observe the polarities *'not nested'* and *'A precedes B'* in any tree, and we restrict the

testing to the single contrast *'B precedes A'* versus *'Simultaneous'*.

The pairwise polarity inference is applied to each pair of traits and summarized

in a trait-pair polarity matrix. We again apply the FDR correction, this time over all trait-

pairs and polarity contrasts. In the present study, with 25 traits, we apply FDR over

3,480 tests. To derive an ordering of the traits, we apply 'Topological sort'[60] to the

significant polarities of type *'A precedes B'*, or vice versa. In the present study the

significant polarities form a partial order, i.e., there are no self-contradicting precedence

cycles, and the topological sorting order is used to order the polarities matrix and to

reduce it to a feed-forward network (Fig. 2).

Data availability

The alignments and trees are available on our webpage at www.uni-

kiel.de/genomik/ressourcen

References

1. Buss, L. W. *The evolution of individuality*. (Princeton University Press, 1987).

2. Smith, J. M. & Szathmáry, E. *The major transitions in evolution*. (Oxford University Press, 1995).

3. Okasha, S. *Evolution and the levels of selection*. (Oxford University Press, 2006).

4. Godfrey-Smith, P. *Darwinian populations and natural selection*. (OUP Oxford, 2009).

5. Bonner, J. T. The origins of multicellularity. *Integr. Biol.* **1,** 27–36 (1998).

6. Nowak, M. A. Five rules for the evolution of cooperation. *Science* **314,** 1560–1563 (2006).

7. Rainey, P. B. & De Monte, S. Resolving conflicts during the evolutionary transition to multicellular life. *Annu. Rev. Ecol. Evol. Syst.* **45,** 599–620 (2014).

8. Kirk, D. L. A twelve-step program for evolving multicellularity and a division of labor. *BioEssays* **27,** 299–310 (2005).

9. Michod, R. E. Evolution of individuality during the transition from unicellular to multicellular life. *Proc. Natl. Acad. Sci. USA* **104 Suppl 1,** 8613–8618 (2007).

10. Ispolatov, I., Ackermann, M. & Doebeli, M. Division of labour and the evolution of multicellularity. *Proc. Biol. Sci.* **279,** 1768–1776 (2012).

11. Schopf, J. W. Microfossils of the Early Archean Apex chert: new evidence of the antiquity of life. *Science* **260,** 640–646 (1993).

12. Grosberg, R. K. & Strathmann, R. R. The evolution of multicellularity: a minor major transition? *Annu. Rev. Ecol. Evol. Syst.* **38,** 621–654 (2007).

13. Shapiro, J. Bacteria as multicellular organisms. *Sci. Am.* **258,** 82–89 (1988).

14. Claessen, D., Rozen, D. E., Kuipers, O. P., Søgaard-Andersen, L. & van Wezel, G. P. Bacterial solutions to multicellularity: a tale of biofilms, filaments and fruiting bodies. *Nat. Rev. Microbiol.* **12,** 115–124 (2014).

15. Tarnita, C. E., Taubes, C. H. & Nowak, M. A. Evolutionary construction by staying

together and coming together. *J. Theor. Biol.* **320,** 10–22 (2013).

16. Karatan, E. & Watnick, P. Signals, regulatory networks, and materials that build and

break bacterial biofilms. *Microbiol. Mol. Biol. Rev.* **73,** 310–347 (2009).

17. Vlamakis, H., Aguilar, C., Losick, R. & Kolter, R. Control of cell fate by the formation of

an architecturally complex bacterial community. *Genes Dev.* **22,** 945–953 (2008).

18. van Gestel, J., Vlamakis, H. & Kolter, R. Division of Labor in Biofilms: the Ecology of

Cell Differentiation. *Microbiol. Spec.* **3,** 1–24 (2015).

19. Herrero, A., Stavans, J. & Flores, E. The multicellular nature of filamentous

heterocyst-forming cyanobacteria. *FEMS Microbiol. Rev.* **40,** 831–854 (2016).

20. Frías, J. E., Flores, E. & Herrero, A. Requirement of the regulatory protein NtcA for

the expression of nitrogen assimilation and heterocyst development genes in the

cyanobacterium *Anabaena* sp. PCC 7120. *Mol. Microbiol.* **14,** 823–832 (1994).

21. Zehr, J. P. Nitrogen fixation by marine cyanobacteria. *Trends Microbiol.* **19,** 162–173

(2011).

22. Kasting, J. F. & Siefert, J. L. Biogeochemistry. The nitrogen fix. *Nature* **412,** 26–27

(2001).

23. Gallon, J. R. The oxygen sensitivity of nitrogenase: a problem for biochemists and

micro-organisms. *Trends Biochem. Sci.* **6,** 19–23 (1981).

24. Mitsui, A. *et al.* Strategy by which nitrogen-fixing unicellular cyanobacteria grow

photoautotrophically. *Nature* **323,** 720–722 (1986).

25. Flores, E., Picossi, S., Valladares, A. & Herrero, A. Transcriptional regulation of

development in heterocyst-forming cyanobacteria. *BBA - Gene Regul. Mech.* (2018).

26. D'Souza, G. *et al.* Ecology and evolution of metabolic cross-feeding interactions in

bacteria. *Nat. Prod. Rep.* **35,** 455–488 (2018).

27. Kaiser, D., Kowalski, N., Böttcher, M. E., Yan, B. & Unger, D. Benthic nutrient fluxes

from mangrove sediments of an anthropogenically impacted estuary in Southern

China. *J. Mar. Sci. Eng.* **3,** 466–491 (2015).

28. Flores, E. & Herrero, A. Compartmentalized function through cell differentiation in

filamentous cyanobacteria. *Nat. Rev. Microbiol.* **8,** 39–50 (2010).

29. Rippka, R., Deruelles, J., Waterbury, J. B., Herdman, M. & Stanier, R. Y. Generic

assignments, strain histories and properties of pure cultures of cyanobacteria.

Microbiol. **111,** 1–61 (1979).

30. Herdman, M. & Rippka, R. Cellular differentiation: hormogonia and baeocytes.

Methods Enzymol. **167,** 232–242 (1988).

31. Rainey, P. B. & Kerr, B. Cheats as first propagules: A new hypothesis for the

evolution of individuality during the transition from single cells to multicellularity.

BioEssays **32,** 872–880 (2010).

32. Hammerschmidt, K., Rose, C. J., Kerr, B. & Rainey, P. B. Life cycles, fitness decoupling

and the evolution of multicellularity. *Nature* **515,** 75–79 (2014).

33. Yoon, H.-S. & Golden, J. W. Heterocyst pattern formation controlled by a diffusible

peptide. *Science* **282,** 935–938 (1998).

34. Flores, E., Herrero, A., Wolk, C. P. & Maldener, I. Is the periplasm continuous in

filamentous multicellular cyanobacteria? *Trends Microbiol.* **14,** 439–443 (2006).

35. Mariscal, V., Herrero, A. & Flores, E. Continuous periplasm in a filamentous,

heterocyst-forming cyanobacterium. *Mol. Microbiol.* **65,** 1139–1145 (2007).

36. Mullineaux, C. W. *et al.* Mechanism of intercellular molecular exchange in

heterocyst-forming cyanobacteria. *EMBO J.* **27,** 1299–1308 (2008).

37. Bornikoel, J. *et al.* Role of two cell wall amidases in septal junction and nanopore

formation in the multicellular cyanobacterium *Anabaena* sp. PCC 7120. *Front. Cell.*

Infect. Microbiol. **7,** 92–15 (2017).

38. Waterbury, J. B. & Stanier, R. Y. Patterns of growth and development in
pleurocapsalean cyanobacteria. *Microbiol. Rev.* **42,** 2–44 (1978).

39. Dagan, T. *et al.* Genomes of stigonematalean cyanobacteria (subsection V) and the
evolution of oxygenic photosynthesis from prokaryotes to plastids. *Genome Biol.*
Evol. **5,** 31–44 (2013).

40. Shih, P. M. *et al.* Improving the coverage of the cyanobacterial phylum using
diversity-driven genome sequencing. *Proc. Natl. Acad. Sci. USA* **110,** 1053–1058
(2013).

41. Giovannoni, S. J. *et al.* Evolutionary relationships among cyanobacteria and green
chloroplasts. *J. Bacteriol.* **170,** 3584–3592 (1988).

42. Baracaldo, P. S., Hayes, P. K. & Blank, C. E. Morphological and habitat evolution in the
cyanobacteria using a compartmentalization approach. *Geobiol.* **3,** 145–165 (2005).

43. Schirrmeister, B. E., Antonelli, A. & Bagheri, H. C. The origin of multicellularity in
cyanobacteria. *BMC Evol. Biol.* **11,** 45 (2011).

44. Uyeda, J. C., Harmon, L. J. & Blank, C. E. A comprehensive study of cyanobacterial
morphological and ecological evolutionary dynamics through deep geologic time.
PLoS ONE **11,** e0162539–32 (2016).

45. Tomitani, A., Knoll, A. H., Cavanaugh, C. M. & Ohno, T. The evolutionary
diversification of cyanobacteria: Molecular–phylogenetic and paleontological
perspectives. *Proc. Natl. Acad. Sci. USA* **103,** 5442–5447 (2006).

46. Shi, T. & Falkowski, P. G. Genome evolution in cyanobacteria: The stable core and the
variable shell. *Proc. Natl. Acad. Sci. USA* **105,** 2510–2515 (2008).

47. David, L. A. & Alm, E. J. Rapid evolutionary innovation during an Archaean genetic
expansion. *Nature* **469,** 93–96 (2010).

48. Tria, F. D. K., Landan, G. & Dagan, T. Phylogenetic rooting using minimal ancestor

deviation. *Nat. Ecol. Evol.* **1,** 0193 (2017).

49. Tomitani, A., Knoll, A. H., Cavanaugh, C. M., & Ohno, T. The evolutionary

diversification of cyanobacteria: molecular–phylogenetic and paleontological

perspectives. *Proc. Natl. Acad. Sci. USA* **103**, 5442-5447 (2006).

50. Wolk, C. P., Ernst, A. & Elhai, J. in *The molecular biology of cyanobacteria* 769–823

(Springer, Dordrecht, 1994).

51. O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current status,

taxonomic expansion, and functional annotation. *Nucleic Acids Res.* **44,** D733–45

(2016).

52. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment

search tool. *J. Mol. Biol.* **215,** 403–410 (1990).

53. Tatusov, R. L., Koonin, E. V. & Lipman, D. J. A genomic perspective on protein families.

Science **278,** 631–637 (1997).

54. Rice, P., Longden, I. & Bleasby, A. EMBOSS: the European Molecular Biology Open

Software Suite. *Trends Gen.* **16,** 276–277 (2000).

55. Enright, A. J., Van Dongen, S. & Ouzounis, C. A. An efficient algorithm for large-scale

detection of protein families. *Nucleic Acids Res.* **30,** 1575–1584 (2002).

56. Katoh, K. & Standley, D. M. MAFFT Multiple Sequence Alignment Software version 7:

improvements in performance and usability. *Mol. Biol. Evol.* **30,** 772–780 (2013).

57. Guindon, S. *et al.* New algorithms and methods to estimate maximum-likelihood

phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* **59,** 307–321

(2010).

58. Zahr, J. H. *Biostatistical analysis.* (2014).

59. Benjamini, Y. & Yekutieli, D. The control of the false discovery rate in multiple testing

under dependency. Ann. Stat. 29, 1165–1188 (2001).

60. Cormen, T. H., Leiserson, C. E., Rivest, R. L. & Stein, C. in *Section 22.4: Topological sort*

549–552 (2001).

Acknowledgments

We thank Tom Williams, Anne Kupczok, Tanita Wein, and Caroline Rose for comments

on the manuscript. KH received funding from the European Union's Framework

Programme for Research and Innovation Horizon 2020 (2014–2020) under the Marie

Skłodowska-Curie Grant Agreement No. 657096 and from the German Academic

Exchange Service (DAAD). F.D.K.T. was supported by CAPES (Coordination for the

Improvement of Higher Education Personnel–Brazil).

Author contributions

K.H. and T.D. conceived the study. K.H. collected the traits data. G.L., T.D. and F.D.K.T.

developed and implemented the method. G.L. performed all analyses. K.H., G.L., and T.D.

wrote the manuscript with contributions from F.D.K.T..

Author information

The authors declare no competing financial interests and no conflict of interest.

Correspondence and requests for materials should be addressed to K.H.

Table 1. Description of cyanobacterial cell types, morphological and physiological traits, and their habitat and life style.

Cell types	
Vegetative cells	Photosynthetic cells.
HORMOGONIA*	Motile reproductive cells that result from repeated rounds of fission without intermittent growth phases. They break of the mother filament, ensuring the reproduction and dispersal of benthic species.
Necridia*	Dead cells resulting from PCD for hormogonia release.
HETEROCYSTS*	Thick-walled cells that are specialized in fixing N_2.
AKINETES*	Thick-walled, spore-like cells that provide reproduction, dormancy, and resilience.
BAEOCYTES*	Reproductive cells that result from repeated rounds of fission without intermittent growth phases.

Morphological and physiological traits	
UNICELLULAR	Single-celled morphology. After cell division cells separate.
FILAMENTOUS*	Multi-celled morphology. Cells remain attached after cell division.
NITROGEN FIXATION	Fixation of N_2 into ammonium.
SHEATH	Part of the cell envelope, located outside the cell wall.
MUCILAGE	Part of the envelope, located outside the cell wall, comprised of EPS, without a defined structure.
GAS VESICLES*	Intracellular gas-filled chambers for regulating buoyancy in the water column.
MOTILITY	Movement across surfaces or through a liquid medium.
FISSION IN MULTIPLE PLANES	Cell division in two or three perpendicular planes.
TRUE BRANCHING*	Fission in multiple planes leads to branches that remain attached to the main filament.

Habitat and life style	
FRESHWATER	Aquatic environments with salinity between 0-0.5ppt.
MARINE	Aquatic environments with salinity between 30–50ppt.
THERMOPHILIC	Optimal growth temperature above 45°.
PLANKTONIC	Organism that lives in the plankton (not attached).
SESSILE/ BENTHIC	Attachment to a substrate.
MATS	Growth inside thick, laminated, microbial structures.
FREE-LIVING	Organism that lives autonomously, in contrast to:
NOT FREE-LIVING	Organism that lives in a symbiotic relationship.
EPILITHIC/ ENDOLITHIC	Growth on or inside rocky substrates.
EPIPHYTIC	Growth on plants.
PERIPHYTIC	Attachment to underwater substrates.

* Multicellularity markers: traits that are adaptations on the level of the filament.

SMALL CAPS indicate the traits that have been used in the analysis.

460 **Table 2. Resolution of on-going debates regarding trait precedencies in**
461 **cyanobacterial evolution.**

Debate and resolution	Description of debate
Unicellular – Filamentous Trait-pair polarity tests show that the cyanobacterial ancestor was UNICELLULAR and that the FILAMENTOUS morphology arose later (FDR adjusted U-test p-value 2.3×10^{-110}).	Whereas early work[41] suggested that both morphologies had multiple origins, a subsequent analysis found the ancestor to be unicellular and the filamentous morphology to arise in independent lineages of the cyanobacterial tree[42]. Another view is that the filamentous morphology evolved early during cyanobacterial evolution and was subsequently lost and regained several times[43].
Planktonic - Benthic Our results show that the cyanobacteria ancestor traits include PLANKTONIC, MOTILITY and FREE LIVING (all FDR adjusted U-test p-values $<3.2 \times 10^{-69}$, see Supplementary table 2).	Whether the cyanobacteria ancestor was planktonic is a matter of debate and opposing views on the topic have been published[11,42,44].
N$_2$ fixation – No N$_2$ fixation Trait-pair polarity tests show that N FIXATION is a derived trait and that the ancestor of cyanobacteria was lacking the ability to fix N$_2$ (FDR adjusted U-test p-value 2.6×10^{-125}).	Whereas some studies claimed the last cyanobacterial common ancestor to fix N$_2$[45], there are others that concluded that it could not fix N$_2$ and that cyanobacteria must have acquired this trait several times independently[42,46,47].
Freshwater – Marine Our results show that there is no evidence for either MARINE or FRESHWATER environments as ancestral or derived habitat (Simultaneous polarity with FDR adjusted U-test p-value 5.4×10^{-148}).	Some studies suggest that early cyanobacteria lived in freshwater and subsequently diverged into marine environments[39,42,44], whereas others provide evidence in support of a marine origin[48]. The cyanobacteria ancestor most likely inhabited an aquatic environment and colonized both environments early.
Akinetes – Heterocysts Our results show that AKINETES and HETEROCYSTS emerged simultaneously (FDR adjusted U-test p-value 1.8×10^{-124}).	There is a common agreement that these cell types appeared late in cyanobacterial evolution[49], but there is a controversy about whether they shared a common ancestor and appeared simultaneously[50] or successively[42,49].

Fig. 1. Phylogenomic reconstruction of trait emergence chronology. Stages in the procedure are depicted clockwise from top-left. A genome-wide sample of single-copy protein coding genes provides measures of ancestor-descendant relations (Ancestor Deviation, AD)[48]. The AD support is then coupled to traits presence-absence patterns for each gene tree. Considering a specific pair of traits, their emergence polarity is deduced from formal statistical tests that contrast alternative hypothesized ancestor-descendant relations, utilizing paired information from all gene trees. All pairwise comparisons of traits are then combined into an overall trait-pairs polarity matrix, and if partial ordering of the traits (i.e., no polarity cycles) is possible, a feed-forward network can be inferred by topological sorting.

Fig. 2. Trait precedence in the evolution of cyanobacteria. (**a**) Trait-pairs polarity matrix. (**b**) Trait emergence order network. Cells in the matrix are shaded according to the polarity of the row and column traits, and the FDR-corrected significance of the U tests (for trait definition and test results see Supplementary Tables 1 and 2). In this dataset there are no cycles of conflicting ordering among the 300 inferred pairwise polarities. Thus, the traits form a *partial order* that is used to determine the trait order in the matrix (**a**) and is visualized as a feed-forward network (**b**). Shades of green mark the

482   four phases of trait emergence. Traits that can be regarded as multicellularity markers

483   in cyanobacteria are labelled in red (Table 1).