Title: PESCA: A scalable platform for the development of cell-type-specific viral drivers

One sentence summary: Highly paralleled functional evaluation of enhancer activity in single

cells generates new cell-type-specific tools with broad medical and scientific applications.

Authors:

Sinisa Hrvatin[1, *], Christopher P. Tzeng[1], M. Aurel Nagy[1], Hume Stroud[1], Charalampia

Koutsioumpa[1, 2], Oren F. Wilcox[1], Eric C. Griffith[1], Michael E. Greenberg[1, *]

Affiliations:

1. Department of Neurobiology, Harvard Medical School, Boston, Massachusetts, USA

2. BBS Program, Harvard Medical School, 220 Longwood Avenue, Boston, MA 02115,

15   USA

16   *Co-corresponding authors:

17   Sinisa Hrvatin (sinisa_hrvatin@hms.harvard.edu),

18   Michael E. Greenberg (michael_greenberg@hms.harvard.edu)

Abstract:

Enhancers are the primary DNA regulatory elements that confer cell type specificity of gene expression, thus giving rise to the hundreds of different cell types that make up a complex organism. Recent studies characterizing individual enhancers have revealed their potential to direct heterologous gene expression in a highly cell-type-specific manner. However, it has not yet been possible to systematically identify such enhancers for each of the many cell types in an organism. We have developed PESCA, a scalable and generalizable method coupling ATAC and single-cell RNA sequencing protocols, to identify cell-type-specific enhancers that should enable genetic access and perturbation of gene function in virtually any mammalian cell type. Focusing on the highly heterogeneous mammalian cerebral cortex, we apply PESCA to find enhancers and generate viral reagents capable of accessing and manipulating a rare subset of somatostatin-expressing cortical interneurons with high specificity. This study demonstrates the utility of this readily generalizable platform for developing new cell-type-specific viral reagents, with significant implications for both basic and translational research.

Main text:

Introduction

Enhancers are DNA elements that regulate gene expression to produce the unique complement of proteins necessary to establish a specialized function for each cell type in an organism. Large scale efforts to build a definitive catalogue of cell types(*1–4*) based on their gene expression have only recently been met with attempts to map epigenomic regulatory landscapes as well(*5–7*), enabling a mechanistic understanding of the underlying gene expression that is critical for cell-type-specific development, identity, and unique function. Importantly, characterization of individual enhancers has revealed their potential to direct highly cell-type-specific gene expression in both endogenous and heterologous contexts(*8–11*), making them ideal for developing tools to access, study, and manipulate virtually any mammalian cell type.

Indeed, despite being able to transcriptionally catalogue distinct cell subpopulations, our efforts to study their function are fundamentally impeded by our limited ability to specifically access them. Nowhere is this more apparent than in the mammalian cerebral cortex, which is composed of a staggeringly intricate and interconnected ensemble of cell types that together enable the complex computations that the brain must make. Glutamatergic excitatory neurons propagate signal across neural circuits, while GABAergic inhibitory interneurons play an essential role in cortical signal processing by modulating neuronal activity, balancing excitability, and gating information(*12–14*). Although relatively sparse, interneurons are highly diverse; for example, somatostatin-expressing cortical interneurons comprise several anatomically, electrophysiologically, and molecularly defined cell types whose dysfunction is associated with neuropsychiatric and neurological disorders(*2*, *15*, *16*). Given the vast diversity of cell types in the brain, and the inability of our current transgenic tools to access most neuronal

cell types, enhancer-driven viral reagents have the potential to become the next generation of

cell-type-specific transgenic tools enabling, for the first time, facile, inexpensive, cross-species,

and targeted observation and functional study of neuronal cell-types and circuits.

Despite the potential of cell-type-specific enhancers to revolutionize neuroscience

research, cell-type-restricted gene regulatory elements (GREs) have not yet been systematically

identified. Moreover, functional evaluation of candidate GRE-driven viral vectors is currently

laborious, expensive, and low-throughput, typically relying on the production of individual viral

vectors and the assessment of expression across a limited number of cell types by *in situ*

hybridization or immunofluorescence. The lack of a high-throughput platform for rapid

identification and functional testing of enhancers is therefore a critical bottleneck impeding the

generation of new cell-type-specific viral reagents required to elucidate the function of each cell

type in a complex organism.

To address these issues, we developed a scalable Paralleled Enhancer Single Cell Assay

(PESCA) to identify and functionally assess the specificity of hundreds of GREs across the full

complement of cell types present in any tissue of interest. In the PESCA protocol, the expression

of a barcoded pool of AAV vectors harboring GREs is analyzed by single-nucleus RNA

sequencing (snRNA-seq) to evaluate the specificity of each constituent GRE across tens of

thousands of individual cells in the target tissue, through the use of an orthogonal cell-indexed

system of transcript barcoding (**Fig. 1a**).

We validated the efficacy of PESCA in the murine primary visual cortex by identifying

GREs that have the ability to confine AAV expression to a layer-restricted, electrophysiologically

distinct subset of somatostatin (SST)-expressing interneurons, and showed that these vectors can

be used to modulate neuronal activity in a cell-type-specific manner. We chose to focus on SST

neurons in the brain because this population is known to be diverse and to be composed of

relatively rare subpopulations(*2*, *15*, *17*), and thus might serve as a good test case for the efficacy

of PESCA. As described below, these results highlight the utility of PESCA for identifying an

enhancer that drives gene expression selectively in a subset of neurons and establish the use of

PESCA as a platform of broad interest to the research and gene therapy community, enabling the

generation of cell-type-specific AAVs for virtually any cell type.

Results

GRE selection and library construction

To identify candidate SST interneuron-restricted gene regulatory elements (GREs), we

carried out comparative epigenetic profiling of the three largest classes of cortical interneurons,

somatostatin (SST)-, vasoactive intestinal polypeptide (VIP)- and parvalbumin (PV)-expressing

cells. To this end, we employed the recently developed isolation of nuclei tagged in specific cell

types (INTACT)(*5*) method to isolate purified chromatin from of each of these cell types from

the cerebral cortex of adult (6-10-week-old) mice. Assay for transposase-accessible chromatin

using sequencing (ATAC-Seq)(*7*), which marks nucleosome-depleted gene regulatory regions

based on their enhanced accessibility to *in vitro* transposition by the Tn5 transposase, was then

used to identify genomic regions with enhanced accessibility in the SST (n = 279,221), PV (n =

275,631), and VIP (n = 258,646) chromatin samples. These datasets can be used as a resource to

identify putative gene regulatory elements as candidates for driving cell-type-specific gene

expression for the numerous SST, PV or VIP-expressing intraneuronal subtypes across diverse

cortical regions.

Among these putative gene regulatory regions, 16,386 (5.9%) were enriched or uniquely

present in SST cells relative to PV or VIP-expressing interneurons (**Fig. 1b, c**). To enrich for

GREs that thus could be useful reagents to study and manipulate interneurons across mammalian

species including in humans, we subsequently filtered the resulting list to exclude GREs with

poor mammalian sequence conservation (**Methods**, **Supplementary Fig. 1**). Remaining

elements were ranked based on cell-type-enrichment (**Methods**), with the top 287 SST-enriched

GREs selected for screening to identify enhancers that drive gene expression selectively in SST

interneurons of the primary visual cortex (**Fig. 1d**).

A PCR-based strategy was used to simultaneously amplify and barcode each GRE from

mouse genomic DNA (**Methods**). To minimize sequencing bias due to the choice of barcode

sequence, each GRE was paired with three unique barcode sequences. The resulting library of

861 GRE-barcode pairs was pooled and cloned into an AAV-based expression vector, with the

GRE element inserted 5' to a minimal promoter driving a GFP expression cassette and the GRE-

paired barcode sequences inserted into the 3' untranslated region (UTR) of the GRE-driven

transcript (**Methods, Fig. 2a, Supplementary Fig. 2)**. This configuration was chosen to

maximize the retrieval of the barcode sequence during single-cell RNA sequencing. The library

was packaged into AAV9, which exhibits broad neural tropism(*18*). The complexity of the

resulting rAAV-GRE library was then confirmed by Next Generation Sequencing, detecting 802

of the 861 barcodes (93.2%), corresponding to 285 of the 287 GREs (99.3%) (**Fig. 2b**).

PESCA screen identifies GREs highly enriched for SST$^+$ interneurons

To quantify the expression of each rAAV-GRE vector across the full complement of cell types in

the mouse visual cortex, we used a modified single-nucleus RNA-Seq (snRNA-Seq) protocol to

first determine the cellular identity of each nucleus and then quantify the abundance of the GRE-

paired barcodes in the transcriptome of nuclei assigned to each cell type. Two injections (800 nL

each) of the pooled AAV library (1×10^{13} viral genomes/mL) were first administered to the

primary visual cortex (V1) of two 6-week-old C57BL/6 mice. Twelve days following injection,

the injected cortical regions were dissected and processed to generate a suspension of nuclei for

snRNA-Seq using the inDrops platform([19, 20]) (**Methods**). A total of 32,335 nuclei were

subsequently analyzed across the two animals, recovering an average of 866 unique non-viral

transcripts per nucleus, representing 610 unique genes (**Supplementary Fig. 3a, b**).

Since droplet-based high-throughput snRNA-Seq samples the nuclear transcriptome with

low sensitivity([19]), viral-derived transcripts were initially detected in only 3.9% of sampled

nuclei. We therefore designed a modified PCR-based approach to enrich for barcode-containing

viral transcripts, which yielded deep coverage of AAV-derived transcripts with simultaneous

shallow coverage of the non-viral transcriptome. PCR enrichment increased the viral transcript

recovery 382-fold in the sampled nuclei, to an average of 15.6 unique viral transcripts, 6.0

unique GRE-barcodes, and 5.7 unique GREs per cell **(Fig. 2b, Supplementary Fig. 3c)**. Using

this modified protocol, viral transcripts were identified across 86% of cells (**Supplementary Fig.**

3d, e), with a high correlation (r = 0.9, $p < 2.2 \times 10^{-16}$) observed between the abundance of each

barcoded AAV in the library and the number of cells infected by that AAV (**Supplementary**

Fig. 3f).

Nuclei were classified into ten cell types using graph-based clustering and expression of

known marker genes **(Methods, Fig. 2c, d, Supplementary Fig. 4)**. The average expression of

each viral-derived barcoded transcript was analyzed across all ten cell types, and an enrichment

score was calculated from the ratio of expression in Sst^+ nuclei compared to all Sst^- nuclei. As

expected, sets of three barcodes associated with the same GRE showed highly statistically

correlated enrichment scores (r = 0.53 ± 0.03, p < 2.2×10⁻¹⁶) (**Fig. 2e, f, Supplementary Fig. 5**),

which were abolished when barcodes were randomly shuffled (shuffled r = 0.002 ± 0.06; Wilcox

test between data and shuffled data, p = 0.003).

Having confirmed a robust, non-random correlation in enrichment scores among the three

barcodes associated with each GRE, we next computed a single expression value for each of the

287 viral drivers by aggregating expression data from barcodes associated with the same GRE,

and carried out differential gene expression analysis between Sst^+ and Sst^- cells for each rAAV-

GRE. Differential gene expression analysis between Sst^+ and Sst^- cells for each rAAV-GRE

revealed a marked overall enrichment of viral-derived transcripts in the Sst^+ subpopulation

(**Supplementary Fig. 6a**). Indeed, multiple viral drivers were identified that promoted highly

specific reporter expression in the Sst^+ subpopulation (q < 0.01, fold-change > 7, **Fig. 2g-i,**

Supplementary Fig. 6b).

In situ **characterization of rAAV-GRE reporter expression**

We next sought to validate the cell-type-specificity of the resulting hits using methods

that do not rely on single-cell sequencing-based approaches. To this end, we selected three of the

top five viral drivers (GRE12, GRE22, GRE44), as well as a control viral construct lacking the

GRE element (ΔGRE), for injection into V1 of adult transgenic Sst-Cre; Ai14 mice, in which

SST⁺ cells express the red fluorescent marker tdTomato. Fluorescence analysis twelve days

following injection with rAAV-GRE12/22/44-GFP revealed strong yet sparse GFP labeling

centered around cortical layers IV and V (**Fig. 3a-c**). By contrast, the control rAAV-ΔGRE-GFP

showed a strikingly different pattern of GFP expression concentrated around the sites of

injection, with expression in a larger number of cells (**Fig. 3d**). Many rAAV-GRE12/22/44-GFP

virally infected cells were indeed SST-positive, marked by the high degree of overlapping GFP

and tdTomato expression: $90.7\% \pm 2.1\%$ for rAAV-GRE12-GFP (170 cells, 4 animals); $72.9 \pm$

4.2% for rAAV-GRE22-GFP (1164 cells, 3 animals), and $95.8 \pm 0.6\%$ for rAAV-GRE44-GFP

(759 cells, 4 animals). (**Fig. 3e, f, Supplementary Fig. 7**). By contrast, we observed that only

$27.2 \pm 1.9\%$ of GFP$^+$ cells following rAAV-ΔGRE-GFP infection were also positive for

tdTomato expression (2066 cells, 3 animals, **Fig. 3e, f**), indicating that the tested GREs serve to

effectively restrict AAV payload expression to SST$^+$ interneurons. It is notable that the GREs

seemingly not only promote expression in SST$^+$ cells but also reduce background expression in

SST$^-$ cells, indicating the tested GREs confer both enhancer and insulator functionality.

Consistent with this hypothesis, the incorporation of the GREs into the rAAV both increased the

number of SST$^+$/GFP$^+$ cells (1.7-2-fold) and dramatically (3-32-fold) decreased the number of

SST$^-$ cells that expressed GFP (**Fig. 3g, Supplementary Fig. 8**). To further investigate the

specificity of our viral drivers among cortical interneuron cell types we injected each construct

into Vip-Cre; Ai14$^+$ mice in which all VIP$^+$ cells express tdTomato, or used fluorescence

antibody staining to label PV-expressing cells (**Supplementary Fig. 9**). Fluorescent signal

analysis indicated the percentage of GFP$^+$ cells that were either VIP$^+$ or PV$^+$ (rAAV-SST12-

GFP$^+$ [$2.6 \pm 2.6\%$], rAAV-GRE22-GFP$^+$ [$3.5 \pm 2.0\%$] and rAAV-GRE44-GFP$^+$ [$6.0 \pm 2.7\%$],

Fig. 3h). This confirms that among major interneuron cell classes, all three vectors are highly

SST-specific.

Because at least five subtypes of cortical SST$^+$ interneurons have been identified based

on the laminar distribution of their cell bodies and projections(*15*, *21*), we also investigated the

laminar distribution of GFP-expressing cells for the three Sst-enriched viral drivers. Intriguingly,

the majority of rAAV-GRE12-GFP$^+$ and rAAV-GRE44-GFP$^+$ SST$^+$ cells were found to reside in

layers IV and V, distinct from the distribution observed for the full SST$^+$ cell population in visual

cortex (p = 1.3×10^{-6}, p < 2.2×10^{-16}, respectively, Mann-Whitney U test, two-tailed, **Fig. 3i, j**),

raising the possibility that these constructs may preferentially label a specific subtype(s) of SST$^+$

interneuron. Consistent with this hypothesis, we observed that these two viral drivers mediated

reporter expression in only a relatively small fraction of all SST$^+$ cells within the region of

infection (44.5 ± 12.0% for rAAV-GRE12-GFP and 35.9 ± 6.2% for rAAV-GRE44-GFP)

compared to rAAV-GRE22-GFP (**Fig. 3k, Supplementary Fig. 10**). Together, these findings

suggest that PESCA may support the isolation of viral drivers capable of discriminating between

fine-grained cell-types within a given interneuron cell class.

Electrophysiological characterization of rAAV-GRE-GFP-expressing SST subtypes

In addition to variability in laminar distribution, different electrophysiological

phenotypes have also been observed in cortical SST interneurons(*22*, *23*). To determine whether

AAV-GRE reporters can be used to distinguish electrophysiologically distinct SST subtypes, we

injected rAAV-GRE44-GFP into the visual cortex of adult Sst-Cre; Ai14 mice and performed

whole-cell current-clamp recordings from double GFP and tdTomato positive neurons (rAAV-

GRE44-GFP$^+$), as well as immediately nearby tdTomato positive but GFP negative cells (rAAV-

GRE44-GFP$^-$). Our recordings indicate that both rAAV-GRE44-GFP$^+$ and rAAV-GRE44-GFP$^-$

neurons are adapting SST interneurons with high input resistances and features consistent with

deep layer cortical SST neurons(*22*, *24*) (**Fig. 4a, b**).

However, rAAV-GRE44-GFP$^+$ SST neurons are distinct in a several electrophysiological

parameters. Action potentials of rAAV-GRE44-GFP$^+$ SST neurons are significantly broader

than those of rAAV-GRE44-GFP$^-$ SST neurons (**Fig. 4c, d**), perhaps due to differences in

expression of certain channels in these subgroups of SST neurons, such as voltage-activated

potassium channels, and BK calcium-activated potassium channels(*25*, *26*). Furthermore,

rAAV-GRE44-GFP$^+$ SST neurons have a lower rheobase, and fire action potentials with a slower

rising phase, and at lower maximal frequencies compared to rAAV-GRE44-GFP$^-$ SST neurons

(Fig. 4a, d, Supplementary Table 1). Our electrophysiology experiments further illuminate the

potential of PESCA to identify novel, functionally distinct subgroups of previously defined

interneuron types.

Modulation of neuronal activity with rAAV-GREs

240   Finally, we evaluated whether the identified viral drivers support sufficiently high and

persistent levels of payload expression to effectively modulate SST$^+$ cell physiology. Designer

receptors exclusively activated by designer drugs (DREADDs) are a commonly employed viral

payload to dynamically regulate neuronal activity in response to the synthetic ligand clozapine-

N-oxide (CNO)(*27*). We therefore injected the visual cortex of adult wild-type mice (6-8-week-

old) with rAAV-GRE12-Gq-DREADD-tdTomato and performed electrophysiological recordings

from tdTomato$^+$ cells of acute cortical slices in a whole-cell, current-clamp configuration two

weeks post-injection. All recordings from tdTomato$^+$ cells evoked with depolarizing current

steps showed striking sensitivity to CNO, as shown by significantly increased firing rates and

depolarized resting membrane potentials during bath application of CNO (**Fig. 4e-g**). To ensure

that increases in firing rate upon CNO application were specific to infected SST$^+$ neurons, we

performed recordings from nearby uninfected pyramidal neurons that were identified by

morphology and found that there was no statistically significant increase in firing rate upon CNO

application (**Fig. 4h-j**). These data demonstrate the ability of these reagents to robustly and

specifically modulate the activity of SST$^+$ cells in non-transgenic animals.

Discussion

The PESCA platform merges the principle of massively paralleled reporter assays (MPRA)(*28,*

29) with scRNA-seq and represents a significant advancement in current approaches to viral

vector design, as it enables the rapid screening of hundreds of viral permutations for enhanced

cell-type-specificity. In this study, we applied PESCA to screen putative enhancer elements for

drivers that robustly and specifically target a rare SST$^+$ population of GABAergic interneurons in

the mouse central nervous system. Given our understanding of the generality of enhancer

function across tissues, this approach could be readily applied to virtually any other neuronal or

non-neuronal cells types, diverse model organisms, tissues, and viral types. Moreover, PESCA is

not limited to GRE screening; the method can be easily adapted to assess the cell-type-specificity

of viral capsid variants or other mutable aspects of viral design. This study addresses the urgent

need for new tools to access, study, and manipulate specific cell types across complex tissues,

organ systems, and animal models, demonstrating the importance and broad utility of PESCA

beyond neuroscience research. Moreover, as the promise of gene therapy to treat and cure a

broad range of diseases is being realized, PESCA has the potential to pave the way for a new

generation of targeted gene therapy vehicles for diseases with cell-type-specific etiologies, such

as congenital blindness, deafness, cystic fibrosis, and spinal muscular atrophy.

Figures and figure legends

Figure 1 – Experimental strategy and GRE selection

(**a**) Paralleled Enhancer Single Cell Assay (PESCA). Comparative ATAC-Seq is used to

identify candidate GREs. A library of gene regulatory elements (GREs) is inserted upstream of a

minimal promoter-driven GFP. The viral barcode sequence is inserted in the 3'UTR, and the

vector packaged into rAAVs. Following *en masse* injection of the rAAV library, the specificity

of the constituent GREs for various cell types *in vivo* is determined by single-nucleus RNA

sequencing, measuring expression of the barcoded transcripts in tens of thousands of individual

cells in the target tissue. Finally, bioinformatic analysis determines the most cell-type-specific

barcode-associated rAAV-GRE-GFP constructs. pA = polyA tail.

(**b**) Area-proportional Venn diagram of the number of putative GREs identified by

ATAC-Seq of purified PV, SST, and VIP interneuron chromatin. Overlapping areas indicate

shared putative GREs. Non-overlapping areas represent GREs that are either unique or strongly

enriched in a single cell type.

(**c**) Representative ATAC-seq genome browser traces of a putative GRE enriched in SST-

, PV-, or VIP-interneurons. Sequence conservation across the Placental mammalian clade is also

shown.

(**d**) Putative GREs (n = 323,369) are plotted based on average sequence conservation

(phyloP, 60 placental mammals) and SST-specificity (ratio of the average ATAC-Seq signal

intensity between SST samples and non-SST samples). Dashed vertical line indicates the

minimal conservation value cutoff (0.5). Green coloring indicates the 287 most SST-specific

GREs selected for PESCA screening.

Figure 2 – PESCA screen identifies GREs highly enriched for SST⁺ interneurons

(**a**) PESCA library plasmid map. ITR, inverted terminal repeats; GRE, gene regulatory

element; pr, HBB minimal promoter; int, intron; GFP, green fluorescent protein; WPRE,

Woodchuck Hepatitis Virus post-transcriptional regulatory element; BAR, 10-mer sequence

barcode associated with each GRE; pA, polyadenylation signal.

(**b**) Library complexity plotted as distribution of the abundance of the 861 barcodes and

287 GREs in the AAV library. Barcodes and GREs were binned by number of sequencing reads

attributed to each barcode or GRE within the library.

(**c**) Transcript count per nucleus (n = 32,335 nuclei). Sequencing libraries were prepared

with or without PCR-enrichment for viral transcripts. PCR enrichment resulted in a 382-fold

increase in the number of recovered viral transcripts (p = 0) to an average of 15.6 unique viral

transcripts per nucleus. Displayed as Log10(Count+1).

(**d**) t-SNE plot of 32,335 nuclei from V1 cortex of two animals**.** Colors denote main cell

types: Exc (Excitatory neurons), Pv (PV Interneurons), Sst (SST Interneurons), Vip (VIP

interneurons), Npy (NPY Interneurons), Astro (Astrocytes), Vasc (Vascular-associated cells),

Micro (Microglia), Olig (Oligodendrocytes), OPCs (Oligodendrocyte precursor cells).

(**e**) Marker gene expression across cell types. Color denotes mean expression across all

nuclei normalized to the highest mean across cell types. Size represents the fraction of nuclei in

which the marker gene was detected.

(**f**) Three-dimensional dot plot with each dot representing one GRE (n = 287). The values

on each axis represent the SST fold-enrichment calculated for each GRE based on the three

barcodes paired with that GRE. Plane of correlation between the enrichment values calculated

from three sets of barcodes associated with 287 GREs (r = 0.53 ± 0.03, p < 2.2×10^{-16}). Color

gradient indicates the average enrichment between the three barcodes.

(**g**) Pairwise Pearson correlation between the enrichment values calculated from three sets

of barcodes associated with 287 GREs for experimental data (Exp. Data, r = 0.53 ± 0.03, p <

2.2×10^{-16}) and after random shuffling of enrichment values (Shuffled Data, r = 0 ± 0.06).

(**h**) GREs ranked by average expression specificity for SST interneurons. Shading

indicates the minimal and maximal specificity calculated by analyzing each of the three barcodes

associated with a GRE. Blue indicates the five top hits that also passed a statistical test for SST

interneuron enrichment (FDR-corrected q < 0.01).

(**i**) Expression of the top five hits: GRE12, GRE19, GRE22, GRE44, GRE80. For each

GRE, expression values are split into two animals, and, for each animal, into the three barcodes

associated with that GRE. Color denotes mean expression across all nuclei normalized to the

highest mean across cell types. Size represents the fraction of nuclei in which the marker gene

was detected.

(**j**) t-SNE plot of 32,335 nuclei from V1 cortex of two animals, showing expression of

GRE12, GRE19, GRE22, GRE44, and GRE80**.** Plot is pseudocolored based on the mean GRE

expression in each cell type.

Figure 3 – *In situ* characterization of rAAV-GRE reporter expression

(**a-d**) Fluorescent images from adult Sst-Cre; Ai14 mouse visual cortex twelve days following injection with rAAV-GRE-GFP as indicated. Scale bars 100 μm.

(**e**) Identification of rAAV-GRE-GFP$^+$ cells that express tdTomato (SST$^+$). Each dot represents a GFP$^+$ cell (n = 2066, 172, 1164, and 765, for AAV-[ΔGRE, GRE12, GRE22, GRE44]-GFP, respectively). Cyan indicates tdTomato$^+$ (SST$^+$) cells. Distribution of cell frequency across tdTomato intensity is plotted on the right for each construct.

(**f**) Quantification of the fraction of GFP$^+$ cells that are SST$^+$. Each dot represents one animal. Box plot represents mean ± standard error of the mean (s.e.m). Values are 27.2 ± 1.9%, 90.7 ± 2.1, 72.9 ± 4.2%, and 95.8 ± 0.6% for AAV-[ΔGRE, GRE12, GRE22, GRE44]-GFP, respectively.

(**g**) Quantification of the number of GFP$^+$ SST$^-$ cells normalized for area of infection. Each dot represents one animal. Box plot represents mean ± standard error of the mean (s.e.m). Values are 198.0 ± 46.0, 16.4 ± 6.2, 56.0 ± 17.3 and 6.1± 2.1 cells/mm^2 for AAV-[ΔGRE, GRE12, GRE22, GRE44]-GFP, respectively.

(**h**) Quantification of the fraction of GFP$^+$ cells that are PV$^+$ or VIP$^+$. Box plot represents mean ± standard error of the mean (s.e.m). Fraction of AAV-GRE-GFP$^+$ cells that are PV$^+$ is 1.4 ± 1.4%, 2.2 ± 0.7, and 4.3 ± 1.7% for AAV-[GRE12, GRE22, GRE44]-GFP, respectively. Similarly, the fraction of AAV-GRE-GFP$^+$ cells that are VIP$^+$ is 1.2 ± 1.2%, 1.3 ± 1.3%, and 1.7 ± 1.0% for AAV-[GRE12, GRE22, GRE44]-GFP$^+$ cells, respectively.

(**i**) Distribution of the location of GFP-expressing cells as function of distance from the pia. Gray represents SST$^+$ cells (n = 2648); Colored lines represents GFP$^+$ SST$^+$ cells (n = 2066,

172, 1164, and 765, respectively, for AAV-[ΔGRE, GRE12, GRE22, GRE44]-GFP). Shading

represents the 95% confidence interval.

(**j**) Quantification of the fraction of GFP$^+$ cells that are present it each cortical layer. Each

dot represents one animal. Box plot represents mean $\pm$ standard error of the mean (s.e.m). Gray

represents all SST$^+$ cells, colored plots represent GFP$^+$ SST$^+$ cells respectively, for AAV-

[GRE12, GRE22, GRE44]-GFP).

(**k**) Quantification of the fraction of Sst-Cre; Ai14$^+$ cells within the infection area that are

GFP$^+$. Each dot represents one animal. Box plot represents mean $\pm$ standard error of the mean

(s.e.m). Values are 44.5 $\pm$ 12.0%, 73.4 $\pm$ 9.4%, and 35.9 $\pm$ 6.2% for AAV-[GRE12, GRE22,

GRE44]-GFP, respectively.

Figure 4 – Electrophysiology of neurons expressing an rAAV-GRE-driven reporter and modulation of neuronal activity with rAAV-GREs

(**a**) Representative current-clamp recordings from SST neurons in the visual cortex of

Sst-Cre; Ai14 mice injected with rAAV-GRE44-GFP. *Top*: Representative traces from a cortical

SST neuron with Cre-dependent expression of red fluorescent protein (RFP), in response to

1000ms depolarizing current injections as indicated in black ("GRE44-"). *Bottom*: Traces from a

RFP+ SST neuron with GRE44-driven expression of GFP ("GRE44+"). GRE44- SST neurons

were only recorded in the immediate vicinity of GRE44+ SST neurons.

(**b**) Recordings from GRE44+ and GRE44- neurons in response to hyperpolarizing,

1000ms currents. Asterisks indicate the sag likely due to the hyperpolarization-activated current

I_h. Rebound action potentials following recovery from hyperpolarization, likely due to low-

threshold calcium spikes mediated by T-type calcium channels, were also present in cells of both

groups. Same scale as (a).

(**c**) Broader action potentials in GRE44+ SST neurons (bottom) compared to GRE44-

SST neurons (top). Same vertical scale as (a) and (b).

(**d**) Electrophysiological properties that differ between GRE44+ (n=16 cells from 5 mice)

and GRE44- (n=16 cells from 4 mice) SST neurons, including rheobase (minimal amount of

current necessary to elicit a spike), maximal rate of rise during the depolarizing phase of the

action potential, the initial and steady state firing frequencies (both measured at the maximal

current step before spike inactivation), and spike width (measured as the width at half-maximal

spike amplitude).

(**e**) Representative current-clamp recordings from AAV-GRE12-Gq-tdTomato$^+$ cells

before and during CNO application.

(**f**) Increased firing rates of AAV-GRE12-Gq-tdTomato$^+$ cells evoked by depolarizing current injections upon bath application of CNO (3 animals, 6-7 cells). * $p < 0.05$; ** $p < 0.01$, paired t-test, two-tailed.

(**g**) Robust depolarization of AAV-GRE12-Gq-tdTomato$^+$ cells upon bath application of CNO (3 animals, 6-7 cells). *** $p < 0.001$, paired t-test, two-tailed.

(**h**) Representative recordings from nearby uninfected pyramidal neurons in the visual cortex of mice that were injected with AAV-GRE-12-Gq-tdTomato$^+$, before (top) and during CNO application (bottom).

(**i**) Firing rates of pyramidal neurons during CNO application remain unchanged (3 animals, 5 cells). ns, $p > 0.05$, paired t-test, two-tailed.

(**j**) Representative image of nearby uninfected pyramidal neuron that was recorded from

Acknowledgements

We thank D. Tom for help with image analysis, Drs. W. Renthal, J. Green, and members of the

Greenberg lab for discussions, the HMS Single Cell Core for single-nucleus RNA-seq sample

processing, Boston Children's Hospital Viral Core for AAV packaging, Dr. B. Bean for feedback

on electrophysiology, and the Harvard NeuroDiscovery Center Enhanced Neuroimaging Core for

imaging and image analysis. This work was supported by the National Institute of Health

BRAIN Initiative grant R01 MH114081-01, NIH grant T32GM007753 to M.A.N. and NIH

Training in the Molecular Biology of Neurodegeneration grant 5T32AG000222-23 to S.H.

Author Contributions

S.H. conceived the study and designed the experiments. H.S. performed ATAC-Seq. M.A.N.

and S.H. analyzed ATAC-seq data. S.H. cloned the library and performed the PESCA screen.

M.A.N. mapped sequencing data and viral barcodes. S.H. analyzed snRNA-seq data. S.H., C.K.

and O.F.W. cloned individual viral constructs. O.F.W. performed stereotactic injections and

tissue sectioning. S.H. performed imaging and image analysis. C.P.T. performed and analyzed

electrophysiology recordings. M.E.G. and E.C.G. advised on all aspects of the study. S.H.,

C.P.T., M.A.N, E.C.G. and M.E.G. wrote the manuscript.

Competing Financial Interests

The authors declare no competing financial interests.

Correspondence

Correspondence and requests for materials should be addressed to S.H.

(sinisa_hrvatin@hms.harvard.edu) or M.E.G. (michael_greenberg@hms.harvard.edu).

References

1.   A. Zeisel *et al.*, Cell types in the mouse cortex and hippocampus revealed by single-cell
RNA-seq. *Science*. **347**, 1138–1142 (2015).

2.   B. Tasic *et al.*, Shared and distinct transcriptomic cell types across neocortical areas. *Nature*.
563, 72–78 (2018).

3.   J. Cao *et al.*, Comprehensive single-cell transcriptional profiling of a multicellular organism.
Science. **357**, 661–667 (2017).

4.   A. B. Rosenberg *et al.*, Single-cell profiling of the developing mouse brain and spinal cord
with split-pool barcoding. *Science*. **360**, 176–182 (2018).

5.   A. Mo *et al.*, Epigenomic Signatures of Neuronal Diversity in the Mammalian Brain.
Neuron. **86**, 1369–1384 (2015).

6.   D. A. Cusanovich *et al.*, Multiplex single-cell profiling of chromatin accessibility by
combinatorial cellular indexing. *Science*. **348**, 910–914 (2015).

7.   J. D. Buenrostro *et al.*, Single-cell chromatin accessibility reveals principles of regulatory
variation. *Nature*. **523**, 486–490 (2015).

8.   J. Dimidschstein *et al.*, A viral strategy for targeting and manipulating interneurons across
vertebrate species. *Nat. Neurosci.* **19**, 1743–1749 (2016).

9.   J. Juettner *et al.*, Targeting neuronal and glial cell types with synthetic promoter AAVs in
mice, non-human primates, and humans. *bioRxiv*, 434720 (2018).

10. L. T. Graybuck *et al.*, Prospective, brain-wide labeling of neuronal subclasses with enhancer-
driven AAVs. *bioRxiv*, 525014 (2019).

11. J. K. Mich *et al.*, Epigenetic landscape and AAV targeting of human neocortical cell classes.
bioRxiv, 555318 (2019).

12. A. Kepecs, G. Fishell, Interneuron cell types are fit to function. *Nature*. **505**, 318–326
(2014).

13. O. Marín, Interneuron dysfunction in psychiatric disorders. *Nat. Rev. Neurosci.* **13**, 107–120
(2012).

14. B. Rudy, G. Fishell, S. Lee, J. Hjerling-Leffler, Three groups of interneurons account for
nearly 100% of neocortical GABAergic neurons. *Dev. Neurobiol.* **71**, 45–61 (2011).

15. W. Muñoz, R. Tremblay, D. Levenstein, B. Rudy, Layer-specific modulation of neocortical
dendritic inhibition during active wakefulness. *Science.* **355**, 954–959 (2017).

16. X. Jiang *et al.*, Principles of connectivity among morphologically defined cell types in adult
neocortex. *Science.* **350**, aac9462 (2015).

17. B. Tasic *et al.*, Adult mouse cortical cell taxonomy revealed by single cell transcriptomics.
Nat. Neurosci. **19**, 335–346 (2016).

18. C. N. Cearley, J. H. Wolfe, Transduction characteristics of adeno-associated virus vectors
expressing cap serotypes 7, 8, 9, and Rh10 in the mouse brain. *Mol. Ther. J. Am. Soc. Gene*
Ther. **13**, 528–537 (2006).

19. A. M. Klein *et al.*, Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic
Stem Cells. *Cell.* **161**, 1187–1201 (2015).

20. R. Zilionis *et al.*, Single-cell barcoding and sequencing using droplet microfluidics. *Nat.*
Protoc. **12**, 44–73 (2017).

21. J. Urban-Ciecko, A. L. Barth, Somatostatin-expressing neurons in cortical networks. *Nat.*
Rev. Neurosci. **17**, 401–409 (2016).

22. Y. Ma, H. Hu, A. S. Berrebi, P. H. Mathers, A. Agmon, Distinct subtypes of somatostatin-
containing neocortical interneurons revealed in transgenic mice. *J. Neurosci. Off. J. Soc.*
Neurosci. **26**, 5069–5082 (2006).

23. R. Tremblay, S. Lee, B. Rudy, GABAergic Interneurons in the Neocortex: From Cellular
Properties to Circuits. *Neuron.* **91**, 260–292 (2016).

24. H. Xu, H.-Y. Jeong, R. Tremblay, B. Rudy, Neocortical somatostatin-expressing GABAergic
interneurons disinhibit the thalamorecipient layer 4. *Neuron.* **77**, 155–167 (2013).

25. B. P. Bean, The action potential in mammalian central neurons. *Nat. Rev. Neurosci.* **8**, 451–
465 (2007).

26. T. Kimm, Z. M. Khaliq, B. P. Bean, Differential Regulation of Action Potential Shape and
Burst-Frequency Firing by BK and Kv2 Channels in Substantia Nigra Dopaminergic
Neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **35**, 16404–16417 (2015).

27. B. N. Armbruster, X. Li, M. H. Pausch, S. Herlitze, B. L. Roth, Evolving the lock to fit the
key to create a family of G protein-coupled receptors potently activated by an inert ligand.
Proc. Natl. Acad. Sci. **104**, 5163–5168 (2007).

28. R. P. Patwardhan *et al.*, Massively parallel functional dissection of mammalian enhancers in
vivo. *Nat. Biotechnol.* **30**, 265–270 (2012).

29. A. Melnikov *et al.*, Systematic dissection and optimization of inducible enhancers in human
cells using a massively parallel reporter assay. *Nat. Biotechnol.* **30**, 271–277 (2012).

Methods

Mice

Animal experiments were approved by the National Institute of Health and

Harvard Medical School Institutional Animal Care and Use Committee, following ethical

guidelines described in the US National Institutes of Health Guide for the Care and Use of

Laboratory Animals. For INTACT we crossed Sst-IRES-Cre (The Jackson Laboratory Stock #

013044), Vip-IRES-Cre (The Jackson Laboratory Stock # 010908) and Pv-Cre (The Jackson

Laboratory Stock # 017320) with SUN1-2xsfGFP-6xMYC (The Jackson Laboratory Stock #

021039) and used adult (6-12 wk old) male and female F1 progeny. For PESCA screening we

used adult (6-10 wk) C57BL/6J (The Jackson Laboratory, Stock # 000664) mice. For

confirmation of hits we crossed Sst-IRES-Cre (The Jackson Laboratory Stock # 013044), Vip-

IRES-Cre (The Jackson Laboratory Stock # 031628) and Gad2-IRES-Cre (The Jackson

Laboratory Stock # 028867) mice with Ai14 mice (The Jackson Laboratory Stock # 007914) and

used adult (6-12 wk old) male and female F1 progeny. All mice were housed under a standard 12

hr light/dark cycle.

INTACT purification and *in vitro* transposition

INTACT employs a transgenic mouse that expresses a cell-type-specific Cre and a Cre-

dependent SUN1-2xsfGFP-6xMYC (SUN1-GFP) fusion protein. Nuclear purifications were

performed from whole cortex of adult mice as previously described using anti-GFP antibodies

(Fisher G10362)(*1*, *2*). Isolated nuclei were gently resuspended in cold L1 buffer (50 mM Hepes

pH 7.5, 140 mM NaCl, 1 mM EDTA, 1 mM EGTA, 0.25% Triton X-100, 0.5% NP40, 10%

Glycerol, protease inhibitors), and pelleted at 800g for 5 minutes at 4°C. DNA libraries were

prepared from the nuclei using the Nextera DNA Library Prep Kit (Illumina) according to manufacturer's protocols. The final libraries were purified using the Qiagen minelute kit and sequenced on a Nextseq 500 benchtop DNA sequencer (Illumina).

ATAC-seq mapping:

All ATAC-seq libraries were sequenced on the Nextseq 500 benchtop DNA sequencer (Illumina). Seventy-five base pair (bp) single-end reads were obtained for all datasets. ATAC-seq experiments were sequenced to a minimum depth of 20 million (M) reads. Reads for all samples were aligned to the mouse genome (GRCm38/mm10, December 2011) using default parameters for the Subread (subread-1.4.6-p3, (Liao et al., 2013)) alignment tool after quality trimming with Trimmomatic v0.33 (Bolger et al., 2014) with the following command: java -jar trimmomatic-0.33.jar SE -threads 1 -phred33 [FASTQ_FILE] ILLUMINACLIP:[ADAPTER_FILE]:2:30:10 LEADING:5 TRAILING:5 SLIDINGWINDOW:4:20 MINLEN:45. Nextera adapters were trimmed out for ATAC-seq data. Duplicates were removed with samtools rmdup. To generate UCSC genome browser tracks for ATAC-seq visualization, BEDtools was used to convert output bam files to BED format with the bedtools bamtobed command. Published mm10 blacklisted regions (Consortium, 2012) were filtered out using the following command: bedops –not-element-of 1 [BLACKLIST_BED]. Filtered BED files were scaled to 20 M reads and converted to coverageBED format using the BEDtools genomecov command. bedGraphToBigWig (UCSC-tools) was used to generate bigWIG files for the UCSC genome browser.

ATAC-seq peak calling and quantification:

Two independent peak calling algorithms were employed to ensure robust, reproducible peak calls. First, tag directories were created using HOMER makeTagDirectory for each replicate, and peaks were called using default parameters for findPeaks with -style factor. MACS2 was also called using default parameters on each replicate. The summit files output by MACS2 were converted to bed format and each summit extended bidirectionally to achieve a total length of 300 bp. As the ATAC-seq peak calls would ultimately be used to narrow down potential regulatory elements for screening of a limited subset, we applied the overly stringent requirement that a peak be called by both approaches in a given replicate for its inclusion in the final peak list for that sample. Peaks identified in any sample in this way were aggregated to produce a final superset of 323,369 regulatory elements called as accessible in at least one cell type. The featurecounts package was used to obtain ATAC-seq read counts for each of these accessible putative GREs.

Identification of SST-enriched GREs

We used genomic coordinates of a superset of 323,369 genomic regions identified as a union of ATAC-Seq peaks across various cell types in the mouse cortex (manuscript in preparation) as a list of reference coordinates over which to quantify the ATAC-Seq signal from SST+, VIP+ and PV+ cells. A matrix was constructed representing the mean ATAC-Seq signal in SST+, VIP+ and PV+ cells for each of the 323,369 GREs and normalized such that the total ATAC-Seq signal from each cell population was scaled to 10^7. Fold-enrichment was calculated for each region/GRE as [(Signal in cell type A)+1] / [mean(signal in cell types B and C)+1]. GREs were subsequently ranked based on fold-enrichment score.

Identification of conserved GREs

To identify GREs whose sequence is highly conserved across mammals, we first needed to identify an appropriate conservation score to use as a threshold for high conservation. We reasoned that by analyzing the conservation of DNA sequences of the same length, but an arbitrary distance of 100,000 bases away from each identified GRE, we would generate a set of DNA sequences whose conservation can be used to determine this threshold.

To this end, conservation scores for GREs and corresponding GRE-distal sequences were calculated using the bigWigAverageOverBed command to determine the average PhyloP score of each sequence based on mm10.60way.phyloP60wayPlacental.bw PhyloP scores (http://hgdownload.cse.ucsc.edu/goldenpath/mm10/phyloP60way/)(*3*) . After plotting the conservation score (phyloP, 60 placental mammals) of 323,369 GRE-distal sequences, we determined the conservation score of the 95[th] percentile of this distribution (PhyloP score = 0.5) and chose it as a minimal conservation score needed to classify any GRE as conserved.

Viral barcode design

Viral barcode sequences were chosen to be at least 3 insertions, deletions, or substitutions apart from each other to minimize the effects of sequencing errors on the correct identification of each barcode. The R library "DNAbarcodes" and following functions were used:

initialPool = create.dnabarcodes(10, dist=3, heuristic="ashlock");

finalPool = create.dnabarcodes(10, pool = initialPool, metric="seqlev");

The result was a list of 1164 10-base barcodes that fit our initial criteria.

Amplification of GREs and barcoding

Genomic PCR:

PCR primers were designed using primer3 2.3.7.(*4*) such that a 150-400 bp flanking sequence was added to each side of the GRE. The forward primers contained a 5' overhang sequence for downstream in-Fusion (Clonetech) cloning into the AAV vector (5'-GCCGCACGCGTTTAAT). The reverse primers contained a 5' overhang sequence containing the recognition sites for AsiSI and SalI restriction enzymes(5'-GCGATCGCTTGTCGAC). Hot Start High-Fidelity Q5 polymerase (NEB) was used according to manufacturer's protocol with mouse genomic DNA as template.

Barcoding PCR:

The unpurified PCR products from the genomic PCR were used as templates for the barcoding PCR. A forward primer containing the sequence for downstream in-Fusion (Clonetech) cloning into the AAV vector (5'-CTGCGGCCGCACGCGTTTA) was used in all reactions. Reverse primers were constructed featuring (in the 5' → 3'direction): 1) a sequence for downstream in-Fusion (Clonetech) cloning into the AAV vector (5'-GCCGCTATCACAGATCTCTCGA), 2) a unique 10-base barcode sequence, and 3) sequence complementary with the AsiSI and SalI restriction enzyme recognition sites that were introduced during the first PCR (5'-GCGATCGCTTGTCGAC). Three different reverse primers were used for each of the GREs amplified during the genomic PCR. Hot Start High-Fidelity Q5 polymerase (NEB) was used according to the manufacturer's protocol.

PESCA Library cloning

All PCR reactions were pooled and the amplicons purified using Agencourt AMPure XP. The pAAV-mDlx-GFP-Fishell-1 was a gift from Gordon Fishell (Addgene plasmid # 83900). The

plasmid was digested with PacI and XhoI, leaving the ITRs and the polyA sequence. in-Fusion was used to shuttle the pool of GRE PCR products into the vector. Following transformation into High Efficiency NEB 5-alpha Competent E.coli and recovery, SalI and AsiSI were used to linearize the AAV vector containing the GREs. The expression cassette containing the human HBB promoter and intron followed by GFP and WPRE was isolated by PCR amplification from pAAV-mDlx-GFP-Fishell-1. The expression cassette was ligated with the linearized GRE-library-containing vector using T4 ligase and transformed into High Efficiency NEB 5-alpha Competent E.coli to yield the final library. 50 colonies were Sanger sequenced to determine the correct pairing between GRE and barcode and the correct arrangement of the AAV vector.

AAV preparation

The pooled PESCA library or individual AAV constructs (100 µg) were packed into AAV9 at the Boston Children's Hospital Viral Core. The titers (2-50 x 10^{13} genome copies/mL) were determined by qPCR. Next generation sequencing using the NextSeq 500 platform was used to determine the complexity of the pooled PESCA library (Fig. 2a).

V1 cortex injections

Animals were anesthetized with isofluorane (1–3% in air) and placed on a stereotactic instrument (Kopf) with a 37°C heated pad. The PESCA library (AAV9, 1.9 x 10^{13} genome copies/mL) was stereotactically injected in V1 (800 nL per site at 25 nL/min) using a sharp glass pipette (25-45 µm diameter) that was left in place for 5 min prior to and 10 min following injection to minimize

backflow. Two injections were performed per animal at coordinates 3.0 and 3.7 mm posterior, 2.5 mm lateral relative to bregma, and 0.6 mm ventral relative to the brain surface. Individual rAAV-GRE constructs were stereotactically injected at a titer of 1 x 10[11] genome copies/mL. (250 nL per site at 25 nL/min). All injections were performed at two depths (0.4 and 0.7 mm ventral relative to the brain surface) to achieve broader infection across cortical layers. The injection coordinates relative to bregma were 3.0 or 3.7 mm posterior, 2.5 or -2.5 mm lateral.

Nuclear isolation

Single-nuclei suspensions were generated as described previously(1) , with minor modifications. V1 was dissected and placed into a Dounce with homogenization buffer (0.25 M sucrose, 25 mM KCl, 5 mM MgCl$_2$, 20 mM Tricine-KOH, pH 7.8, 1 mM DTT, 0.15 mM spermine, 0.5 mM spermidine, protease inhibitors). The sample was homogenized using a tight pestle with 10 stokes. IGEPAL solution (5%, Sigma) was added to a final concentration of 0.32%, and 5 additional strokes were performed. The homogenate was filtered through a 40-μm filter, and OptiPrep (Sigma) added to a final concentration of 25% iodixanol. The sample was layered onto an iodixanol gradient and centrifuged at 10,000g for 18 minutes as previously described[1,2] . Nuclei were collected between the 30% and 40% iodixanol layers and diluted to 80,000-100,000 nuclei/mL for encapsulation. All buffers contained 0.15% RNasin Plus RNase Inhibitor (Promega) and 0.04% BSA.

snRNA-Seq library preparation and sequencing

Single nuclei were captured and barcoded whole-transcriptome libraries prepared using the inDrops platform as previously described(5, 6), collecting five libraries of approximately 3,000 nuclei from each animal. Briefly, single nuclei along with single primer-carrying hydrogels were captured into droplets using a microfluidic platform. Each hydrogel carried oligodT primers with a unique cell-barcode. Nuclei were lysed and the cell-barcode containing primers released from the hydrogel, initiating reverse transcription and barcoding of all cDNA in each droplet. Next, the emulsions were broken and cDNA across ~3000 nuclei pooled into the same library. The cDNA was amplified by second strand synthesis and *in vitro* transcription, generating an amplified RNA intermediate which was fragmented and reverse transcribed into an amplified cDNA library.

For enrichment of virally-derived transcripts, a fraction (3 μL) of the amplified RNA intermediate was reverse transcribed with random hexamers without prior fragmentation. PCR was next used to amplify virally derived transcripts. The forward primer was designed to introduce the R1 sequence and anneal to a sequence uniquely present 5' of the viral-barcode sequence present in the viral transcripts (5'- GCATCGATACCGAGCGC). The reverse primer was designed to anneal to a sequence present 5' of the cell-barcode (5'- GGGTGTCGGGTGCAG). The result of the PCR is preferential amplification of the viral-derived transcripts, while simultaneously retaining the cell-barcode sequence necessary to assign each transcript to a particular cell/nucleus. Following PCR amplification (18 cycles, Hot Start High-Fidelity Q5 polymerase) all the libraries were indexed, pooled, and sequenced on a Nextseq 500 benchtop DNA sequencer (Illumina).

inDrop sample mapping and viral barcode deconvolution by cell:

The published inDrops mapping pipeline (github.com/indrops/indrops) was used to assign reads to cells. To map viral sequences, a custom annotated transcriptome was generated using the indrops pipeline build_index command supplied with the following newly generated reference files: a custom genome with one additional contig comprising a shared 5' sequence (gcatcgataccgagcgcgcgatcgc), the given 10 bp barcode, and a shared 3' sequence (tcgagagatctgtgatagcggc) was appended to the GRCm38.dna_sm.primary_assembly.fa genome file for each cloned GRE. These sequences were also appended GRCm38.88.gtf gene annotation file, with all sequences assigned the same gene_id and gene_name, but unique transcript_id, transcript_name, and protein_id. After inDrops pipeline mapping and cell deconvolution, the pysam package was used to extract the 'XB' and 'XU' tags, which contain cell barcode and UMI sequences, respectively, from every read that mapped uniquely to any one of the custom viral contigs (i.e. requiring the read map to the 10 bp barcode with at most 1 mismatch) in the inDrops pipeline-output bam files. These barcode-UMI combinations were condensed to generate a final cell x GRE barcode UMI counts table for each sample.

Embedding and identification of cell types

Data from all nuclei (two animals, 5 libraries of ~3,000 nuclei per animal) were analyzed simultaneously. Viral-derived sequences were removed for the purposes of embedding clustering and cell type identification. The initial dataset contained 32,335 nuclei, with more than 200 unique non-viral transcripts (UMIs) assigned to each nucleus. The R software package Seurat(*7*, *8*) was used to cluster cells. First, the data were log-normalized and scaled to 10,000 transcripts per cell. Variable genes were identified using the FindVariableGenes() function. The following

parameters were used to set the minimum and maximum average expression and the minimum dispersion: x.low.cutoff=0.0125, x.high.cutoff=3, y.cutoff=0.5. Next, the data was scaled using the ScaleData() function, and principle component analysis (PCA) was carried out. The FindClusters() function using the top 30 principal components (PCs) and a resolution of 1.5 was used to determine the initial 29 clusters. Based on the expression of known marker genes we merged clusters that represented the same cell type. Our final list of cell types was: Excitatory neurons, PV Interneurons, SST Interneurons, VIP interneurons, NPY Interneurons, Astrocytes, Vascular-associated cells, Microglia, Oligodendrocytes, and Oligodendrocyte precursor cells.

Enrichment calculation

Viral vector expression for each of the 861 barcodes across the ten cell types was calculated by averaging the expression of barcoded transcripts across all the individual nuclei that were assigned to that cell type. The relative fold-enrichment in expression toward *Sst+* cells was computed as the ratio of the mean expression in *Sst+* cells and the mean expression in *Sst-* cells: (mean(*Sst+* cells)+0.01)/ (mean(*Sst-* cells)+0.01).

Viral GRE expression for each of the 287 barcodes was calculated at the single-nucleus level as a sum of the expression of the three barcodes that were paired with that GRE. Average GRE-driven expression across the ten cell types was calculated by averaging the expression of the GRE transcripts across all the individual nuclei that were assigned to that cell type. The relative fold-enrichment in GRE expression toward *Sst+* cells was determined as the ratio of the mean expression in *Sst+* cells and the mean expression in *Sst-* cells: (mean(*Sst+* cells)+0.01)/ (mean(*Sst-* cells)+0.01).

Differential gene expression

To identify which of the GRE-driven transcripts were statistically enriched in *Sst+* vs. *Sst-* cells, we carried out differential gene expression analysis using the R package Monocle2(*9*) . The data were modeled and normalized using a negative binomial distribution, consistent with snRNA-seq experiments. The functions estimateSizeFactors(), estimateDispersions() and differentialGeneTest() were used to identify which of the GRE-derived transcripts were statistically enriched in *Sst+* cells. GREs whose false discovery rate (FDR) was less than 0.01 were considered enriched.

Fluorescence microscopy

Sample preparation:

Mice were sacrificed and perfused with 4% PFA followed by PBS. The brain was dissected out of the skull and post-fixed with 4% PFA for 1-3 days at 4°C. The brain was mounted on the vibratome (Leica VT1000S) and coronally sectioned into 100 μm slices. Sections containing V1 were arrayed on glass slides and mounted using DAPI Fluoromount-G (Southern Biotech).

Sample imaging:

Sections containing V1 were imaged on a Leica SPE confocal microscope using an ACS APO 10x/0.30 CS objective (Harvard NeuroDiscovery Center). Tiled V1 cortical areas of ~1.2 mm by ~0.5 mm were imaged at a single optical section to avoid counting the same cell across multiple optical sections. Channels were imaged sequentially to avoid any optical crosstalk.

Immunostaining

To identify parvalbumin (PV)+ cells, coronal sections were washed three times with PBS containing 0.3% TritonX-100 (PBST) and blocked for 1 h at room temperature with PBST containing 5% donkey serum. Section were incubated overnight at 4°C with mouse anti-PVALB antibody 1:2000 (Milipore), washed again three times with PBST, and incubated for 1 h at room temperature with 1:500 donkey anti-mouse 647 secondary antibody (Life Technologies). After washing in PBST and PBS, samples were mounted onto glass slides using DAPI Fluoromount-G.

Quantification of the percentage of GFP+ cells that were SST+, VIP+, and PV+

Across all images, coordinates were registered for each GFP+ cell that could be visually discerned. An automated ImageJ script was developed to quantify the intensity of each acquired channel for a given GFP+ cell. We created a circular mask (radius = 5.7 μm) at each coordinate representing a GFP positive cell, background subtracted (rolling ball, radius = 72 μm) each channel, and quantified the mean signal of the masked area. To identify the threshold intensity used to classify each GFP+ cell as either SST+, VIP+ or PV+, we first determined the background signal in the channel representing SST, VIP or PV by selecting multiple points throughout the area visually identified as background. These background points were masked as small circular areas (radius = 5.7 μm), over which the mean background signal was quantified. The highest mean background signal for SST, VIP and PV was conservatively chosen as the threshold for classifying GFP+ cells as SST+, VIP+ or PV+, respectively.

Quantification of the distribution of cells as a function of distance from pia

A semiautomated ImageJ algorithm was developed to trace the pia in each image, generate a Euclidean Distance Map (EDM), and calculate the distance from the pia to each GFP+ cell.

Quantification of the percentage of SST+ cells that were GFP+

An automated algorithm was developed to identify SST+ cells after appropriate background subtraction, image thresholding, masking and filtering for all objects of appropriate size and circularity. The number of SST+ objects (cells) was then counted within a minimal polygonal area that encompassed all GFP+ cells in that image. The ratio of the number of GFP+ cells and SST+ cells within the area of infection (here identified as area with discernable GFP+ cells) was calculated.

Slice Preparation

Acute, coronal brain slices containing visual cortex of 250 – 300 μm thickness were prepared using a sapphire blade (Delaware Diamond Knives, Wilmington, DE) and a VT1000S vibratome (Leica, Deerfield, IL). Mice were anesthetized though inhalation of isoflurane, then decapitated. The head was immediately immersed in an ice-cold solution containing (in mM): 130 K-gluconate, 15 KCl, 0.05 EGTA, 20 HEPES, and 25 glucose (pH 7.4 with NaOH; Sigma). The brains were quickly dissected and cut in the same ice-cold, gluconate based solution while oxygenated with 95% O_2/5% CO_2. Slices then recovered at 32ºC for 20-30 minutes in oxygenated artificial cerebrospinal fluid (ACSF) in mM: 125 NaCl, 26 NaHCO3, 1.25 NaH2PO4, 2.5 KCl, 1.0 MgCl2, 2.0 CaCl2, and 25 glucose (Sigma), adjusted to 310-312 mOsm with water.

Electrophysiological Recordings

Using an Olympus BX51WI microscope equipped with a 60x water immersion objective, we used fluorescence illumination to identify rAAV-GRE44-GFP$^+$ (red and green) and rAAV-GRE44-GFP$^-$ (only red) SST neurons in the area of injection/ AAV infection (Fig. 4a-d). rAAV-GRE44-GFP$^-$ neurons were recorded if they were in the same field of view as rAAV-GRE44-GFP$^+$ neurons under 60x. For rAAV-GRE12-Gq-DREADD-tdTomato experiments (Fig. 4e-j), tdTomato$^+$ cells and morphologically identified pyramidal neurons in the same field of view under 60x were recorded. Whole-cell current clamp recordings of these neurons in coronal visual cortex slices of P50 to P80 wild-type mice were performed using borosilicate glass pipettes (3-6 MOhms, Sutter Instrument, Novato, CA) filled with an internal solution (in mM): 116 KMeSO$_3$, 6 KCl, 2 NaCl, 0.5 EGTA, 20 HEPES, 4 MgATP, 0.3 NaGTP, 10 NaPO$_4$ creatine (pH 7.25 with KOH; Sigma). Neurobiotin (1.5%) was occasionally included in the internal solution to allow for post-hoc morphological reconstruction of recorded cells. All experiments were performed at room temperature in oxygenated ACSF. Series resistance was compensated by at least 60% in a voltage-clamp configuration before switching to current-clamp ("I Clamp Normal"). After break-in, a systematic series of 1 second current injections ranging from -100 pA to 500 pA were applied to each cell using the User List function in the "Edit Waveform" tab of pClamp. After such baseline firing rates were calculated, CNO (2 μM, Sigma) was bath applied. An average of at least three trials for each current injection was calculated before and during CNO application.

Data Acquisition and Analysis

For electrophysiology, data acquisition of current-clamp experiments was performed using Clampex10.2, an Axopatch 200B amplifier, filtered at 2 kHz and digitized at 20 kHz with a DigiData 1440 data acquisition board (Molecular Devices, Sunnyvale, CA). Analysis of electrophysiological parameters was done using Clampfit (Molecular Devices, Sunnyvale, CA), Prism 7 (GraphPad Software, La Jolla, CA), Excel (Microsoft, Redmond, WA), and custom software written and generously shared by Dr. Bruce Bean in Igor Pro version 6.1.2.1 (WaveMetrics, Lake Oswego, OR). Membrane potentials in this study were not corrected for the liquid junction potential and are thus positively biased by 8 mV. For analysis of action potential waveform in Fig. 4a-d and Supplementary Table 1, the first action potential that appeared during a current injection equivalent to the rheobase was analyzed, as well as the first action potential of the subsequent two current injections. For example, if the rheobase were 20 pA, then all the parameters defined in the next section were also analyzed for the first action potential elicited with 20, 25, and 30 pA of injected current, and averaged.

Definition of Electrophysiological Parameters

AP Height (in millivolts): the difference between the peak of the action potential and the most negative voltage during the afterhyperpolarization immediately following the spike.

AP Peak (in millivolts): the most depolarized (positive) potential of the spike.

AP Trough (in millivolts): the most negative voltage reached during the afterhyperpolarization immediately following the spike.

F_{max} initial (in Hertz): the average of the reciprocal of the first three interstimulus intervals, measured at the maximal current step injected before spike inactivation.

F_{max} steady-state (in Hertz): the average of the reciprocal of the last three interstimulus intervals, measured at the maximal current step injected before spike inactivation.

Rate of rise (in volts per second): maximal voltage slope (*dV/dt*) during the upstroke (rising phase) of the action potential.

Rheobase (in picoamperes): the minimal 1000 ms current step (in increments of 5 pA) needed to elicit an action potential.

R_{in} (in megaohms, MΩ): input resistance, determined by using Ohm's law to measure the change in voltage in response to a -50 pA, 1000ms hyperpolarizing current at rest.

Spike adaptation ratio: the ratio of F_{max} steady-state to F_{max} initial

Spike width (in milliseconds, used interchangeably with spike half-width): the width at half-maximal spike height as defined above.

τ_m (in milliseconds): membrane time constant, determined by fitting a monoexponential curve to the voltage chance in response to a -50pA, 1000ms hyperpolarizing current at rest.

Threshold (in millivolts): the membrane potential at which dV/dt = 5 V/s.

V_{rest} (in millivolts): resting membrane potential a few minutes after breaking in without any current injection.

Supplementary figures

Supplementary figure 1

Identification of conserved GREs. Left: For each of the 323,369 genomic regions that were identified by ATAC-Seq as GREs in either SST[+], VIP[+] or PV[+] cells, a region of the same size was chosen exactly 100,000 bases away from the GRE. The mean sequence conservation score (phyloP, 60 placental mammals) for each of these GRE-distal regions was calculated and plotted. A vertical line at the conservation score of 0.5 indicates the 95th percentile of that distribution and was chosen as a minimal conservation score needed to consider a GRE sequence as conserved. Right: The mean sequence conservation score (phyloP, 60 placental mammals) for each of the 323,369 GREs was calculated and plotted. A vertical line indicates the minimal conservation score of 0.5. 36,215 GREs (11%) had a mean conservation greater than 0.5 and were deemed conserved.

| Genomic PCR | Barcoding PCR | Cloning GRE pool into pAAV | Insertion of expression cassette |

Supplementary figure 2

(**a**) Schema for PESCA library construction. PCR is used to amplify GREs from the genomic DNA and to introduce appropriate restriction enzyme sites and, subsequently, a 10 bp barcode sequence. Each GRE is amplified three times using three different barcode sequences. The amplified GREs are pooled and cloned into an AAV vector. Restriction enzyme sites between the GRE and the barcode are used to insert an expression cassette consisting of a minimal promoter, intron, GFP and WPRE sequences. See methods for details.

Supplementary figure 3

(**a**) Dot plot of the number of unique molecular identifiers (UMIs) and the number of genes for each nucleus that was analyzed.

(**b**) Plot showing the density distribution of number of UMIs and genes per nucleus.

(**c**) Distribution of the number of unique barcodes and unique GREs detected per nucleus, displayed as Log10(Count+1).

(**d**) Quantification of the fraction of cells within each defined cell type in which we detected barcoded viral transcripts. Each dot represents one animal. Box plot represents mean ± standard error of the mean (s.e.m) for Exc (86.6 ± 4.6%), Int_Pv (91.6 ± 3.1%), Int_Sst (93.9 ± 2.1%), Int_Vip (90.4 ± 3.5%), Int_Npy (87.1 ± 3.8%), Astro (82.0 ± 5.0%), Vasc (76.9 ± 7.1%), Microglia (78.6 ± 7.1%), Olig (75.4 ± 6.6%), and OPCs (77.4 ± 8.3%).

(**e**) t-SNE plot of 32,335 nuclei from V1 cortex of two injected animals**.** Color gradient denotes number of unique viral transcripts per nucleus displayed as Log10(Count+1).

(**f**) Dot plot of the number of viral genomes in the AAV library and the number of infected cells recovered after snRNA-Seq. Each dot represents one barcode (n = 861). Line of linear fit with 95% confidence intervals (shaded). Pearson correlation r = 0.9, $p < 2.2 \times 10^{-16}$.

Supplementary figure 4

t-SNE plots of 32,335 nuclei from V1 cortex of two analyzed animals. Color gradient denotes number of unique transcripts per nucleus of the indicated cellular marker gene.

Supplementary figure 5

Dot plot of pairwise comparison between SST fold-enrichment values across three sets of barcodes. The values on each axis represent the SST fold-enrichment calculated for each GRE based on one of the three barcodes paired with that GRE. Blue line indicates linear fit with 95% confidence intervals (shaded). Correlation and p-values are indicated for each plot. Color gradient indicates the average enrichment between all three barcodes.

Supplementary figure 6

(**a**) t-SNE plot of 32,335 nuclei from V1 cortex of two analyzed animals showing the mean viral expression across all GREs. Plot is pseudocolored based on the mean expression in each cell type.

(**b**) Volcano plots for identified SST-enriched GREs (Fold-enrichment > 7 and FDR < 0.01). Colored dots represent the five SST-enriched GREs that were considered hits.

Supplementary Figure 7

Fluorescent images from adult Sst-Cre; Ai14 mouse visual cortex twelve days following

injection with rAAV-GRE-GFP as indicated.

Supplementary Figure 8

Quantification of the number of GFP[+] SST[+] cells normalized for area of infection. Each dot represents one animal. Box plot represents mean ± standard error of the mean (s.e.m). Values are 73.0 ± 17.2, 146.9 ± 19.7, 144.8 ± 38.6 and 125.6± 26.4 cells/mm^2 for AAV-[ΔGRE, GRE12, GRE22, GRE44]-GFP, respectively.

Supplementary Figure 9

Fluorescent images from adult Vip-Cre; Ai14 mouse visual cortex immunostained for PVALB

twelve days following injection with rAAV-GRE-GFP as indicated.

	GRE44- (n=16)	GRE44+ (n=16)	p value (2-tailed unpaired t-test)
Supplementary Table 1 Electrophysiological Parameters of GRE44- and GRE44+ SST Neurons in Visual Cortex			
V_{rest} (mV)	-62.4 ± 1.51	-60.6 ± 1.63	0.41
R_{in} (MΩ)	304 ± 54.8	391 ± 47.3	0.24
τ_m (ms)	14.2 ± 2.35	22.8 ± 4.32	0.094
Threshold (mV)	-45.6 ± 1.24	-48.1 ± 1.26	0.17
AP Peak (mV)	13 ± 3.38	11.6 ± 3.19	0.76
AP Trough (mV)	-63.6 ± 1.36	-63.7 ± 1.43	0.96
AP Height (mV)	76.5 ± 4.46	75.2 ± 4.06	0.83
Rate of Rise (V/s)	**122 ± 11.7**	**85 ± 7.34**	**0.013***
Rheobase (pA)	**43.5 ± 10.4**	**20.3 ± 4.64**	**0.044***
Spike Half-Width (ms)	**1.25 ± 0.0819**	**2.52 ± 0.307**	**0.0004***
F_{max}, steady-state (Hz)	**83.4 ± 9.8**	**34.5 ± 4.32**	**0.0002***
F_{max}, initial (Hz)	**111 ± 8.35**	**67.3 ± 7.48**	**0.0007***
Spike adaptation ratio	0.763 ± 0.0839	0.561 ± 0.0699	0.08

Values are shown as mean ± SEM.

Supplementary Table 1. Electrophysiological Parameters of GRE44- and GRE44+ SST

Neurons in Visual Cortex

References

1. A. Mo *et al.*, Epigenomic Signatures of Neuronal Diversity in the Mammalian Brain. *Neuron*. **86**, 1369–1384 (2015).

2. H. Stroud *et al.*, Early-Life Gene Expression in Neurons Modulates Lasting Epigenetic States. *Cell*. **171**, 1151-1164.e16 (2017).

3. K. S. Pollard, M. J. Hubisz, K. R. Rosenbloom, A. Siepel, Detection of nonneutral substitution rates on mammalian phylogenies. *Genome Res.* **20**, 110–121 (2010).

4. A. Untergasser *et al.*, Primer3--new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115 (2012).

5. A. M. Klein *et al.*, Droplet Barcoding for Single-Cell Transcriptomics Applied to Embryonic Stem Cells. *Cell*. **161**, 1187–1201 (2015).

6. R. Zilionis *et al.*, Single-cell barcoding and sequencing using droplet microfluidics. *Nat. Protoc.* **12**, 44–73 (2017).

7. R. Satija, J. A. Farrell, D. Gennert, A. F. Schier, A. Regev, Spatial reconstruction of single-cell gene expression data. *Nat. Biotechnol.* **33**, 495–502 (2015).

8. A. Butler, P. Hoffman, P. Smibert, E. Papalexi, R. Satija, Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411–420 (2018).

9. C. Trapnell *et al.*, The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32**, 381–386 (2014).