The Cdc14 phosphatase controls resolution of recombination intermediates and crossover formation during meiosis

Paula Alonso-Ramos,[1,*] David Álvarez-Melo,[1,*] Katerina Strouhalova,[1] Carolina Pascual-Silva,[1] George B. Garside,[2] Meret Arter,[3] Teresa Bermejo,[1] Rokas Grigaitis,[3] Rahel Wettstein,[3] Marta Fernández-Díaz,[1] Joao Matos,[3] Marco Geymonat,[4] Pedro A. San-Segundo,[5] Jesús A. Carballo[1,6]

[1] Department of Cellular and Molecular Biology, Centro de Investigaciones Biológicas Margarita Salas, CSIC, Ramiro de Maeztu 9, 28040, Madrid, Spain

[2] Leibniz Institute for Age Research/Fritz Lipmann Institute (FLI) Beutenbergstr. 11; D-07745 Jena; Germany

[3] Max Perutz Labs, University of Vienna, Dr. Bohr-Gasse 9, 1030 Vienna, Austria

[4] Department of Genetics, University of Cambridge, Downing Street, Cambridge, CB2 3EH, United Kingdom

[5] Instituto de Biología Funcional y Genómica. Consejo Superior de Investigaciones Científicas and University of Salamanca. 37007-Salamanca, Spain

* These authors contributed equally to this work

[6]Corresponding author: j.carballo@ csic.es

Abstract

Meiotic defects derived from incorrect DNA repair during gametogenesis can lead to mutations, aneuploidies and infertility. Coordinated resolution of meiotic recombination intermediates is required for crossover formation, ultimately necessary for accurate completion of both rounds of chromosome segregation. Numerous master kinases orchestrate the correct assembly and activity of the repair machinery. Although much less is known, reversal of phosphorylation events in meiosis must also be key to coordinate the timing and functionality of repair enzymes. Cdc14 is an evolutionarily conserved phosphatase required for the dephosphorylation of multiple CDK1 targets. Mutations that inactivate this phosphatase lead to meiotic failure, but until now it was unknown if Cdc14 plays a direct role in meiotic recombination. Here, we show that elimination of Cdc14 leads to severe defects in the processing and resolution of recombination intermediates, causing a drastic depletion of crossovers when other repair pathways are compromised. We also show that Cdc14 is required for correct activity and localization of the Holliday Junction resolvase Yen1/GEN1. We reveal that Cdc14 regulates Yen1 activity from meiosis I onwards, and this function is essential for crossover resolution in the absence of other repair pathways. We also demonstrate that Cdc14 and Yen1 are required to safeguard sister

36 chromatid segregation during the second meiotic division, a late action that is independent of the earlier

37 role in crossover formation. Thus, this work uncovers previously undescribed functions of Cdc14 in the

38 regulation of meiotic recombination.

Introduction

Meiotic recombination is initiated by the conserved Spo11 transesterase, which introduces numerous

DNA Double-Strand Breaks (DSBs) into the genome (Keeney et al. 1997). Association of single-strand

DNA binding proteins, including RPA, Rad51 and the meiosis specific recombinase, Dmc1, with the

resected DSB ends promotes strand invasion into the intact homologous non-sister chromatid template,

which culminates in the formation of a displacement loop (D-loop). Nascent D-loops can be processed

through the Synthesis-Dependent Strand Annealing (SDSA) repair pathway to generate non-crossover

(NCO) repair products (Bishop 2006). Alternatively, stabilization of the D-loop followed by second-

end capture gives rise to even more stable structures, such as double Holliday Junction (dHJ)

intermediates. In budding yeast meiosis, dHJs are most frequently resolved into crossovers (COs)

through the combined action of Mlh1-Mlh3 (MutLγ) and Exo1 (Zakharyevich et al. 2010; Zakharyevich

et al. 2012; Sanchez et al. 2020). A second class of COs arises from the resolution of recombination

intermediates via the Structure-Selective Endonucleases (SSEs) Mus81/Mms4, Slx1/Slx4 and Yen1 (de

los Santos et al. 2001; Jessop and Lichten 2008; Oh et al. 2008; Matos et al. 2011; Arter et al. 2018).

Finally, Sgs1, the ortholog of the Bloom's syndrome helicase (BLM), can also promote dissolution of

JMs via the Sgs1-Top3-Rmi1 (STR) complex, which would exclusively generate NCOs products. In

addition, multiple rounds of strand exchange over a stabilized D-loop, can give rise to three and four

interconnected duplexes, also known as multichromatid joint molecules (mc-JMs) (Oh et al. 2007).

Further, both DSB-ends might also participate in multi-invasion events on two or three chromatids,

creating heavily branched DNA structures (Piazza et al. 2017). These complex DNA species must be

efficiently processed to prevent the risk of becoming potential hazards for the genome integrity due to

their capacity to form aberrant crossovers, or other forms of toxic repair products. Correct processing

of all types of JMs requires the orchestrated action of a set of helicases, topoisomerases and

endonucleases, capable of dislodging multi-branched DNA assemblies. The BLM/Sgs1, in addition to

be involved in SDSA repair, plays a prominent role in eliminating aberrant "off-pathway" JMs,

including mc-JMs (Jessop and Lichten 2008; Oh et al. 2008; De Muyt et al. 2012; Copsey et al. 2013;

Xaver et al. 2013; Grigaitis et al. 2018).

In budding yeast, it is not fully clear which enzymatic complexes take over the function of

removing any unprocessed branched DNA intermediates outside prophase I, due to the existence of

numerous SSEs capable of processing such substrates (Machin 2020; San-Segundo and Clemente-

Blanco 2020). Upon initiation of *NDT80*-dependent transcription, and activation of the budding yeast

polo-like kinase Cdc5, cells abandon the pachytene stage concomitantly with a surge of enzymatic

activity that culminates in the resolution of ZMM-dependent and ZMM-independent dHJs to give rise

to COs and some NCO products (Clyne et al. 2003; Sourirajan and Lichten 2008). Cdc5

phosphorylation is required to stimulate the function of a set of SSEs, most notoriously Mus81-Mms4

(Matos et al. 2011; Wild et al. 2019). The Mus81-Mms4 complex acts on branched DNA substrates that

have not been resolved by the canonical ZMM-dependent Class-I CO pathway (De Muyt et al. 2012;

Zakharyevich et al. 2012). On the other hand, Yen1[GEN1], a conserved member of the Rad2/XPG family

of SSEs, appears to act later in meiosis, despite the fact that Yen1 is considered a prototypical HJ

resolvase (Ip et al. 2008). In meiotic cells, this discrepancy is easily explained since the activity of Yen1

is negatively regulated by CDK1-dependent phosphorylation, preventing the activation of the nuclease

during prophase I (Matos et al. 2011). Elimination of CDK-phosphorylation sites in Yen1, turns on its

enzymatic activity prematurely and promotes its nuclear localization. Therefore, counteracting the

phosphorylation of Yen1 generates the fully active form of the nuclease (Matos et al. 2011; Arter et al.

2018). In mitotic cells, dephosphorylation of Yen1 is carried out by the Cdc14 phosphatase (Blanco et

al. 2014; Eissler et al. 2014). Activation of Yen1 during anaphase by Cdc14 allows the resolution of

persistent repair intermediates that otherwise would impose a physical impediment to chromosome

segregation. It is currently unknown whether Cdc14 also activates Yen1 in meiosis (Matos et al. 2011;

Zakharyevich et al. 2012; Grigaitis et al. 2020). Furthermore, it is possible that Cdc14 dephosphorylates

other substrates important for the timely resolution of different types of JMs (Grigaitis et al. 2020).

Cdc14 is a well-conserved dual-specificity phosphatase that has been defined as a key

component in the reversal of CDK phosphorylation during exit from mitosis (Visintin et al. 1998). In

budding yeast, Cdc14 activity is essential and cells lacking this phosphatase remain arrested at the end

of mitosis (Culotti and Hartwell 1971; Hartwell et al. 1973). Additionally, Cdc14 regulates

transcriptional silencing at the rDNA and other repeated sequences (Machin et al. 2006; Clemente-

Blanco et al. 2009; Clemente-Blanco et al. 2011). Several pathways regulate Cdc14 localization and

activity. The RENT complex retains Cdc14 at the nucleolus until anaphase through its interaction with

the Cif1/Net1 protein (Shou et al. 1999; Visintin et al. 1999). Cdc14 is released from the nucleolus by

the sequential action of two pathways. The FEAR (Cdc Fourteen Early Anaphase Release) pathway

promotes the early release of Cdc14 through the phosphorylation of Cif1/Net1 and Cdc14 (Yoshida and

Toh-e 2002; Visintin et al. 2003; Azzam et al. 2004; Queralt et al. 2006; Rahal and Amon 2008;

Rodriguez-Rodriguez et al. 2016; Zhou et al. 2021). Later in anaphase, the Mitotic Exit Network (MEN)

keeps Cdc14 in its released state, allowing the dephosphorylation of additional substrates, important for

the full termination of mitosis and cytokinesis (Stegmeier et al. 2002; Visintin et al. 2008).

In addition to curtailing CDK activity at the end of mitosis, Cdc14 plays critical roles promoting

chromosome segregation (Hartwell and Smith 1985). In *S. cerevisiae*, it is required for the correct

segregation of the ribosomal gene array (rDNA) and telomeres (D'Amours et al. 2004; Sullivan et al.

2004; Torres-Rosell et al. 2004). Lack of transcriptional silencing at these regions prevents the loading

of condensins at the rDNA and subtelomeric regions (D'Amours et al. 2004; Wang et al. 2004;

Clemente-Blanco et al. 2011), which leads to unresolved linkages, intertwined sister chromatids, and non-disjunction during mid-anaphase. Notably, a number of functions have been allocated to Cdc14 before its anaphase-triggered release. In particular, Cdc14 has been involved in completion of replication of late-replicating regions, such as the rDNA locus and other parts of the genome (Dulev et al. 2009). Additionally, anaphase-independent release of Cdc14 has been observed upon induction of DNA damage; generation of DSBs promotes the transitory release of Cdc14 from the nucleolus, targeting the SPB component Spc110 for dephosphorylation (Villoria et al. 2017).

Cdc14 is highly conserved and orthologs have been identified in other organisms, including fission yeast, nematodes, insects, and vertebrates (Mocciaro and Schiebel 2010). The human genome contains three Cdc14 paralogs, hCDC14A, hCDC14B and hCDC14C (Li et al. 1997; Rosso et al. 2008). Depletion of Cdc14A, which is required for mitotic CDK inactivation, leads to defects in centrosome duplication, mitosis and cytokinesis (Trautmann and McCollum 2002). In vertebrates, CDC14B exits the nucleolus in response to DNA damage (Bassermann et al. 2008; Mocciaro et al. 2010), a process conserved in fission yeast (Diaz-Cuervo and Bueno 2008), and more recently also observed in budding yeast (Villoria et al. 2017). Thus, in addition to playing important roles in the DNA Damage Response (DDR) (Bassermann et al. 2008) it appears that both Cdc14A and Cdc14B may be required for efficient repair of damaged DNA (Mocciaro et al. 2010).

Cdc14 is also critically required for the completion of meiosis (Sharon and Simchen 1990). In budding yeast, Cdc14 is released from the nucleolus in two waves; the first during anaphase I and the second in anaphase II (Buonomo et al. 2003; Marston et al. 2003; Yellman and Roeder 2015). Contrary to mitotic cells, Cdc14 release from the nucleolus requires the essential function of the FEAR pathway, but MEN appears to be dispensable (Kamieniecki et al. 2005; Pablo-Hernando et al. 2007; Attner and Amon 2012). Additionally, some components of the MEN pathway, such as Cdc15, have functionally differentiated to fulfil a role in spore morphogenesis (Kamieniecki et al. 2005; Pablo-Hernando et al. 2007). Curiously, inactivation of the FEAR pathway in meiosis allows exclusively one round of nuclear division to take place, culminating with the formation of asci carrying two diploid spores (dyads) (Kamieniecki et al. 2000; Zeng and Saunders 2000; Buonomo et al. 2003; Marston et al. 2003; Kamieniecki et al. 2005). Premature activation of FEAR blocks spindle assembly during meiosis, a process normally averted by PP2A^{Cdc55} (Queralt et al. 2006; Bizzari and Marston 2011; Kerr et al. 2011; Nolt et al. 2011). On the other hand, inactivation of Cdc14 function by means of FEAR mutants, or by employing conditional temperature-sensitive alleles of *CDC14*, impairs the second round of chromosome segregation (Buonomo et al. 2003; Marston et al. 2003). Furthermore, the absence of the phosphatase creates chromosome entanglements, a problem that can be reverted by deleting *SPO11* (Marston et al. 2003). Occasionally, in the absence of Cdc14, the second meiotic division takes place over non-tetrapolar meiosis II spindles and, subsequently, these cells carry out aberrant chromosome segregation (Bizzari and Marston 2011). These interesting observations prompted the identification of another critical function of Cdc14 at the meiosis I to meiosis II transition, which is the licensing of SPB

re-duplication/half-bridge separation by asymmetrical enrichment of Cdc14 on a single SPB during

anaphase I (Fox et al. 2017).

Here, we describe a previously undefined role of Cdc14 in budding yeast meiosis. We reveal

that Cdc14 is involved in ensuring correct meiotic recombination by gradually implementing Yen1

activity following Cdc5 activation. We show that, at anaphase I, Yen1 already shows a prominent

nuclear localization promoted by Cdc14. Yen1 reaches its maximum activity during meiosis II, and

Cdc14 is strictly required for such activation. A constitutively active form of Yen1, termed Yen1^{ON},

which is insensitive to CDK downregulation, compensates for Cdc14 deficiency, suggesting that Cdc14

normally reverses CDK phosphorylation in Yen1 during meiosis. Unexpectedly, Cdc14 also promotes

Yen1 activity in *CDC20*-depleted cells, allowing JM processing and CO formation when other repair

pathways are absent. Thus, Yen1 plays a more prominent role during meiotic recombination than

originally anticipated, which can be manifested even before meiosis I division is normally completed.

Notably, Yen1 complements other endonucleases whose activities are mainly constrained to the time-

window between the end of prophase I and before anaphase I. We propose that Yen1, controlled by

Cdc14, implements its resolvase function when other enzymatic activities begin to decline outside of

their optimum operative time-window by the end of prophase I, and that Yen1 covers a wider gap

extending its action until the end of the second meiotic division.

Results

Identification of a separation-of-function allele of *CDC14* defective in sporulation

Reversal of CDK1 phosphorylation is carried out in budding yeast by the conserved Cdc14

phosphatase. This essential protein can only be studied utilizing conditional *cdc14* alleles that fully or

partially impair its functionality. Thermosensitive mutants have been widely used in the past for mitotic

and meiotic studies. However, meiotic recombination is sensitive to variations of temperature (Borner

et al. 2004), thus we considered using other types of conditional alleles to further examine the role of

Cdc14 in meiosis. During the course of a different study, we generated the *cdc14-HA* allele carrying an

endogenous C-terminal tagged version of *CDC14* with three copies of the hemagglutinin epitope (HA).

The *cdc14-HA* strain was highly defective for sporulation, displaying only 10% of asci containing one

or two spores (Fig 1A), and lacking dityrosine autofluorescence, an indicator of spore wall maturation

(Briza et al. 1986) (Fig 1B & F and figure S2B). Unlike other temperature-sensitive alleles of *cdc14*

used in many studies (see introduction), mitotic growth was not affected in diploid cells carrying the

cdc14-HA allele incubated at different temperatures (Fig 1C). Thus, we conclude that *cdc14-HA* cells

cannot efficiently complete sporulation despite their apparently normal mitotic growth and may be a

useful tool to explore additional aspects of Cdc14 meiotic function.

Distinct sporulation-defective alleles of *CDC14* block meiotic progression at different stages

To determine the cause(s) of the sporulation defect of the *cdc14-HA* mutant we analyzed the progression of the meiotic program by studying the kinetics of meiotic DNA replication and nuclear divisions. During the mitotic cell cycle, released Cdc14 dephosphorylates replication factors, such as Sld2 and Dpb2 (Bloom and Cross 2007), and the lack of Cdc14 activity leads to problems during termination of DNA replication (Dulev et al. 2009). However, FACS analysis of DNA content revealed that S-phase progression occurred with nearly identical kinetics in both control and *cdc14-HA* strains (Fig 1D), suggesting that meiotic S-phase is not altered in *cdc14-HA*. Next, meiotic divisions were monitored in synchronous cultures of *CDC14* and *cdc14-HA*. We found that *cdc14-HA* cells displayed slightly slower and less efficient kinetics of nuclear divisions; nevertheless, most *cdc14-HA* cells exited the one nuclei stage, although with a ~60 min delay compared to wild-type cells (Fig 1E; Fig. S1A). Previous work has shown that most meiotic cells in *cdc14* temperature-sensitive mutants are blocked with two nuclei, due to the inability to assemble fully functional tetrapolar spindles at meiosis II (Schild and Byers 1980; Sharon and Simchen 1990; Marston et al. 2003; Bizzari and Marston 2011). However, we noticed that the *cdc14-HA* mutant progressed beyond the two-nuclei stage in a high proportion of cells (Fig S1B-C), distinguishing this *cdc14-HA* allele from temperature-sensitive mutants used in other studies.

Next, we generated an additional meiosis-specific allele of *CDC14* by replacing its endogenous promoter with the *CLB2* promoter, which is strongly repressed in meiotic cells (Lee and Amon 2003). In this case, a 3HA tag was located at the N-terminal region of the protein. Cells expressing P_{CLB2}-*3HA-CDC14* (hereafter, *cdc14-md,* for meiotic depletion) recapitulated the sporulation defect of *cdc14-HA* (Fig 3G). The most striking difference was that although the first meiotic nuclear division took place in *cdc14-md* as it occurs in *cdc14-HA,* most cells remained binucleated in *cdc14-md* even at late time points in meiosis (Fig S1A). Thus, the terminal meiotic phenotype of *cdc14-md* appears to be different from that of *cdc14-HA*.

Cdc14 protein levels are depleted during meiosis in *cdc14-HA* cells

We next used western blot to analyze Cdc14 protein levels during meiosis in *CDC14, cdc14-md* and *cdc14-HA* strains. The *cdc14-HA* mutant displayed reduced levels of the tagged protein, even at time zero, prior to meiotic induction, and they were further reduced after 6-8 hours into meiosis (Fig 2A and Fig S2A-B). Depletion of Cdc14 was more effective in the *cdc14-md* allele, while in *cdc14-HA* it occurred more gradually (Fig 2A). This is not related to delayed entry into the meiotic program, since all strains underwent meiotic S-phase with similar kinetics as determined by FACS analysis (Fig 2A). To directly ascertain if Cdc14 protein levels varied from mitotically dividing cells to meiotic cells, we also analyzed extracts from exponentially growing cultures compared to extracts prepared from cultures

immediately after meiotic induction (Fig 2B). The wild-type strain showed similar amounts of Cdc14
protein in both mitotic and meiotic cultures. The levels of Cdc14-HA were lower in mitotic cells
compared to the wild type, and they were further reduce upon meiotic entry. On the other hand, Cdc14
protein levels in the mitotically dividing *cdc14-md* mutant were similar to the wild type, but they
showed a drastic depletion with meiosis onset (Fig 2B). Increasing the copy number of different *cdc14*
alleles, as well as overproduction of the Cdc14-HA protein restored sporulation to a large extent (Fig
1F and Fig S2C-D). These results indicate that the reduction of Cdc14 phosphatase levels, but not its
full depletion, is causing the meiotic phenotype in the hypomorphic *cdc14-HA* mutant. On the other
hand, the almost complete absence of the protein throughout the entire meiotic process likely provokes
the more stringent block in nuclear divisions observed in *cdc14-md* cells. Thus, both alleles are useful
tools to investigate different meiotic events that could be regulated by Cdc14.

SPB integrity is compromised in *cdc14-HA* cells after the second meiotic division
As mentioned above, a great proportion of *cdc14-HA* cells eventually passed the two-nuclei
stage observed as terminal meiotic phenotype in *cdc14-md* (this study) and *cdc14* temperature-sensitive
mutants (Schild and Byers 1980; Sharon and Simchen 1990; Marston et al. 2003). Thus, we monitored
the kinetics and morphology of the spindle pole bodies (SPBs) and spindles during the first and the
second meiotic divisions (Fig 2C-D). We observed that by eight hours into meiosis, *CDC14* and *cdc14-*
HA cultures contained around 75% and 60% of cells with more than one SPB, respectively. Among the
population of cells that showed multiple SPBs, 56% of wild-type cells presented more than two;
therefore, they had initiated meiosis II. However, only 40% of *cdc14-HA* cells displayed more than two
SPBs, confirming a slight reduction of cells forming meiosis II spindles carrying four separated SPBs.
This observation correlates with the slightly lower frequency of *cdc14-HA* cells containing more than
two DAPI masses (Fig 2D; Fig S1B). As expected, after 24 hours of meiotic induction, the wild type
showed over 90% of cells with three or four SPBs. In marked contrast, most *cdc14-HA* meiotic mutant
cells lacked a single SPB, suggesting that, in the absence of Cdc14, the structural integrity of SPBs
becomes compromised once both meiotic divisions take place (Fig 2D; Fig S3A). On the other hand,
most cells of the *cdc14-md* mutant, contained only 2 SPBs for the entire length of the time course (Fig
S3D), confirming earlier studies revealing a role for the phosphatase in SPB reduplication/half-bridge
separation (Fox et al. 2017). Additionally, the average time taken to complete both meiotic divisions
was not significantly different in the *cdc14-HA* mutant compared to the wild type (Fig S4A-C). Thus,
we can conclude that, albeit with some difficulties, the *cdc14-HA* mutant is still capable of undergoing
both meiotic divisions, but it is unable to form spores. On the other hand, the *cdc14-md* mutation, which
produces a more drastic meiosis-specific depletion of Cdc14, prevents the separation of duplicated SPBs
after the first division; therefore, these cells are unable to assemble proper tetrapolar spindles at meiosis
II. Such behaviour resembles more closely the phenotype(s) previously described for other non-meiosis
specific *cdc14* mutants (Marston et al. 2003; Fox et al. 2017).

Impeded homolog separation and sister missegregation are common hallmarks of *cdc14* mutations in meiosis

We took advantage of the hypomorphic *cdc14-HA* allele, which allows us to study the second division in the absence, or with very little amount, of the Cdc14 protein to analyze chromosome segregation. In the *cdc14-HA* mutant, it was frequent to observe cells with more than four DAPI-stained masses late in meiosis (Fig S1B and Fig S3C). This might be due to impediments during chromosome segregation in the first and/or the second meiotic division. To test this possibility, we analyzed nuclear morphology, along with the meiotic spindle, in individual cells undergoing the first and second divisions (Fig 3Ai-v). Cells presenting meiosis I (MI) spindles with a length of ≥5μm were considered to be in late anaphase I, and therefore, two DAPI masses on each pole should be distinguishable (Fig 3Ai). On the other hand, cells presenting DAPI-stained bridges might reflect some form of chromosome entanglement (Fig 3Aii-iii; (Marston et al. 2003)). Wild-type cells showed a clear separation of nuclear content in 74% of anaphase I cells analyzed (Fig 3A). By contrast, in both *cdc14-HA* and *cdc14-md* mutants, only ~20% of anaphase I cells displayed two fully resolved nuclei (Fig 3A). A large proportion (~80%) of anaphase I mutant cells presented a chromosomal bridge, suggesting that despite having undergone MI spindle elongation correctly, some form of DNA entwining was not properly resolved.

A subtle delay in cohesin removal during anaphase I onset was observed in a small proportion of *cdc14-HA* cells (Fig 3B-C and Fig S4D), which may be responsible for at least a fraction of these bridged DAPI masses (Marston et al. 2003). To test if these entanglements were eventually eliminated or persisted until the second meiotic division, we examined metaphase II cells and scored the configuration of their nuclei (Fig 3Aiv-v). The wild type presented over 76% of metaphase II cells with single individualized nuclei. By contrast, 75% of metaphase II cells in the *cdc14-HA* mutant showed DNA threads connecting both nuclear masses. As mentioned above, the severity of the SPB re-duplication/half-bridge separation phenotype in the *cdc14-md* strain precluded us to identify metaphase II cells with 4-SPBs in *cdc14-md* cells. We conclude that diminished levels of Cdc14 in meiosis results in the abnormal formation of nuclear bridges after the first meiotic division and that these bridges persist at least until metaphase II.

To determine whether the DNA bridges present at metaphase II in *cdc14-HA* mutant cells were due to problems arising during chromosome segregation, we constructed strains carrying fluorescent markers on specific chromosomes and followed their segregation (Straight et al. 1996). We made use of *tetO* repeats integrated at the interstitial *lys2* locus of chromosome II, together with the presence of the *tetR* repressor gene fused with GFP, which associates to these repeats (Michaelis et al. 1997; Katis et al. 2010). First, we analyzed strains labelled at chromosome II on both homologs to determine chromosome segregation fidelity in cells that had completed both meiotic divisions. We found that chromosome II segregated correctly in 95% and 78% of wild-type and *cdc14-HA* tetranucleated cells,

respectively. The remaining 22% of *cdc14-HA* mutant cells displayed at least one of the four nuclei lacking a GFP focus (Fig 3D; Fig S5B type1, 2 and 3 only). We further analyzed the first and the second division separately using strains carrying *lacO* repeats at the *trp1* locus, which is linked to the centromere of chromosome IV (*CENIV*), and expressing the *lacI* repressor fused to GFP (Shonn et al. 2002) (Fig S5A). In order to assess homolog disjunction, we first examined binucleated cells homozygous for the *CENIV-GFP*, and we did not find any defect in the segregation of homologs during the first meiotic division in *cdc14-HA* (Fig 3E). Very similar results were previously reported for homozygous *CENV-GFP* markers in *cdc14-1* mutants (Bizzari and Marston 2011). Next, we analyzed sister chromatid segregation in the second meiotic division using heterozygous strains for the *CENIV-GFP* marker. In this case, 50% of tetranucleated *cdc14-HA* cells displayed problems segregating sister centromeres (Fig 3F). Taken all results together, we conclude that accurate sister chromatid segregation in meiosis requires unperturbed Cdc14 activity.

Preventing recombination alleviates defective events in meiosis-specific *cdc14* mutants

It has been shown that elimination of meiotic DSBs alleviates problems with chromosome segregation in FEAR and *cdc14-1* mutants (Marston et al. 2003). Therefore, we considered that the presence of anaphase I nuclear bridges and sister chromatid missegregation in *cdc14-HA* cells could be, at least in part, originated from problems caused by defective recombination. If this were the case, these problems should be alleviated by eliminating meiotic recombination. To test this hypothesis, we combined the *cdc14-HA* allele with the *spo11-Y135F* mutation, which prevents DSB formation (Keeney et al. 1997). The absence of DSBs improved the separation of DAPI masses in anaphase I in both *cdc14-HA* and *cdc14-md* mutants (Fig 3A), indicating that recombination leads to the formation of anaphase I bridges in both meiosis-defective *cdc14* mutant strains. Accordingly, inactivation of Spo11 catalytic activity also improved sporulation in *cdc14-HA* and *cdc14-md* (Fig 3G). Thus, these results suggest that during meiosis Cdc14 plays a role in promoting accurate repair of Spo11-dependent DSBs.

Meiotic recombination is impaired in *cdc14* mutants

To gain insight into the function of Cdc14 during meiotic recombination, we performed DNA physical assays using the well-characterized *HIS4LEU2* recombination hotspot (Fig 4A; (Cao et al. 1990; Schwacha and Kleckner 1994; Hunter and Kleckner 2001; Borner et al. 2004)) in wild type, *cdc14-HA* and *cdc14-md* strains. First, we measured total levels of DSBs, CO and NCOs products (Fig 4C-D). Levels of DSBs and COs were very similar in the three strains analyzed, albeit a small, not significant, reduction in NCO levels was observed in *cdc14-HA* and *cdc14-md* mutants (Fig 4C-D). Although a strong effect in the formation of recombination products was not perceived in *cdc14* mutant cells, shorter, or longer, lifespan of JMs could alter the ratio of NCO to CO to compensate for lower, or higher, efficiency in the processing of these intermediates (Martini et al. 2006; Oh et al. 2007; Shinohara

et al. 2008). To address this possibility, we analyzed JM formation in the *ndt80Δ* mutant background

(Allers and Lichten 2001). Accumulation of JMs takes place in *ndt80Δ* cells due to the lack of *CDC5*-

dependent JM-resolution promoting activity (Sourirajan and Lichten 2008). DSBs, NCOs and JMs

formed efficiently in the *cdc14-HA* mutant (Fig. 4E-F). Thus, under unchallenged conditions, Cdc14

appears to play little or no effect in the initiation and processing of meiotic recombination during

prophase I, at least at the *HIS4LEU2* hotspot.

Next, we wanted to monitor if CO and/or NCO formation was affected in *cdc14* mutants when

DSB repair is moderately compromised by eliminating one of the two main resolution pathways

described in budding yeast, such as the Mus81/Mms4 pathway (de los Santos et al. 2001). Thus, we

proceeded to evaluate the efficiency of JM resolution in a *mus81Δ ndt80Δ* mutant background. In order

to obtain an accurate reading of repair product formation we first confirmed that total levels of JMs

were similar in both *mus81Δ ndt80Δ* and *mus81Δ cdc14-HA ndt80Δ* mutant combinations (Fig 4B; Fig

S6). To promote JM resolution in the absence of Ndt80, we took advantage of the widely-employed

CDC5-inducible expression system (*CDC5-IN*), where the expression of the *CDC5*, which is required

for the resolution of JMs into COs, can be induced at any point required in the meiotic time-course by

the addition of β-estradiol (ES) to the media (Carlile and Amon 2008; Sourirajan and Lichten 2008).

We added ES after 7 hours in meiosis and monitored CO and NCO formation. Induced expression of

CDC5 in the *mus81Δ* mutant efficiently led to the formation of COs only one hour after ES addition to

the medium (Fig 5A-B); CO levels did not increase further after the 8-hour time-point. On the other

hand, NCOs steadily increased reaching their maximum at 24h. In contrast, fewer COs were detected

in the *mus81Δ cdc14-HA* mutant. NCOs levels were also lower in *mus81Δ cdc14-HA*, and total levels

did not rise over the period when *CDC5* was expressed (Fig 5B). Thus, Cdc14 is required for efficient

formation of COs and NCOs, at least when meiotic DSB repair takes place in the absence of the

Mus81/Mms4-dependent resolution pathway.

To confirm that Cdc14 plays a relevant role in the repair of recombination intermediates at least

when certain resolution pathways are compromised, we decided to study the efficiency of repair in a

NDT80 background using 2D-gels that provide more precise information of the different DNA species

involved during the JM resolution process (Fig 4B). To this end, we took advantage of the well-known

negative effect on DSB-repair when depleting Sgs1 in combination with mutations in one or more of

the SSEs, such as *mms4/mus81* and *yen1/slx1/slx4* (De Muyt et al. 2012; Zakharyevich et al. 2012). In

this case, we used the *cdc14-md* allele, to prevent any residual activity of the Cdc14 phosphatase. As

expected, elimination of *YEN1* in the *sgs1-md mms4Δ* background led to increased accumulation of

unprocessed JMs compared to the *sgs1-md mms4Δ* double mutant (Fig 5C-D). Strikingly, after 24 hours

in meiosis, the *sgs1-md mms4Δ cdc14-md* triple mutant also accumulated great levels of unresolved

JMs compared to the *sgs1-md mms4Δ* double mutant (Fig 5C-D). Interestingly, the accumulation of

unresolved JMs in *sgs1-md mms4Δ cdc14-md* was similar to that of the *sgs1-md mms4Δ yen1Δ* triple

mutant (Fig 5C-D). Furthermore, CO levels at the *HIS4LEU2* hotspot were highly reduced in the *sgs1-*

md mms4Δ cdc14-md strain; a phenotype that was nearly identical to that of *sgs1-md mms4Δ yen1Δ* (Fig 5E-F). Thus, in the absence of the Sgs1 and Mus81/Mms4-dependent pathways for processing recombination intermediates, the lack of the Cdc14 phosphatase leads to the same defects as the absence of the Yen1 resolvase, strongly suggesting that Cdc14 and Yen1 are involved in the same repair pathway during meiosis.

Yen1 nuclear localization and activity during meiosis requires Cdc14 function.

In mitotic cells, Yen1 localization is cytoplasmic when CDK activity is high, but release of Cdc14 during anaphase I reverts CDK-dependent phosphorylation on Yen1 promoting its nuclear entry and activity (Matos et al. 2011; Blanco et al. 2014; Eissler et al. 2014; Garcia-Luis et al. 2014). In meiotic cells, Yen1 subcellular localization is also regulated in a phosphorylation-dependent and cell-cycle stage-specific manner (Matos et al. 2011). To determine whether Cdc14 also regulates Yen1 subcellular distribution during meiosis, we examined nuclear localization of Yen1 in anaphase I cells, and we found that it was markedly reduced in *cdc14-HA* compared to the wild type (Fig 6A). We also tested if the enzymatic activity of Yen1 is controlled by Cdc14 during meiosis (Fig 6B-C). Samples were collected from synchronized meiotic cultures of wild-type and *cdc14-HA* strains carrying a functional 9MYC-tagged version of *YEN1* (Arter et al. 2018). From each time-point taken, Yen1 was immunoprecipitated from the meiotic extracts and the beads were incubated with a synthetic HJ substrate (Grigaitis et al. 2018). Active Yen1 should be able to cleave the HJ substrate whereas the phosphorylated inactive form of the nuclease would not (Blanco et al. 2014). In wild-type cells, Yen1 did not display significant nuclease activity during early stages of the meiotic time course prior to the accumulation of Cdc5 (Fig 6B), but approximately 2 hr after Cdc5 induction, we detected robust HJ processing, consistent with activation of Yen1 during meiotic divisions (Fig 6B-C; (Matos et al. 2011)). Notably, the resolvase activity displayed by Yen1 in the *cdc14-HA* mutant was extremely low for the duration of the whole time course (Fig 6B-C). Meiotic S-phase was completed with similar timing in wild-type and *cdc14-HA* cells suggesting that the initiation of the meiotic program occurred with similar kinetics in both strains (Fig 6D). Moreover, the Cdc5 protein (marking the exit from prophase I) was detected at the same time, and at the same levels, in both wild-type and *cdc14-HA* strains (Fig. 6B). Thus, these results strongly suggest that Cdc14 promotes nuclear accumulation of Yen1 during anaphase I and is required for its enzymatic activity at times when Cdc5 activity is high.

Constitutively active Yen1 supresses the accumulation of aberrant recombination intermediates in *cdc14* mutants

To determine whether the activation of Yen1 by Cdc14 is exerted by controlling its phosphorylation status, we next used a phosphorylation-resistant version of the nuclease (*YEN1^ON^*) in which mutation of nine of the CDK consensus sites of phosphorylation in Yen1 renders the protein

constitutively active (Blanco et al. 2014; Arter et al. 2018). We combined the *cdc14-HA* allele with

YEN1^{ON} and examined sporulation efficiency. Similar to the elimination of Spo11-dependent DSBs,

unrestrained activity of Yen1 partly rescued the sporulation defect of *cdc14-HA* (Fig 6E). We also

directly tested the effect of *YEN1*^{ON} in chromosome segregation during both meiotic divisions (Fig 6F).

First, we checked if the presence of Yen1^{ON} promotes resolution of the DNA bridges observed at late

anaphase I in *cdc14-HA* mutants. Indeed, a two-fold improvement in *cdc14-HA YEN1*^{ON} over the *cdc14-*

HA single mutant was observed (Fig 6F, left graph). Furthermore, missegregation of sister chromatids

was also fully rescued to the levels of the *YEN1*^{ON} single mutant in the *cdc14-HA YEN1*^{ON} double mutant

(Fig 6F, right graph). Moreover, to confirm if the improvement in chromosome segregation by *YEN1*^{ON}

was due to the reestablishment of Yen1 resolvase activity, we employed two types of assays. First, we

checked using 2D-gels whether persistent accumulation of JMs and/or CO formation observed at the

HIS4LEU2 hotspot in the repair-deficient *sgs1-md mms4Δ cdc14-md* triple mutant was alleviated by

the constitutively active Yen1^{ON} (Fig 5C-F). Notably, the introduction of *YEN1*^{ON} efficiently prevented

the accumulation of unresolved JMs after 24 hours in meiosis (Figs 5C and D). Furthermore, JMs were

efficiently resolved giving rise to high levels of COs in the *sgs1-md mms4Δ cdc14-md YEN1*^{ON}

quadruple mutant (Figs 5E and F). Finally, we used the *in vitro* resolution assay to determine whether

Yen1^{ON} was capable of restoring resolvase activity in the *cdc14-HA* mutant. We found that, unlike

Yen1, the constitutively active Yen1^{ON} displayed potent resolvase activity in the absence of Cdc14 (Fig

6G). Altogether, these results indicate that Yen1 activation by Cdc14 is important for the repair of

unprocessed recombination intermediates in meiosis.

Cdc14 and Yen1 promote JM resolution during the first meiotic division

In the meiotic program, Yen1 activity appears to reach its maximum level during the second

meiotic division (Matos et al. 2011; Arter et al. 2018). Nevertheless, it is possible that, during the first

release of Cdc14 at meiosis I, Yen1 can be activated to act over remnants of unprocessed or complex

JMs. In support of this idea we observed that 52% of wild-type cells displayed nuclear Yen1 in anaphase

I (Fig 6A), indicating that a high proportion of Yen1 molecules have been translocated to the nucleus

during the first meiotic division. To test if Cdc14 is acting over Yen1 already in meiosis I, or by contrast,

Yen1 is exclusively activated during the second release of Cdc14 at meiosis II, we used the *cdc20-md*

mutant to deplete the APC component Cdc20 and arrest cells in meiosis I (Carlile and Amon 2008). We

analyzed the effect of Cdc14 and Yen1 on JM resolution in *cdc20-md* cells blocked at the first meiotic

division when the Sgs1 and Mus81/Mms4 repair pathways, normally acting at the end of prophase I,

have been removed. Remarkably, the *cdc20-md sgs1-md mms4Δ yen1Δ* and *cdc20-md sgs1-md mms4Δ*

cdc14-md quadruple mutants exhibited much higher levels of unprocessed JMs than the *cdc20-md sgs1-*

md mms4Δ triple mutant (Fig. 7A-B). In addition, this JM resolution defect was accompanied by a

markedly reduced capacity of CO formation in both *cdc20-md sgs1-md mms4Δ yen1Δ* and *cdc20-md*

sgs1-md mms4Δ cdc14-md quadruple mutants compared to the *cdc20-md sgs1-md mms4Δ* triple mutant

(Fig. 7C-D). Altogether, we conclude that, when the action of other relevant JM processing pathways

is compromised, Cdc14 and Yen1 are required for proper JM resolution as early as in the first meiotic

division.

Discussion

Meiotic recombination is fundamental for sexual reproduction to ensure that homologous

chromosomes are accurately distributed to the gametes as well as to facilitate distinct allele

combinations that sustain evolution. Homologous recombination is initiated by the introduction of

programmed DSBs followed by resection, homology search, and DNA strand exchange. Those early

steps in the recombination process lead to the formation of stable JMs, which are ultimately resolved

into two main classes of HR repair products, known as COs and NCOs (Fig 8). Recombining

chromosomes may also contain intermediates consisting of three- and four-armed DNA structures, such

as mc-JMs, where three and four DNA duplexes stay connected (Schwacha and Kleckner 1995; Hunter

and Kleckner 2001; Liberi et al. 2005; Cromie et al. 2006; Jessop and Lichten 2008; Oh et al. 2008;

Bzymek et al. 2010; Mankouri et al. 2011). In some cases, unresolved recombination intermediates can

persist until the metaphase to anaphase transition where a subset of late-acting nucleases take charge of

their processing in order to safeguard genome integrity (San-Segundo and Clemente-Blanco 2020).

Unrestrained activity of such nucleases can interfere with the correct allotment of COs and NCOs in

meiosis (Arter et al. 2018); thus, they are tightly controlled by several layers of regulation (Fig. 8).

Phosphorylation events carried out by key cell cycle kinases, like CDK, DDK, and Polo Kinase, are the

most extended form of regulation of these enzymatic activities (Matos et al. 2011; Wild et al. 2019);

therefore, it is expected that phosphatases can play also crucial roles in this regulation. In *S. cerevisiae*

mitotic cells, at least one of the late acting nucleases, Yen1, is modulated by the highly-conserved Cdc14

phosphatase (Blanco et al. 2014; Eissler et al. 2014; Garcia-Luis et al. 2014). Although a number of

studies have unveiled the relevance of Cdc14 during meiotic chromosome segregation and

SPB/centrosome dynamics, a role for Cdc14 in the regulation of meiotic recombination has not been

established so far. Given the essentiality of *CDC14* in the budding yeast, most previous meiotic studies

have used conditional temperature-sensitive alleles of *cdc14* that are blocked in meiosis after the first

round of chromosome segregation (Culotti and Hartwell 1971; Schild and Byers 1980; Marston et al.

2003; Fox et al. 2017). Despite the large amount of valuable data that has been collected over the years

using these *ts* alleles, other less conspicuous functions of Cdc14, for example those affecting meiosis

II, might have been precluded from being discovered. Here, using two different meiosis-specific alleles

of *CDC14* we have been able to identify previously undetected functions of the phosphatase during

meiosis, particularly those affecting meiotic recombination (Fig 8). Strikingly, we found the

unprecedented requirement for Cdc14 to promote JM resolution at the transition between prophase I

and the first meiotic division, suggesting that Cdc14 activity is not strictly ligated to its full release from

the nucleolus.

475   Novel insights into Cdc14 meiotic functions using different *cdc14* alleles

In the present work we describe an allele of *CDC14* (*cdc14-HA*) that displays no obvious defects

during unchallenged mitotic divisions, but it is strongly deficient in sporulation. Protein levels of the

phosphatase are reduced in both mitotic and meiotic *cdc14-HA* cells, but this reduction is more dramatic

during meiosis (Fig 2A-D). However, the amount of the phosphatase in *cdc14-HA* meiotic cells is

sufficient to complete meiotic DNA replication and both nuclear divisions with rather normal kinetics.

Nonetheless, the *cdc14-HA* mutant fails to form spores once cells completed both nuclear divisions. It

is possible that the impaired structural integrity of the SPBs observed in *cdc14-HA* can contribute to

this phenotype. Similar defects in sporulation have been also described in the absence of components

of the MEN pathway (Pablo-Hernando et al. 2007; Arguello-Miranda et al. 2017). Importantly, correct

SPB re-duplication and separation occur in a high proportion of *cdc14-HA* cells, but not in the more

restrictive *cdc14-md* mutant that contains negligible levels of Cdc14 during meiosis (Fig 2E-F; S3 Fig).

Furthermore, many *cdc14-HA* cells are able to assemble functional tetrapolar spindles at

metaphase/anaphase II confirming that problems arising in the *cdc14-HA* mutant somewhat differ from

those occurring in *cdc14-md* as well as in the widely-employed, *cdc14-1, cdc14-3* or other FEAR

mutants (Marston et al. 2003; Bizzari and Marston 2011; Fox et al. 2017). This observation suggests

that the small amount of the phosphatase still present in *cdc14-HA* is sufficient to carry out those roles

required for meiotic SPB dynamics, but not to maintain SPB integrity after divisions. The *cdc14-HA*

mutant also displays a short delay in cohesin removal when entering anaphase I (Fig 3B-C and Fig

S4D), sharing this particular meiotic phenotype with *spo12*, *slk19* and *cdc14-1* mutants (Marston et al.

2003). It is possible that delayed Rec8 removal from chromosomes could contribute to the presence of

DNA bridges in late anaphase I. However, the proportion of *cdc14-HA* cells presenting anaphase I

bridges is consistently higher than that of cells displaying persistent Rec8 signal during anaphase I;

thus, it is unlikely that inefficient Rec8 removal is the only source of most DNA bridges, but instead a

contributing factor. The fact that the formation of chromatin bridges in *cdc14-HA* and *cdc14-md* is

substantially alleviated in the absence of meiotic DSBs strongly suggests that they may arise from

chromosome entanglements resulting from unresolved recombination intermediates, and raises the

possibility of a direct role for Cdc14 in the regulation of meiotic recombination that has not been

previously explored. Here we demonstrate that, in some circumstances, Cdc14 is critically required for

proper resolution of meiotic JMs through the regulation of the Yen1 nuclease.

506   Multistage *CDC14*-dependent processing of recombination intermediates

Meiotic cells are endowed with a battery of enzymatic activities, including dissolvases and

resolvases, that are tightly regulated to ensure efficient and timely processing of recombination

intermediates (San-Segundo and Clemente-Blanco 2020). Processing of most meiotic JMs occurs at the

end of pachytene (Borner et al. 2004), far earlier than the reported first release of Cdc14 at anaphase I.

Nevertheless, this is not the first time that the requirement of Cdc14 has been linked to cell cycle stages

preceding its bulk release from the nucleolus at anaphase. For example, DNA damage caused in

vegetative cells triggers the transitory release of Cdc14 from the nucleolus to the nucleoplasm, allowing

the phosphatase to act on components of the SPBs to stabilize them at metaphase (Villoria et al. 2017).

Cdc14 is involved in completion of late-replicating regions in the rDNA, and other parts of the genome

(Dulev et al. 2009). Although the *cdc14-HA* mutant does not show a discernible mitotic phenotype

under unchallenged conditions, it is formally possible that *cdc14-HA* cells, which contain reduced

protein levels, may accumulate aberrant DNA structures during preceding mitoses prior to meiosis

entry. However, this is unlikely the case for the *cdc14-md* mutant, whose mitotic protein levels are very

similar to those of the wild type. Therefore, we favor a role for Cdc14 in the direct regulation of

substrates required for correct processing of meiotic JMs (Fig 8). Interestingly, the conserved SSE,

Yen1^{GEN1} is a critical substrate of Cdc14 during budding yeast mitosis, and it exhibits a phosphorylation-

regulated nucleocytoplasmic shuttling behaviour (Kosugi et al. 2009). CDK-dependent phosphorylation

restricts Yen1 from entering the nucleus and becoming active, whereas reversal of that phosphorylation

by Cdc14 allows Yen1 to enter the nucleus and resolve entangled DNA structures (Blanco et al. 2014;

Eissler et al. 2014; Garcia-Luis et al. 2014). Nonetheless, Yen1 nuclear import and activation appears

to be concomitant with bulk Cdc14 release from the nucleolus during anaphase. This is why a role for

Yen1 in safeguarding chromosome segregation has been proposed, especially, during the second

meiotic division (Matos et al. 2011; Blanco et al. 2014; Arter et al. 2018). Our results support this

conclusion, since high frequencies of aneuploidy and missegregation events are detected during meiosis

II in the *cdc14-HA* meiotic mutant when the other repair pathways are intact (Fig 3F). This is consistent

with the requirement for Cdc14 to activate Yen1 (Fig 6A-C). Puzzlingly, we also observe Yen1-

dependent processing of JMs in cells arrested during MI by Cdc20 depletion (Fig 7). This effect is

manifested more prominently when the Mus81/Mms4 endonuclease and the Sgs1 helicase are missing.

Such resolvase activity is not observed in *cdc14* mutants (Fig 7), suggesting that Cdc14 can activate

Yen1 in cells that are about to initiate anaphase I. Taking into account that the massive Yen1 nuclear

transport takes place during the ensuing anaphase (Arter et al. 2018), our results open the question of

the origin of the Yen1 population acting in *CDC20*-depleted cells, which are stably arrested at the

metaphase I to anaphase I transition in meiotic cells (Lee and Amon 2003; Katis et al. 2004; Carlile and

Amon 2008; Arguello-Miranda et al. 2017). We speculate about other possible scenarios, like the

existence of a different fraction of Yen1 arising from a different subcellular compartment to perform

this earlier role. In support of this possibility, it has been recently postulated that a subpopulation of

Yen1 localizes at the nucleolus in mitotic cells (Talhaoui et al. 2018). Whether the nuclear Yen1

observed in MI is sourced from the cytoplasm or from other sub-nuclear compartment will be a matter

of future studies.

Alternative repair pathways to process JMs during meiosis I

An intriguing possibility is that different SSEs might be required for processing different types of JMs that may not be necessarily good substrates for the canonical MutLγ complex. There is a risk that spindle forces originated during the *cdc20-md* arrest could generate tension at a number of still unprocessed JMs eventually forcing them towards conformations that cannot be easily processed by prophase-specific type of resolution/dissolution enzymes, for example affecting branch migration kinetics or directionality (Tang et al. 2015; Machin 2020). In such perilous scenario, Cdc14 could be an ideal regulator, together with Cdc5, to promote the switch between different endonucleases, by activating some while inhibiting others, once spindle forces start becoming dominant under circumstances where the SC no longer exists to hold homologs in close association (Shodhan et al. 2019; Cannavo et al. 2020; Kulkarni et al. 2020).

In budding yeast meiosis, the timing of JM resolution and CO formation is coordinated with cell-cycle progression through the *NDT80*-dependent expression of the Polo-like kinase Cdc5 (Clyne et al. 2003; Sourirajan and Lichten 2008; Matos et al. 2011). Thus, the requirement of Cdc5 for the resolution of dHJ at the pachytene to MI transition, upon Ndt80 activation, might too involve its ability to regulate the interaction of Cdc14 with Cif1/Net1 in the nucleolus. In opposition to mitotic cells, the action of Cdc5 could temporally counteract the negative regulatory effect of PP2A^{Cdc55} allowing some Cdc14 molecules to escape from its captor (Bizzari and Marston 2011; Kerr et al. 2011). It is tempting to speculate that Cdc5 might also play a relevant role during meiosis in promoting an early, metastable, partial release of a Cdc14 population at the transition from the pachytene stage to metaphase I in order to modulate the activity of a number of safeguarding enzymes required for correct chromosome segregation (Tsuchiya et al. 2014). Understanding the regulatory pathway/s which control Cdc14 activity previous to its full release during anaphase will be important to decipher its full contribution to the regulation of meiotic recombination.

In recent years, human orthologs of Cdc14 phosphatase have received increased attention due to their involvement in key processes like DDR, DNA repair and cell cycle control. Furthermore, recent findings point to recessive variants of the phosphatase to be directly responsible of human diseases, like deafness and male infertility (Imtiaz et al. 2018). Thus, in order to comprehend the underlying factors that trigger those conditions, a deeper understanding of the genetic and molecular mechanisms that are responsible for the countless functionalities of the Cdc14 phosphatase during gametogenesis and HR repair will be required.

Materials and Methods

Yeast strains and plasmids

All strains were SK1, as detailed in Table S1. Standard yeast manipulation procedures and growth media

were utilized. To introduce the 3HA tag at the C-terminal end of Cdc14, the CDC14ha3-pKan^R plasmid

containing the last ~500bps of the *CDC14* gene with 3xHA inserted at the NotI restriction site and

containing the *CLB1* terminator, was used. The plasmid was linearized using a unique restriction site

located within the *CDC14* sequence and transformed into a SK1 haploid strain. Alternatively, *CDC14*

was tagged using the PCR-based method described in (Longtine et al. 1998) using the plasmid pFA6a-

3HA-kanMX6. The phenotype of *cdc14-HA* strains obtained from both tagging methods was checked

and the sporulation defect was identical. Transformants containing correct tag integration were

identified and tested by western blot for the presence of the tag. Southern blot analysis and/or PCR was

performed to confirm the integration at the endogenous locus. Multi-copy plasmids used in Figure S2C

were originally described in (Jaspersen et al. 1998).

Synchronous meiotic time courses

592   Induction of synchronous meiosis was carried out according to the established protocols for standard

593   assays (Padmore et al. 1991). All pre-growth and meiotic time courses were carried out at 30°C unless

594   specified otherwise. For *cdc14-1* meiosis, the culture was kept at 23°C and shifted to 30°C 2 hours after

595   transferring into sporulation medium (SPM). Aliquots were removed at the specified times and

596   subjected to various analyses.

DNA manipulation, extraction and southern blot detection

598   Standard DNA extraction was performed as in (Carballo et al. 2013). For studies at the *HIS4LEU2*

599   recombination hotspot, the protocol described in (Ahuja and Borner 2011) was followed. For 2D gel

600   agarose electrophoresis, cell cultures were photo-crosslinked with trioxalen (Merck) using long-wave

601   UV light before DNA extraction, in order to visualize recombination intermediates by standard southern

602   blotting techniques at the *HIS4LEU2* hotspot (Ahuja and Borner 2011).

Time-lapse imaging, immunofluorescence, microscopy, and image analysis

Time-lapse experiments were performed as in (Newnham et al. 2013), with small variations. In brief, 1

ml aliquots from synchronous meiotic cultures were taken at specific times and diluted 1:9 in fresh SPM

(kept at 30°C). 300μl of diluted cells were placed in suitable multi-well slides (80821 uncoated, ibidi).

Slides were placed in a temperature-controlled incubation chamber from a Multidimensional

Microscopy System Leica AF6000 LX. Images were taken at multiple positions and channels every 5,

10 or 15 minutes, depending on the experiment. Image acquisition was carried out using a CCD

Hamamatsu 9100-02 camera. The system was controlled by the proprietary software, Leica Application

Suite X, version 3.30. For preparations of fixed cells and immunofluorescence, aliquots were fixed and

prepared as described in (Carballo et al. 2013) and (Arter et al. 2018), respectively. Chromosomal DNA

was stained with 1μg/ml 4,6-diamino-2-phenylimide (DAPI). Images were recorded and analyzed using

a Deltavision (DV3) workstation from Applied Precision Inc. with a Photometrics CoolSnap HQ (10-20MHz) air cooled CCD camera and controlled by SoftWorx image acquisition and deconvolution software.

Protein extraction, Western blot analysis and antibodies

Whole-cell extracts were prepared from cell suspensions in 20% trichloroacetic acid by agitation with glass beads. Precipitated proteins were solubilized in SDS-PAGE sample buffer, and appropriate dilutions were analyzed by SDS-PAGE and western blotting (Gimenez-Abian et al. 2004). Antibodies used for western blotting were mouse monoclonal anti-MYC (1:1000, Abcam), mouse monoclonal anti-HA (1:1000) from S. Ley (NIMR), goat polyclonal anti-Cdc14 (yE-17; 1:1000; Santa Cruz Biotechnology), goat polyclonal anti-Cdc5 (1:1000; Santa Cruz Biotechnology), mouse monoclonal anti-Pgk1 (1:5000; Invitrogen), goat anti-mouse IgG conjugated to horseradish peroxidase (1:10000; Sigma-Aldrich), and chicken anti-rabbit IgG conjugated to horseradish peroxidase (1:10000; Sigma-Aldrich).

Nuclease assays

For nuclease assays, myc9-tagged Yen1 and Yen1[ON] were immuno-affinity purified from yeast using anti-Myc agarose beads (9E10) and washed extensively. The beads (approx. volume 10 µl) were then mixed with 10 µl cleavage buffer (50 mM Tris-HCl pH 7.5, 3 mM $MgCl_2$) and 15 ng of 5'-Cy3-end-labeled synthetic Holliday junction X0 DNA. After 1 h incubation at 37°C with gentle rotation, reactions were stopped by addition of 2.5 µl of 10 mg/ml proteinase K and 2% SDS, followed by incubation for 30 min at 37°C. Loading buffer (3 µl) was then added and fluorescently-labelled products were separated by 10% native PAGE and analyzed using a Typhoon scanner and ImageQuant software. Resolution activity was calculated by determining the fraction of nicked duplex DNA product relative to the sum of the intact substrate and resolution product. The protein input was estimated by western blot.

Data analysis and Biostatistics

Data was compiled and analyzed using Excel, LibreOffice Calc, and SPSS Statistical Data Editor. For multiple comparisons, analysis of variance (one-way ANOVA) was performed. For pairwise comparisons, two-tailed unpaired *t*-tests were used using IBM SPSS Statistics and SigmaPlot.

Acknowledgements

We would like to thank C.R. Vázquez de Aldana, M. Knop, E. Winter, A.M. Neiman, E. Hoffmann, J.L. Santos, G.V. Börner, A. Marston, M.G. Blanco and F. Machín for kindly providing us with reagents, strains, antibodies and/or advice. We are especially thankful to the Molecular Biology of the Chromosome lab in CIB for sharing lab space, reagents and equipment. We would like to thank to the

647 microscopy facility at the CIB for valuable technical advice in designing and analysing time-lapse, to

648 P. Jalón for UV photography acquisition and to J.S. Ahuja, for advice in 2D gel experiments.

Figure legends

Figure 1. *cdc14-HA* behaves as a specific sporulation-deficient separation-of-function allele of

CDC14. (A) Homozygous *cdc14-HA* diploids do not form asci containing spores under standard

conditions for sporulation. (B) *cdc14-HA* diploid cells divide mitotically but lack di-tyrosine

autofluorescence induced by UV light after several days of incubation in SPM media at 30°C. (C) The

cdc14-HA allele does not perturb normal growth conditions when cultivated at 25°C, 30°C and 37°C in

rich media (YPD). (D) FACS analysis of DNA content shows that *cdc14-HA* cells complete meiotic

DNA replication with identical kinetics as *CDC14* diploid cells. (E) *cdc14-HA* cells undergo meiotic

nuclear divisions at subtly slower kinetics, and reduced frequencies, than *CDC14* cells. Error bars

represent the SEM over the mean values plotted. (F) Di-tyrosine autofluorescence of different mutant

combinations as well as the control strains grown and sporulated on plates at 25°C. *cdc14-1*

homozygous diploids (JCY2353) sporulate at high efficiency under semi-permissive temperature

forming preferentially tetrads whereas *cdc14-HA* homozygous diploids (JCY840) do not sporulate.

Combinations, and variable copy number, of the mutant genes can rescue the sporulation defect to

different degrees (JCY2365/JCY2354/JCY2356).

Figure 2. Diminished Cdc14 protein levels in *cdc14-HA* meiotic cells compromise SPB integrity

following meiotic divisions. (A) Dynamics of Cdc14 protein levels throughout meiosis in synchronous

time-courses for *CDC14* (JCY2232), *cdc14-HA* (JCY2231), and *cdc14-md* (JCY2389). Higher contrast

was applied to visualize residual protein levels for all three strains. Immunodetection of Pgk1 was used

as loading control. FACS histograms for each time-point are depicted at the bottom to show the degree

of synchrony reached in each culture. (B) Variability of protein levels in mitotic versus meiotic cells in

different *CDC14* alleles. Exponentially growing cells show similar Cdc14 levels except for *cdc14-HA*,

which has lower overall protein levels (left panel). Direct comparison between the three alleles *CDC14*

(JCY2232); *cdc14-HA* (JCY2231) and *cdc14-md* (JCY2389*)* in exponentially growing mitotic cells

(Exp.) and meiotic cultures immediately after they were transferred to SPM (Mei$_0$). Loading control for

each lane was determined using Pgk1. (C) Visualization of Spc29-CFP at SPBs, tubulin, and DNA, in

wild-type (JCY892) and *cdc14-HA* (JCY893) cells fixed at different stages of meiosis at 30°C. (D)

cdc14-HA cells develop meiosis I and II spindles with near wild-type kinetics. Lack of spore formation

in *cdc14-HA* meiotic cells parallels the loss of SPB's structural integrity.

Figure 3. Missegregation of sister chromatids at MII in meiosis-deficient *cdc14* mutant cells. (A; i-iii) Frequency of cells presenting connected DAPI masses at anaphase I in wild type (JCY902), *cdc14-HA* (JCY904), *cdc14-md* (PRY132), *spo11* (PRY53) and *spo11 cdc14-HA* (PRY55). (A; iv-v) Frequency of cells presenting connected nuclei at metaphase II. Meiotic-deficient *cdc14-HA* mutants (GGY54) present higher frequencies of unresolved nuclear divisions at late anaphase I and at metaphase II than wild type (GGY53). (B) Representative images of distinct morphological patterns for Rec8-GFP in meiosis. Full: bright nuclear localization. Diffuse: faint staining within the stretched nucleus. Peri: Rec8-GFP only visible at peri-centromeric locations. Absent: no Rec8-GFP signal. Pds1-tdTomato and CNM67-tdTomato were used to follow Pds1 accumulation/degradation and SPB number/location. (C) Temporal distribution of the distinct morphological categories of Rec8-GFP shown in (B) for *CDC14* (JCY2406; n=57 cells) and *cdc14-HA* (JCY2404; n=52 cells). (D) Unequal distribution of homozygous chromosome II-linked GFP markers at *lys2* locus in *cdc14-HA* mutant cells (JCY2330) denotes increased chromosome missegregation in comparison with wild-type cells (JCY2331). (E) Similar distribution of homozygous *CENIV*-linked GFP markers at *trp1* locus in meiosis-deficient *cdc14-HA* mutant cells (JCY2286) in comparison with wild-type cells (JCY2284) denotes correct homolog disjunction in anaphase I. (F) Unequal distribution of heterozygous *CENIV-GFP* marker in meiosis-deficient *cdc14-HA* tetranucleated mutant cells (JCY2327) denotes increased sister chromatid missegregation in comparison with wild-type cells (JCY2326). Statistical significance of differences was calculated by two-tailed *t*-test, assuming unequal variances ($^*p < 0.05$; $^{**}p < 0.01$; $^{***}p < 0.001$; $^{****}p < 0.0001$; n.s.: not significant). (G) Sporulation defects in *cdc14-HA* (JCY844) or *cdc14-md* (JCY2376) can be partly alleviated (JCY840) by eliminating recombination (JCY2270/JCY2280/PRY151).

Figure 4. Analysis of meiotic recombination in meiotic-deficient *cdc14* mutants. (A) Schematic illustration of recombination analysis and intermediates at the *HIS4LEU2* hotspot (see Results for more details). (B) Schematic representation of DNA species commonly observed after analysing the *HIS4LEU2* hotspot by 2D-gel electrophoresis. (C; top) Representative 1D-gel Southern blot for analysis of DSBs and COs at the *HIS4LEU2* hotspot using *XhoI* as the restriction site. Wild-type strain (JCY2232), *cdc14-HA* (JCY2231), and *cdc14-md* (JCY2389). (C; bottom) Analysis of COs and NCOs at the *HIS4LEU2* hotspot using *XhoI* and *NgoMVI* in the same strains as in (C). (D) Quantification of DSBs, COs and NCOs from the Southern blots shown in (C). (E) Representative 1D-gel Southern blot image for analysis of DSBs, COs (top) and NCOs (bottom) at the *HIS4LEU2* hotspot in *CDC14 ndt80Δ* strain (JCY2390) and *cdc14-HA ndt80Δ* (JCY2385) strains. Asterisk indicates meiosis-specific non-characterized recombination products. (F) Quantification of DSBs, NCOs and JMs from the Southern blots shown in (E).

Figure 5. Analysis of recombination intermediates in *cdc14* mutants shows early and late accumulation of JMs. (A) Representative Southern blots depicting 1D-gels at the *HIS4LEU2* hotspot following induction of *CDC5* in prophase-arrested *ndt80Δ mus81Δ cdc14-HA* cells (JCY2440) leads to inefficient CO and NCO formation compared to *ndt80Δ mus81Δ* (JCY2442). (B) Quantification of COs and NCOs from Southern blots shown in (A). (C) Representative Southern blots depicting 2D-gels at the *HIS4LEU2* hotspot in *sgs1-md mms4Δ* (JCY2444), *sgs1-md mms4Δ cdc14-md* (JCY2446), *sgs1-md mms4Δ yen1Δ* (JCY2448) and *sgs1-md mms4Δ cdc14-md YEN1^ON* (JCY2529) 24 h into meiosis. Arrows mark IH-dHJs and lines mark mc-JMs (see Fig 4A-B). (D) Quantification of total JMs in the strains shown in (C) from at least two independent gels. Error bars represent SEM over the calculated mean value. (E) Representative 1D-gel Southern blot images for analysis of crossovers at the *HIS4LEU2* hotspot for all strains shown in (C) and for *cdc14-md* (JCY2389). (F) Quantification of COs from at least three different image acquisitions like the one depicted in (E). Error bars represent SEM over the calculated mean value.

Figure 6. The absence of Cdc14 prevents activation of Yen1 during meiotic divisions. (A) Quantification of distinct localization patterns of Yen1 at meiosis I. (B) Analysis of expression levels and nuclease activity of Yen1 in *CDC14* (YJM7692) and *cdc14-HA* (YML7693) meiotic cells. Soluble extracts were prepared from *YEN1-Myc9* strains at 2-hr intervals after transfer into sporulation medium (SPM). Following anti-Myc immunoaffinity purification (IP), the IPs were analyzed by western blotting and tested for nuclease activity using Cy3-labeled Holliday junction DNA as a substrate. Upper panel: western blots of the cell extracts, with detection of Yen1-myc9, Cdc5, and Pgk1 (loading control). Lower panel: HJ resolution assay. The experiment shown is representative of two independent experiments. (C) Percentage of HJ cleavage in Yen1 IPs during meiosis in wild type and *cdc14-HA*. (D) Evolution of DNA content during meiosis from strains in (B). (E) Unrestrained resolution of recombination intermediates by Yen1^ON improves sporulation in *cdc14-HA* (JCY2164) and *cdc14-md* (PRY182) cells. Frequency of asci containing one, two, three and four spores in the strains of the indicated genotypes. (F) Frequency of cells presenting connected DAPI masses at anaphase I (left panel) or sister chromatid missegregation (right panel) in *YEN1^ON* (PRY123/PRY99), *cdc14-HA* (JCY2327/PRY55) and *cdc14-HA YEN1^ON* (PRY121/PRY108). (G) Poor HJ resolution in *cdc14-HA* (YML7693) is efficiently restored by the presence of Yen1^ON (JCY2421). Western blot analysis of Yen1/Yen1^ON immunoprecipitates is shown on the left panels. HJ resolution assay is shown on the right panel. Quantification of resolution efficiency is displayed at the bottom. Resolution data arise from two independent experiments.

Figure 7. Analysis of recombination intermediates in the *cdc20-md* mutant reveals an early meiotic role for Cdc14 and Yen1. (A) Representative Southern blots depicting 2D-gels at the

HIS4LEU2 hotspot in *CDC14 cdc20-md sgs1-md* (JCY2480), *cdc20-md sgs1-md yen1* (JCY2469), *cdc20-md sgs1-md cdc14-md* (JCY2508), *cdc20-md sgs1-md mms4* (JCY2480), *cdc20-md sgs1-md mms4 yen1* (JCY2478) and *cdc20-md sgs1-md mms4 cdc14-md* (JCY2502). Arrows mark IH-dHJs and lines mark mc-JMs (see Fig 4A-B). (B) Quantification of total JMs in the strains shown in (A) from several independent gels. Error bars display SEM over the mean values plotted. (C) Representative 1D-gel Southern blot images for analysis of crossovers at the *HIS4LEU2* hotspot for all strains shown in (A). (D) Quantification of COs from at least three different image acquisitions like the one depicted in (C). Error bars represent SEM over the calculated mean value.

Figure 8. Model for *CDC14*-dependent resolution of recombination intermediates via multiple mechanisms. Contribution of Cdc14 to the correct disjunction of recombining chromosomes during meiosis. Early in prophase I, chromosomes initiate homologous recombination. 3'-resected ssDNA overhangs invade an intact template with the help of recombinases. Displacement of the intact strand from the template allows the formation of D-loops, which can be stabilized allowing DNA synthesis. Next, stabilized branched DNA molecules might be disrupted by the action of the anti-recombinases. Reannealing of the extended 3'-ssDNA overhang with the resected complementary strand, followed by further DNA synthesis will lead to the repair and ligation of the broken DNA duplex, giving rise to NCOs via SDSA. ZMM stabilization of JMs is followed by the resolution of dHJs through the MutLγ, class I, dedicated CO pathway. Ndt80-dependent expression, and activation, of Cdc5 triggers MutLγ resolvase activity. Unresolved linkages between bivalents that persist until metaphase/anaphase I are mostly resolved by the action of the SSEs, Mus81-Mms4. Slx1-Slx4, Top3-Rmi1 as well as Cdc14/Yen1 also contribute to the correct resolution of chromosomal entanglements between homologs during MI. Residual chromatid intertwining between sister chromatids during the second meiotic division will be removed by the action of Cdc14/Yen1. Gradual implementation of Yen1 activity during both divisions by Cdc14 will transfer Yen1 inactive population to its active/nuclear-enriched form.

Supplemental material legends

Figure S1. Cdc14 is required during meiosis for correct nuclear division. (A) *cdc14-HA* cells (JCY844) sustain both nuclear divisions although with subtly slower kinetics than wild type cells (JCY840) predominantly during the second round of nuclear segregation. *cdc14-md* cells (JCY2376) do not undergo the second nuclear division (B) Quantification of the number of nuclei as well as other aberrant nuclear structures during meiosis in *CDC14* (JCY840) and *cdc14-HA* (JCY844) cells. (C) Representative images of all nuclear structures quantified in (B).

Figure S2. Reduced levels of Cdc14 causes the sporulation defect observed in the *cdc14-HA* mutant. (A) Analysis of Cdc14 protein levels in wild-type (JCY902), and *cdc14-HA* (JCY904) cells. Top panel shows immunodetection using a polyclonal α-Cdc14 (yE-17) antibody. The same blot was reanalyzed using α-HA (12CA5) (bottom panel). (*) denotes residual signal from the earlier α-Cdc14 detection. *SMK1-3HA* is also expressed in both *CDC14* and *cdc14-HA* strains. (B) Quantification of Cdc14 protein levels shown in (B). Values were normalized against the maximum signal corresponding to the 12 h time-point in the wild type (C) Sporulation efficiency of *cdc14-HA* diploids transformed with multi-copy plasmids carrying MEN genes (GGY102/GGY103/GGY104/GGY105) and *cdc14-HA* (GGY93). Only overproduction of the Cdc14 phosphatase rescues the sporulation defect. (D) Di-tyrosine autofluorescence of different mutant combinations as well as the control strains grown and sporulated on plates at 25ºC and 30ºC. *cdc14-1* homozygous diploids (JCY2353) sporulate at high efficiency under semi-permissive temperature forming preferentially tetrads whereas *cdc14-HA* homozygous diploids do not sporulate at any temperature (JCY840). Combinations, and variable copy number, of the mutant genes can rescue the sporulation defect at different degrees (JCY2365/JCY2354/JCY2356).

Figure S3. Meiotic *cdc14* mutant cells re-duplicate their SPBs and assemble meiosis I and meiosis II spindles. (A) Representative images of fixed cells at different time-points during synchronous parallel time courses for both *CDC14* (JCY892) and *cdc14* mutant (JCY893) cells. *cdc14-HA* cells initiate both meiotic divisions, as visualized by different fluorescence markers, but they do not complete sporulation. (B) Some *cdc14-HA* (JCY893) cells display abnormal number of SPBs and atypical spindle conformations. (C) Example of the terminal phenotype of a post-meiotic *cdc14-HA* (JCY893) cell. (D) *cdc14-md* cells display strong SPB re-duplication defects and arrest with two nuclei.

Figure S4. Meiotic spindle lifespan and dynamics are not altered in the meiosis-deficient *cdc14-HA* mutant. (A) Live-imaging of wild-type and *cdc14-HA* cells (GGY53 and GGY54, respectively) undergoing the first meiotic division carrying GFP-tubulin (red) and Spc29-CFP (green). (B) Visualization of spindle and SPB dynamics in the same strains (GGY53 and GGY54) undergoing the second meiotic division. For both A and B, microscopy fields from at least four different positions and from two separate wells were analyzed. A minimum of three time-courses per strain were run, acquired images were processed and movies generated (Video S1 and S2). Representative cells are shown. (C) Quantification of spindle lifespan from cells undergoing MI or MII. Meiotic deficient *cdc14-HA* cells (GGY54) show similar spindle dynamics compared to the wild type when completing both meiotic divisions. Box plots depict median number of spindle lifespan with whiskers representing upper and lower 1.5 interquartile range. Black dots represent outliers. Statistical test was performed using a two-tailed unpaired *t*-test. (D) Example of a meiotic *cdc14-HA* (JCY2404) cell transiting from prophase I to complete both meiotic divisions. Rec8-GFP (yellow) and Pds1-tdTomato/Cnm67-tdTomato (magenta).

Time 0 (min) was considered the last frame where Pds1 was visualized and SPBs stayed in close association. Data obtained from Video S3 and S4.

Figure S5. Lack of Cdc14 in meiosis causes missegregation of chromosomes at both meiotic divisions. (A) Scheme depicting the chromosome GFP-tagging system placed at an interstitial *locus* within chromosome II (left) and at a centromere proximal *locus* within chromosome IV (right). (B) Representative images and frequencies of the observed types of distribution of interstitial chromosome II GFP dots in homozygosis of both *cdc14-HA* mutant (JCY2230) and wild-type (JCY2231) meiotic cells. Arrowheads indicate diffuse GFP in the nucleus. For chromosome segregation fidelity analysis (Fig 3D), different percentages from depicted were calculated from considering only cells with four nuclei (type 1, 2 and 3). . Cells were fixed in formaldehyde and nuclei stained with DAPI. Bar=5µm.

Figure S6. Accumulation of JMs is analogous in *cdc14-HA* in the presence or absence of Mus81. (A; left) Representative 2D gels showing recombination intermediates in *mus81 ndt80*, and *mus81 cdc14-HA ndt80* prophase-arrested cells 8 h after meiotic induction. (A; right) Quantification of DNA species from (2D-gels). (B) Comparison of JM abundance and resolution from the Southern blot shown in (Fig 5A). Asterisk indicates meiosis-specific non-characterized recombination products.

Table S1. *S. cerevisiae* strains used in this study. SK1 background strains were used throughout the study, unless otherwise specified.

Video S1. Meiotic spindle lifespan and dynamics in *CDC14* cells. Live imaging of cells (GGY53) undergoing both meiotic divisions carrying GFP-tubulin (red) and Spc29-CFP (green).

Video S2. Meiotic spindle lifespan and dynamics in *cdc14-HA* cells. Live imaging of cells (GGY54) undergoing both meiotic divisions carrying GFP-tubulin (red) and Spc29-CFP (green).

Video S3. Kinetics of Rec8-GFP association/removal in *CDC14* cells. Sequence of Rec8-GFP wild-type cells (JCY2406) at 15-minute intervals displaying kinetics of cohesin assembly and removal.

Video S4. Kinetics of Rec8-GFP association/removal in *cdc14-HA* cells. Sequence of Rec8-GFP *cdc14-HA* cells (JCY2404) at 15 minutes interval displaying kinetics of cohesin removal.

References

Ahuja JS, Borner GV. 2011. Analysis of meiotic recombination intermediates by two-dimensional gel
electrophoresis. *Methods Mol Biol* **745**: 99-116.

Allers T, Lichten M. 2001. Differential timing and control of noncrossover and crossover recombination
during meiosis. *Cell* **106**: 47-57.

Arguello-Miranda O, Zagoriy I, Mengoli V, Rojas J, Jonak K, Oz T, Graf P, Zachariae W. 2017. Casein
Kinase 1 Coordinates Cohesin Cleavage, Gametogenesis, and Exit from M Phase in Meiosis II.
Dev Cell **40**: 37-52.

Arter M, Hurtado-Nieves V, Oke A, Zhuge T, Wettstein R, Fung JC, Blanco MG, Matos J. 2018.
Regulated Crossing-Over Requires Inactivation of Yen1/GEN1 Resolvase during Meiotic
Prophase I. *Dev Cell* **45**: 785-800 e786.

Attner MA, Amon A. 2012. Control of the mitotic exit network during meiosis. *Mol Biol Cell* **23**: 3122-
3132.

Azzam R, Chen SL, Shou W, Mah AS, Alexandru G, Nasmyth K, Annan RS, Carr SA, Deshaies RJ.
2004. Phosphorylation by cyclin B-Cdk underlies release of mitotic exit activator Cdc14 from
the nucleolus. *Science* **305**: 516-519.

Bassermann F, Frescas D, Guardavaccaro D, Busino L, Peschiaroli A, Pagano M. 2008. The Cdc14B-
Cdh1-Plk1 axis controls the G2 DNA-damage-response checkpoint. *Cell* **134**: 256-267.

Bishop DK. 2006. Multiple mechanisms of meiotic recombination. *Cell* **127**: 1095-1097.

Bizzari F, Marston AL. 2011. Cdc55 coordinates spindle assembly and chromosome disjunction during
meiosis. *J Cell Biol* **193**: 1213-1228.

Blanco MG, Matos J, West SC. 2014. Dual control of Yen1 nuclease activity and cellular localization
by Cdk and Cdc14 prevents genome instability. *Mol Cell* **54**: 94-106.

Bloom J, Cross FR. 2007. Novel role for Cdc14 sequestration: Cdc14 dephosphorylates factors that
promote DNA replication. *Mol Cell Biol* **27**: 842-853.

881 Borner GV, Kleckner N, Hunter N. 2004. Crossover/noncrossover differentiation, synaptonemal
882  complex formation, and regulatory surveillance at the leptotene/zygotene transition of meiosis.
883  *Cell* **117**: 29-45.

884 Briza P, Winkler G, Kalchhauser H, Breitenbach M. 1986. Dityrosine is a prominent component of the
885  yeast ascospore wall. A proof of its structure. *J Biol Chem* **261**: 4288-4294.

886 Buonomo SB, Rabitsch KP, Fuchs J, Gruber S, Sullivan M, Uhlmann F, Petronczki M, Toth A, Nasmyth
887  K. 2003. Division of the nucleolus and its release of CDC14 during anaphase of meiosis I
888  depends on separase, SPO12, and SLK19. *Dev Cell* **4**: 727-739.

889 Bzymek M, Thayer NH, Oh SD, Kleckner N, Hunter N. 2010. Double Holliday junctions are
890  intermediates of DNA break repair. *Nature* **464**: 937-941.

891 Cannavo E, Sanchez A, Anand R, Ranjha L, Hugener J, Adam C, Acharya A, Weyland N, Aran-Guiu
892  X, Charbonnier JB et al. 2020. Regulation of the MLH1-MLH3 endonuclease in meiosis.
893  *Nature* **586**: 618-622.

894 Cao L, Alani E, Kleckner N. 1990. A pathway for generation and processing of double-strand breaks
895  during meiotic recombination in S. cerevisiae. *Cell* **61**: 1089-1101.

896 Carballo JA, Panizza S, Serrentino ME, Johnson AL, Geymonat M, Borde V, Klein F, Cha RS. 2013.
897  Budding yeast ATM/ATR control meiotic double-strand break (DSB) levels by down-
898  regulating Rec114, an essential component of the DSB-machinery. *PLoS Genet* **9**: e1003545.

899 Carlile TM, Amon A. 2008. Meiosis I is established through division-specific translational control of a
900  cyclin. *Cell* **133**: 280-291.

901 Clemente-Blanco A, Mayan-Santos M, Schneider DA, Machin F, Jarmuz A, Tschochner H, Aragon L.
902  2009. Cdc14 inhibits transcription by RNA polymerase I during anaphase. *Nature* **458**: 219-
903  222.

904 Clemente-Blanco A, Sen N, Mayan-Santos M, Sacristan MP, Graham B, Jarmuz A, Giess A, Webb E,
905  Game L, Eick D et al. 2011. Cdc14 phosphatase promotes segregation of telomeres through
906  repression of RNA polymerase II transcription. *Nat Cell Biol* **13**: 1450-1456.

Clyne RK, Katis VL, Jessop L, Benjamin KR, Herskowitz I, Lichten M, Nasmyth K. 2003. Polo-like
kinase Cdc5 promotes chiasmata formation and cosegregation of sister centromeres at meiosis
I. *Nat Cell Biol* **5**: 480-485.

Copsey A, Tang S, Jordan PW, Blitzblau HG, Newcombe S, Chan AC, Newnham L, Li Z, Gray S,
Herbert AD et al. 2013. Smc5/6 coordinates formation and resolution of joint molecules with
chromosome morphology to ensure meiotic divisions. *PLoS Genet* **9**: e1004071.

Cromie GA, Hyppa RW, Taylor AF, Zakharyevich K, Hunter N, Smith GR. 2006. Single Holliday
junctions are intermediates of meiotic recombination. *Cell* **127**: 1167-1178.

Culotti J, Hartwell LH. 1971. Genetic control of the cell division cycle in yeast. 3. Seven genes
controlling nuclear division. *Exp Cell Res* **67**: 389-401.

D'Amours D, Stegmeier F, Amon A. 2004. Cdc14 and condensin control the dissolution of cohesin-
independent chromosome linkages at repeated DNA. *Cell* **117**: 455-469.

de los Santos T, Loidl J, Larkin B, Hollingsworth NM. 2001. A role for MMS4 in the processing of
recombination intermediates during meiosis in Saccharomyces cerevisiae. *Genetics* **159**: 1511-
1525.

De Muyt A, Jessop L, Kolar E, Sourirajan A, Chen J, Dayani Y, Lichten M. 2012. BLM helicase
ortholog Sgs1 is a central regulator of meiotic recombination intermediate metabolism. *Mol
Cell* **46**: 43-53.

Diaz-Cuervo H, Bueno A. 2008. Cds1 controls the release of Cdc14-like phosphatase Flp1 from the
nucleolus to drive full activation of the checkpoint response to replication stress in fission yeast.
Mol Biol Cell **19**: 2488-2499.

Dulev S, de Renty C, Mehta R, Minkov I, Schwob E, Strunnikov A. 2009. Essential global role of
CDC14 in DNA synthesis revealed by chromosome underreplication unrecognized by
checkpoints in cdc14 mutants. *Proc Natl Acad Sci U S A* **106**: 14466-14471.

Eissler CL, Mazon G, Powers BL, Savinov SN, Symington LS, Hall MC. 2014. The Cdk/cDc14 module
controls activation of the Yen1 holliday junction resolvase to promote genome stability. *Mol
Cell* **54**: 80-93.

Fox C, Zou J, Rappsilber J, Marston AL. 2017. Cdc14 phosphatase directs centrosome re-duplication
at the meiosis I to meiosis II transition in budding yeast. *Wellcome open research* **2**: 2.

Garcia-Luis J, Clemente-Blanco A, Aragon L, Machin F. 2014. Cdc14 targets the Holliday junction
resolvase Yen1 to the nucleus in early anaphase. *Cell Cycle* **13**: 1392-1399.

Gimenez-Abian MI, Rozalen AE, Carballo JA, Botella LM, Pincheira J, Lopez-Saez JF, de la Torre C.
2004. HSP90 and checkpoint-dependent lengthening of the G2 phase observed in plant cells
under hypoxia and cold. *Protoplasma* **223**: 191-196.

Grigaitis R, Ranjha L, Wild P, Kasaciunaite K, Ceppi I, Kissling V, Henggeler A, Susperregui A, Peter
M, Seidel R et al. 2020. Phosphorylation of the RecQ Helicase Sgs1/BLM Controls Its DNA
Unwinding Activity during Meiosis and Mitosis. *Dev Cell* **53**: 706-723 e705.

Grigaitis R, Susperregui A, Wild P, Matos J. 2018. Characterization of DNA helicases and nucleases
from meiotic extracts of S. cerevisiae. *Methods Cell Biol* **144**: 371-388.

Hartwell LH, Mortimer RK, Culotti J, Culotti M. 1973. Genetic Control of the Cell Division Cycle in
Yeast: V. Genetic Analysis of cdc Mutants. *Genetics* **74**: 267-286.

Hartwell LH, Smith D. 1985. Altered fidelity of mitotic chromosome transmission in cell cycle mutants
of S. cerevisiae. *Genetics* **110**: 381-395.

Hunter N, Kleckner N. 2001. The single-end invasion: an asymmetric intermediate at the double-strand
break to double-holliday junction transition of meiotic recombination. *Cell* **106**: 59-70.

Imtiaz A, Belyantseva IA, Beirl AJ, Fenollar-Ferrer C, Bashir R, Bukhari I, Bouzid A, Shaukat U,
Azaiez H, Booth KT et al. 2018. CDC14A phosphatase is essential for hearing and male fertility
in mouse and human. *Human molecular genetics* **27**: 780-798.

Ip SC, Rass U, Blanco MG, Flynn HR, Skehel JM, West SC. 2008. Identification of Holliday junction
resolvases from humans and yeast. *Nature* **456**: 357-361.

Jaspersen SL, Charles JF, Tinker-Kulberg RL, Morgan DO. 1998. A late mitotic regulatory network
controlling cyclin destruction in Saccharomyces cerevisiae. *Mol Biol Cell* **9**: 2803-2817.

Jessop L, Lichten M. 2008. Mus81/Mms4 endonuclease and Sgs1 helicase collaborate to ensure proper
recombination intermediate metabolism during meiosis. *Mol Cell* **31**: 313-323.

Kamieniecki RJ, Liu L, Dawson DS. 2005. FEAR but not MEN genes are required for exit from meiosis
I. *Cell Cycle* **4**: 1093-1098.

Kamieniecki RJ, Shanks RM, Dawson DS. 2000. Slk19p is necessary to prevent separation of sister
chromatids in meiosis I. *Curr Biol* **10**: 1182-1190.

Katis VL, Lipp JJ, Imre R, Bogdanova A, Okaz E, Habermann B, Mechtler K, Nasmyth K, Zachariae
W. 2010. Rec8 phosphorylation by casein kinase 1 and Cdc7-Dbf4 kinase regulates cohesin
cleavage by separase during meiosis. *Dev Cell* **18**: 397-409.

Katis VL, Matos J, Mori S, Shirahige K, Zachariae W, Nasmyth K. 2004. Spo13 facilitates monopolin
recruitment to kinetochores and regulates maintenance of centromeric cohesion during yeast
meiosis. *Curr Biol* **14**: 2183-2196.

Keeney S, Giroux CN, Kleckner N. 1997. Meiosis-specific DNA double-strand breaks are catalyzed by
Spo11, a member of a widely conserved protein family. *Cell* **88**: 375-384.

Kerr GW, Sarkar S, Tibbles KL, Petronczki M, Millar JB, Arumugam P. 2011. Meiotic nuclear
divisions in budding yeast require PP2A(Cdc55)-mediated antagonism of Net1 phosphorylation
by Cdk. *J Cell Biol* **193**: 1157-1166.

Kosugi S, Hasebe M, Tomita M, Yanagawa H. 2009. Systematic identification of cell cycle-dependent
yeast nucleocytoplasmic shuttling proteins by prediction of composite motifs. *Proc Natl Acad*
Sci U S A **106**: 10171-10176.

Kulkarni DS, Owens SN, Honda M, Ito M, Yang Y, Corrigan MW, Chen L, Quan AL, Hunter N. 2020.
PCNA activates the MutLgamma endonuclease to promote meiotic crossing over. *Nature* **586**:
623-627.

Lee BH, Amon A. 2003. Role of Polo-like kinase CDC5 in programming meiosis I chromosome
segregation. *Science* **300**: 482-486.

Li L, Ernsting BR, Wishart MJ, Lohse DL, Dixon JE. 1997. A family of putative tumor suppressors is
structurally and functionally conserved in humans and yeast. *J Biol Chem* **272**: 29403-29406.

Liberi G, Maffioletti G, Lucca C, Chiolo I, Baryshnikova A, Cotta-Ramusino C, Lopes M, Pellicioli A,
Haber JE, Foiani M. 2005. Rad51-dependent DNA structures accumulate at damaged
replication forks in sgs1 mutants defective in the yeast ortholog of BLM RecQ helicase. *Genes*
Dev **19**: 339-350.

Longtine MS, McKenzie A, 3rd, Demarini DJ, Shah NG, Wach A, Brachat A, Philippsen P, Pringle JR.
1998. Additional modules for versatile and economical PCR-based gene deletion and
modification in Saccharomyces cerevisiae. *Yeast* **14**: 953-961.

Machin F. 2020. Implications of Metastable Nicks and Nicked Holliday Junctions in Processing Joint
Molecules in Mitosis and Meiosis. *Genes* **11**.

Machin F, Torres-Rosell J, De Piccoli G, Carballo JA, Cha RS, Jarmuz A, Aragon L. 2006.
Transcription of ribosomal genes can cause nondisjunction. *J Cell Biol* **173**: 893-903.

Mankouri HW, Ashton TM, Hickson ID. 2011. Holliday junction-containing DNA structures persist in
cells lacking Sgs1 or Top3 following exposure to DNA damage. *Proc Natl Acad Sci U S A* **108**:
4944-4949.

Marston AL, Lee BH, Amon A. 2003. The Cdc14 phosphatase and the FEAR network control meiotic
spindle disassembly and chromosome segregation. *Dev Cell* **4**: 711-726.

Martini E, Diaz RL, Hunter N, Keeney S. 2006. Crossover homeostasis in yeast meiosis. *Cell* **126**: 285-
295.

Matos J, Blanco MG, Maslen S, Skehel JM, West SC. 2011. Regulatory control of the resolution of
DNA recombination intermediates during meiosis and mitosis. *Cell* **147**: 158-172.

Michaelis C, Ciosk R, Nasmyth K. 1997. Cohesins: Chromosomal Proteins that Prevent Premature
Separation of Sister Chromatids. *Cell* **91**: 35-45.

Mocciaro A, Berdougo E, Zeng K, Black E, Vagnarelli P, Earnshaw W, Gillespie D, Jallepalli P,
Schiebel E. 2010. Vertebrate cells genetically deficient for Cdc14A or Cdc14B retain DNA
damage checkpoint proficiency but are impaired in DNA repair. *J Cell Biol* **189**: 631-639.

Mocciaro A, Schiebel E. 2010. Cdc14: a highly conserved family of phosphatases with non-conserved
functions? *J Cell Sci* **123**: 2867-2876.

Newnham L, Jordan PW, Carballo JA, Newcombe S, Hoffmann E. 2013. Ipl1/Aurora kinase suppresses S-CDK-driven spindle formation during prophase I to ensure chromosome integrity during meiosis. *PLoS One* **8**: e83982.

Nolt JK, Rice LM, Gallo-Ebert C, Bisher ME, Nickels JT. 2011. PP2A (Cdc)(5)(5) is required for multiple events during meiosis I. *Cell Cycle* **10**: 1420-1434.

Oh SD, Lao JP, Hwang PY, Taylor AF, Smith GR, Hunter N. 2007. BLM ortholog, Sgs1, prevents aberrant crossing-over by suppressing formation of multichromatid joint molecules. *Cell* **130**: 259-272.

Oh SD, Lao JP, Taylor AF, Smith GR, Hunter N. 2008. RecQ helicase, Sgs1, and XPF family endonuclease, Mus81-Mms4, resolve aberrant joint molecules during meiotic recombination. *Mol Cell* **31**: 324-336.

Pablo-Hernando ME, Arnaiz-Pita Y, Nakanishi H, Dawson D, del Rey F, Neiman AM, Vazquez de Aldana CR. 2007. Cdc15 is required for spore morphogenesis independently of Cdc14 in Saccharomyces cerevisiae. *Genetics* **177**: 281-293.

Padmore R, Cao L, Kleckner N. 1991. Temporal comparison of recombination and synaptonemal complex formation during meiosis in S. cerevisiae. *Cell* **66**: 1239-1256.

Piazza A, Wright WD, Heyer WD. 2017. Multi-invasions Are Recombination Byproducts that Induce Chromosomal Rearrangements. *Cell* **170**: 760-773 e715.

Queralt E, Lehane C, Novak B, Uhlmann F. 2006. Downregulation of PP2A(Cdc55) phosphatase by separase initiates mitotic exit in budding yeast. *Cell* **125**: 719-732.

Rahal R, Amon A. 2008. The Polo-like kinase Cdc5 interacts with FEAR network components and Cdc14. *Cell Cycle* **7**: 3262-3272.

Rodriguez-Rodriguez JA, Moyano Y, Jativa S, Queralt E. 2016. Mitotic Exit Function of Polo-like Kinase Cdc5 Is Dependent on Sequential Activation by Cdk1. *Cell reports* **15**: 2050-2062.

Rosso L, Marques AC, Weier M, Lambert N, Lambot MA, Vanderhaeghen P, Kaessmann H. 2008. Birth and rapid subcellular adaptation of a hominoid-specific CDC14 protein. *PLoS Biol* **6**: e140.

San-Segundo PA, Clemente-Blanco A. 2020. Resolvases, Dissolvases, and Helicases in Homologous
Recombination: Clearing the Road for Chromosome Segregation. *Genes* **11**.

Sanchez A, Adam C, Rauh F, Duroc Y, Ranjha L, Lombard B, Mu X, Wintrebert M, Loew D, Guarne
A et al. 2020. Exo1 recruits Cdc5 polo kinase to MutLgamma to ensure efficient meiotic
crossover formation. *Proc Natl Acad Sci U S A* **117**: 30577-30588.

Schild D, Byers B. 1980. Diploid spore formation and other meiotic effects of two cell-division-cycle
mutations of Saccharomyces cerevisiae. *Genetics* **96**: 859-876.

Schwacha A, Kleckner N. 1994. Identification of joint molecules that form frequently between
homologs but rarely between sister chromatids during yeast meiosis. *Cell* **76**: 51-63.

-. 1995. Identification of double Holliday junctions as intermediates in meiotic recombination. *Cell* **83**:
783-791.

Sharon G, Simchen G. 1990. Mixed segregation of chromosomes during single-division meiosis of
Saccharomyces cerevisiae. *Genetics* **125**: 475-485.

Shinohara M, Oh SD, Hunter N, Shinohara A. 2008. Crossover assurance and crossover interference
are distinctly regulated by the ZMM proteins during yeast meiosis. *Nat Genet* **40**: 299-309.

Shodhan A, Medhi D, Lichten M. 2019. Noncanonical Contributions of MutLgamma to VDE-Initiated
Crossovers During Saccharomyces cerevisiae Meiosis. *G3 (Bethesda, Md)* **9**: 1647-1654.

Shonn MA, McCarroll R, Murray AW. 2002. Spo13 protects meiotic cohesin at centromeres in meiosis
I. *Genes Dev* **16**: 1659-1671.

Shou W, Seol JH, Shevchenko A, Baskerville C, Moazed D, Chen ZW, Jang J, Shevchenko A,
Charbonneau H, Deshaies RJ. 1999. Exit from mitosis is triggered by Tem1-dependent release
of the protein phosphatase Cdc14 from nucleolar RENT complex. *Cell* **97**: 233-244.

Sourirajan A, Lichten M. 2008. Polo-like kinase Cdc5 drives exit from pachytene during budding yeast
meiosis. *Genes Dev* **22**: 2627-2632.

Stegmeier F, Visintin R, Amon A. 2002. Separase, polo kinase, the kinetochore protein Slk19, and
1065   Spo12 function in a network that controls Cdc14 localization during early anaphase. *Cell* **108**:
1066   207-220.

1067   Straight AF, Belmont AS, Robinett CC, Murray AW. 1996. GFP tagging of budding yeast
1068   chromosomes reveals that protein-protein interactions can mediate sister chromatid cohesion.
1069   *Curr Biol* **6**: 1599-1608.

1070   Sullivan M, Higuchi T, Katis VL, Uhlmann F. 2004. Cdc14 phosphatase induces rDNA condensation
1071   and resolves cohesin-independent cohesion during budding yeast anaphase. *Cell* **117**: 471-482.

1072   Talhaoui I, Bernal M, Mullen JR, Dorison H, Palancade B, Brill SJ, Mazon G. 2018. Slx5-Slx8 ubiquitin
1073   ligase targets active pools of the Yen1 nuclease to limit crossover formation. *Nat Commun* **9**:
1074   5016.

1075   Tang S, Wu MKY, Zhang R, Hunter N. 2015. Pervasive and essential roles of the Top3-Rmi1
1076   decatenase orchestrate recombination and facilitate chromosome segregation in meiosis. *Mol*
1077   *Cell* **57**: 607-621.

1078   Torres-Rosell J, Machin F, Jarmuz A, Aragon L. 2004. Nucleolar segregation lags behind the rest of
1079   the genome and requires Cdc14p activation by the FEAR network. *Cell Cycle* **3**: 496-502.

1080   Trautmann S, McCollum D. 2002. Cell cycle: new functions for Cdc14 family phosphatases. *Curr Biol*
1081   **12**: R733-735.

1082   Tsuchiya D, Yang Y, Lacefield S. 2014. Positive feedback of NDT80 expression ensures irreversible
1083   meiotic commitment in budding yeast. *PLoS Genet* **10**: e1004398.

1084   Villoria MT, Ramos F, Duenas E, Faull P, Cutillas PR, Clemente-Blanco A. 2017. Stabilization of the
1085   metaphase spindle by Cdc14 is required for recombinational DNA repair. *EMBO J* **36**: 79-101.

1086   Visintin C, Tomson BN, Rahal R, Paulson J, Cohen M, Taunton J, Amon A, Visintin R. 2008. APC/C-
1087   Cdh1-mediated degradation of the Polo kinase Cdc5 promotes the return of Cdc14 into the
1088   nucleolus. *Genes Dev* **22**: 79-90.

1089   Visintin R, Craig K, Hwang ES, Prinz S, Tyers M, Amon A. 1998. The phosphatase Cdc14 triggers
1090   mitotic exit by reversal of Cdk-dependent phosphorylation. *Mol Cell* **2**: 709-718.

Visintin R, Hwang ES, Amon A. 1999. Cfi1 prevents premature exit from mitosis by anchoring Cdc14
phosphatase in the nucleolus. *Nature* **398**: 818-823.

Visintin R, Stegmeier F, Amon A. 2003. The role of the polo kinase Cdc5 in controlling Cdc14
localization. *Mol Biol Cell* **14**: 4486-4498.

Wang BD, Yong-Gonzalez V, Strunnikov AV. 2004. Cdc14p/FEAR pathway controls segregation of
nucleolus in S. cerevisiae by facilitating condensin targeting to rDNA chromatin in anaphase.
Cell Cycle **3**: 960-967.

Wild P, Susperregui A, Piazza I, Dorig C, Oke A, Arter M, Yamaguchi M, Hilditch AT, Vuina K, Chan
KC et al. 2019. Network Rewiring of Homologous Recombination Enzymes during Mitotic
Proliferation and Meiosis. *Mol Cell* **75**: 859-874 e854.

Xaver M, Huang L, Chen D, Klein F. 2013. Smc5/6-Mms21 prevents and eliminates inappropriate
recombination intermediates in meiosis. *PLoS Genet* **9**: e1004067.

Yellman CM, Roeder GS. 2015. Cdc14 Early Anaphase Release, FEAR, Is Limited to the Nucleus and
Dispensable for Efficient Mitotic Exit. *PLoS One* **10**: e0128604.

Yoshida S, Toh-e A. 2002. Budding yeast Cdc5 phosphorylates Net1 and assists Cdc14 release from
the nucleolus. *Biochem Biophys Res Commun* **294**: 687-691.

Zakharyevich K, Ma Y, Tang S, Hwang PY, Boiteux S, Hunter N. 2010. Temporally and biochemically
distinct activities of Exo1 during meiosis: double-strand break resection and resolution of
double Holliday junctions. *Mol Cell* **40**: 1001-1015.

Zakharyevich K, Tang S, Ma Y, Hunter N. 2012. Delineation of joint molecule resolution pathways in
meiosis identifies a crossover-specific resolvase. *Cell* **149**: 334-347.

Zeng X, Saunders WS. 2000. The Saccharomyces cerevisiae centromere protein Slk19p is required for
two successive divisions during meiosis. *Genetics* **155**: 577-587.

Zhou X, Li W, Liu Y, Amon A. 2021. Cross-compartment signal propagation in the mitotic exit
network. *Elife* **10**.

1116

A

B

C

D

A

	Anaphase I			Metaphase II	
Genotype	i	ii	iii	iv	v
WT	74.3%	22.6%	3.1%	76.3%	24.7%
cdc14-HA	19.6%	67.3%	13.1%	24.6%	75.3%
cdc14-md	20.5%	71.1%	8.4%	N.A.	N.A.
spo11-Y135F	77.1%	20.0%	2.9%	N.A.	N.A.
spo11-Y135F cdc14-HA	61.4%	34.5%	4.1%	N.A.	N.A.

Table S1. *S. cerevisiae* strains used in this study

Strain	Genotype
JCY840	*MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", trp1::hisG/", arg4-nsp/", his4x/"*
JCY844	*MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", trp1::hisG/", arg4-nsp/", his4x/", cdc14-3HA::KanMX4/"*
GGY92	*MAT a/α ho::hisG/LYS2, lys2, ura3(Dsma-pst::hisG)/ura3, HIS4::LEU2-(NBam;ori)/his4x, Zip1-GFP(at AA700)/", leu2::hisG/", cdc14-3HA::KanMX4/CDC14*
GGY91	*MAT a/α ho::hisG/LYS2, lys2, ura3(Dsma-pst::hisG)/ura3, HIS4::LEU2-(NBam;ori)/his4X or LEU2(NgomIV;ori)-URA3/his4x, Zip1-GFP(at AA700), leu2::hisG/", cdc14-3HA::KanMX4/"*
JCY902	*MAT a/α ho::hisG/", ura3/", leu2::hisG/+, lys2/", trp1/", SPC29-CFP::TRP1/", DON1-dGFP-KanMX/", pSMK1-3HA-URA3*
JCY904	As JCY902 but *cdc14-3HA::KanMX4/"*
GGY53	*MAT a/α ho::hisG/", ura3/", leu2::hisG/+, lys2/", trp1/", SPC29-CFP::TRP1/", TUB-dGFP-URA3/"*
GGY54	*MAT a/α ho::hisG/", ura3/", leu2::hisG/+, lys2/", trp1/", SPC29-CFP::TRP1/", TUB-dGFP-URA3/", cdc14-3HA::KanMX4/"*
JCY2196	*MAT a/α ho::hisG/", lys2/", ura3/", trp1/", his4/", leu2::hisG/", cdc14-3HA::KanMX4/"*
PRY151	*MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", trp1::hisG/", spo11-Y135F-HA::URA3/", pCLB2::3HA-CDC14::KanMX4/"*
JCY2247	As JCY2235 but *cdc14-3HA::KanMX4/"*
JCY2270	*MAT a/α ho::LYS2/", lys2/", leu2::hisG/", ura3/", arg4-nsp/", trp1::hisG/", his4x/", spo11-Y135F-HA::URA3/"*
JCY2280	As JCY2270 but *cdc14-3HA::KanMX4/"*
JCY2232	*MAT a/α ho::hisG/", lys2/", ura3(Δsma-pst::hisG)/", leu2::hisG/", his4X::LEU2-(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori)*
JCY2231	As JCY2232 but *cdc14-3HA::KanMX4/"*
YJM7692	*MAT a/α ho::LYS2/" ura3/" leu2::hisG/" trp1::hisG/" his3::hisG/" YEN1-myc9::URA3/"*
YML7693	As YML3952 but *cdc14-3HA::KanMX4/"*

JCY2164   *MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", trp1/",*
               his4x::LEU2-URA3/", YEN1-9A-myc18::URA3/", cdc14-
               3HA::KanMX4/", ZIP1-eGFP(at AA700)/"

GGY102   JCY844 with pSJ103 (*CDC15HA3-URA3)*

GGY93   JCY844 with pPD.2 (*CDC14HA3-URA3)*

GGY103   *JCY844 with pSJ57 (HA3-DBF2-URA3)*

GGY104   JCY844 with pSJ56 (*TEM1HA3-URA3)*

GGY105   JCY844 with pSJ29 (*HACDC5-URA3)*

JCY2353   *MAT a/α ho::LYS2/", lys2/", his4x/", leu2::hisG/", ura3/",*
               cdc14Δ::KanMX4/", trp1::cdc14-1::TRP1::LEU2/"

JCY2365   *MAT a/α ho::LYS2/", lys2/", his4x/", leu2::hisG/", ura3/",*
               cdc14Δ::KanMX4/cdc14-3HA::KanMX4, trp1::cdc14-
               1::TRP1::LEU2/trp1 arg4-nsp/ARG4

JCY2354   *MAT a/α ho::LYS2/", lys2/", his4x/", leu2::hisG/", ura3/", cdc14-*
               3HA::KanMX4/cdc14Δ::KanMX4, trp1::cdc14-1::TRP1::LEU2/"

JCY2356   *MAT a/α ho::LYS2/", lys2/", his4x/", leu2::hisG/", ura3/", cdc14-*
               3HA::KanMX4/cdc14-3HA::KanMX4, trp1::cdc14-1::TRP1::LEU2/"

JCY2222   *MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", his4/", Cyc1p-LacI-GFP-*
               URA3/", trp1::LacO-TRP1 (at CENIV)/"

JCY2224   As JCY2222 but *cdc14-3HA::KanMX4/"*

JCY2284   *MAT a/α ho::LYS2/", lys2/", ura3/", leu2::hisG/", his4/", Cyc1p-LacI-GFP-URA3/ura3,*
               trp1::LacO-TRP1 (at CENIV)/trp1

JCY2286   As JCY2284 but *cdc14-3HA::KanMX4/"*

JCY2327   *MATa/α ho::LYS2/" his4/" ura3::Cyc1p-LacI-GFP-URA3/" trp1::LacO-TRP1/"*
               CNM67-mCherry::natMX4/CNM67

JCY2326   *MATa/α ho::LYS2/" his4/" ura3::Cyc1p-LacI-GFP-URA3/" trp1::LacO-TRP1/"*
               CNM67-mCherry::natMX4/CNM67

JCY2251   *MAT a/α ho::LYS2/", his4/", trp1/", lys2::TetOx240:URA3/lys2, leu2::LEU2 tetR-*
               GFP/leu2

JCY2248   As JCY2251 but *cdc14-3HA::KanMX4/"*

JCY2331   *MAT a/α ho::LYS2/", his4/", trp1/", lys2::TetOx240:URA3/", leu2::LEU2 tetR-*
               GFP/", CNM67-mCherry::natMX4/CNM67

JCY2330   As JCY2331 but *cdc14-3HA::KanMX4/"*

PRY53   *MAT a/α ho::LYS2/" leu::hisG/" spo11-Y135F-HA::URA3/",TUB-dGFP-URA3/"*
SPC29-CFP::TRP1/" HIS4X/hisx4x

PRY55   *MAT a/α ho::LYS2/" leu::hisG/" spo11-Y135F-HA::URA3/" TUB-dGFP-URA3/"*
SPC29-CFP::TRP1/" cdc14-3HA::KanMX4/" arg4/" HIS4/his4x

PRY99   *MAT a/α ho::LYS2/" leu2/" his4x/" trp1-ura3/LacO-TRP1 (at CENIV))- Cyc1p-*
LacI-GFP-URA3, YEN1-9A/"

PRY108   *MAT a/α ho::LYS2/" leu2/" his4x/" trp1-ura3/LacO-TRP1 (at CENIV)- Cyc1p-LacI-*
GFP-URA3 cdc14-3HA::KanMX4/" YEN1-9A/"

PRY121   *MAT a/α ho::LYS/" leu2/" SPC29-CFP::TRP1/" YEN1-9A/" TUB-dGFP-URA3/"*
his3/", cdc14-3HA::KanMX/"

PRY123   *MAT a/α ho::LYS/" leu2/" SPC29-CFP::TRP1/" YEN1-9A/" TUB-dGFP-URA3/"*
his3/"

PRY132   *MAT a/α ho::LYS2/ho::hisG, lys2/" leu2::hisG/" TUB-dGFP-URA3/" SPC29-*
CFP::TRP1/", pCLB2:3HA-CDC14::KanMX4/", his3/"

PRY182   *MATa/α ho:.LYS2/" his3 or his4/" leu2/" LacOTRP1 (at CENIV)-Cyc1p-LacI-GFP-*
URA3/" YEN1-9A/" lys2/" pCLB2::3HA-CDC14::KanMX4/"

JCY2389   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) pCLB2::3HA-CDC14::KanMX4/"

JCY2385   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) ndt80::hphMX6/" cdc14-
3HA::KanMX4/"

JCY2390   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) ndt80::hphMX6/"

JCY2399   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) ndt80::hphMX6/" pCLB2::3HA-
CDC14::KanMX4/"

JCY2440   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) ndt80::hphMX6/" mus81::hphMX4/"
cdc14-3HA::KanMX4/" pCDC5-Gal4-ER-pGAL1-CDC5-natMX4/"

JCY2442   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*
(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) ndt80::hphMX6/" mus81::hphMX4/"
pCDC5-Gal4-ER-pGAL1-CDC5-natMX4/"

JCY2444   *MAT a/α ho::hisG/" lys2/" ura3(Δsma-pst::hisG)/" leu2::hisG/" his4X::LEU2-*

(NgoMIV;ori)-URA3/HIS4::LEU2-(NBam;ori) mms4::hphMX6/" pCLB2::3HA-SGS1::KanMX4/"

JCY2446	As JCY2444 but *pCLB2::3HA-CDC14::KanMX4/"*
JCY2448	As JCY2444 but *yen1::natMX4/"*
JCY2529	*MATa/α ho::LYS2/" mms4Δ:.hphMX4/" HIS4B::LEU2/leu2 his4x::LEU2-URA3 pCLB2::3HA-CDC14::TRP1/ pCLB2::3HA-CDC14::KanMX4 trp1 pCLB2-3HA-CDC20 (KANMX6) sgs1::CLB2p-HA3-SGS1::KanMX5/" YEN1-9A-18MYC::URA3/YEN1 his3::hisG*
JCY2421	*MATa/α ho::LYS2/" ura3/" leu2::hisG/" trp1::hisG/" his3::hisG/" YEN1-9A-myc18::URA3/"*
JCY2376	*MATa/α ho::LYS2/" lys2/" ura3/" trp1/" his4/" leu2::hisG/" pCLB2::3HA-CDC14::KanMX4/"*
JCY2478	*MATa/α ho::LYS2/" lys2/" trp1/" HIS4B::LEU2/hix4x::LEU2-URA3 ura3/" yen1Δ::CNAT/" sgs1::CLB2p-HA3-SGS1::KanMX5/" mms4Δ:.hphMX4/" pCLB2-3HA-CDC20 (KANMX6)/"*
JCY2480	*MATa/α ho::LYS2/" ura3/" leu2::hisG/"trp1::hisG/" his3::hisG?/? sgs1::CLB2p-HA3-SGS1::NatMX4/sgs1::CLB2p-HA3-SGS1::KanMX ms4::hphMX4/" yen1::NAT/YEN1 pCLB2-HA3-CDC20::KanMX6/" HIS4(NewBamH)-LEU2/his4X-LEU2-URA3*
JCY2469	*MATa/α ho::LYS2/" lys2/" trp1/TRP1 his4x::LEU2-URA3/HIS4B::LEU2 ura3/" sgs1::CLB2p-HA3-SGS1::KanMX5/" mms4::hphMX4/MMS4 yen1Δ::NatMX4/"*
JCY2502	*MATa/α ho::LYS2" his4x::LEU2-URA3/HIS4B::LEU2 sgs1::CLB2p-HA3-SGS1::KanMX5/" pCLB2-3HA-CDC20 (KANMX6)/" pCLB2::3HA-CDC14::TRP1/" his3::hisG/? mms4Δ::hphMX4/MMS4*
JCY2508	*MATa/α ho::LYS2/" sgs1::CLB2p-HA3-SGS1::KanMX5/" HIS4B::LEU2/his4x::LEU2-URA3*
JCY2406	*MATa/α ho::LYS2/" ura3/" leu2::hisG/" trp1::hisG/" his3::hisG/" REC8-GFP::LEU2::KanMX4/", PDS1-tdTomato::KiTRP1/", CNM67-tdTomato::NatMX4/"*
JCY2404	As JCY2406 but *cdc14-3HA::KanMX4/"*
JCY892	*MATa/α ho::hisG/" ura3/" leu2::hisG/LEU2 lys2/" trp1/" SPC29-CFP::TRP1/" TUB-dGFP-URA3/"*
JCY893	*MATa/α ho::hisG/" ura3/" leu2::hisG/LEU2 lys2/" trp1/" SPC29-CFP::TRP1/" TUB-dGFP-URA3/" cdc14-3HA::KanMX4/"*

A

B

C

D

A

B

<u>ChrII (*lys2*)</u>

	TetR-GFP	DAPI	MERGE	DIC	*CDC14*	*cdc14-HA*
Type 1					92.0%	42.4%
Type 2					2.4%	8.5%
Type 3					2.1%	3.3%
Type 4					0.3%	5.7%
Type 5					1.4%	13.2%
Type 6					0.3%	5.7%
Type 7					1.0%	14.5%
Type 8					0.5%	6.7%

A

B