***Leishmania major* degrades murine CXCL1 – an immune evasion strategy**

Matthew S. Yorek[1,2], Barun Poudel[1,2], Lalita Mazgaeen[1,2,4], R. Marshall Pope[5], Mary E.

Wilson[2,3,6,7,8], Prajwal Gurung[1,2,3,4,*]

[1] Inflammation Program, University of Iowa, Iowa City, IA, USA

[2] Department of Internal Medicine, University of Iowa, Iowa City, IA, USA

[3] Immunology Graduate Program, University of Iowa, Iowa City, IA, USA

[4] Interdisciplinary Graduate Program in Human Toxicology, University of Iowa, Iowa City,

IA, USA

[5]The Proteomics Facility, University of Iowa, Iowa City, IA, USA.

[6]Department of Microbiology and Immunology, University of Iowa, Iowa City, IA, USA

[7]Department of Epidemiology, University of Iowa, Iowa City, IA, USA

[8]Iowa City VA Medical Center, Iowa City, IA, USA

*Correspondence should be addressed to:

Prajwal Gurung

Inflammation Program, Department of Internal Medicine

University of Iowa

431 Newton Road, 429 EMRB

Iowa City, IA 52242

Tel: (319) 335-4536; Fax: (319) 335-4194

E-mail: prajwal-gurung@uiowa.edu

Running title: *Leishmania spp.* degrades CXCL1

Keywords: Leishmania, CXCL1, leishmaniasis, metalloprotease

Summary sentence: *L. major* metalloproteases can specifically degrade murine

CXCL1 to subvert the host immune response and establish infection.

Abstract

Leishmaniasis is a global health problem with an estimated report of 2 million new cases

every year and more than 1 billion people at risk of contracting this disease in endemic

areas. The innate immune system plays a central role in controlling *L. major* infection by

initiating a signaling cascade that results in production of pro-inflammatory cytokines and

recruitment of both innate and adaptive immune cells. Upon infection with *L. major*,

CXCL1 is produced locally and plays an important role in the recruitment of neutrophils

to the site of infection. Herein, we report that *L. major* specifically targets murine CXCL1

for degradation. The degradation of CXCL1 is not dependent on host factors as *L. major*

can directly degrade recombinant CXCL1 in a cell-free system. Using mass

spectrometry, we discovered that the *L. major* protease cleaves at the C-terminal end of

murine CXCL1. Finally, our data suggest that *L. major* metalloproteases are involved in

the direct cleavage and degradation of CXCL1, and a synthetic peptide spanning the

CXCL1 cleavage site can be used to inhibit *L. major* metalloprotease activity. In

conclusion, our study has identified an immune evasion strategy employed by *L. major*

to evade innate immune responses in mice, likely reservoirs in the endemic areas, and

further highlights that targeting these *L. major* metalloproteases may be important in

controlling infection within the reservoir population and transmittance of the disease.

Authors' summary

Our study discovered a highly specific role for *L. major* metalloprotease in cleaving and

degrading murine CXCL1. Indeed, *L. major* metalloprotease did not cleave murine

CXCL2 or human CXCL1, CXCL2 and CXCL8. CXCL1 is a critical chemokine required

for neutrophil recruitment to the site of infection; thus, we propose that this

metalloprotease may have evolved to evade immune responses specifically in the

murine host. We have further identified that the C-terminal end on CXCL1 is targeted for

cleavage by the *L. major* metalloprotease. Finally, this cleavage site information was

used to design peptides that are able to inhibit CXCL1 degradation by *L. major*. Our

study highlights an immune evasion strategy utilized by *L. major* to establish infection

within a murine host.

Introduction

Leishmania spp. are unicellular eukaryotic protozoan parasites that are transmitted to

mammalian hosts by sandfly (*Phlebotomine and Lutzomyia spp.*) bites [1]. Upon

transmission of *L. major* promastigotes (the infectious stage for mammalian hosts with a

long slender body and an anterior flagellin), the promastigotes are quickly taken up by

neutrophils, macrophages and keratinocytes [2-6]. Within the macrophages, *Leishmania*

spp. promastigotes hijack the phagocytic vacuole and transform into amastigotes (round

body lacking an anterior flagellin) [7]. The *Leishmania spp.* amastigotes then proliferate

within the vacuole and establish infection within the host [8, 9]. While a mammalian

host-vector system is the major mode of *Leishmania spp.* transmission, several studies

have reported a vertical transmission of these parasites in mammalian hosts, from a

pregnant female to its offspring [10-13]. Specifically, *Leishmania spp.* infection has been

found to be endemic in foxhound dog populations in the United States, where the

vectors are not present [14-17]. Given that *Leishmania spp.* can infect several hosts,

including rodents and dogs (in addition to humans) [18], these studies demonstrate how

these parasites can remain endemic even with strategies to eradicate sandflies.

Our immune system is extremely efficient in killing pathogens. Professional phagocytes

such as macrophages and neutrophils phagocytose and kill the invading pathogen in

the intracellular phagosome-lysosome compartment. In addition, these phagocytes also

respond to the foreign pathogens by secreting pro-inflammatory cytokines and

chemokines to recruit neutrophils to the site of infection [19-21]. *Leishmania spp.* have

evolved to evade the host immune response by using its armada of virulence factors to

avoid host killing [22]. The two major virulence factors of *Leishmania spp.* include

leishmanolysin metalloprotease glycoprotein 63 (GP63) and lipophosphoglycan (LPG),

and both have been extensively studied for their roles in immune evasion [23, 24].

GP63 and LPG inhibit the formation of the membrane attack complex to evade

complement-mediated lysis [25, 26], inhibit acidification of leishmania containing

vacuoles [27, 28], and dampen host immune signaling pathways [29, 30] for

establishing infection within the mammalian host. In addition, *Leishmania spp.* can

hijack the host immune responses to establish infections as shown by *Leishmania*

chagasi (L. chagasi)-mediated activation of TGF-β and *Leishmania major (L. major)*-

induced activation of NLRP3 inflammasome, events that promote *L. major* survival and

pathology [31-33]. Here, we have identified one such immune evasion mechanism

employed by *L. major.*

L. major is the most common *Leishmania spp.* and a major cause of cutaneous

leishmaniasis that affects 600,000-1,000,000 people globally [34]. Once deposited at

the site of a sandfly bite, *L. major* promastigotes are taken up by the resident

macrophages and keratinocytes, which then secrete essential cytokines and

chemokines to elicit an innate immune response [35]. One of the earliest chemokines

that are produced in the skin in response to *L. major* include CXCL1 [36]. CXCL1 is a

functional homolog of human IL-8 in mice and a potent neutrophil chemoattractant [37-

41]. Keratinocytes [3], neutrophils [42] and macrophages [43, 44] have all been

suggested to produce CXCL1 (or IL-8 in humans) in response to *Leishmania spp.*

infection. Several studies have shown that neutrophils infiltrate the site of *Leishmania*

spp. infection as early as 1-hour post-infection and are important for optimal resolution

of the infection [21, 45]. Given the importance of CXCL1 in modulating the early innate

immune response, we reasoned that *L. major* targets CXCL1 to evade early host innate

immune responses.

In this current study, we have identified murine CXCL1 as a highly specific substrate for

L. major metalloprotease and a possible immune evasion strategy employed by this

parasite to establish a successful infection within the murine host. We further report that

L. major promotes proteolytic cleavage of murine CXCL1 at the C-terminal end to

initiate its degradation. Finally, we have designed a synthetic peptide spanning the

CXCL1 cleavage site that inhibits *L. major* protease activity. In conclusion, our study

has uncovered a specific mechanism employed by *L. major* to degrade murine CXCL1

which may help the parasite to establish infection within the murine host.

Results

L. major infection abrogates LPS-induced CXCL1 production by bone marrow-derived macrophages.

To investigate how _L. major_ impacts innate immune responses elicited by macrophages, we stimulated bone marrow-derived macrophages (BMDM) with lipopolysaccharide (LPS) in the presence or absence of _L. major_ (WHOM/IR/-173) infection following established protocol [32]. As detailed in the experimental outline (**Fig. 1A**), BMDM were stimulated with 20 ng/ml LPS in the presence or absence of 20 MOI _L. major_ promastigotes for 48 hours. LPS stimulation of BMDM induced robust production of IL-6, TNF and CXCL1. Simultaneous LPS stimulation and _L. major_ infection did not impact the production of IL-6 and TNF by BMDM whereas CXCL1 levels were significantly blunted (**Fig. 1A**). These data suggest that _L. major_ specifically targets LPS-induced CXCL1 production by BMDM.

Previous studies have shown that _L. major_ targets signaling pathways to evade immune responses and establish infection [[29]]. To examine whether _L. major_ specifically targeted CXCL1 expression and production, we designed an experiment whereby BMDM were infected with 20 MOI _L. major_ for 6 hours, extracellular _L. major_ were washed, then stimulated with 20 ng/ml LPS for the next 42 hours (**Fig. 1B**). Interestingly, _L. major_-infected BMDM produced equal levels of CXCL1 when compared to controls (**Fig. 1B**). These results show that inhibition of CXCL1 by _L. major_ may not occur through modulation of intracellular signaling pathway that promotes CXCL1 expression and/or production by BMDM.

CXCL1 detection is significantly reduced by *L. major* in a cell free system

Given that *L. major* pre-infection of BMDM prior to LPS stimulation did not affect CXCL1

production, we posited that *L. major* regulates secreted CXCL1 in the extracellular

milieu. To this end, we used supernatants derived from LPS-stimulated BMDM as a

source of CXCL1 and cultured 500 μl of these cell-free supernatants with 20×10^6 *L.*

major promastigotes for 24 hours (**Fig. 2A**). While levels of IL-6 and TNF detected in the

supernatants remained unchanged, levels of CXCL1 were significantly reduced after

addition of *L. major* to the supernatants. The reduced levels of CXCL1 were not due to

its short half-life in the culture, as the levels of CXCL1 in the cell free supernatants were

stable up to 48 hours (**Supplemental Fig. 1**). Thus, *L. major* directly dampens CXCL1

detection.

We next examined whether the observed effect of *L. major* on CXCL1 detection was

dependent on live parasites. To this end, we cultured supernatants from LPS-stimulated

BMDM with *L. major* lysates (*Lm* lysate: generated by 3× freeze-thaw cycles of *L. major*)

or supernatants from *L. major* culture (*Lm* sup: supernatant collected from stationary

phase of *L. major* growth; **Fig. 2B**). Similar to the addition of live *L. major*, addition of

Lm lysate or *Lm* sup also reduced CXCL1 in the supernatants while IL-6 and TNF

remained unaffected (**Fig. 2B**). These results demonstrate that: first, *L. major* need not

be alive to dampen CXCL1 detection; second, *L. major* lysate can dampen CXCL1

detection; and finally, *L. major* secreted factors dampen CXCL1 detection. Boiling *Lm*

lysate or *Lm* sup for 20 minutes at 100°C rescued CXCL1 detected in the supernatants,

suggesting that the responsible *L. major* components are susceptible to heat treatment

(**Supplemental Fig. 2**). Given the sensitivity to heat treatment, we propose that the

CXCL1 regulating *L. major* components are proteinaceous.

***L. major* reduces recombinant murine CXCL1 (rm-CXCL1) levels in a cell free**

system

Macrophages secrete several hundreds of different proteins that may indirectly alter our

observed effect of *L. major* components on CXCL1 [46]. To this end, we obtained

recombinant murine CXCL1 (rm-CXCL1, Tonbo Biosciences, San Diego, CA) which

was stable in culture up to 48 hours (**Supplemental Fig. 3**). Importantly, *Lm* sup

addition to rm-CXCL1 dampened its detection by ELISA in a time-dependent manner,

and boiling *Lm* sup rescued rm-CXCL1 detection (**Supplemental Fig. 3**).

To further investigate the specificity of *L. major* in reducing rm-CXCL1 levels, we

examined rm-CXCL2, rh-CXCL1, rh-CXCL2 and rh-CXCL8 sequence homology by

CLUSTAL-W alignment and found significant homology between these recombinant

proteins. Despite the significant homology, *L. major* failed to reduce levels of rm-

CXCL2, rh-CXCL1, rh-CXCL2 or rh-CXCL8, demonstrating specific regulation of rm-

CXCL1 (**Fig. 3, B-F**). As expected, *L. major* did not inhibit rm-TNF levels (**Fig. 3G**).

Given the specific nature of *L. major* (WHOM/IR/-170) in dampening rm-CXCL1 levels,

we next determined whether this activity was specific to the WHOM/IR/-170 isolate.

However, supernatant from *L. major* (IA0, isolated from a patient in Iowa who acquired

L. major in Iraq [47]) was also able to inhibit rm-CXCL1 detection when compared to the

WHOM/IR/-170 strain (**Fig. 3H**). Furthermore, exosomes from the *L. major* supernatants

from both isolates were able to reduce rm-CXCL1 levels, suggesting that the active

components are also present in the *L. major*-derived vesicles (**Fig. 3H**).

***L. major* inactivates biological activity of CXCL1**

Our data show that *L. major* reduced CXCL1 levels as demonstrated by ELISA (**Figs. 1-**

3). Possible reasons why CXCL1 levels were reduced when incubated with *L. major*

include: 1) *L. major* proteins bind to CXCL1, limiting the ability of anti-CXCL1 antibody in

ELISA to interact with CXCL1 or 2) *L. major* proteases cleaves and degrades CXCL1.

Considering both outcomes, it is possible that CXCL1, either masked or cleaved by *L.*

major components, could still be biologically active. To this end, we performed a

functional assay where rm-CXCL1 or rm-CXCL1+*Lm* lysate was used to stimulate

BMDM (**Fig. 4A**). As demonstrated previously, when rm-CXCL1 is incubated with *L.*

major, rm-CXCL1 levels are significantly reduced (**Fig. 4B**). *Lm* lysate alone has very

little stimulatory activity as demonstrated by mRNA induction of *Cxcl1*, *Tnf* and *Il6* (**Fig.**

4, C-E). As expected, rm-CXCL1 induced modest increase of *Cxcl1*, *Tnf* and *Il6* but rm-

CXCL1+*Lm* lysate failed to induce *Cxcl1*, *Tnf* and *Il6* mRNA (**Fig. 4, C-E**). These results

altogether demonstrate that *L. major* inactivates biological activity of rm-CXCL1.

***L. major* cleaves CXCL1 at the C-terminal end to promote its degradation**

To determine whether CXCL1 is degraded by *L. major*, we incubated rm-CXCL1 with

Lm lysate for various time periods and examined rm-CXCL1 by silver staining (**Fig. 5A**).

Rm-CXCL1 was detected at approximately 10KDa by silver stain. Interestingly, when

incubated with *Lm* lysate rm-CXCL1 showed two bands as early as 30 minutes after

incubation, suggesting *Lm* lysate-mediated cleavage of rm-CXCL1 (**Fig. 5A and**

Supplemental Fig. 4). By 2 hours, only the cleaved rm-CXCL1 was observed and this

cleaved band intensity decreased over time, suggesting further degradation (**Fig. 5A**).

L. major-mediated cleavage of rm-CXCL1 was specific because similar experiments

done with rm-CXCL2, rh-CXCL1 and rh-CXCL2 did not result in cleavage or

degradation of these recombinant proteins (**Fig. 5B**).

Based on the cleavage pattern of rm-CXCL1 (less than 1KDa shift), cleavage either at

the N-terminal or C-terminal end would result in a large fragment (detected by silver

stain as the lower band, **Fig. 5A**) and a smaller fragment (which could not be detected

by silver stain; **Fig. 5C**). To identify the cleavage product and site, we processed the

full-length rm-CXCL1 and *Lm* lysate-cleaved rm-CXCL1 bands by trypsin digestion and

performed mass spectrometry to determine the sequence of the bands (**Supplemental**

Fig. 5). The tryptic peptide sequence readouts of full-length rm-CXCL1 covered the

whole rm-CXCL1 peptide sequence (**Supplemental Fig. 5B**), while the peptide

coverage of cleaved rm-CXCL1 covered all of the rm-CXCL1 peptide sequence except

the last 7 amino acids (MLKGVPK) at the C-terminal end (**Supplemental Fig. 5C**).

Thus, *L. major* cleaves rm-CXCL1 after lysine 65 (K65) residue that results in a large N-

terminal rm-CXCL1 fragment lacking 7 amino acids at the C-terminal end (**Fig. 5D**).

***L. major* metalloprotease cleaves CXCL1**

L. major has several proteases that enable it to survive within a cell and establish

infection [48]. Metalloproteases and cysteine, serine and aspartic proteases are the

major proteases described in *L. major* [48]. To examine whether cleavage activity was

dependent on proteases, we first treated our rm-CXCL1 + *Lm* sup culture with pan

protease inhibitor (Roche and Sigma) (**Fig.6, A-C**). Roche cOmplete protease inhibitor

is a broad inhibitor of proteases including serine, cysteine and metalloproteases, and

Sigma P8340 inhibitor is reported to inhibit serine, cysteine, acid proteases and

aminopeptidases. However, the presence of protease inhibitors from Roche (cOmplete)

or Sigma (P8340) did not inhibit *Lm* sup-mediated degradation of rm-CXCL1 as

demonstrated by ELISA (**Fig. 6, B and C, and Supplemental Fig. 6**). Marimastat, a

broad inhibitor of matrix metalloproteases [49], did not rescue the degradation of rm-

CXCL1 by *Lm* sup (**Supplemental Fig. 6B**). EDTA treatment, which chelates metal ions

such as Ca^{2+} and Fe^{3+} (and thus can inhibit certain metalloprotease), did not rescue rm-

CXCL1 cleavage by *Lm* sup (**Fig. 6, D and E**). Interestingly, 1,10-Phenanthroline, a

Zn^{2+} metalloprotease inhibitor [50], rescued rm-CXCL1 degradation by *Lm* sup (**Fig. 6, F**

and G). While the addition of 1,10-Phenanthroline resulted in the upward shift of rm-

CXCL1 bands in SDS-PAGE gels; importantly, no cleavage or degradation of rm-

CXCL1 was observed (**Fig. 6G**).

Several studies have shown that 1,10-Phenanthroline inhibits GP63, a Zn^{2+}

metalloprotease present on all *Leishmania spp* and is often used in biochemical assays

to inhibit non-specific proteolytic activity of GP63 [51]. To determine whether GP63 was

involved in specific cleavage of rm-CXCL1, we immunoprecipitated (IP) GP63 from *Lm*

lysate using anti-GP63 antibody (**Fig. 6H**). Immunoblotting the GP63+ve and GP63-ve

IP fractions for GP63 showed that GP63 was exclusively present in the IP+ve fraction

and not present in the IP-ve fraction demonstrating successful immunoprecipitation of

GP63 (**Fig. 6H**). More importantly, when these fractions were cultured with rm-CXCL1,

the IP-ve (*Lm* lysate lacking GP63) but not the IP+ve (i.e. GP63) fraction degraded rm-

CXCL1 suggesting a role for GP63-independent metalloprotease in the specific

degradation of rm-CXCL1 (**Fig. 6I**).

Synthetic peptide spanning the CXCL1 cleavage site inhibits *L. major* proteolytic

activity against rm-CXCL1

Our data suggest that *L. major* Zn^{2+} metalloprotease specifically cleaves rm-CXCL1

after the K65 residue leaving a 7-mer amino acid sequence (MLKGVPK). To further

examine the specificity of this *L. major* metalloprotease, we designed a blocking peptide

that covered this cleavage site (i.e. the last 15 amino acid sequences of rm-CXCL1; **Fig.**

7A). The addition of blocking peptide was able to rescue *Lm* lysate-mediated

degradation of rm-CXCL1 in a dose-dependent manner (**Fig. 7B**). The peptide

sequence from the signal peptide region of rm-CXCL1 (peptide #1 and peptide #2, Fig.

7A) did not inhibit *Lm* lysate-mediated degradation of rm-CXCL1, demonstrating the

specificity of the blocking peptide (**Fig. 7B**). These data demonstrate that a synthetic

peptide spanning the murine CXCL1 cleavage site can competitively inhibit *L. major*-

mediated rm-CXCL1 degradation.

Discussion

CXCL1 is a potent neutrophil chemoattractant and is rapidly upregulated following

Leishmania spp. infection in the skin [36]; however, the role of CXCL1 during

Leishmania spp. infection of mice in vivo has been understudied. Charmoy et al showed

that mice deficient in CXCL1 have a slight increase in lesion size and reduced numbers

of neutrophil infiltrates in chronic lesions, but the overall pathology remained similar

between WT and *Cxcl1*[-/-] mice [33]. However, it should be noted that the study used the

L. major Seidman strain (*Lm*Sd) that causes a non-healing infection in C57BL/6 mice

[33]. Thus, more thorough analyses are needed to determine the precise role of CXCL1

during *L. major* infection in mice and humans.

Importantly, even depletion of neutrophils using anti-Gr1 neutralizing antibody does not

always lead to increased parasitic burden, lesion or pathology [21]. While some studies

with acute depletion of neutrophils reported worsened pathology during *L. major*

infections, other studies have shown no role or even amelioration of disease pathology

[5, 52-60]. Thus, mouse strain, *L. major* strain, route of infection and timing of CXCL1

release may all contribute to the fate of the infection and overall pathology. In line with

this thought, two independent *L. major* strains were tested in our study, and both strains

degraded CXCL1 in our experimental systems. Whether this degradation of murine

CXCL1 is a specific feature of *L. major* or a more general feature of all *Leishmania spp.*

will need further investigation.

Our initial observation demonstrated that *L. major* in culture with macrophages

effectively inhibited LPS-induced CXCL1 production. Further examination showed that

intracellular *L. major* within the macrophages did not inhibit LPS-induced CXCL1

production. These experiments suggest that *L. major* does not interfere with signaling

pathways that promote CXCL1 production but directly regulates secreted CXCL1 in the

extracellular milieu. *L. major* are present as metacyclic promastigotes in sandflies [61].

When infected into the mammalian host, *L. major* infect macrophages and resides

within the vacuole where they transform into amastigotes, multiply and establish

infection [62]. Because *L. major*-infected macrophages which contain amastigotes did

not inhibit CXCL1 production, it could be posited that only the promastigote form of *L.*

major degrades CXCL1. However, genome microarray analysis shows that the majority

of genes are constitutively expressed in both *L. major* promastigotes and amastigotes

(>90%) and therefore the major genes and virulent factors expressed by these different

stages of *L. major* may not be different [63]. Thus, both *L. major* promastigotes and

amastigotes may similarly cleave and degrade murine CXCL1 in an in vitro assay. In

addition, infected macrophages can secrete exosomes containing *L. major* proteins [64,

65], which could degrade murine CXCL1. We have consistently shown that *L. major*

secreted microvesicles (i.e. exosomes) contain these metalloproteases that can

degrade murine CXCL1. Physiologically, *L. major* promastigotes released during a

sandfly bite may act locally to limit acute CXCL1 produced in response to the infection.

Once *L. major* are phagocytosed and transformed into amastigotes, they are physically

separated from the CXCL1 due to their location (amastigotes are in the vacuole while

CXCL1 are secreted by the macrophages and are extracellular) and may only impact

CXCL1 production through the *L. major* component laden exosomes.

Our studies suggest that a yet-unknown metalloprotease from *L. major* cleaves murine

CXCL1. Given that 1,10-Phenanthroline (Zn^{2+} chelator) but not EDTA (Ca^{2+} and Fe^{3+}

metal ion chelator) or Marimastat (Matrix metalloprotease inhibitor) rescued *L. major*-

mediated CXCL1 degradation, it is possible that the unknown metalloprotease is a Zn^{2+}

metalloprotease. Our studies also demonstrate that the *L. major* metalloprotease is

highly specific for murine CXCL1 in that its closest murine homolog, CXCL2, or human

homologs, CXCL1, CXCL2 or CXCL8, were not degraded. We hypothesize that this

particular *L. major* metalloprotease may have evolved to specifically evade host immune

response in rodents. Given that human CXCL1 homologs are not susceptible to this

metalloprotease, one can argue that our results may not have any importance from a

public health standpoint, but we reason otherwise. Our results are highly relevant to

public health because rodents are ubiquitous, serve as a reservoir for *Leishmania spp.*

and we highlight a rodent-specific *L. major* evolution in targeting murine CXCL1 [66].

Our studies demonstrate that the murine CXCL1 is first cleaved at the C-terminal end

after K65 releasing a 7aa residue. CXCL1 cleavage occurs as early as 30 minutes of

incubation with *L. major* and by 4 hours the cleavage is complete in that only the

cleaved bands are observed. Interestingly, we did not observe any accumulation of the

cleaved CXCL1 band, suggesting continuous degradation of the cleaved form. While

our data clearly demonstrate that the unknown *L. major* metalloprotease mediates

cleavage of murine CXCL1, how the cleaved CXCL1 is further degraded will be the

subject of future investigation. The cleaved murine CXCL1 (lacking the C-terminal 7 aa)

may be highly unstable and undergo spontaneous degradation overtime. Alternatively,

the cleaved murine CXCL1 may be susceptible to additional *L. major* proteases which

promote its subsequent degradation.

In conclusion, we have identified an immune evasion strategy utilized by *L. major* that is

highly specific to murine CXCL1 (**Fig. 7C**). Specifically, *L. major*-associated

metalloprotease cleaved murine CXCL1 at K65 residue and released a C-terminal 7

amino acid fragment to promote its degradation. Finally, we have designed a peptide

spanning the cleavage site of CXCL1 that inhibited murine CXCL1 cleavage by *L.*

major. Our study altogether uncovered an immune evasion strategy employed by *L.*

major that may have evolved in rodents and highlights how parasites may utilize diverse

immune evasion strategy to establish infection within its diverse mammalian hosts.

Methods

Ethics statement

Experimental procedures that utilized mice were all approved by the University of Iowa

Animal Care and Use Committee (Approved Animal Protocol # 7042004 – PI Dr.

Prajwal Gurung) and performed in accordance to the Office of Laboratory Animal

Welfare guidelines and the PHS Policy on Humane Care and Use of Laboratory

Animals.

BMDM culture

BMDMs were prepared as described previously (32). Briefly, bone marrow cells were

harvested from the hind limbs of BALB/c mice (Jackson Laboratory, Stock No. 000651)

and cultured in L cell-conditioned IMDM medium supplemented with 10% FBS, 1%

nonessential amino acid, and 1% penicillin-streptomycin for 5-7 d to differentiate into

macrophages. BMDMs were counted and seeded at 1×10^6 cells in 12-well cell culture

plates in IMDM media containing 10% FBS, 1% nonessential amino acids, and 1%

penicillin-streptomycin. BMDMs were primed with LPS (20 ng/ml) and infected with 20

MOI *L. major* promastigotes for 24 and 48 hours (Supplemental Figure 1). For some

experiments, BMDMs were LPS primed to generate supernatant containing cytokines

for in vitro biochemical analysis with *L. major* supernatant (*Lm* sup) and *L. major* lysate

(*Lm* lysate). BMDMs were also treated with rm-CXCL1 (20 ng/ml) or rm-CXCL1+*Lm* sup

preparations in some experiments.

L. major culture

L. major strains WHOM/IR/-173 [67] and IA0 [47] were grown in T-25 flasks with M199

media supplemented with 10% FBS, 5% HEPES and 1% penicillin-streptomycin at room

temperature. BMDMs were infected with 20 MOI of *L. major* promastigotes for 48 hours.

Conditioned *L. major* supernatant (*Lm* sup) was prepared after *L. major* reached the

stable growth phase at approximately 20×10^6 / ml. Following centrifugation (3000×g) to

pellet *L. major*, supernatants were collected and filtered using a 0.2μm vacuum filter to

harvest *Lm* sup. *Lm* lysate was prepared by collecting the *L. major* pellet, washing 3

times with PBS followed by 3 freeze-thaw cycles of 7×10^9 *L. major* /ml in PBS. For the

preparation of *L. major* exosomes, promastigotes were grown to stable phase and

changed to serum-free media and incubated at room temperature overnight.

Supernatant was obtained as above and then separated using a 100kDa molecular

weight Amicon ultra-15 centrifugal filter (MilliporeSigma, Burlington, MA). The

concentrated vesicles were washed twice with PBS by ultracentrifugation at 110,000xg

at 4°C for 1 hour (Beckman Optma MAX-XP Ultracentrifuge, rotor TLA 120.2; Beckman

Coulter Inc., USA). The final pellet (exosomes) was resuspended in PBS and used in in

vitro biochemistry assays.

Electrophoresis and in vitro biochemistry

For in vitro reactions, 125ng of total recombinant protein was separated on 4-20% or

10-20% Novex WedgeWell Tris-Glycine precast gels (Thermofisher, Waltham, MA).

For silver stain, the Pierce Mass Spec compatible silver stain kit was used per

manufacturer's instructions (Thermofisher). Synthetic peptides were generated and

purchased from Pierce Biotechnology (Thermofisher) and are composed of the

following sequences: peptide#1: CAALLLLATSRLA; peptide#2: RLATGAPIAN;

blocking peptide: LVQKIVQKMLKGVPK. For Western blot analysis a semi-dry transfer

was completed using trans-blot turbo system (Bio-Rad, Hercules, CA) with Immobilon

PVDF membrane. The membranes were blocked in 5% non-fat milk for 1 hour at room

temperature. Membranes were probed with sheep polyclonal antibody Sp180 raised

against *L. chagasi* promastigote GP63 at a concentration 1:1500 incubated overnight at

4°C. The membranes were then probed with HRP-tagged secondary antibodies at room

temperature for 1 hour and developed using Immobilon Forte Western HRP Substrate

(MilliporeSigma) and imaged with an Odyssey Fc Infrared Imaging System (LI-COR

Bioscience, Lincoln, NE). Immunoprecipitation was performed using Protein G

Dynabeads and Leishmania GP63 monoclonal antibody clone 96-126 (Thermofisher).

As instructed by the manufacturer, 50μL beads were washed in PBS with 0.02% Tween

20 and bound to 5μg antibody for at least 10 minutes. Following a wash, the antibody

bound beads were incubated with *Lm* lysate for another 10 minutes and finally either

antigen-antibody binding was negated by acidic (pH 2.8) 50mM Glycine treatment or

bead bound antibody-antigen complexes were used directly for biochemistry.

Mass spectrometry

In-gel trypsin digestion: The gel was stained using a Pierce mass spec compatible silver

stain kit (Thermofisher) per manufacturer directions. A procedure slightly modified than

that described by Yu et al [68] was used for in-gel digestion. Briefly, the targeted protein

bands from SDS-PAGE gel were manually excised, cut into 1mm^3 pieces, and washed

in 100mM ammonium bicarbonate:acetonitrile (1:1, v/v) and 25mM ammonium

bicarbonate /acetonitrile (1:1, v/v), respectively, to achieve complete de-staining. The

gel pieces were further treated with acetonitrile (ACN), to effectively "dry" the gel

segments and then reduced in 50μl of 10mM DTT at 56 °C for 60 min. Gel-trapped

proteins were alkylated with 55mM chloroacetamide (CAM) for 30 min at room

temperature. The gel pieces were washed with 25mM ammonium bicarbonate:

acetonitrile (1:1, v/v) twice to remove excess DTT and CAM. 50μl of cold trypsin solution

at 10ng/μl in 25mM ammonium bicarbonate was then added to the gel pieces and they

were allowed to swell on ice for 60 min. Digestion was conducted at 37°C for 16 h.

Peptide extraction was performed three times, adding 100µL of 50% acetonitrile/0.1%

formic acid for 0.5 h, combining the supernatants. The combined extracts were

concentrated in a lyophilizer and rehydrated with 15 μl of mobile phase A.

LC-MS/MS: Mass spectrometry data were collected using an Orbitrap Fusion Lumos

mass spectrometer (Thermofisher) coupled to an Easy-nLC-1200™ System (Proxeon

P/N LC1400). The autosampler is set to aspirate 3μl (estimated 0.3ug) of reconstituted

digest and load the solution on a 2.5cm C18 trap (New Objective, P/N IT100-25H002)

coupled to waste, HV or analytical column through a microcross assembly (IDEX, P/N

UH-752). Peptides are desalted on the trap using 16μl mobile phase A for 4 min. The

waste valve is then blocked and a gradient begins flowing at 0.4μl/min through a self-

packed analytical column, 10cm in length × 75μm id. The fused silica column was

tapered from 100μm ID (Polymicro) to ~8μm at the tip using a Sutter P-2000 laser puller

then packed with 2.7micron Halo C18 particles using a He-pressurized SS cylinder.

Peptides were separated in-line with the mass spectrometer using a 70 min gradient

composed of linear and static segments wherein buffer A is 0.1% formic acid and B is

95%ACN, 0.1% formic acid. The gradient begins first hold at 4% for 3 min then makes

the following transitions (%B, min): (2, 0), (35, 46), (60, 56), (98, 62), (98, 70).

Tandem mass spectrometry on the Thermo Q-Exactive hf: Data acquisitions begin with

a survey scan (m/z 380 -1800) acquired on a Q-Exactive Orbitrap mass spectrometer

(Thermofisher) at a resolution of 120,000 in the off-axis Orbitrap segment (MS1) with

automatic gain control (AGC) set to 3E06 and a maximum injection time of 50 ms. MS1

scans were acquired every 3 sec during the 70-min gradient described above. The most

abundant precursors were selected among 2-6 charge state ions at a 1E05 AGC and 70

ms maximum injection time. Ions were isolated with a 1.6 Th window using the multi-

segment quadrupole and subjected to dynamic exclusion for 30 sec if they were

targeted twice in the prior 30 sec. Selected ions were then subjected to high energy

collision-induced dissociation (HCD) in the ion routing multipole (IRM). Targeted

precursors were fragmented by (HCD) at 30% collision energy in the IRM. HCD

fragment ions were analyzed using the Orbitrap (AGC 1.2E05, maximum injection time

110 ms, and resolution set to 30,000 at 400 Th). Both MS2 data were recorded as

centroid and the MS1 survey scans were recorded in profile mode.

Proteomic searches: Initial spectral searches were performed with both Mascot version

2.6.2 (MatrixScience) and Byonic search engines (Protein Metrics ver. 2.8.2). Search

databases were composed of the Uniprot KB for species 10090 (mouse) downloaded

February 6, 2016 containing 58436 sequences. In either search, an equal number of

decoy entries were created and searched simultaneously by reversing the original

entries in the target database. Precursor mass tolerance was set to 5 ppm and

fragments were searched at 10 ppm. A fixed 57 Da modification was assumed for

cysteine residues while variable oxidation was allowed at methionine. A variable GG

modification at lysine was set to monitor ubiquitylation and potential phosphorylation

was accessed at Ser and Thr residues. The false discovery rate was maintained at 1%

by tracking matches to the decoy database. Both Mascot and Byonic search results

were combined and validated using Scaffold ver. 4.8.5 (Proteome Software). Protein

assignments required a minimum of two peptides established at 70% probability (local

FDR algorithm) and an overall 95% protein probability (assigned by Protein Prophet).

Approximately 300 protein families (including common contaminants) were assigned at

a total FDR to 1.2%. Proteins were annotated with GO terms from goa_uniprot_all.gaf

downloaded on May 3, 2017.

Immunoassays and recombinant proteins

Cytokine ELISAs and multiplex immunoassays were performed according to

manufacturer instructions. Mouse and human CXCL1 and CXCL2 as well as human IL-

8 (CXCL8) were obtained from R&D Systems (Minneapolis, MN) and multiplex CXCL1,

IL-6, and TNF were completed using ProcartaPlex assays (Thermofisher) which were

run on the BioRad Bio-Plex (Luminex, Austin, TX). We obtained our recombinant

proteins from Tonbo biosciences which included mouse CXCL1, CXCL2, and TNF as

well as human CXCL1, CXCL2, and CXCL8.

PCR methods

RNA was extracted from BMDM cultures in TRIzol reagent (Thermofisher) followed by

chloroform extraction and isopropanol precipitation. The extracted RNA was reverse-

transcribed into cDNA by using qScript Supermix (Quanta Biosciences, Beverly, MA).

Real-time quantitative PCR was performed on Eppendorf realplex EPgradient S

Mastercycler (Eppendorf, Germany) using PerfeCTa SYBR Green SuperMix ROX

(Quanta Biosciences) and the appropriate primers. The sequences of the quantitative

RT-PCR primers are as follows: *mCXCL1:* 5'-CAATGAGCTGCGCTGTCAGTG-3', 5'-

CTTGGGGACACCTTTTAGCATC-3'; *mTnf*: 5′-CATCTTCTCAAAATTCGAGTGACAA-

3′, 5′-TGGGAGTAGACAAGGTACAACCC-3′; *mIL-6:* 5'-

CAAGAAAGACAAAGCCAGAGTC-3', 5'-GAAATTGGGGTAGGAAGGAC-3'.

Threshold cycle Ct values were normalized to GAPDH, and gene fold change was

determined by the relative comparison method, relative to the 0 h time point.

Statistical analysis and alignment

GraphPad Prism 8.0 software was used for data analysis and figure presentation. Data

are shown as mean ± SEM. Statistical significance was determined by *t* tests (two-tailed

and Mann-Whitney) for two groups, and one-way ANOVA (with Dunnett's or Tukey's

multiple comparisons tests) for three or more groups. For the purpose of alignment

presentation, sequences were obtained from the protein manufacturer or the NCBI

protein database and figures were prepared using Jalview 2 [69].

Authorships

M.S.Y, B.P., L.M. performed the experiments. M.S.Y, M.E.W., and P.G. analyzed the

data. R.M.P. performed the mass spectrometry and analysis of CXCL1 cleavage bands.

M.S.Y. and P.G. wrote the manuscript with input from all authors. P.G. designed and

conceptualized the study.

Acknowledgements

We thank the University of Iowa Proteomics Core for the comparative analysis

performed in the University of Iowa Proteomic facility, Directed by RMP and supported

by an endowment from the Carver Foundation and by Thermo Lumos provided through

an HHMI grant to Dr. Kevin Campbell. We thank Kristina Greiner for her help with

scientific editing of the manuscript. We are also thankful to the members of the

Inflammation Program for critical insights provided during the development of the

project. This work was supported by NIAID K22AI127836 Grant, NIEHS P30ES005605

Pilot Grant, and the University of Iowa Startup funds to Dr. Prajwal Gurung.

Conflict of Interest

The authors have no conflict of interest to declare.

References

1.   Elnaiem DE. Ecology and control of the sand fly vectors of Leishmania donovani in East Africa, with special emphasis on Phlebotomus orientalis. J Vector Ecol. 2011;36 Suppl 1:S23-31. Epub 2011/03/05. doi: 10.1111/j.1948-7134.2011.00109.x. PubMed PMID: 21366778.

2.   Ehrchen JM, Roebrock K, Foell D, Nippe N, von Stebut E, Weiss JM, et al. Keratinocytes determine Th1 immunity during early experimental leishmaniasis. PLoS Pathog. 2010;6(4):e1000871. Epub 2010/05/06. doi: 10.1371/journal.ppat.1000871. PubMed PMID: 20442861; PubMed Central PMCID: PMCPMC2861693.

3.   Scorza BM, Wacker MA, Messingham K, Kim P, Klingelhutz A, Fairley J, et al. Differential Activation of Human Keratinocytes by Leishmania Species Causing Localized or Disseminated Disease. J Invest Dermatol. 2017;137(10):2149-56. Epub 2017/06/26. doi: 10.1016/j.jid.2017.05.028. PubMed PMID: 28647347; PubMed Central PMCID: PMCPMC5786447.

4.   Ribeiro-Gomes FL, Moniz-de-Souza MC, Alexandre-Moreira MS, Dias WB, Lopes MF, Nunes MP, et al. Neutrophils activate macrophages for intracellular killing of Leishmania major through recruitment of TLR4 by neutrophil elastase. J Immunol. 2007;179(6):3988-94. Epub 2007/09/06. PubMed PMID: 17785837.

5.   Ribeiro-Gomes FL, Otero AC, Gomes NA, Moniz-De-Souza MC, Cysne-Finkelstein L, Arnholdt AC, et al. Macrophage interactions with neutrophils regulate Leishmania major infection. J Immunol. 2004;172(7):4454-62. Epub 2004/03/23. PubMed PMID: 15034061.

6.   Liu D, Uzonna JE. The early interaction of Leishmania with macrophages and

dendritic cells and its influence on the host immune response. Front Cell Infect

Microbiol. 2012;2:83. Epub 2012/08/25. doi: 10.3389/fcimb.2012.00083. PubMed PMID:

22919674; PubMed Central PMCID: PMCPMC3417671.

7.   Gurung P, Kanneganti TD. Innate immunity against Leishmania infections. Cell

Microbiol. 2015;17(9):1286-94. Epub 2015/08/08. doi: 10.1111/cmi.12484. PubMed

PMID: 26249747; PubMed Central PMCID: PMCPMC4698274.

8.   Kaye P, Scott P. Leishmaniasis: complexity at the host-pathogen interface. Nat

Rev Microbiol. 2011;9(8):604-15. Epub 2011/07/13. doi: 10.1038/nrmicro2608. PubMed

PMID: 21747391.

9.   Naderer T, Heng J, Saunders EC, Kloehn J, Rupasinghe TW, Brown TJ, et al.

Intracellular Survival of Leishmania major Depends on Uptake and Degradation of

Extracellular Matrix Glycosaminoglycans by Macrophages. PLoS Pathog.

2015;11(9):e1005136. Epub 2015/09/04. doi: 10.1371/journal.ppat.1005136. PubMed

PMID: 26334531; PubMed Central PMCID: PMCPMC4559419.

10.   Avila-Garcia M, Mancilla-Ramirez J, Segura-Cervantes E, Farfan-Labonne B,

Ramirez-Ramirez A, Galindo-Sevilla N. Transplacental transmission of cutaneous

Leishmania mexicana strain in BALB/c mice. Am J Trop Med Hyg. 2013;89(2):354-8.

Epub 2013/06/27. doi: 10.4269/ajtmh.12-0716. PubMed PMID: 23798582; PubMed

Central PMCID: PMCPMC3741259.

11.   Ben Slimane T, Chouihi E, Ben Hadj Ahmed S, Chelbi I, Barhoumi W, Cherni S,

et al. An investigation on vertical transmission of Leishmania infantum in experimentally

infected dogs and assessment of offspring's infectiousness potential by xenodiagnosis.

Vet Parasitol. 2014;206(3-4):282-6. Epub 2014/12/04. doi:

10.1016/j.vetpar.2014.10.020. PubMed PMID: 25468025.

12.   da Silva SM, Ribeiro VM, Ribeiro RR, Tafuri WL, Melo MN, Michalick MS. First

report of vertical transmission of Leishmania (Leishmania) infantum in a naturally

infected bitch from Brazil. Vet Parasitol. 2009;166(1-2):159-62. Epub 2009/09/08. doi:

10.1016/j.vetpar.2009.08.011. PubMed PMID: 19733439.

13.   Naucke TJ, Lorentz S. First report of venereal and vertical transmission of canine

leishmaniosis from naturally infected dogs in Germany. Parasit Vectors. 2012;5:67.

Epub 2012/04/03. doi: 10.1186/1756-3305-5-67. PubMed PMID: 22463789; PubMed

Central PMCID: PMCPMC3350385.

14.   Gibson-Corley KN, Hostetter JM, Hostetter SJ, Mullin K, Ramer-Tait AE,

Boggiatto PM, et al. Disseminated Leishmania infantum infection in two sibling

foxhounds due to possible vertical transmission. Can Vet J. 2008;49(10):1005-8. Epub

2009/01/03. PubMed PMID: 19119370; PubMed Central PMCID: PMCPMC2553493.

15.   Rosypal AC, Troy GC, Duncan RB, Zajac AM, Lindsay DS. Utility of diagnostic

tests used in diagnosis of infection in dogs experimentally inoculated with a North

American isolate of Leishmania infantum infantum. J Vet Intern Med. 2005;19(6):802-9.

Epub 2005/12/17. PubMed PMID: 16355672.

16.   Swenson CL, Silverman J, Stromberg PC, Johnson SE, Wilkie DA, Eaton KA, et

al. Visceral leishmaniasis in an English foxhound from an Ohio research colony. J Am

Vet Med Assoc. 1988;193(9):1089-92. Epub 1988/11/01. PubMed PMID: 3198462.

17.   Petersen CA, Barr SC. Canine leishmaniasis in North America: emerging or

newly recognized? Vet Clin North Am Small Anim Pract. 2009;39(6):1065-74, vi. Epub

2009/11/26. doi: 10.1016/j.cvsm.2009.06.008. PubMed PMID: 19932363; PubMed

Central PMCID: PMCPMC2824922.

18.   Ashford RW. Leishmaniasis reservoirs and their significance in control. Clin

Dermatol. 1996;14(5):523-32. Epub 1996/09/01. PubMed PMID: 8889331.

19.   Oghumu S, Lezama-Davila CM, Isaac-Marquez AP, Satoskar AR. Role of

chemokines in regulation of immunity against leishmaniasis. Exp Parasitol.

2010;126(3):389-96. Epub 2010/03/09. doi: 10.1016/j.exppara.2010.02.010. PubMed

PMID: 20206625; PubMed Central PMCID: PMCPMC3661772.

20.   Charmoy M, Auderset F, Allenbach C, Tacchini-Cottier F. The prominent role of

neutrophils during the initial phase of infection by Leishmania parasites. J Biomed

Biotechnol. 2010;2010:719361. Epub 2009/11/04. doi: 10.1155/2010/719361. PubMed

PMID: 19884987; PubMed Central PMCID: PMCPMC2768872.

21.   Ribeiro-Gomes FL, Sacks D. The influence of early neutrophil-Leishmania

interactions on the host immune response to infection. Front Cell Infect Microbiol.

2012;2:59. Epub 2012/08/25. doi: 10.3389/fcimb.2012.00059. PubMed PMID:

22919650; PubMed Central PMCID: PMCPMC3417510.

22.   Chang KP, McGwire BS. Molecular determinants and regulation of Leishmania

virulence. Kinetoplastid Biol Dis. 2002;1(1):1. Epub 2002/09/18. PubMed PMID:

12234388; PubMed Central PMCID: PMCPMC119322.

23.   Joshi PB, Kelly BL, Kamhawi S, Sacks DL, McMaster WR. Targeted gene

deletion in Leishmania major identifies leishmanolysin (GP63) as a virulence factor. Mol

Biochem Parasitol. 2002;120(1):33-40. Epub 2002/02/19. PubMed PMID: 11849703.

24.   Spath GF, Lye LF, Segawa H, Sacks DL, Turco SJ, Beverley SM. Persistence

without pathology in phosphoglycan-deficient Leishmania major. Science.

2003;301(5637):1241-3. Epub 2003/08/30. doi: 10.1126/science.1087499. PubMed

PMID: 12947201.

25.   Brittingham A, Morrison CJ, McMaster WR, McGwire BS, Chang KP, Mosser

DM. Role of the Leishmania surface protease gp63 in complement fixation, cell

adhesion, and resistance to complement-mediated lysis. J Immunol. 1995;155(6):3102-

11. Epub 1995/09/15. PubMed PMID: 7673725.

26.   Puentes SM, Sacks DL, da Silva RP, Joiner KA. Complement binding by two

developmental stages of Leishmania major promastigotes varying in expression of a

surface lipophosphoglycan. J Exp Med. 1988;167(3):887-902. Epub 1988/03/01.

PubMed PMID: 3280727; PubMed Central PMCID: PMCPMC2188887.

27.   Desjardins M, Descoteaux A. Inhibition of phagolysosomal biogenesis by the

Leishmania lipophosphoglycan. J Exp Med. 1997;185(12):2061-8. Epub 1997/06/16.

PubMed PMID: 9182677; PubMed Central PMCID: PMCPMC2196352.

28.   Vinet AF, Fukuda M, Turco SJ, Descoteaux A. The Leishmania donovani

lipophosphoglycan excludes the vesicular proton-ATPase from phagosomes by

impairing the recruitment of synaptotagmin V. PLoS Pathog. 2009;5(10):e1000628.

Epub 2009/10/17. doi: 10.1371/journal.ppat.1000628. PubMed PMID: 19834555;

PubMed Central PMCID: PMCPMC2757729.

29.   Gupta G, Oghumu S, Satoskar AR. Mechanisms of immune evasion in

leishmaniasis. Adv Appl Microbiol. 2013;82:155-84. Epub 2013/02/19. doi:

10.1016/B978-0-12-407679-2.00005-3. PubMed PMID: 23415155; PubMed Central

PMCID: PMCPMC3697132.

30.   Isnard A, Shio MT, Olivier M. Impact of Leishmania metalloprotease GP63 on

macrophage signaling. Front Cell Infect Microbiol. 2012;2:72. Epub 2012/08/25. doi:

10.3389/fcimb.2012.00072. PubMed PMID: 22919663; PubMed Central PMCID:

PMCPMC3417651.

31.   Gantt KR, Schultz-Cherry S, Rodriguez N, Jeronimo SM, Nascimento ET,

Goldman TL, et al. Activation of TGF-beta by Leishmania chagasi: importance for

parasite survival in macrophages. J Immunol. 2003;170(5):2613-20. Epub 2003/02/21.

PubMed PMID: 12594289.

32.   Gurung P, Karki R, Vogel P, Watanabe M, Bix M, Lamkanfi M, et al. An NLRP3

inflammasome-triggered Th2-biased adaptive immune response promotes

leishmaniasis. J Clin Invest. 2015;125(3):1329-38. Epub 2015/02/18. doi:

10.1172/JCI79526. PubMed PMID: 25689249; PubMed Central PMCID:

PMCPMC4362229.

33.   Charmoy M, Hurrell BP, Romano A, Lee SH, Ribeiro-Gomes F, Riteau N, et al.

The Nlrp3 inflammasome, IL-1beta, and neutrophil recruitment are required for

susceptibility to a nonhealing strain of Leishmania major in C57BL/6 mice. Eur J

Immunol. 2016;46(4):897-911. Epub 2015/12/23. doi: 10.1002/eji.201546015. PubMed

PMID: 26689285; PubMed Central PMCID: PMCPMC4828310.

34.   WHO. Leishmaniasis. https://wwwwhoint/news-room/fact-

sheets/detail/leishmaniasis. 2018;World Health Orgnaization.

35.   Teixeira MJ, Teixeira CR, Andrade BB, Barral-Netto M, Barral A. Chemokines in

host-parasite interactions in leishmaniasis. Trends Parasitol. 2006;22(1):32-40. Epub

2005/11/29. doi: 10.1016/j.pt.2005.11.010. PubMed PMID: 16310413.

36.   Muller K, van Zandbergen G, Hansen B, Laufs H, Jahnke N, Solbach W, et al.

Chemokines, natural killer cells and granulocytes in the early course of Leishmania

major infection in mice. Med Microbiol Immunol. 2001;190(1-2):73-6. Epub 2002/01/05.

PubMed PMID: 11770115.

37.   Modi WS, Yoshimura T. Isolation of novel GRO genes and a phylogenetic

analysis of the CXC chemokine subfamily in mammals. Mol Biol Evol. 1999;16(2):180-

93. Epub 1999/02/24. doi: 10.1093/oxfordjournals.molbev.a026101. PubMed PMID:

10028286.

38.   Bozic CR, Kolakowski LF, Jr., Gerard NP, Garcia-Rodriguez C, von Uexkull-

Guldenband C, Conklyn MJ, et al. Expression and biologic characterization of the

murine chemokine KC. J Immunol. 1995;154(11):6048-57. Epub 1995/06/01. PubMed

PMID: 7751647.

39.   Chintakuntlawar AV, Chodosh J. Chemokine CXCL1/KC and its receptor CXCR2

are responsible for neutrophil chemotaxis in adenoviral keratitis. J Interferon Cytokine

Res. 2009;29(10):657-66. Epub 2009/08/01. doi: 10.1089/jir.2009.0006. PubMed PMID:

19642907; PubMed Central PMCID: PMCPMC2956677.

40.   Frevert CW, Huang S, Danaee H, Paulauskis JD, Kobzik L. Functional

characterization of the rat chemokine KC and its importance in neutrophil recruitment in

a rat model of pulmonary inflammation. J Immunol. 1995;154(1):335-44. Epub

1995/01/01. PubMed PMID: 7995953.

41. Girbl T, Lenn T, Perez L, Rolas L, Barkaway A, Thiriot A, et al. Distinct

Compartmentalization of the Chemokines CXCL1 and CXCL2 and the Atypical

Receptor ACKR1 Determine Discrete Stages of Neutrophil Diapedesis. Immunity.

2018;49(6):1062-76 e6. Epub 2018/11/18. doi: 10.1016/j.immuni.2018.09.018. PubMed

PMID: 30446388; PubMed Central PMCID: PMCPMC6303217.

42. van Zandbergen G, Hermann N, Laufs H, Solbach W, Laskay T. Leishmania

promastigotes release a granulocyte chemotactic factor and induce interleukin-8 release

but inhibit gamma interferon-inducible protein 10 production by neutrophil granulocytes.

Infect Immun. 2002;70(8):4177-84. Epub 2002/07/16. PubMed PMID: 12117926;

PubMed Central PMCID: PMCPMC128123.

43. Badolato R, Sacks DL, Savoia D, Musso T. Leishmania major: infection of human

monocytes induces expression of IL-8 and MCAF. Exp Parasitol. 1996;82(1):21-6. Epub

1996/01/01. doi: 10.1006/expr.1996.0003. PubMed PMID: 8617327.

44. Racoosin EL, Beverley SM. Leishmania major: promastigotes induce expression

of a subset of chemokine genes in murine macrophages. Exp Parasitol. 1997;85(3):283-

95. Epub 1997/03/01. doi: 10.1006/expr.1996.4139. PubMed PMID: 9085925.

45. Wilson ME, Innes DJ, Sousa AD, Pearson RD. Early histopathology of

experimental infection with Leishmania donovani in hamsters. J Parasitol.

1987;73(1):55-63. Epub 1987/02/01. PubMed PMID: 3572666.

46. Koppenol-Raab M, Sjoelund V, Manes NP, Gottschalk RA, Dutta B, Benet ZL, et

al. Proteome and Secretome Analysis Reveals Differential Post-transcriptional

Regulation of Toll-like Receptor Responses. Mol Cell Proteomics. 2017;16(4 suppl

1):S172-S86. Epub 2017/02/27. doi: 10.1074/mcp.M116.064261. PubMed PMID:

28235783; PubMed Central PMCID: PMCPMC5393387.

47.   Clay GM, Valadares DG, Graff JW, Ulland TK, Davis RE, Scorza BM, et al. An

Anti-Inflammatory Role for NLRP10 in Murine Cutaneous Leishmaniasis. J Immunol.

2017;199(8):2823-33. Epub 2017/09/22. doi: 10.4049/jimmunol.1500832. PubMed

PMID: 28931602; PubMed Central PMCID: PMCPMC5679237.

48.   Silva-Almeida M, Souza-Silva F, Pereira BA, Ribeiro-Guimaraes ML, Alves CR.

Overview of the organization of protease genes in the genome of Leishmania spp.

Parasit Vectors. 2014;7:387. Epub 2014/08/22. doi: 10.1186/1756-3305-7-387. PubMed

PMID: 25142315; PubMed Central PMCID: PMCPMC4158035.

49.   Rasmussen HS, McCann PP. Matrix metalloproteinase inhibition as a novel

anticancer strategy: a review with special focus on batimastat and marimastat.

Pharmacol Ther. 1997;75(1):69-75. Epub 1997/01/01. PubMed PMID: 9364582.

50.   Day TA, Chen GZ. The metalloprotease inhibitor 1,10-phenanthroline affects

Schistosoma mansoni motor activity, egg laying and viability. Parasitology. 1998;116 (

Pt 4):319-25. Epub 1998/05/20. PubMed PMID: 9585934.

51.   Chaudhuri G, Chaudhuri M, Pan A, Chang KP. Surface acid proteinase (gp63) of

Leishmania mexicana. A metalloenzyme capable of protecting liposome-encapsulated

proteins from phagolysosomal degradation by macrophages. J Biol Chem.

1989;264(13):7483-9. Epub 1989/05/05. PubMed PMID: 2708373.

52.   Chen L, Zhang ZH, Watanabe T, Yamashita T, Kobayakawa T, Kaneko A, et al.

The involvement of neutrophils in the resistance to Leishmania major infection in

susceptible but not in resistant mice. Parasitol Int. 2005;54(2):109-18. Epub 2005/05/04.

doi: 10.1016/j.parint.2005.02.001. PubMed PMID: 15866472.

53. Lima GM, Vallochi AL, Silva UR, Bevilacqua EM, Kiffer MM, Abrahamsohn IA.

The role of polymorphonuclear leukocytes in the resistance to cutaneous

Leishmaniasis. Immunol Lett. 1998;64(2-3):145-51. Epub 1998/12/31. PubMed PMID:

9870666.

54. McFarlane E, Perez C, Charmoy M, Allenbach C, Carter KC, Alexander J, et al.

Neutrophils contribute to development of a protective immune response during onset of

infection with Leishmania donovani. Infect Immun. 2008;76(2):532-41. Epub

2007/12/07. doi: 10.1128/IAI.01388-07. PubMed PMID: 18056477; PubMed Central

PMCID: PMCPMC2223441.

55. Novais FO, Santiago RC, Bafica A, Khouri R, Afonso L, Borges VM, et al.

Neutrophils and macrophages cooperate in host resistance against Leishmania

braziliensis infection. J Immunol. 2009;183(12):8088-98. Epub 2009/11/20. doi:

10.4049/jimmunol.0803720. PubMed PMID: 19923470.

56. Peters NC, Egen JG, Secundino N, Debrabant A, Kimblin N, Kamhawi S, et al. In

vivo imaging reveals an essential role for neutrophils in leishmaniasis transmitted by

sand flies. Science. 2008;321(5891):970-4. Epub 2008/08/16. doi:

10.1126/science.1159194. PubMed PMID: 18703742; PubMed Central PMCID:

PMCPMC2606057.

57. Rousseau D, Demartino S, Ferrua B, Michiels JF, Anjuere F, Fragaki K, et al. In

vivo involvement of polymorphonuclear neutrophils in Leishmania infantum infection.

BMC Microbiol. 2001;1:17. Epub 2001/10/10. PubMed PMID: 11591218; PubMed

Central PMCID: PMCPMC57739.

58.   Smelt SC, Cotterell SE, Engwerda CR, Kaye PM. B cell-deficient mice are highly

resistant to Leishmania donovani infection, but develop neutrophil-mediated tissue

pathology. J Immunol. 2000;164(7):3681-8. Epub 2000/03/22. PubMed PMID:

10725726.

59.   Tacchini-Cottier F, Zweifel C, Belkaid Y, Mukankundiye C, Vasei M, Launois P, et

al. An immunomodulatory function for neutrophils during the induction of a CD4+ Th2

response in BALB/c mice infected with Leishmania major. J Immunol.

2000;165(5):2628-36. Epub 2000/08/18. PubMed PMID: 10946291.

60.   Thalhofer CJ, Chen Y, Sudan B, Love-Homan L, Wilson ME. Leukocytes infiltrate

the skin and draining lymph nodes in response to the protozoan Leishmania infantum

chagasi. Infect Immun. 2011;79(1):108-17. Epub 2010/10/13. doi: 10.1128/IAI.00338-

10. PubMed PMID: 20937764; PubMed Central PMCID: PMCPMC3019875.

61.   Inbar E, Hughitt VK, Dillon LA, Ghosh K, El-Sayed NM, Sacks DL. The

Transcriptome of Leishmania major Developmental Stages in Their Natural Sand Fly

Vector. MBio. 2017;8(2). Epub 2017/04/06. doi: 10.1128/mBio.00029-17. PubMed

PMID: 28377524; PubMed Central PMCID: PMCPMC5380837.

62.   Sacks D, Noben-Trauth N. The immunology of susceptibility and resistance to

Leishmania major in mice. Nat Rev Immunol. 2002;2(11):845-58. Epub 2002/11/05. doi:

10.1038/nri933. PubMed PMID: 12415308.

63.   Leifso K, Cohen-Freue G, Dogra N, Murray A, McMaster WR. Genomic and

proteomic expression analysis of Leishmania promastigote and amastigote life stages:

the Leishmania genome is constitutively expressed. Mol Biochem Parasitol.

2007;152(1):35-46. Epub 2006/12/26. doi: 10.1016/j.molbiopara.2006.11.009. PubMed

PMID: 17188763.

64.   Cronemberger-Andrade A, Aragao-Franca L, de Araujo CF, Rocha VJ, Borges-

Silva Mda C, Figueira CP, et al. Extracellular vesicles from Leishmania-infected

macrophages confer an anti-infection cytokine-production profile to naive macrophages.

PLoS Negl Trop Dis. 2014;8(9):e3161. Epub 2014/09/19. doi:

10.1371/journal.pntd.0003161. PubMed PMID: 25232947; PubMed Central PMCID:

PMCPMC4169249.

65.   Silverman JM, Clos J, Horakova E, Wang AY, Wiesgigl M, Kelly I, et al.

Leishmania exosomes modulate innate and adaptive immune responses through effects

on monocytes and dendritic cells. J Immunol. 2010;185(9):5011-22. Epub 2010/10/01.

doi: 10.4049/jimmunol.1000541. PubMed PMID: 20881185.

66.   Roque AL, Jansen AM. Wild and synanthropic reservoirs of Leishmania species

in the Americas. Int J Parasitol Parasites Wildl. 2014;3(3):251-62. Epub 2014/11/27. doi:

10.1016/j.ijppaw.2014.08.004. PubMed PMID: 25426421; PubMed Central PMCID:

PMCPMC4241529.

67.   Baguet A, Epler J, Wen KW, Bix M. A Leishmania major response locus

identified by interval-specific congenic mapping of a T helper type 2 cell bias-controlling

quantitative trait locus. J Exp Med. 2004;200(12):1605-12. Epub 2004/12/15. doi:

10.1084/jem.20040334. PubMed PMID: 15596523; PubMed Central PMCID:

PMCPMC2211989.

68.   Yu CL, Summers RM, Li Y, Mohanty SK, Subramanian M, Pope RM. Rapid

identification and quantitative validation of a caffeine-degrading pathway in

Pseudomonas sp. CES. J Proteome Res. 2015;14(1):95-106. Epub 2014/10/29. doi:

10.1021/pr500751w. PubMed PMID: 25350919.

69.   Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. Jalview Version 2-

-a multiple sequence alignment editor and analysis workbench. Bioinformatics.

2009;25(9):1189-91. Epub 2009/01/20. doi: 10.1093/bioinformatics/btp033. PubMed

PMID: 19151095; PubMed Central PMCID: PMCPMC2672624.

820
821
822

Figure legends

Figure 1: *L. major* diminishes LPS-induced CXCL1 release by BMDMs. (A) BMDMs were treated with LPS (20 ng/ml) in the presence or absence of *L. major* (20 MOI) for 48h and cell culture supernatants were analyzed for the indicated cytokines. (B) BMDMs were treated with PBS or *L. major* (20 MOI) for 6h. Then, the cells were washed to remove extracellular *L. major* and stimulated with LPS for next 42h and cell culture supernatants were subjected to cytokines analysis as in (A). Data are representative of at least three independent experiments. Results are represented as mean $\pm$ SEM. ***P<0.001.

Figure 2: *L. major* inhibits CXCL1 detection in a cell free system. (A) Conditioned supernatants from 24h LPS (20 ng/ml)-stimulated BMDMs were treated with or without *L. major* for 24h and cytokines were analyzed by ELISA. (B) 24h LPS treated conditioned media from BMDMs were treated with or without *L. major*, *L. major* lysate (*Lm* lysate), or *L. major* culture supernatant (*Lm* sup) for 24h and cytokines were measured as in (A). Data are representative of at least three independent experiments. Results are represented as mean $\pm$ SEM. ****P<0.0001.

Figure 3: Selective inhibitory activity of *L. major* on recombinant murine CXCL1 (A) Sequence alignment of recombinant murine (rm) and recombinant human (rh) CXCL1, CXCL2, and CXCL8. (B) Rm-CXCL1, (C) rh-CXCL1, (D) rm-CXCL2, (E) rh-CXCL2, (F) rh-CXCL8, and (G) rm-TNF were incubated with or without *Lm* sup for indicated time-points and ELISA was performed to quantify the mentioned cytokine/chemokines. (H)

Rm-CXCL1 was incubated with or without *Lm* sup or *Lm* exosomes derived from indicated

strains of *L. major* for 24h, and CXCL1 quantification was performed as in (G). Data are

representative of at least three independent experiments. Results are represented as

mean $\pm$ SEM. ****$P<0.0001$.

Figure 4: Biological activity of *L. major*-treated CXCL1 is diminished (A) Schematic

representation of experimental design. (B) Recombinant murine CXCL1 was incubated

with *Lm* lysate for 24h and CXCL1 levels was determined by ELISA. (C) BMDMs were

treated *Lm* lysate, rm-CXCL1, or *Lm* lysate + rm-CXCL1 from (B) for indicated time-points

and mRNA expression of pro-inflammatory genes were evaluated by qPCR. Data are

representative of at least three independent experiments.

Figure 5: *L. major* cleaves CXCL1 at the C-terminal end. (A) A silver-stained SDS-

PAGE demonstrate time-dependent cleavage of rm-CXCL1. Rm-CXCL1 were left alone

or treated with *Lm* lysate for the indicated time points. Rm-CXCL1 bands were enlarged

to show the cleavage products on right. (B) Indicated chemokines were treated with *Lm*

lysate for indicated time-points and subjected to SDS-PAGE followed by silver-staining.

(C) Hypothesized *L. major* cleavage site on rm-CXCL1 based on the observed cleaved

rm-CXCL1 bands in (A and B). Arrow denotes possible cleavage site on CXCL1 (D)

Identification of CXCL1 cleavage site by Mass Spectrometry analysis (raw data presented

in Supplemental Fig. 4). Arrow represents cleavage site on rm-CXCL1 targeted by *L.*

major. Data in A and B are representative of at least three independent experiments.

Figure 6: *L. major* metalloprotease cleaves CXCL1. (A) Recombinant CXCL1 was

incubated with or without *Lm* sup containing (B) Roche cOmplete protease inhibitor, or

(C) Sigma P8340 protease inhibitor for indicated time-points and subjected to CXCL1

quantification using ELISA. (D) Recombinant CXCL1 was treated with *Lm* sup in th

presence of (E) EDTA or (F) 1,10-Phenanthroline for indicated time-points and CXCL1

degradation determined by ELISA. (G) Recombinant CXCL1 was treated with *Lm* sup in

the presence of 1,10-Phenanthroline for indicated time-points and CXCL1 degradation

determined by silver staining. (H) *Lm* lysate was immunoprecipitated with GP63 antibody

and Western blot was carried out to evaluate the levels of GP63 protein. (I) The fractions

from (H) were incubated with rm-CXCL1 for indicated time-points and ELISA was

performed to quantify the levels of CXCL1. Data are representative of at least three

independent experiments. Results are represented as mean $\pm$ SEM. ***P<0.001,

****P<0.0001.

Figure 7: Synthetic peptide spanning rm-CXCL1 cleavage site inhibits *L. major*-

mediated CXCL1 degradation. (A) Schematic of synthetic peptide generation.

Alignment of full length murine CXCL1 with mature murine CXCL1 demonstrating the

schematic for generation of peptide#1, peptide#2 and blocking peptide. Peptide#1 and

peptide#2 were generated from the N-terminal region of full length murine CXCL1.

Blocking peptide spanning the *L. major* cleavage site on murine CXCL1 were generated.

Lm lysate-mediated degradation of rm-CXCL1 in the presence of (B) Activity of synthetic

peptides from (A) in inhibiting L. major-mediated degradation of rm-CXCL1. (C) Model of

L. major metalloprotease function on CXCL1 during infection. Data are representative of

at least three independent experiments. Results are represented as mean $\pm$ SEM.

****P<0.0001.

Supplemental Figure 1: CXCL1, IL-6, and TNF secreted by LPS-activated BMDMs

are stable in cell culture conditions. Conditioned supernatants from 24h LPS (20

ng/ml) stimulated BMDMs were collected and further cultured for 0, 24, or 48h and stability

of CXCL1, IL-6 and TNF in the culture were determined by ELISA. Data are representative

of at least three independent experiments. Results are represented as mean $\pm$ SEM.

Supplemental Figure 2: Heat inactivation of *Lm* lysate or *Lm* sup rescues CXCL1

detection. (A) Experimental Design. (B) Conditioned supernatants from 24h LPS (20

ng/ml) stimulated BMDMs were collected and subjected to 24h treatment with control or

boiled (20 min at 100ºC) *Lm* lysate or *Lm* sup. Levels of CXCL1, IL-6 and TNF were

determined by ELISA. Results are represented as mean $\pm$ SEM. ****P<0.0001.

Supplemental Figure 3: *L. major* reduces rm-CXCL1 levels in culture. Rm-CXCL1

were left alone or treated with *Lm* sup (control or boiled) for up to 48 hours in culture. The

quantity of rm-CXCL1 in the culture at 0, 24 and 48 h were determined by ELISA. Data

are representative of at least three independent experiments. Results are represented as

mean $\pm$ SEM.

Supplemental Figure 4: Acute time course of *L. major*-induced CXCL1 cleavage. Rm-

CXCL1 were left alone or treated with *Lm* lysate for acute time points as indicated to

determine the precise time course of rm-CXCL1 cleavage. Silver staining was used to

visualize rm-CXCL1. Data are representative of at least three independent experiments.

Supplemental Figure 5: Identification of CXCL1 cleavage site and cleaved products.

(A) A silver-stained SDS-PAGE showing cleavage of full length (BAND 1) and cleaved

rm-CXCL1 (BAND 2). (B) Mass spectrometry analysis of BAND 1 (full length CXCL1) and

tryptic peptide coverage analysis confirm a full length rm-CXCL1. (C) Mass spectrometry

analysis of BAND 2 (cleaved CXCL1) and tryptic peptide coverage analysis confirm a C-

terminal end cleavage after K65 residue.

Supplemental Figure 6: Proteolytic activity of _L. major_ on CXCL1 is not prevented

by Roche cOmplete Inhibitor or Marimastat (A) Conditioned supernatants from 24h

LPS (20 ng/ml)-stimulated BMDMs were collected and treated with _Lm_ sup in the

presence or absence of Roche cOmplete protease inhibitor. Levels of CXCL1, IL-6 and

TNF were determined by ELISA. (B) Rm-CXCL1 was treated with _Lm_ sup in the presence

or absence of Marimastat (10μM) for indicated time-points and ELISA was performed to

evaluate the levels of CXCL1. Data are representative of at least three independent

experiments. Results are represented as mean $\pm$ SEM. ***$P<0.001$, ****$P<0.0001$.

934