TITLE

Transfer of *Xanthomonas campestris* pv. *arecae,* and *Xanthomonas campestris* pv. *musacearum* to

Xanthomonas vasicola (Vauterin) as *Xanthomonas vasicola* pv. *arecae* comb. nov., and *Xanthomonas*

vasicola pv. *musacearum* comb. nov. and description of *Xanthomonas vasicola* pv. *vasculorum* pv.

nov.

AUTHORS

David J. Studholme, Emmanuel Wicker, Adam Bogdanove, Kirk Broders, Zoe Dubrow, Murray Grant,
Jeffrey B. Jones, Georgina Karamura, Jillian Lang, Jan Leach, Sadik Muzemil, George Mahuku, Gloria
Valentine Nakato, Teresa Coutinho, Julian Smith, Carolee T. Bull

Corresponding author: David J. Studholme (d.j.studholme@exeter.ac.uk)

ABSTRACT

Recent DNA sequence and other data indicated that several important bacterial pathogens should

be transferred into the species *Xanthomonas vasicola* Vauterin 1995. The first objective of this letter

is to propose the transfer of *X. campestris* pv. *musacearum* (Yirgou and Bradbury 1968) Dye 1978 to

X. vasicola Vauterin 1995. The second objective is to give a clear overview of the different

evolutionary lineages that constitute the species *X. vasicola*, in the light of recent genomics analyses.

These analyses also indicate that strains described as [*X. campestris* pv. *zeae*] (Qhobela, Claflin, and

Nowell 1990; Coutinho and Wallis 1991) fall within the species *X. vasicola* Vauterin 1995.

Furthermore, the sequence of its *gyrB* gene suggested that *X. campestris* pv. *arecae* (Rao and Mohan

1970) Dye 1978 is closely related to the type strain of *X. vasicola*. Note that in this manuscript

pathovar names that have no valid standing in nomenclature are presented with square brackets as

is standard.

AUTHOR ADDRESSES

• David J. Studholme: Biosciences, University of Exeter, Exeter, United Kingdom
• Emmanuel Wicker: UMR IPME, Univ Montpellier, CIRAD, IRD, Montpellier, France
• Adam Bogdanove: Plant Pathology and Plant-Microbe Biology Section, School of Integrative
Plant Science, Cornell University, 334 Plant Science Building, Ithaca, NY 14853, USA
• Kirk Broders: Department of Bioagricultural Sciences and Pest Management, Colorado State
University
• Zoe Dubrow: Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant
Science, Cornell University, 334 Plant Science Building, Ithaca, NY 14853, USA
• Murray Grant: School of Life Sciences, Gibbet Hill, University of Warwick, Coventry, CV4 7AL,
UK
• Jeffrey B. Jones: University of Florida, Plant Patholgoy Dept. , 1453 Fifield Hall , Gainesville,
Florida, United States , 32611
• Georgina Karamura: Bioversity International, Uganda
• Jillian Lang: Department of Bioagricultural Sciences and Pest Management, Colorado State
University, Fort Collins, CO 80523
• Jan Leach: Bioagricultural Sciences and Pest Management, Colorado State University, Ft
Collins, CO 80523
• Sadik Muzemil: Southern Agricultural Research Institute (SARI), Areka Agricultural Research
Center, Areka, Ethiopia
• George Mahuku: International Institute of Tropical Agiculture (IITA), East Africa Hub, IITA-
Tanzania P.O. Box 34441, Dar es Salaam, Tanzania
• Gloria Valentine Nakato: International Institute of Tropical Agiculture (IITA), Plot 15B,
Naguru East Road, Upper Naguru, P.O. Box 7878, Kampala, Uganda
• Teresa Coutinho: Department of Microbiology and Plant Pathology, Centre for Microbial
Ecology and Genomics (CMEG), Forestry and Agricultural Biotechnology Institute (FABI),
University of Pretoria, Private Bag X28, Pretoria 0028, South Africa
• Julian Smith: Fera Science Ltd. York, UK
• Carolee T. Bull: Department of Plant Pathology and Environmental Microbiology, Penn State
University, University Park, Pennsylvania, USA

LETTER TO THE EDITOR

The genus *Xanthomonas* comprises gamma-Proteobacteria that collectively cause disease on more

than 400 species of plants (Hayward 1993), though some species and strains are apparently non-

pathogenic (Vauterin et al. 1996) and some have been isolated from clinical samples such as skin

microbiota (Seité, Zelenkova, and Martin 2017). Historically, taxonomy of *Xanthomonas* was tied to

the plant host of isolation (Vauterin et al. 1995), with the genus being split into large numbers of

species, each defined by this single phenotypic feature (Dye 1962). Subsequently, most of the

species were transferred (lumped) into a single species, *X. campestris*, because the organisms could

not be distinguished from one another by phenotypic and physiological tests (Lapage et al. 1992). As

a temporary solution, and to help to maintain a connection with the historical and plant pathological

literature, these nomenspecies were designated as pathovars within *X. campestris*, each defined by

host range or disease syndrome (Dye et al. 1980). More recently, DNA sequence comparisons and

biochemical approaches revealed that some of the host ranges of pathovars of *X. campestris* were

not correlated with inferred phylogenies (Parkinson et al. 2007, 2009; Rodriguez-R et al. 2012).

There have been heroic advances to improve the taxonomy of the genus as a whole (Vauterin et al.

1990; Vauterin, Rademaker, and Swings 2000; Rademaker et al. 2005; Vauterin et al. 1995) and

individual taxa (da Gama et al. 2018; Constantin et al. 2016), based on phenotypic, chemotaxonomic

and genotypic analyses, but in a number of taxa there remain issues not fully resolved.

The bacterial pathogen *X. campestris* pv. *musacearum* (Yirgou and Bradbury 1968) Dye 1978

presents a major threat to cultivation of banana and enset crops in central and eastern Africa, where

is causes banana Xanthomonas wilt (BXW) and enset Xanthomonas wilt (EXW). Originally described

as *X. musacearum* (Yirgou and Bradbury 1968), this pathogen was first isolated from enset and

banana in Ethiopia in the late 1960s and early 1970s, respectively (Yirgou and Bradbury 1968, 1974),

although symptoms consistent with EXW were reported for Ethiopia as early as the 1930s (Castellani

1939). However, only in the 21st century did the disease establish in the banana-growing areas of

Burundi, Democratic Republic of Congo, Kenya, Rwanda, Tanzania and Uganda, (Biruma, Pillay, and

Tripathi 2007; Tushemereirwe et al. 2004; Ndungo et al. 2006; Reeder et al. 2007; Carter et al. 2010).

In this region around the Great Lakes of eastern and central Africa, BXW disease severely challenges

livelihoods and food security (Blomme et al. 2017; Shimwela et al. 2016; Tinzaara et al. 2016;

Blomme et al. 2013; Biruma et al. 2007; Nakato, Mahuku, and Coutinho 2018).

There is confusion in the literature about the taxonomy of this bacterium. Since its assignment to *X.*

campestris (Young et al. 1978), molecular sequence and biochemical data indicate that this pathogen

is more closely related with *X. vasicola* (Parkinson et al. 2007; Aritua et al. 2007) as detailed below.

Thus, the first objective of this letter is to propose the transfer of *X. campestris* pv. *musacearum*

(Yirgou and Bradbury 1968) Dye 1978 to *X. vasicola* Vauterin 1995. The second objective is to give a

clear overview of the different evolutionary lineages that constitute the species *X. vasicola*, in the

light of recent genomics analyses. These analyses also indicate that strains described as [*X.*

campestris pv. *zeae*] (Qhobela, Claflin, and Nowell 1990; Coutinho and Wallis 1991) fall within a

clade of *X. campestris* pv. *vasculorum* (Cobb 1894) Dye 1978 that belongs within the species *X.*

vasicola Vauterin 1995. Furthermore, the sequence of its *gyrB* gene suggested that *X. campestris* pv.

arecae (Rao and Mohan 1970) Dye 1978 is closely related to the type strain of *X. vasicola* (Parkinson

et al. 2009). Note that in this manuscript pathovar names that have no valid standing in

nomenclature are presented with square brackets as is standard (Bull et al. 2012).

The species *X. vasicola* (Vauterin et al. 1995) was created to encompass *X. campestris* pv. *holcicola*

(Elliott 1930) Dye 1978 and a subset of strains (not including the pathotype) of *X. campestris* pv.

vasculorum (Cobb 1894) Dye 1978 (Young et al. 1978; Vauterin et al. 1995). Taxonomic studies

revealed that *X. campestris* pv. *vasculorum* contained groups of strains that are clearly

distinguishable from the pathotype strain (of *X. campestris* pv. *vasculorum*) by phenotypic and

molecular traits despite their shared host ranges (Vauterin et al. 1992; Péros et al. 1994; Dookun,

Stead, and Autrey 2000; Stead 1989; Vauterin et al. 1995; Destéfano et al. 2003). Vauterin's type-B

strains are distinguished from type-A strains (to which the pathotype strain of *X. campestris* pv.

vasculorum belongs) by SDS-PAGE of proteins, gas chromatography of fatty acid methyl esters and

DNA-DNA hybridization (Yang et al. 1993). Type-A and type-B strains can also be distinguished by

PCR-RFLP analysis (Destéfano et al. 2003). Table 1 lists examples of *X. campestris* pv. *vasculorum*

(Cobb 1894) Dye 1978 strains that were classified in one or more of those studies. Vauterin and

colleagues assigned type-A strains, along with the pathotype, to *X. axonopodis* pv. *vasculorum*

(Cobb) Vauterin, Hoste, Kersters & Swings and type-B to [*X. vasicola* pv. *vasculorum*] (Vauterin et al.

1995). However, we note that this pathovar is invalid because of the lack of a formal proposal

differentiating it from other pathovars (Young et al. 2004) and no designation of a pathotype strain.

For a given organism, competing classifications and invalid names result in three different valid

species names, *X. campestris*, *X. axonopodis* and *X. vasicola,* having been used in the literature. For

example, various authors have referred to strain NCPPB 1326 as *X. campestris* pv. *vasculorum*, *X.*

axonopodis pv. *vasculorum* (to which the strain clearly does not belong) or [*X. vasicola* pv.

vasculorum] (Wasukira et al. 2014; Lewis Ivey, Tusiime, and Miller 2010; Qhobela, Claflin, and Nowell

1990; Qhobela and Claflin 1992). Type-B strains NCPPB 702, NCPPB 1326 and NCPPB 206 were

erroneously described as *X. axonopodis* pv. *vasculorum* (Lewis Ivey, Tusiime, and Miller 2010) though

they are clearly members of *X. vasicola*. However, we acknowledge that examples of mistakes such

as these will not likely be resolved by transfer of the pathovars from *X. campestris* into *X. vasicola*.

A further source of confusion is the status of strains isolated from maize for which some authors use

the invalid name [*X. campestris* pv. *zeae*] (Qhobela, Claflin, and Nowell 1990; Coutinho and Wallis

1991). Adding to the confusion, at least one strain of *X. campestris pv. vasculorum* (NCPPB 206)

isolated from maize has the fatty-acid type characteristic of *X. vasicola* (Dookun, Stead, and Autrey

2000); consistent with this, on the basis of phylogenetic analysis of DNA sequence, NCPPB 206 falls

among strains assigned to Vauterin's invalid [*X. vasicola* pv. *vasculorum*] (Wasukira et al. 2014). A

useful nomenclature for this group has become more pressing since the recent outbreak of leaf

streak on corn in the USA, caused by bacteria very closely related to strains previously described as

[*X. campestris* pv. *zeae*]. One of these strains, NCPPB 4614 (=SAM119), has been suggested to be the

eventual pathotype strain of *X. vasicola* pv. *vasculorum* though no valid proposal has been made

(Lang et al. 2017; Korus et al. 2017). Although [*X. vasicola* pv. *vasculorum*] (Vauterin et al. 1995) is

invalid, this name has come to be understood by the community to represent a meaningful

biological reality; that is a set of *X. campestris* pv. *vasculorum* strains that are biochemically and

phylogenetically similar to *X. vasicola*.   Therefore, below we propose a formal description of *X.*

vasicola pv. *vasculorum* pv. nov. that should be considered valid to harmonize the formal

nomenclature with that which is in use. Further, consistent with the previous suggestion (Lang et al.

2017) that strains classified to [*X. vasicola* pv. *vasculorum*] and [*X. campestris* pv. *zeae*] (Vauterin et

al. 1995) are insufficiently distinct to warrant separate pathovars, we therefore propose that [*X.*

vasicola pv. *vasculorum*] group B, [*X. campestris* pv. *zeae*] and phylogenetically closely related strains

isolated from sugarcane and maize be assigned into the newly described *X. vasicola* pv. *vasculorum*

pv. nov.

Vauterin et al. (1995) designated the pathotype strain of X. *vasicola* pv. *holicola* (LMG 736, NCPPB

2417, ICMP 3103 and CFBP 2543) as the type strain of *X. vasicola*, although they did not use the

pathovar epithet for the specific epithet of the species as is most appropriate to indicate this

relationship. The natural host range of *X. vasicola* pv. *holicola* includes the cereal crops millet and

sorghum on which it causes bacterial leaf streak (Table 2). The host range of the strains that Vauterin

et al. (1995) called [*X. vasicola* pv. *vasculorum*] is less well defined because in most of the relevant

pre-1995 literature it is impossible to distinguish between type-A and type-B of *X. campestris* pv.

vasculorum and therefore between *X. axonopodis* pv. *vasculorum* and strains belonging to *X.*

vasicola.   However, *X. campestris* pv. *vasculorum* type-B strains (that is, members of *X. vasicola*)

have been isolated from sugarcane and maize and shown to infect these hosts on artificial

inoculation (Vauterin et al. 1995; Karamura et al. 2015).

Previous studies suggested a close relationship between *X. campestris* pv. *musacearum* (Yirgou and

Bradbury 1968) Dye 1978b and *X. vasicola* pv. *holcicola* (Elliott 1930) Vauterin et al. 1995 based on

fatty acid methyl ester analysis, genomic fingerprinting using rep-PCR and partial nucleotide

sequencing of the *gyrB* gene (Aritua et al. 2007; Parkinson et al. 2009). Draft or complete sequence

assemblies are now available for more than a thousand *Xanthomonas* genomes, including those of

type strains for most species and pathotypes for most pathovars. Genome-wide sequence data can

offer some advantages, such as generally applicable threshold values for species delineation (Glaeser

and Kämpfer 2015; Meier-Kolthoff et al. 2013; Meier-Kolthoff, Klenk, and Göker 2014; Richter and

Rosselló-Móra 2009). Therefore, we further explored relationships among these organisms using

whole genome sequences. We calculated pairwise average nucleotide identity (ANI) between *X.*

campestris pv. *musacearum* and representative *Xanthomonas* strains, including all available species

type strains and relevant pathotype strains. A representative subset of these pairwise ANI

percentages is tabulated in Figure 1. This revealed that the pathotype strain of *X. campestris* pv.

musacearum (Yirgou and Bradbury 1968) Dye 1978b shares 98.43 % ANI with the type strain of *X.*

vasicola but only 87.27 % with the type strain of *X. campestris*. As expected, strains of *X. vasicola* pv.

holcicola share high ANI (> 99.6 %) with the *X. vasicola* type strain NCPPB 2417, which is also the

pathotype strain of X. *vasicola* pv. *holcicola* (Elliott 1930) Vauterin et al. 1995. Also as expected,

strains of *X. campestris* pv. *vasculorum* previously called [*X. vasicola* pv. *vasculorum*] or [*X.*

campestris pv. *zeae*], including sequenced strain SAM119 (=NCPPB 4614) from corn isolated by T.

Coutinho (Qhobela, Claflin, and Nowell 1990), share > 98.5 % ANI with the type strain of *X. vasicola*,

supporting the need to transfer these strains to this species. Furthermore, unclassified strains NCPPB

902, NCPPB 1394, NCPPB 1395 and NCPPB 1396, from *Tripsacum laxum* (Mulder 1961) and the

pathotype strain of X. *campestris* pv. *arecae* (Rao and Mohan 1970) Dye 1978 (NCPPB 2649) all share

more than 98 % ANI with the type strain of *X. vasicola*, which places them unambiguously within *X.*

vasicola. The next-nearest species to *X. vasicola* is *X. oryzae*; ANI between the respective type strains

of these two species is 91.7%. It has been proposed that the boundary of a prokaryotic species can

be delimited by 95 to 96% (Richter and Rosselló-Móra 2009). By this criterion, *X. campestris* pv.

arecae, *X. campestris* pv. *musacearum* and strains from corn that are referred to by the invalid name

[*Xanthomonas vasicola* pv. *zeae*] clearly fall within *X. vasicola* and outside *X. campestris*.

The high ANI levels clearly delineate a genomo-species that includes the type strain *X. vasicola*

NCPPB 2417. Nevertheless, despite the usefulness of ANI for delimiting species boundaries, it does

not include any model of molecular evolution and thus is unsuited for phylogenetic reconstruction.

Therefore, we used RaxML via the RealPhy pipeline (Bertels et al. 2014; Stamatakis, Ludwig, and

Meier 2005) to elucidate phylogenetic relationships, based on genome-wide sequencing data. This

approach has the additional advantage of being based on sequence reads rather than on genome

assemblies, where the latter may be of variable quality and completeness (Bertels et al. 2014).

Figure 2 depitcs the phylogeny of *X. vasicola* based on RealPhy analysis of genome-wide sequence

data. Pathovars *X. vasicola* pv. *holcicola* and *X. campestris* pv. *musacearum* are monophyletic,

comprising well supported clades within the *X. vasicola* genomospecies. A third well supported clade

includes the four "*Xanthomonas* sp." strains originating from *Tripsacum laxum*. A fourth clade

consists of mostly *X. campestris* pv. *vasculorum* strains isolated from sugarcane but also includes *X.*

campestris pv. *vasculorum* strain NCPPB 206 isolated from maize and several strains from maize

attributed to the invalid name [*X. campestris* pv. *zeae*]. This indicates that sequenced strains of [*X.*

campestris pv. *zeae*] from corn (Sanko et al. 2018; Lang et al. 2017; Qhobela, Claflin, and Nowell

1990; Coutinho and Wallis 1991) are monophyletic and fall within the clade containing type-B strains

of *X. campestris* pv. *vasculorum* (Figure 2). The single sequenced pathotype strain of *X. campestris*

pv. *arecae* falls immediately adjacent to the *X. vasicola* clade containing strains from corn and *X.*

campestris pv. *vasculorum* type B strains (Figure 2).

Overall, our molecular sequence analyses strongly point to the existence of a phylogenetically

coherent species *X. vasicola* that includes strains previously assigned to *X. campestris* pathovars

musacearum, arecae, and some strains of *X. campestris* pv. *vasculorum* and strains collected from

corn and *T. laxum* grass that have not been previously assigned to species nor pathovar. Here we

propose that the pathovar *Xanthomonas vasicola* pv. *vasculorum* pv. nov. includes strains formerly

classified as *X. campestris* pv. *vasculorum* but distinguishable from *X. axonopodis* pv. *vasculorum*

(Cobb) Vauterin, Hoste, Kersters & Swings by protein SDS-PAGE, fatty acid methyl esterase (FAME)

analysis and DNA hybridisation (Vauterin et al. 1992; Yang et al. 1993; Vauterin et al. 1995). Our

analyses also support the transfer of *X. campestris* pv. *arecae* (Rao and Mohan 1970) Dye 1978 to *X.*

vasicola. Although only a single genome of this pathovar has been sequenced, that genome belongs

to the pathotype strain for this pathovar (Rao and Mohan 1970; C.T. Bull et al. 2010).

Our results are consistent with previous evidence for similarity between *X. campestris* pv.

musacearum (Yirgou and Bradbury 1968) Dye 1978 and strains of *X. vasicola*, based on FAME,

genomic fingerprinting with rep-PCR and *gyrB* sequencing (Aritua et al. 2007; Parkinson et al. 2007).

The formal species description for *X. vasicola* states that this species can be clearly distinguished by

its FAME profiles (Vauterin et al. 1995). Pathogenicity studies demonstrated phenotypic

distinctiveness of *X. campestris* pv. *musacearum* (Yirgou and Bradbury 1968) Dye 1978 on banana; *X.*

campestris pv. *musacearum* produces severe disease whereas X. *vasicola* pv. *holcicola* NCPPB 2417

and *X. campestris* pv. *vasculorum* NCPPB 702 (which belongs to *X. vasicola*) showed no symptoms

(Aritua et al. 2007). The species description (Vauterin et al. 1995) also states that *X. vasicola* is

characterised by metabolic activity on the carbon substrates D-psicose and L-glutamic acid, and by a

lack of metabolic activity on the carbon substrates N-acetyl-D-galactosamine, L-arabinose, a-D-

lactose, D-melibiose, P-methyl-D- glucoside, L-rhamnose, D-sorbitol, formic add, D-galactonic acid

lactone, D-galacturonic acid, D-gluconic acid, D-glucuronic acid, p-hydroxyphenylacetic acid, a-

ketovaleric acid, quinic acid, glucuronamide, L-asparagine, L-histidine, L-phenylalanine, urocanic

acid, inosine, uridine, thymidine, DL-a-glycerol- phosphate, glucose 1-phosphate, and glucose 6-

phosphate. We are not aware that these metabolic activities have been tested for *X. campestris* pv.

arecae, *X. campestris* pv. *musacearum* and strains referred to as [*X. campestris* pv. *zeae*]; it is

possible that the species description may need to be amended to accommodate any deviation from

this definition among the repositioned pathovars.

Overall, it seems that the species *X. vasicola* (including *X. vasicola* pv. *holcicola*, *X. campestris* pv.

vasculorum type-B strains, [*X. campestris* pv. *zeae*] strains, *X. campestris* pv. *arecae* and some strains

isolated from *T. laxum*) is almost exclusively associated with monocot plants of the families Palmae

and Gramineae. In this respect, it is similar to its closest sibling species *X. oryzae*, whose host range is

limited to Gramineae (Bradbury 1986). The exception is a report of leaf blight and dieback in

Eucalyptus caused by *X. vasicola* (Coutinho et al. 2015), remarkable given the phylogenetic distance

between this dicot plant and the usual monocot hosts of *X. vasicola*; the infected South African

plantation was in an area where sugarcane is grown.

In conclusion, analysis of available genome sequence data, combined with published pathogenicity

and biochemical data, strongly support the transfer of *X. campestris* pathovars *arecae* and

musacearum to the species *X. vasicola* as *X. vasicola* pv. *musacearum* comb. nov. with NCPPB 2005

as the pathotype strain (being the type strain of *X. musacearum* and pathotype strain of *X.*

campestris pv. *musacearum*) and *X. vasicola* pv. *arecae* comb. nov with NCPPB 2649 as the

pathotype strain (being the type strain of *X. arecae* and pathotype strain of *X. campestris* pv.

arecae). Strains NCPPB 206, NCPPB 702, NCPPB 795, NCPPB 890, NCPPB 895, NCPPB 1326, NCPPB

1381, and NCPPB 4614 form a phylogenetically and phenotypically coherent group with a distinctive

host range causing symptoms on maize and sugarcane but not on banana (Aritua et al. 2007;

Karamura et al. 2015). We designate the pathotype strain for *X. vasicola* pv. *vasculorum* pv. *nov.* as

NCPPB 4614. This strain was previously proposed as the pathotype of *X. vasicola* pv. *vasculorum*

(Lang et al. 2017) and causes disease symptoms on maize and sugarcane (Lang et al. 2017) but not

on banana (unpublished observation, Z. Dubrow and A. Bogdanove). Furthermore, given that strains

from corn formerly described by the invalid name [*X. campestris* pv. *zeae*] are members of *X.*

vasicola and have host ranges that can not be distinguished from the pathotype strain of *X. vasicola*

pv. *vasculorum*, we propose that these strains are members of this pathovar. Phylogenetic data

support this as the corn strains represent a sub-clade within strains of *X. campestris* pv. *vasculorum*

that fall within the emended *X. vasicola*.

EMENDED DESCRIPTION OF *XANTHOMONAS VASICOLA* VAUTERIN
ET. AL., 1995.

The characteristics are as described for the genus and the species (Vauterin et al., 1995) extended

with phylogenetic data from this study. The species can be clearly distinguished from other

xanthomonads by MLSA and whole genome sequence analysis with members having more than 98

% ANI with the type strain. SDS-PAGE protein and FAME profiles have been shown to be

distinguishing for some pathovars (Yang et al. 1993; Vauterin et al. 1992; Aritua et al. 2007), by the

presence of metabolic activity on the carbon substrates D-psicose and L-glutamic acid, and by a lack

of metabolic activity on the carbon substrates N-acetyl-D-galactosamine, L-arabinose, a-D-lactose, D-

melibiose, P-methyl-D- glucoside, L-rhamnose, D-sorbitol, formic add, D-galactonic acid lactone, D-

galacturonic acid, D-gluconic acid, D-glucuronic acid, p-hydroxyphenylacetic acid, a-ketovaleric acid,

quinic acid, glucuronamide, L-asparagine, L-histidine, L-phenylalanine, urocanic acid, inosine, uridine,

thymidine, DL-a-glycerol- phosphate, glucose 1-phosphate, and glucose 6-phosphate. The G+C

content is between 63.1 and 63.6 mol % as calculated from whole-genome sequence data. The type

strain is *X. vasicola* pv. *holcicola* LMG 736 (= NCPPB 2417 = ICMP 3103 = CFBP 2543).

***X. vasicola* pv. *holcicola* Vauterin et al., 1995.**

= *X. campestris* pv. *holcicola* (Elliott) Dye 1978.

Description is as presented by Vauterin et al., (1995). The pathovar is distinguished on the

basis of phytopathogenic specialization. As shown here and elsewhere (Lang et al. 2017), the

282 pathovar is distinct from other pathovars by MLSA and genome-wide sequence analysis.

283 According to Bradbury (1986), gelatin and starch are hydrolysed by most isolates examined.

284 The natural host range includes: *Panicum miliaceum, Sorghum* spp., *S. almum, S. bicolor* (*S.*

285 *vulgare*), *S. caffrorum, S. durra, S. halepense, S. sudanense, S. technicum* (*S. bicolor* var.

286 *technicus*), *Zea mays.* The artificial host range (by inoculation) includes *Echinochloa*

287 *frumentacea, Pennisetum typhoides, Setaria italica.*

288 Pathotype strain: PDDCC 3103; NCPPB 2417.

289 ***X. vasicola* pv. *vasculorum* pv. nov.**

290 Description as for the species and this pathovar is distinguished on the basis of

291 phytopathogenic specialization and includes the strains of the former taxon *X. campestris*

292 pv. *vasculorum* type B and pathogens from corn. The pathovar is identified to species and

293 distinguished from other pathovars by its *gyrB* gene sequence (Parkinson et al. 2009) and

294 genome-wide sequence analysis. It is not known whether the strains being transferred to

295 this taxon conform to the species description for metabolic activity. According to previously

296 published work (Coutinho et al. 2015; Aritua et al. 2007; Karamura et al. 2015; Hayward

297 1962)(unpublished observation, S. Dubrow and A. Bogdanove) the natural host range

298 includes: *Saccharum* spp., *Zea mays, Eucalyptus grandis.*

299 Pathotype strain: NCPPB 4614; SAM119.

300 ***X. vasicola* pv. *arecae* (Rao &Mohan) Dye 1978 *comb. nov.***

301 = *X. campestris* pv. *arecae* (Rao &Mohan) Dye 1978.

302 Description as for the species and this pathovar is distinguished on the basis of

303 phytopathogenic specialization. The pathovar is identified to species and distinguished from

304 other pathovars by its *gyrB* gene sequence (Parkinson et al. 2009) and by genome-wide

305 sequence analysis. According to Bradbury (1980) the natural host range includes: *Areca*

306 *catechu* (areca nut). Bradbury (1986) reports the artificial host range to include: *Cocos*

307 *nucifera* (coconut). Needle prick into sugar cane produced limited streaks, but the bacteria

308 did multiply to some extent and could be re-isolated. Disease: leaf stripe. Long, narrow

309 water-soaked lesions, becoming dark brown or black with age. It is not known if the strains

310 being transferred to this taxon conform to the species description for metabolic activity.

311 Pathotype strain: NCPPB 2649; PDDCC 5791.

312 ***X. vasicola* pv. *musacearum* (Yirgou & Bradbury) Dye 1978 *comb. nov.***

313 = *X. campestris* pv. *musacearum* (Yirgou & Bradbury) Dye 1978.

Description as for the species and this pathovar is identified to species and distinguished on

the basis of phytopathogenic specialization and is distinct from other pathovars by its *gyrB*

gene sequence (Parkinson et al. 2009) and genome-wide sequence analysis. Gelatin slowly

liquefied, starch not hydrolysed. Growth quite rapid and very mucoid. According to Bradbury

(1986), the natural hosts include: *Ensete ventricosum* (enset), *Musa* spp. (banana).

Additional hosts by inoculation: *Saccharum* sp. (sugarcane), *Zea mays* (maize) and disease is

exhibited as a bacterial wilt where leaves wilt and wither; yellowish bacterial masses are

found in vascular tissue and parenchyma. It is not known if the strains being transferred to

this taxon conform to the species description for metabolic activity.

Pathotype strain: NCPPB 2005; PDDCC 2870.

REFERENCES

Table 1. Classification of strains previously assigned to X. campestris pv. vasculorum.

Strain	Vauterin (Vauterin et al. 1992, 1995)	Dookun (Dookun, Stead, and Autrey 2000)	Péros (Péros et al. 1994)	Current species assignation
NCPPB 186	Type A	Group A	n/a	X. axonopodis
NCPPB 891	Type A	Group A	G1	X. axonopodis
NCPPB 892	n/a	Group A	n/a	X. axonopodis
NCPPB 893	n/a	Group A	n/a	X. axonopodis
NCPPB 181	Type A	Group B	n/a	X. axonopodis
NCPPB 796 PT	Type A	Group B	n/a	X. axonopodis
NCPPB 899	n/a	Group D	n/a	X. axonopodis
NCPPB 900	n/a	Group D	n/a	X. axonopodis
NCPPB 795	Type B	Group C	n/a	X. vasicola
NCPPB 889	Type B	Group C	n/a	X. vasicola
NCPPB 206	n/a	Group C	n/a	X. vasicola
NCPPB 702	n/a	Group C	n/a	X. vasicola
NCPPB 795	n/a	Group C	n/a	X. vasicola
NCPPB 889	n/a	Group C	n/a	X. vasicola
NCPPB 890	n/a	Group C	n/a	X. vasicola
NCPPB 895	n/a	Group C	n/a	X. vasicola
NCPPB 1326	n/a	Group C	n/a	X. vasicola
NCPPB 1381	n/a	Group C	n/a	X. vasicola

329 [z]*In this table, the superscript PT indicates the pathotype strain of X. campestris pv. vasculorum*

331   ***Table 2. Host ranges of the taxa discussed in this letter.***

Current taxon	Proposed taxon	Pathotype or Type strains	Additional strains in NCPPB known to be part of the newly proposed taxon	Natural hosts	Hosts by inoculation
X. campestris pv. arecae (Rao and Mohan 1970) Dye 1978	X. vasicola pv. arecae pv. nov.	NCPPB 2649 = ICMP 5719 = LMG 533	None	Areca catechu (Bradbury 1986; Kumar 1993, 1983)	Cocos nucifera, Saccharum sp. (Bradbury 1986)
X. campestris pv. musacearum (Yirgou and Bradbury 1968) Dye 1978	X. vasicola pv. musacearum pv. nov	NCPPB 2005 = ATCC 49084 = CFBP 7123 = ICMP 2870 = LMG 785	NCPPB 2251; NCPPB 4378; NCPPB 4379; NCPPB 4380; NCPPB 4381; NCPPB 4383; NCPPB 4384; NCPPB 4386; NCPPB 4387; NCPPB 4388; NCPPB 4389; NCPPB 4390; NCPPB 4391; NCPPB 4392; NCPPB 4393; NCPPB 4394; NCPPB 4395; NCPPB 4433; NCPPB 4434	Ensete ventricosum, Musa sp. (Bradbury 1986), Tripsacum sp. (Unpublished observation, E. Wicker),	Saccharum sp., (Karamura et al. 2015), Zea mays (Karamura et al. 2015; Aritua et al. 2007)
[Xanthomonas vasicola pv. zeae Coutinho and Wallis 1990] [Xanthomonas vasicola pv. zeae Qhobela et al 1990]	X. vasicola pv. vasculorum pv. nov.	NCPPB 4614 = SAM119	None	Zea mays (Coutinho and Wallis 1991)	Sorghum sp. (Lang et al. 2017)
X. vasicola pv. holcicola (Elliott 1930) (Elliott 1930) Vauterin et al. 1995 (synonym of X. campestris pv. holcicola)	X. vasicola pv. holcicola (Elliott 1930) Vauterin et al. 1995	NCPPB 2417 = CFBP 2543 = ICMP 3103 = LMG 736	NCPPB 989; NCPPB 1060; NCPPB 1241; NCPPB 2417; NCPPB 2930; NCPPB 3162	Panicum miliaceum, Sorghum spp., Zea mays (Bradbury 1986)	Echinochloa frumentacea, Pennisetum typhoides, Setaria italica (Bradbury 1986)
X. campestris pv. vasculorum type B = [X. vasicola pv. vasculorum (Vauterin et al., 1995)]	X. vasicola pv. vasculorum pv. nov.	NCPPB 4614 = SAM119	NCPPB 206; NCPPB 702; NCPPB 795; NCPPB 889; NCPPB 890; NCPPB 895; NCPPB 1326; NCPPB 1381; NCPPB 4614	Saccharum spp., Zea mays, Eucalyptus grandis (Coutinho et al. 2015; Bradbury 1986; Vauterin et al. 1995)	Saccharum spp., Zea mays (Karamura et al. 2015)
Xanthomonas sp.	The taxonomic placement of these strains requires further study.	Not applicable	NCPPB 1394; NCPPB 1395; NCPPB 1396; NCPPB 902	Tripsacum laxum (Mulder 1961), Vetiveria zizanoides (Kumar 1993, 1983)	Not known

Figure 1. Average nucleotide identity (ANI) with type strains of Xanthomonas species. *Genome sequence assemblies were obtained from GenBank and aligned against each other and ANI was calculated using the dnadiff function in MUMmer version 4 (Marçais et al. 2018). Accession numbers of the genome assemblies: GCA_000774005.1, GCA_000772705.1, GCA_000277875.1, GCA_000770355.1, GCA_000277995.1, GCA_000159795.2, GCA_000278035.1, GCA_003111865.1, GCA_002191965.1, GCA_002191955.1, GCA_003111905.1, GCA_003111825.1, GCA_000007145.1, GCA_001660815.1, GCA_002939755.1, GCA_001401595.1, GCA_002939725.1, GCA_000724905.2, GCA_000192045.3, GCA_000488955.1, GCA_001401605.1, GCA_002018575.1, GCA_000482445.1 and GCA_002846205.1 (Studholme et al. 2010; Wasukira et al. 2014, 2012; Lang et al. 2017; Sanko et al. 2018; da Silva et al. 2002; Vicente et al. 2017; Harrison and Studholme 2014; Potnis et al. 2011; Jacques et al. 2013).*

Figure 2. Maximum-likelihood phylogenetic tree based on genomic sequencing reads. The maximum likelihood tree was generated using RealPhy (Bertels et al. 2014) and RaxML (Stamatakis, Ludwig, and Meier 2005). Bootstrap values are expressed as percentages of 500 trials. Type and pathotype strains are indicated by 'P' and 'PT' respectively. Whole-genome shotgun sequence reads were obtained from the Sequence Read Archive (Leinonen, Sugawara, and Shumway 2011) via BioProjects PRJNA73853, PRJNA163305, PRJNA163307, PRJNA31213, PRJNA374510, PRJNA374557, PRJNA439013, PRJNA439327, PRJNA439328, PRJNA439329 and PRJNA449864 (Lang et al. 2017; Wasukira et al. 2014, 2012; Sanko et al. 2018).