The type VII secretion system protects *Staphylococcus aureus* against

antimicrobial host fatty acids

Arnaud Kengmo Tchoupa[1], Kate E. Watkins[1], Rebekah A. Jones[1], Agnès Kuroki[2],

Mohammad Tauqeer Alam[1], Sebastien Perrier[1,2,3], Yin Chen[4], Meera Unnikrishnan[1*]

[1]Warwick Medical School, University of Warwick, Coventry, United Kingdom

[2]Department of Chemistry, University of Warwick, Coventry, United Kingdom

[3]Faculty of Pharmacy and Pharmaceutical Sciences, Monash University, Parkville,

Victoria 3052, Australia

[4]School of Life Sciences, University of Warwick, Coventry, United Kingdom

Running title: *S. aureus* T7SS and fatty acid toxicity

[*]Corresponding author

Email: M.Unnikrishnan@warwick.ac.uk.

Summary

The *Staphylococcus aureus* type VII secretion system (T7SS) exports several proteins

that are pivotal for bacterial virulence. The mechanisms underlying T7SS-mediated

staphylococcal survival during infection nevertheless remain unclear. Here we show

that the absence of EsxC, a small secreted effector implicated in bacterial persistence,

results in cell membrane defects in *S. aureus*. Interestingly, isogenic mutants lacking

EsxC, other T7SS effectors EsxA and EsxB, or the membrane-bound ATPase EssC,

are more sensitive to killing by the host-derived antimicrobial fatty acid, linoleic acid

(LA), compared to the wild-type (WT). LA induces more cell membrane damage in the

T7SS mutants compared to the WT. Although WT and mutant strains did not differ in

their ability to bind labelled LA, membrane lipid profiles show that T7SS mutants are

less able to incorporate LA into their membrane phospholipids. Furthermore,

proteomic analyses of WT and mutant cell fractions reveal that, in addition to

compromising membranes, T7SS defects induce oxidative stress and hamper their

response to LA challenge. Thus, our findings indicate that T7SS is crucial for *S. aureus*

membrane integrity and homeostasis, which is critical when bacteria encounter

antimicrobial fatty acids.

Keywords: *Staphylococcus aureus*, Type VII secretion system, long-chain

unsaturated free fatty acids.

Introduction

Staphylococcus aureus is a facultative pathogen that can colonize the skin and nares of healthy individuals. The asymptomatic carriage of *S. aureus* is a major risk for subsequent infections (von Eiff *et al.*, 2001). *S. aureus* infections, which can be healthcare or community-associated, range from benign impetigo to life-threatening bacteraemia (Tong *et al.*, 2015). Clinical management of staphylococcal infections is complicated by the increasing prevalence of multidrug resistant strains (Lee *et al.*, 2018).

The success of *S. aureus* as a deadly pathogen is attributed to an array of virulence factors that facilitate host tissue adhesion and immune response evasion (Gordon & Lowy, 2008). One of these virulence factors is the type VII secretion system (T7SS), also known as the ESAT-6 secretion system (ESS). The orthologous ESX-1 system was initially discovered in *Mycobacterium tuberculosis*, where it is essential for bacterial virulence (Conrad *et al.*, 2017). T7SSs (T7SSb) are found in both Gram-positive and Gram-negative bacteria, although these systems and their secretion machineries appear to be distinct to their mycobacterial counterparts (Unnikrishnan *et al.*, 2017).

In extensively studied strains (COL, RN6390, USA300 and Newman), the T7SS consists of four integral membrane proteins (EsaA, EssA, EssB and EssC), two cytosolic proteins (EsaB and EsaG), five secreted substrates (EsxA, EsxB, EsxC, EsxD and EsaD), and EsaE, which interacts with the T7SS substrates to target them to the secretion apparatus (Cao *et al.*, 2016). A peptidoglycan hydrolase, EssH, was reported to mediate T7SS transport across the bacterial cell wall envelope (Bobrovskyy *et al.*, 2018).

The molecular architecture of the staphylococcal T7SS has not yet been fully

characterized. T7SS integral membrane proteins EsaA, EssA, EssB, and EssC are

thought to be the core of the T7 secretion machinery, with EssC being the central

membrane transporter (Burts *et al.*, 2005, Jager *et al.*, 2018, Zoltner *et al.*, 2016).

Interactions between secreted substrates and co-dependent secretion of substrates

have been demonstrated (Anderson *et al.*, 2013, Cao *et al.*, 2016, Kneuper *et al.*,

2014, Ohr *et al.*, 2017). A recent study showed that the functional assembly of the

T7SS machinery in *S. aureus* is supported by the flotillin homolog FloA, within

functional membrane microdomains (Mielich-Suss *et al.*, 2017).

The *S. aureus* T7SS is pivotal for bacterial virulence. Indeed, S. *aureus* mutants

lacking the entire T7SS (Kneuper *et al.*, 2014) or specific T7SS components (EsxA,

EssB, EssC, EsxC, EsxB, EsaB, EsaD or EsaE) were consistently shown to be less

virulent and/or persistent in various mouse infection models (Anderson *et al.*, 2011,

Anderson *et al.*, 2017, Burts *et al.*, 2008, Burts *et al.*, 2005, Ishii *et al.*, 2014, Lopez *et

al.*, 2017). EsxA is necessary to delay apoptosis of *S. aureus*-infected epithelial and

dendritic cells, while other substrates modulate cytokine production (Anderson *et al.*,

2017, Cruciani *et al.*, 2017, Korea *et al.*, 2014). Although the relevance of T7SS to *S.*

aureus is less understood, a role for the toxin-antitoxin pair EsaD (or EssD) and EsaG

(or EssI) was recently demonstrated in intraspecies competition (Cao *et al.*, 2016, Ohr

et al., 2017).

Interestingly, *S. aureus* T7SS expression is induced in response to host-specific FAs

(Ishii *et al.*, 2014, Kenny *et al.*, 2009, Lopez *et al.*, 2017), although the role of T7SS in

bacterial resistance to antimicrobial FAs remains unclear. In this study, we

demonstrate that *S. aureus* lacking the T7SS substrate EsxC has a defective cell

membrane. Intriguingly, EsxC and other T7SS mutants were more sensitive to an

unsaturated FA, linoleic acid (LA), compared to the wild-type (WT). Although there

were no differences in binding labelled LA, LA induced a more leaky membrane in the

T7SS mutants, and there was less incorporation of LA into membrane phospholipids.

Cellular proteomics revealed that in addition to membrane discrepancies, T7SS

mutants exhibited different redox and metabolic states, which likely result in a distinct

response to LA.

Results

The type VII substrate EsxC is present on the staphylococcal surface and affects

its composition

EsxC, a small 15-kDa protein secreted by the T7SS, is important for *S. aureus*

persistence in mice (Burts *et al.*, 2008). However, mechanisms underlying EsxC- or

T7SS-mediated bacterial survival are not known. In order to understand the role of

EsxC, we generated an isogenic *esxC* mutant as described previously (Bae &

Schneewind, 2006), and confirmed the absence of any secondary site mutations by

whole genome sequencing. Δ*esxC* had a similar growth rate to the WT USA300 JE2

strain (Fig. S1). EsxC is known to be a secreted effector of T7SS (Burts *et al.*, 2008),

although it has been consistently detected in cell membrane (CM) fractions of RN6390

and USA300 (Bobrovskyy *et al.*, 2018, Kneuper *et al.*, 2014). As reported previously,

we detected EsxC in the CM fractions of the WT but not in Δ*esxC* (Fig. S2). Moreover,

we also detected EsxC in the WT cell wall (CW) fraction (Fig. S2). In a deletion mutant

of the membrane-bound major ATPase EssC (a core T7SS component), EsxC was

detected in CM and CW preparations even in the absence of EssC (Fig. S2). While

we were unable to detect EsxC in the cell-free supernatants by immunoblotting, we

detected this protein by secretome analyses. Along with EsxC, other T7SS proteins,

EsxA and EsxD, were detected in the WT but not in Δ*esxC or* Δ*essC* supernatants

(Table S1).

Given the association of EsxC with the cell membrane, we stained the WT, Δ*esxC* and

Δ*essC* mutants with FM-143, a fluorescent cell membrane probe (Pulschen *et al.*,

2017). We observed a mild but statistically significant increase in FM-143 staining in

the mutants as compared to the WT (Fig. 1A). Increased staining of membranes by

FM-143 has been associated previously with membrane blebbing in bacteria (Wood

et al., 2019).

To study if the lack of T7SS proteins affected other surface proteins, a surface

proteome analysis of the WT and mutants was carried out using trypsin treatment to

digest surface proteins (Solis *et al.*, 2014). We found that most proteins (172/218) were

less abundant in Δ*esxC* compared to the WT (Fig. S3). Interestingly, majority of

relatively less abundant proteins (16/20) in the Δ*esxC* surface proteome were

predicted or proven to be cytosolic (Table S2). Metabolic (PfKA, FolD, FabI GatB, and

ImdH) or stress-related (Asp23 and putative universal stress proteins) proteins were

lowest in abundance (log_2 fold change < - 1.0). The T7SS core component EsaA, which

has a prominent extracellular loop (Dreisbach *et al.*, 2010), was also ~50% less

abundant in Δ*esxC*, implying that EsxC absence may affect T7SS assembly.

Identification of cytosolic proteins during surface proteome analysis could be attributed

to non-canonical secretion of cytosolic proteins that bind to bacterial surfaces (Hempel

et al., 2011, Pasztor *et al.*, 2010) or cell lysis during sample preparation (Dreisbach *et*

al., 2010, Solis *et al.*, 2014, Ventura *et al.*, 2010). To study if the T7SS mutants were

more resistant to cell lysis, we compared the Triton X-100 lysis of WT, Δ*esxC* and,

Δ*essC* in PBS in non-growing conditions. While Δ*esxC* was slightly more resistant to

Triton X-100, Δ*essC* was clearly defective in lysis (Fig. 1B), as compared to the WT.

We probed T7SS effects on the membrane further by assessing membrane fluidity of

the WT and Δ*esxC* mutant. We used pyrene decanoic acid, an eximer-forming lipid

(Lopez *et al.*, 2017), to measure the fluidity of WT and Δ*esxC* membranes. Compared

to the WT, the cytosolic membrane of the Δ*esxC* mutant showed a mild but statistically

significant increase in rigidity (Fig. 1C). Furthermore, we stained WT and with a

fluorescent membrane dye DiI12C, which has been used detect regions of increased

fluidity in bacteria (Saeloh *et al.*, 2018). As reported previously for *S. aureus*, we

observe few fluid regions in the WT, and an uneven staining of the membrane. In

contrast, a very weak staining of the membranes was observed for the Δ*esxC* mutant,

(Fig 1 D and E), which may indicate a decrease in membrane fluidity.

Overall, the data indicate that EsxC contributes to *S. aureus* cell surface structure and

membrane fluidity.

***S. aureus esxC* and *essC* mutants are more sensitive to antimicrobial fatty acids**

Multiple studies have reported the activation of the T7SS, when *S. aureus* is grown in

presence of host FAs (Ishii *et al.*, 2014, Kenny *et al.*, 2009, Lopez *et al.*, 2017). Given

our findings that the *esxC* mutant has membrane defects and that the *esxC* deletion

affects the surface abundance of IsdA, which is essential for *S. aureus* resistance to

antimicrobial fatty acids (Clarke *et al.*, 2007), we cultured WT USA300 and its isogenic

essC or *esxC* mutant in the presence of an unsaturated FA (C18:2), linoleic acid (LA),

at a concentration (80 µM) that still allows WT growth. Bacteria were also cultured in

parallel in presence of stearic acid (SA), a saturated C18:0 FA. The T7SS mutants

displayed impaired growth in presence of LA but not SA, as measured by optical

density (OD) (Fig. 2A) or colony forming units (CFU) (Fig. 2B). Importantly, Δ*esxC*

complemented with a plasmid containing the *esxC* gene reverted to the WT phenotype

(Fig. 2C). The increased susceptibility of T7SS mutants to antimicrobial fatty acids was

not restricted to linoleic acid; when cultured in the presence of arachidonic acid,

another unsaturated FA (C20:4), growth of Δ*esxC* and Δ*essC* were inhibited more as

compared to the WT (Fig. S4)

T7SS substrates contribute to *S. aureus* resistance to LA toxicity

Next, we investigated whether T7SS proteins other than *essC* and *esxC* contributed

to *S. aureus* growth in presence of LA. Mutants lacking two other substrates, Δ*esxA*

and Δ*esxB*, were grown in presence of FAs. Both mutants grew slower than the WT

USA300 (Fig. S5A). To ensure that the increased sensitivity observed for the T7SS

mutants was not strain specific, RN6390 Δ*essC* or Δ*esxC* and Newman Δ*esxA* or

Δ*esxB* mutants were tested. Similar to the USA300 mutants, the growth of all these

T7SS mutants was also impacted in the presence of LA (Fig. S5B and C). The growth

defect in Newman Δ*esxA* was abrogated upon complementation (Fig. 2D). Of note,

the Newman WT was readily inhibited by a lower concentration of LA (40 µM), which

is in agreement with the lower T7SS expression levels in this strain compared to

USA300 (Anderson *et al.*, 2013, Kneuper *et al.*, 2014). We conclude that a functional

T7SS plays a role in *S. aureus* resistance to LA toxicity.

T7SS is required for maintaining membrane integrity in presence of LA

To study the mechanisms involved in T7SS mediated protection to LA toxicity, further

studies were performed using strains lacking esxC, a representative T7SS effector, or

essC the main T7SS transporter. As our membrane and surface proteome analyses

with Δ*esxC* suggested that the bacterial envelope is altered upon T7SS defect, we

wanted to test if LA-mediated growth inhibition was due to an increased binding of LA

to T7SS mutants. To do this, we chemically engineered LA to produce an azide

functionalised LA (N^6-diazo-N^2-((9Z,12Z)-octadeca-9,12-dienoyl)lysine, N_3-LA) or

azide-LA (Fig. 3A). After incubating bacteria with azide-LA, click-chemistry with an

alkyne dye (Click-iT™ Alexa Fluor™ 488 sDIBO alkyne) was used to stain azide-LA

associated with bacteria. There were no obvious differences in the fluorescence from

Δ*essC* and Δ*esxC* compared to the WT (Fig. 3B), suggesting that T7SS components

are not involved in binding or sequestering LA. However, when bacteria treated with

azide-LA were stained with propidium iodide (PI), a good indicator of membrane

integrity, Δ*essC* and Δ*esxC* displayed a more intense PI staining compared to the WT

(Fig. 3C and D). Similarly, when WT and mutants were stained with PI and SYTO 9

after treatment with 80 μM unlabelled LA, we observed an increased PI staining (Fig.

4A and B) and therefore lower SYTO 9 / PI (Live/Dead) ratio for both mutants (Figure

4C). These data suggest that an intact T7SS helps *S. aureus* to maintain its membrane

integrity when faced with the detergent-like effects of unsaturated FAs.

LA-incorporation into membrane phospholipids is modulated by T7SS

As the T7SS mutants have more compromised cell membranes in presence of LA, we

next investigated if membrane lipids were altered in the T7SS mutants. Lipids from

WT USA300 and T7SS mutants were analysed by high-performance liquid

chromatography (HPLC)-mass spectrometry (MS) in negative ionisation mode. As

reported previously (Delekta *et al.*, 2018, Parsons *et al.*, 2011), phosphatidylglycerol

(PG) was the major phospholipid present in the membrane of WT grown in TSB (Fig.

S6A). Δ*essC* and Δ*esxC* grown with or without 10 μM LA [(a concentration that has

been previously shown to be sub-inhibitory for USA300 (Lopez *et al.*, 2017)] displayed

lipid profiles similar to that of WT (Fig. S6A and B). Notably, PG molecular species

were significantly altered upon growth in LA-supplemented TSB for WT (Fig. 5A),

Δ*essC* (Fig. S6C) and Δ*esxC* (Fig. 5B). Three new LA-specific PG species with mass

to charge ratios (m/z) 731 (C33:2), 759 (C35:2), and 787 (C37:2) appeared to contain

LA (C18:2) or its elongated C20:2 or C22:2 versions, as revealed by their

fragmentations (Fig. S7A, B and C). PG species containing exogenous, unsaturated

FAs were also present in Δ*essC* and Δ*esxC*. However, LA (C18:2)-containing PG

species (C33:2) were less abundant in the *esxC* mutant compared to WT (Fig. 5C). A

similar trend, although statistically non-significant ($P > 0.05$), was observed for C20:2-

and C22:2-containing PG species (Fig. S7D and E), and when all the unsaturated

exogenous PG species were combined (Fig. 5D). The data suggest that a T7SS defect

may compromise the incorporation and elongation of LA in *S. aureus* membranes.

T7SS mutations affect the total cellular content and *S. aureus* responses to LA

In order to gain further insight into T7SS-mediated modulation of proteins involved in

FA incorporation and membrane homeostasis in presence of LA, we used an unbiased

proteomic approach to study protein profiles of WT USA300, Δ*essC*, and Δ*esxC* grown

exponentially with or without 10 μM LA. Of note, WT and both these T7SS mutants

grew similarly in presence of up to 40 μM LA (Fig. S8).

WT vs T7SS mutants in absence of LA treatment: Interestingly, ΔessC or ΔesxC

cultured in TSB readily displayed proteins with changed abundance when compared

to the WT, with 37 and 24 proteins significantly ($P < 0.05$) altered in ΔessC and ΔesxC,

respectively. Similarly, 14 proteins were differentially abundant in both ΔessC and

ΔesxC (Fig. 6A and B). These included proteins associated with signal transduction

(LytR and ArlR), the CW (acetyltransferase GNAT, FnbB and MazF), DNA repair (MutL

and RadA), nucleotide binding (ATP-grasp domain protein and YqeH), hydrolysis

(amidohydrolase), cell stress response [universal stress protein (Usp) family], or were

uncharacterised (A0A0H2XGJ8, YbbR and lipoprotein) (Fig. 6B and Table 1). Of the

33 proteins changed only in ΔessC (23 proteins) or ΔesxC (10 proteins), nearly 40%

(13 proteins) were associated with oxidation-reduction and other metabolic processes.

Ten membrane proteins were more abundant in ΔessC (Table 1), which included SrrB,

a membrane protein that is activated by impaired respiration (Mashruwala *et al.*, 2017),

and whose gene expression increased 6 times upon growth in presence of LA (Lopez

et al., 2017). SrrB was also detected at higher levels in the *esxC* mutant although the

increase was non-significant ($P = 0.07$).

Table 1. Proteins with changed abundance in Δ*essC* and Δ*esxC* mutants relative to the WT USA300 JE2

Functions	Uniprot ID	Δ*essC*/WT		Δ*esxC*/WT		Description
		Log$_2$ FC	Adjusted P value	Log$_2$ FC	Adjusted P value	
Signal transduction systems	Q2FK09	-3.0	4.90E-13	-3.1	4.90E-13	Sensory transduction protein LytR
	Q2FH23	-2.9	0.002527	-2.1	0.026184	Response regulator ArlR
Membrane proteins	A0A0H2XF42	3.0	1.75E-10	0	1	Cytochrome D ubiquinol oxidase, subunit I
	A0A0H2XDZ5	1.7	1.68E-05	0	1	Uncharacterized membrane protein
	A0A0H2XFJ8	2.0	0.001077	0.5	0.883953	Uracil permease
	A0A0H2XGW7	2.7	0.005757	2.5	0.010784	Putative lipoprotein
	A0A0H2XIA9	1.0	0.006115	0	1	Protein translocase subunit SecY
	Q2FIN2	1.6	0.029451	0.5	0.970614	Prolipoprotein diacylglyceryl transferase LGT
	A0A0H2XKD9	2.0	0.038093	1.8	0.070347	Staphylococcal respiratory response protein SrrB
	A0A0H2XFE1	3.4	0.039814	1.1	0.970614	Peptidase
	A0A0H2XJV8	2.0	0.04444	1.5	0.231285	Cyclic-di-AMP phosphodiesterase
	A0A0H2XGF4	3.0	0.044681	0.9	0.986458	Sodium:dicarboxylate symporter family protein
	A0A0H2XHV2	2.5	0.049168	2.4	0.070182	Glycine betaine transporter OpuD
Stress response	A0A0H2XKH6	2.2	5.80E-05	2.5	1.07E-05	Universal stress protein family
	A0A0H2XIZ0	0.0	1	-3.4	1. 06E-10	OsmC/Ohr family protein
DNA repair	Q2FHE2	-2.3	1.79E-08	-2.3	1.19E-08	DNA mismatch repair protein MutL
	A0A0H2XI63	-2.0	0.004036	-2.0	0.004036	DNA repair protein RadA
	A0A0H2XHT1	-0.3	0.708343	-1.9	0.008379	Formamidopyrimidine-DNA glycosylase MutM
Oxidation-reduction process	A0A0H2XJ90	1.1	0.038093	1.0	0.088334	D-isomer specific 2-hydroxyacid dehydrogenase family protein
	A0A0H2XHE0	2.4	0.039099	-0.1	1	Thiol-disulphide oxidoreductase, DCC family protein
	A0A0H2XGR9	0.0	1	1.4	8.89E-08	Oxidoreductase, Gfo/Idh/MocA family
	A0A0H2XK08	1.0	0.406975	2.9	0.00791	Oxidoreductase, short chain dehydrogenase/reductase family
	A0A0H2XFZ3	-0.8	0.404225	2.1	0.016405	Nitroreductase family protein
Hydrolases	A0A0H2XE49	2.9	1.07E-06	2.9	5.99E-07	Amidohydrolase
	Q2FES9	-2.7	0.003385	-0.3	1	Uncharacterized hydrolase
	A0A0H2XFF2	-0.8	0.016697	-0.5	0.157809	Peptidase, U32 family
	A0A0H2XJH8	0.0	1	2.8	1.06E-10	Peptidase M20 domain-containing protein 2
	A0A0H2XJ54	0.0	1	2.0	0.000949	Hydrolase (HAD superfamily)
	Q2FEG2	-0.2	0.854748	-2.9	0.004615	Formimidoylglutamase
Metabolism	A0A0H2XGU2	-1.6	4.99E-06	-0.1	1	Pseudouridine synthase
	A0A0H2XK15	2.8	0.004424	0.9	0.777138	1-phosphatidylinositol phosphodiesterase
	Q2FEK2	-1.6	0.038093	-0.3	1	Urease accessory protein UreE
	Q2FI05	1.1	0.038093	0.0	1	Bifunctional purine biosynthesis protein PurH
	Q2FIL2	2.9	0.038093	0.8	0.970614	SsrA-binding protein
	A0A0H2XII6	1.8	0.038093	1.5	0.088334	Orn/Lys/Arg decarboxylase
	A0A0H2XJR8	-0.9	0.04444	-0.5	0.61246	RNA methyltransferase, RsmD family
	A0A0H2XKG7	0.0	1	1.4	8.18E-08	Aspartokinase
Cell wall composition	A0A0H2XJQ4	-4.0	4.92E-11	-4.0	3.28E-11	Acetyltransferase, GNAT family
	A0A0H2XKG3	-2.3	0.001238	-1.7	0.016405	Fibronectin binding protein B
	A0A0H2XJC8	-2.3	0.009779	-3.0	0.000819	Phi77 ORF017-like protein (Toxin MazF)
	Q2FE03	0.0	1	2.6	1.18E-12	Fibronectin-binding protein A
Nucleotide binding	A0A0H2XHY5	-2.3	8.58E-06	-2.3	6.01E-06	ATP-grasp domain protein
	A0A0H2XFA5	3.3	0.014259	3.5	0.008379	Putative GTP-binding YqeH protein
Uncharacterised proteins	A0A0H2XGJ8	2.2	0.000372	2.4	9.41E-05	Uncharacterized protein
	A0A0H2XE09	2.1	0.008231	1.9	0.016405	Ybbr-like uncharacterized protein
	Q2FFI4	3.4	0.020136	0.9	0.970614	UPF0316 membrane protein
	A0A0H2XG24	1.1	0.022345	0.8	0.088334	Uncharacterized protein

WT vs T7SS mutants in presence of LA: We then compared the proteomic profiles of

LA-treated strains (WT, Δ*essC* or Δ*esxC*) with their untreated counterparts. Clearly,

the principal component analysis revealed that the differences due to the genetic

makeup (WT or T7SS mutants) were less prominent than the dramatic changes

induced by LA (Fig. S9). These changes are exemplified for the WT; 163/1132 proteins

identified had an altered relative abundance upon growth with LA (Fig. 6C). 167 and

171 proteins were changed ($P < 0.05$) in Δ*essC* and Δ*esxC*, respectively, in response

to LA, of which ~ 40% (68 proteins) were common to these mutants and their WT (Fig.

6D). At least 30% of the significantly changed proteins ($P < 0.05$) were unique to WT

(53 proteins), Δ*essC* (50 proteins), or Δ*esxC* (64 proteins) (Fig. 6D), suggesting that

each strain responds differently to LA. However, almost all proteins (13/14 proteins)

that were similarly deregulated in Δ*essC* and Δ*esxC* grown without LA (Fig. 6B) were

modulated in presence of LA (highlighted in bold in Dataset S1). Proteins that were

less abundant in both mutants were, upon LA treatment, either increased to WT levels

(MutL, acetyltransferase GNAT, Toxin MazF, and ATP-grasp domain protein), or were

unchanged in the mutants and decreased in the LA-treated WT (LytR and FnbB)

(Dataset S1). Likewise, proteins with increased amounts in Δ*essC* or Δ*esxC* were: (i)

downregulated to WT levels in response to LA (putative lipoprotein A0A0H2XGW7),

(ii) unaltered in both mutants and upregulated in WT (Usp, amidohydrolase, and

YbbR), (iii) or further increased in the *essC* mutant and strongly upregulated in WT

(A0A0H2XGJ8) (Dataset S1). In sum, except for ArlR and RadA that were conversely

regulated in all strains after LA treatment, proteins similarly deregulated in Δ*esxC* and

Δ*essC* were returned to similar levels in response to LA. A similar trend was observed

for 15/23 and all 10 proteins exclusively more or less abundant in Δ*essC* and Δ*esxC*,

respectively.

Altered molecular functions in presence of LA: We then used QuickGO (a web-based tool for Gene Ontology searching) (Binns *et al.*, 2009) to retrieve GO terms associated with the ten most significantly upregulated proteins in LA-treated WT (Dataset S1). Strikingly, 9/10 proteins had a hydrolase or an oxidoreductase activity. A comprehensive, statistical analysis showed a clear enrichment of 8 specific molecular functions ($P < 0.05$) in at least one strain (WT or T7SS mutants) (Fig. 7A). Oxidoreductase and hydrolase activities were enhanced in LA-treated WT, while Δ*essC* and Δ*esxC* were less able to upregulate proteins with these molecular functions. Flavin adenine dinucleotide (FAD)-binding, which plays a role in oxidation-reduction and FA metabolic processes, was similarly more enriched in the LA-treated WT. In contrast, transferase activity, which is linked to CW synthesis, was induced more in T7SS mutants compared to the WT. Molecular functions that are decreased upon LA challenge were also determined (Fig. 7B). In agreement with reduced intracellular ATP levels following membrane damage by antimicrobial FAs (Cartron *et al.*, 2014), genes with the ATP-binding function (mainly ATP-binding ABC transporters) were strongly inhibited in the WT. ATP-dependent lyases were also repressed in the WT. On the contrary, T7SS mutants were less able to modulate ATP-binding proteins. Instead, a strong inhibition of ribosomal constituents and other translation-related components was seen (Fig. 7B).

To test the oxidoreductive states of the WT and the mutants, we stained bacteria with dichlorofluorescin (DCF), which detects reactive oxygen species (George *et al.*, 2019). Reflecting the changes seen in the proteomics data, when treated with 10 µM LA there is an increase in the ROS generated in the T7SS mutants compared to the WT (Fig 7C). However, in bacteria grown without LA, the mutants have slightly less or no change in the ROS generated compared to WT (Figure 7D). Taken together, our

proteomic analyses reveal that the lack of T7SS induces altered membrane and

metabolic states indicative of oxidative stress responses. While the WT shows a

multifaceted response to mitigate LA-induced damage on the bacterial membrane,

such responses are clearly altered in the absence of the T7SS.

Discussion

Host fatty acids (FAs) play a crucial role in the host defence to *S. aureus* infections. *S.*

aureus is particularly sensitive to unsaturated FAs, which are abundant in the human

skin (Clarke *et al.*, 2007, Parsons *et al.*, 2012, Arsic *et al.*, 2012, Kelsey *et al.*, 2006,

Kenny *et al.*, 2009). We report here that the T7SS, an important component of *S.*

aureus virulence arsenal, is critical in modulating the response to antimicrobial host

FAs by maintaining the bacterial cell membrane integrity. Specifically, we demonstrate

that a T7SS substrate, EsxC, impacts *S. aureus* membrane properties. A functional

T7SS enables bacteria to mitigate LA-induced toxicity and grow better than mutants

with a compromised T7SS. In the absence of T7SS components, LA is less

incorporated into membrane phospholipids and enhances cell membrane damage,

while bacteria are unable to activate adaptive mechanisms involved in LA resistance,

as indicated by cellular proteomics.

Although several studies have shown multiple interactions between the staphylococcal

T7SS components, the precise molecular architecture of this system remains unclear.

EsxC (previously EsaC) was first described as a secreted protein (Burts *et al.*, 2008).

However, in keeping with prior studies, we found that EsxC can also localize within

staphylococcal membranes (Bobrovskyy *et al.*, 2018, Kneuper *et al.*, 2014). Based on

available data, EsxC is likely to be associated to EsxA, EsaD, or EsaE on the

membrane (Anderson *et al.*, 2013, Anderson *et al.*, 2017, Cao *et al.*, 2016).

Additionally, the reduced EsaA protein levels that we found on the surface of *S. aureus*

USA300 Δ*esxC* combined with prior observations of diminished EsxC protein levels in

RN6390 Δ*esaA* membranes (Kneuper *et al.*, 2014) suggest that EsxC may interact

with EsaA, a key component of the T7SS core (Aly *et al.*, 2017), in *S. aureus*

membranes.

T7SS integral membrane proteins interact with the flotillin homolog FloA within

functional membrane microdomains (FMMs) (Mielich-Suss *et al.*, 2017). FMMs were

recently shown to contain staphyloxanthin derivatives (Garcia-Fernandez *et al.*, 2017),

which are known to increase membrane rigidity (Chamberlain *et al.*, 1991, Sen *et al.*,

2016, Tiwari *et al.*, 2018). Hence, well-structured FMMs may play a role in *S. aureus*

membrane fluidity as reported for *Bacillus subtilis* (Bach & Bramkamp, 2013).

Interestingly, mutations or treatments affecting *S. aureus* membrane fluidity also

modulate T7SS transcription (Lopez *et al.*, 2017, Ishii *et al.*, 2014, Parsons *et al.*,

2014). Hence, there appears to be a link between membrane fluidity and T7SS in *S.*

aureus. Also, the state of the membrane was suggested to trigger the production of

T7SS (Lopez *et al.*, 2017). Remarkably, deletion of *esxC* led to a mild increase in

bacterial membrane rigidity and membrane defects (Fig. 1), suggesting membrane

modulation by the T7SS. Possibly, interactions between FloA and T7SS are perturbed

upon *esxC* deletion and affect *S. aureus* FMMs. We surmise that a functional T7SS

helps *S. aureus* to maintain its membrane architecture.

The current cellular proteomics data reveal that the abundance of more proteins is

altered in Δ*essC* (37) than *esxC* (24) in comparison to *S. aureus* WT, which is in

keeping with the greater importance of EssC as the conserved driving force of the

T7SS (Warne *et al.*, 2016). Importantly, almost 60% of proteins deregulated in Δ*esxC*

are similarly affected in Δ*essC*, strongly suggesting that any modification of the T7SS

core leads to a similar staphylococcal response. Surprisingly, proteins with altered

abundance in USA300 Δ*essC* were distinct to the ferric uptake regulator (Fur)-

controlled genes differentially expressed in RN6390 Δ*essC* (Casabona *et al.*, 2017b).

This discrepancy might be due to strain differences, including *rsbU* defect in RN6390

that impairs SigB activity (Cassat *et al.*, 2006, Giachino *et al.*, 2001). Nevertheless,

given the known role of Fur in oxidative stress resistance (Horsburgh *et al.*, 2001,

Johnson *et al.*, 2011), both mutants may display an altered oxidative status following

essC deletion. *S aureus* RN6390 also differentially expresses redox-sensitive genes

in absence of EsaB (Casabona *et al.*, 2017a). Also, since the T7SS substrate EsxA is

upregulated in response to hydrogen pyroxide (Casabona *et al.*, 2017b), one could

speculate that lack of T7SS stimulates an oxidative stress response. A further

indication of altered physiological states of Δ*essC* and Δ*esxC* was the decreased

abundance of the two-component regulatory system proteins, LytSR, ArlSR and

SrrAB, which was consistent with down-regulation of *lytR* transcription observed

previously in the absence of *arlR* (Liang *et al.*, 2005). Importantly, the *S. aureus*

response to antimicrobial FAs includes downregulation of *lytRS* (Kenny *et al.*, 2009,

Neumann *et al.*, 2015), and upregulation of *srrB* (Lopez *et al.*, 2017). Given that LytSR

is involved in bacterial surface and membrane potential modulation (Patton *et al.*,

2006, Groicher *et al.*, 2000), T7SS defects are likely to result in an altered cell

envelope.

It is striking that the staphylococcal T7SS is strongly upregulated in presence of sub-

inhibitory concentrations of LA (Kenny *et al.*, 2009, Lopez *et al.*, 2017). FAs with more

cis double bonds, which are more toxic toward *S. aureus* (Parsons *et al.*, 2012), are

also more potent T7SS activators (Lopez *et al.*, 2017). Our current study interestingly

suggests a protective role of T7SS against LA toxicity. Previously described *S. aureus*

antimicrobial FA (AFA) resistance mechanisms, including IsdA or wall teichoic acid-

mediated modulation of cellular hydrophobicity (Clarke *et al.*, 2007, Kohler *et al.*, 2009,

Parsons *et al.*, 2012, Moran *et al.*, 2017), and AFA detoxification with the efflux pumps

Tet38 and FarE (Alnaseri *et al.*, 2015, Truong-Bolduc *et al.*, 2014), do not appear to

explain the increased susceptibility of T7SS mutants to LA, as indicated by cellular

proteomics. In line with a role for T7SS in the oxidative stress response, T7SS mutants

were less able to prime their redox-active proteins in response to LA-induced oxidative

stress. Instead, to cope with LA, they appear to rely on strong inhibition of the protein

synthesis machinery, which is reminiscent of the stringent response (Geiger *et al.*,

2012).

The Fak pathway which is required for incorporation of exogenous FA into membrane

phospholipids via a two-component fatty acid kinase (Fak) (Nguyen *et al.*, 2016,

Parsons *et al.*, 2012, Parsons *et al.*, 2014), was reported to be important for T7SS

activation by unsaturated FA (Lopez *et al.*, 2017). FakB1 and FakB2, bind to FAs, and

FakB-bound FAs are phosphorylated by FakA prior to their incorporation (Parsons *et

al.*, 2014). In presence of inhibitory concentrations of unsaturated FA like LA, while FA

is incorporated into the membrane lipids, free LA causes pore formation that leads to

bacterial lysis (Greenway & Dyke, 1979). Our lipidomic analyses revealed that in the

absence of T7SS, bacteria were less able to incorporate LA into their phospholipids

(Figure 5), and displayed an increased membrane permeability in presence of LA.

However, it seems counterintuitive that LA incorporation was impacted more in Δ*esxC*

than in Δ*essC* given the central role of EssC in T7 secretion (Burts *et al.*, 2005, Jager

et al., 2018, Zoltner *et al.*, 2016). It is possible that EsxC that accumulates in the

membrane (Fig. 1A) in the absence of protein secretion by the main transporter, EssC,

mediates partial LA incorporation in the *essC* mutant; secretion per se may not be

required for LA incorporation. It is also worth noting that transcript levels of *esxC*, and

not *essC*, were strongly upregulated in a *S. aureus fakA* mutant (Parsons *et al.*, 2014).

As protein levels of Fak proteins in the T7SS mutants stay unaltered in presence or

absence of LA, suggesting no T7SS-mediated regulatory control of the Fak pathway,

we speculate that EsxC and other interdependent T7SS substrates play an important

role in facilitating Fak function in *S. aureus* membranes, either by mediating

recruitment or targeting of Fak proteins to the membrane. Our findings warrant further

investigations   into   molecular   mechanisms   underlying   T7SS-mediated   FA

incorporation within staphylococcal membranes.

The increased susceptibility of T7SS mutants to LA might explain why they are less

virulent in environments rich in LA and other AFAs like the mouse lungs (Δ*essC*) (Ishii

et al., 2014), abscesses (Δ*esxC* and Δ*esaB*), liver and skin (Δ*essB*) (Wang *et al.*, 2016,

Lopez *et al.*, 2017). Previous research showing T7SS induction by host-derived FAs

further supports the importance of T7SS in such environments (Lopez *et al.*, 2017,

Ishii *et al.*, 2014). Taken together, we conclude that T7SS plays a key role in

modulating the *S. aureus* cell membrane in response to toxic host FAs. Although, at

present, it is unclear how T7SS contributes to staphylococcal membrane architecture,

T7SS interaction with the flotillin homolog FloA within functional membrane

microdomains (Mielich-Suss *et al.*, 2017) corroborates the idea that T7SS proteins

interact with many other proteins to modulate *S. aureus* membranes. Indeed, our data

also suggest that blocking T7SS activity would make *S. aureus* more vulnerable to

AFAs, a key anti-staphylococcal host defence, thus making T7SS a very attractive

drug target.

Experimental procedures:

Bacterial strains and growth conditions. *S. aureus* strains used are listed in Table

S3, and were grown aerobically in tryptic soy broth (TSB) overnight (O/N) at 37ºC for

each experiment unless stated otherwise. For complemented *S. aureus* strains, TSB

was supplemented with 10 µg/mL chloramphenicol.

Construction of bacterial mutants. The primers used are listed in Table S4. In-frame

deletion of *essC* or *esxC* was performed as described previously (Bae & Schneewind,

2006). Briefly, 1-kb DNA fragments up and downstream of the targeted gene sequence

were PCR-amplified from USA300 LAC JE2 chromosomal DNA, and both PCR

products fused via SOEing (splicing by overlap extension)-PCR. The 2-kb DNA

fragment obtained was cloned into pKORI, and used for in-frame deletion. Putative

mutants were screened by PCR-amplification of a fragment including the gene of

interest, whose deletion was confirmed by Sanger sequencing. Further, to confirm that

successful mutants did not have any additional mutations, Illumina whole genome

sequencing was performed on libraries prepared with the Nextera XT kit and an

Illumina MiSeq® instrument following manufacturers' recommendations. For

complementation, full-length *esxC* gene was cloned onto pOS1CK described

previously (Korea *et al.*, 2014).

Membrane fluidity assay. O/N bacterial cultures were diluted to an OD_{600} of 0.15 in

TSB, and were grown to an OD_{600} of 1 (OD1). Bacteria were washed with PBS prior

to treatment for 30 min at 37ºC with 37.5 µg/mL lysostaphin in PBS containing 20%

sucrose. The spheroblasts were then centrifuged at 8000 × *g* for 10 min, and the pellet

resuspended in the labelling solution (PBS, 20% sucrose, 0.01% F-127, 5 µM pyrene

decanoic acid). The incubation in the dark was done for 1h at 25ºC under gentle

rotation. PBS supplemented with 20% sucrose was used to wash the stained

spheroblasts that were afterwards transferred to 96-well plates for fluorescence

measurements as previously described (Lopez *et al.*, 2017).

Triton X-100 lysis assays. Whole cell autolysis assays were performed as described

elsewhere with a few modifications (Mashruwala *et al.*, 2017). Specifically, OD1-grown

S. aureus USA300 JE2 WT, *essC* and *esxC* mutants were extensively washed with

PBS followed by ice-cold water, and resuspended in PBS with 0.1% Triton X-100 to

an OD_{600} of 0.7. Subsequently, the samples were incubated with shaking at 37ºC for

2h, after which bacteria were diluted with PBS and plated for CFU determination.

FM1-43 staining. *S. aureus* WT and mutant strains grown to OD_{600} of 1.0 were

centrifuged, and pellets were resuspended in PBS supplemented with 1 µg/mL FM1-

43 (Invitrogen). After incubation for 15 min in the dark at 37°C with shaking, cells were

washed once with PBS. Fluorescence was quantified using a FLUOstar OMEGA plate

reader (BMG Labtech, UK) at excitation and emission wavelengths of 482 nm and 620

nm, respectively.

DiIC12 staining. O/N bacterial cultures were diluted to an OD_{600} of 0.15 in TSB with

1 µg/mL 1,1′-didodecyl-3,3,3′,3′-tetramethylindocarbocyanine perchlorate (DiIC12)

(Invitrogen). Cultures were grown to an OD_{600} of 1.0, centrifuged and washed twice

with fresh TSB. Samples were spotted on to agarose pads and imaged using a Leica

DMi8 widefield microscope (Leica Microsystems, UK). Acquired images were

analysed with the ImageJ processing package, Fiji.

Growth curves. O/N bacterial cultures were diluted to an OD_{600} of 0.05 in plain TSB

or TSB supplemented with fatty acids. Bacteria were then grown in a 96-well plate with

shaking, and the OD_{600} was measured every 15 mins with a FLUOstar OMEGA plate

reader (BMG Labtech, UK).

Synthesis of azide functionalized linoleic acid. A 2-step synthesis was used to

obtain N^6-diazo-N^2-((9Z,12Z)-octadeca-9,12-dienoyl)lysine, N_3-LA (azide-LA). LA was

first functionalized with *N*-hydroxysuccinimide (NHS) in anhydrous dimethyl

formamide (DMF) in presence of *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide

hydrochloride. The solvent was then removed and replaced by dicholoromethane

(DCM), following which the reaction mixture was washed with water and dried over

magnesium sulphate. The product, 2,5-dioxopyrrolidin-1-yl (9Z,12Z)-octadeca-9,12-

dienoate (NHS-LA), was analysed using ^{1}H nuclear magnetic resonance (NMR)

spectroscopy (Fig. S10A) and mass spectrometry (MS). MS: [M+Na]$^+$ 400.5

(calculated), 400.5 (found).

NHS-LA was left O/N at room temperature (RT) to react with L-azidolysine

hydrochloride in anhydrous DMF, and produce azide-LA.

DMF was then removed, the reaction mixture precipitated in water, and dried under

vacuum to obtain a clear oil. The composition of the oil was confirmed as being a

mixture of azide-LA and unmodified LA (20% and 80%, respectively) based on ^{1}H

NMR (Fig. S10B) and MS data. MS: [LA-H]$^-$ 279.5 (calculated), 279.2 (found), [M-H]$^-$

433.3 (calculated), 433.6 (found).

Binding assays with azide-LA and click chemistry. *S. aureus* USA300 JE2 WT, Δ*essC*, and Δ*esxC*, grown to OD1, were treated with 10 μM azide-LA for 15 min at 37°C with shaking. The samples were then centrifuged, and the bacterial pellets resuspended in PBS supplemented with 4 μg/mL Click-iT™ Alexa Fluor™ 488 sDIBO alkyne (Life Technologies LTD, UK). After incubation at 25°C for 1h with shaking, bacteria were washed with PBS, and binding to azide-LA was quantified by measuring fluorescence using a FLUOstar OMEGA plate reader (BMG Labtech, UK). The samples imaged with a microscope were additionally stained with 3 μM propidium iodide, following click chemistry. Bacteria stained with Click-iT™ Alexa Fluor™ 488 sDIBO alkyne and 3 μM propidium iodide were immobilized on agarose-covered glass slides, and viewed with a Leica DMi8 widefield microscope (Leica Microsystems LTD, UK). Images were analysed with the ImageJ processing package Fiji (Schindelin *et al.*, 2012).

Live/Dead Staining. Bacteria grown to OD_{600} of 1.0, were treated with 80 μM linoleic acid for 15 min at 37°C with shaking. The samples were then centrifuged, and the bacterial pellets resuspended in PBS and supplemented with a 1:1 ratio of 2X LIVE/DEAD solution (6 μM SYTO-9 stain and 30 μM propidium iodide) from LIVE/DEAD® BacLight kit (Invitrogen). After incubation in the dark for 15 min, bacteria were washed with PBS, spotted on to agarose pads and imaged using a Leica DMi8 widefield microscope (Leica Microsystems, UK). Acquired images were analysed with the ImageJ processing package, Fiji.

Lipid extraction and analyses. Lipids were extracted from bacterial cultures as described elsewhere (Smith *et al.*, 2019). Briefly, bacteria were grown to OD1 in TSB or TSB supplemented with 10 μM LA, centrifuged in a 2 mL glass Chromacol vial (Thermo Scientific), and resuspended in 0.5 mL MS grade methanol (Sigma-Aldrich).

MS grade chloroform was then used to extract lipids. The extracted lipids were dried under nitrogen gas with a Techne sample concentrator (Staffordshire, UK), and the lipid pellets resuspended in 1 mL acetonitrile. The samples were then analysed by LC-MS with a Dionex 3400RS HPLC coupled to an amaZon SL quadrupole ion trap mass spectrometer (Bruker Scientific) via an electrospray ionisation interface. Both positive and negative ionisation modes were used for sample analyses. The Bruker Compass software package was utilized for data analyses, using DataAnalysis for peak identification and characterization of lipid class, and QuantAnalysis for quantification of the relative abundance of distinct PG species to total PG species.

Cell shaving for surface proteome analysis. *S. aureus* USA300 JE2 WT grown to OD1 and Δ*esxC* were washed three times before being treated with Proteomics grade trypsin from porcine pancreas (Sigma-Aldrich, UK) for 15 min as described (Solis *et al.*, 2014). The samples were then centrifuged at 1000 × *g* for 15 min, and the bacterial pellets discarded while supernatants were filtered through a 0.2 µM filter. The freshly prepared peptides were frozen (-20ºC) until 2 additional, independent biological replicates per strain were prepared.

Cellular proteomics. *S. aureus* strains were grown O/N at 37ºC on tryptic soy agar plates. The next day, single colonies were used to inoculate 10 mL plain TSB or TSB with 10 µM LA. Cultures were grown at 37ºC with 180-rpm shaking until an OD_{600} of 3.2 ± 0.2 was reached. The bacteria were then centrifuged, washed with PBS, and resuspended in lysis buffer (PBS, 250 mM sucrose, 1 mM EDTA, and 50 µg/mL lysostaphin) supplemented with cOmplete™, mini, EDTA-free protease inhibitor cocktail (Sigma-Aldrich, UK). After 15 min incubation at 37ºC, cells were lysed mechanically with silica spheres (Lysing Matrix B, Fischer Scientific, UK) in a fast-prep shaker as described previously (Mielich-Suss *et al.*, 2017). Samples were then

centrifuged, and the supernatants transferred to fresh tubes, where proteins were

reduced and alkylated for 20 min at 70ºC with 10 mM TCEP (tris(2-

carboxyethyl)phosphine) and 40 mM CAA (2-chloroacetamide), respectively. Next, the

solvent was exchanged first to 8M urea buffer then to 50 mM ammonium bicarbonate

(ABC). Proteins were digested O/N at 37ºC with mass spectrometry grade lysyl

endopeptidase LysC and sequencing grade modified trypsin (Promega LTD, UK).

Preparation of culture supernatant for proteomics analysis. *S. aureus* strains

were grown to an OD_{600} of 3 in TSB. After centrifugation of cultures, supernatants were

sterile-filtered and incubated at 4°C overnight with 10% trichloroacetic acid and 50 μM

sodium deoxycholate in presence of one cOmplete™, Mini, EDTA-free Protease

Inhibitor Cocktail tablet (Sigma-Aldrich, UK). The precipitated proteins (10 000 × *g* at

4ºC for 15 min) were gently washed with acetone and dry at RT for 10 min. After

denaturation, proteins were run on a gel until all the proteins had moved from the

stacking gel to the resolving gel. The gel was then stained with Instant*Blue*™ (Sigma-

Aldrich, UK) for 3 h, after which the protein bands were excised and diced. Proteins

were in-gel digested with trypsin as recently described(Goodman *et al.*, 2018). Briefly,

proteins were reduced and alkylated for 5 min at 70ºC with 10 mM TCEP and 40 mM

CAA, respectively. Tryptic digestion was carried out at O/N at 37°C in 50 mM ABC.

Label-free protein quantification. Peptides prepared for proteome analyses were

desalted and concentrated with a C18 cartridge in 40 μL MS buffer (2% acetonitrile

plus 0.1% trifluoroacetic acid). For each sample, 20 μL were analysed by nanoLC-

ESI-MS/MS using the Ultimate 3000/Orbitrap Fusion instrumentation (Thermo

Scientific), and a 90-minute LC separation on a 50 cm column. The data were used to

interrogate the Uniprot *Staphylococcus aureus* USA300 database UP000001939, and

the common contaminant database from MaxQuant (Cox *et al.*, 2014). MaxQuant

software was used for protein identification and quantification using default settings.

Intensities were $\log_2$-tansformed with the Perseus software, and proteins with one or

no valid value for every sample in triplicate were filtered. For surfome data, the

removeBatchEffect function of the limma R package (Ritchie *et al.*, 2015) was used to

remove differences accounting for variation in shaving efficiency done on three

different days for all the biological replicates. Missing values in cellular proteomics

data were imputed on R. Specifically, for each sample, the imputed value was either

the lowest intensity across all samples if at least two biological replicates had missing

values or the average of two valid values if only one was missing.

ROS measurement. Chemical hydrolysis of 2,7-dichlorofluorescein diacetate (Sigma-

Aldrich) was performed to acquire a final dichlorofluorescein (DCF) reagent yield of 50

µM. Briefly, 0.5 ml of 5 mM DCF-DA (dissolved in 100% ethanol), was reacted with 2

ml of 0.1 N NaOH at RT for 30 min. The reaction was stopped using 7.5 mL 10X PBS

pH 7.4 (without calcium and magnesium; Gibco). Bacteria grown to OD_{600} of 1.0 were

treated with 10µM linoleic acid or left untreated for 15 min at 37°C shaking and 1 mL

of this culture was centrifuged. Cells were resuspended in 100 µL of DCF reagent and

incubated for 40 min in the dark at RT. Fluorescence was measured using a FLUOstar

OMEGA plate reader (BMG Labtech, UK) at an excitation wavelength of 482 nm and

emission wavelength of 520 nm.

Data availability statement. The mass spectrometry proteomics data have been

deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol *et al.*,

2019) partner repository with the dataset identifier PXD013081 and

10.6019/PXD013081. Cellular proteomic and surface proteome samples are labelled

MS18-193 and MS17-185, respectively.

Statistical analyses. Except for the proteomics results, the statistical tests were

performed with GraphPad Prism 8.0 as indicated in the Figure legends, with P values

< 0.05 considered significant. A paired two-tailed Student's t-test or a paired Mann-

Whitney U test was used for pairwise comparisons. An ordinary one-way analysis of

variance (ANOVA) with Dunnett's multiple comparisons test or a Kruskal-Wallis test

with Dunn's multiple comparisons test was applied to data form three or more groups.

The fold changes and P values of the proteomics data were calculated with the R

package limma (Ritchie *et al.*, 2015), with USA300 JE2 WT or bacteria grown without

LA as references. These fold changes and P values were used by the R package

piano (Varemo *et al.*, 2013) to compute the enrichment of gene ontology (GO) terms.

Acknowledgments

We thank Professor Tracy Palmer (Newcastle University) and Professor Olaf

Scheewind (University of Chicago) for providing us *S. aureus* strains and reagents.

We thank GSK, Siena, Italy for providing the *esxA*, *esxB* mutant strains used in this

study. We acknowledge the contribution of the Proteomics Research Technology

Platform, University of Warwick. The authors have no conflict of interest to declare.

References

References

Alnaseri, H., Arsic, B., Schneider, J.E., Kaiser, J.C., Scinocca, Z.C., Heinrichs, D.E., and

McGavin, M.J. (2015) Inducible Expression of a Resistance-Nodulation-Division-Type

Efflux Pump in Staphylococcus aureus Provides Resistance to Linoleic and

Arachidonic Acids. *J Bacteriol* **197**: 1893-1905.

Aly, K.A., Anderson, M., Ohr, R.J., and Missiakas, D. (2017) Isolation of a Membrane Protein

Complex for Type VII Secretion in Staphylococcus aureus. *J Bacteriol* **199**.

Anderson, M., Aly, K.A., Chen, Y.H., and Missiakas, D. (2013) Secretion of atypical protein

substrates by the ESAT-6 secretion system of Staphylococcus aureus. *Mol Microbiol*

90: 734-743.

Anderson, M., Chen, Y.H., Butler, E.K., and Missiakas, D.M. (2011) EsaD, a secretion factor

for the Ess pathway in Staphylococcus aureus. *J Bacteriol* **193**: 1583-1589.

Anderson, M., Ohr, R.J., Aly, K.A., Nocadello, S., Kim, H.K., Schneewind, C.E., Schneewind,

O., and Missiakas, D. (2017) EssE Promotes Staphylococcus aureus ESS-Dependent

Protein Secretion To Modify Host Immune Responses during Infection. *J Bacteriol* **199**.

Arsic, B., Zhu, Y., Heinrichs, D.E., and McGavin, M.J. (2012) Induction of the staphylococcal

proteolytic cascade by antimicrobial fatty acids in community acquired methicillin

resistant Staphylococcus aureus. *PLoS One* **7**: e45952.

Bach, J.N., and Bramkamp, M. (2013) Flotillins functionally organize the bacterial membrane.

Mol Microbiol **88**: 1205-1217.

Bae, T., and Schneewind, O. (2006) Allelic replacement in Staphylococcus aureus with

inducible counter-selection. *Plasmid* **55**: 58-63.

Binns, D., Dimmer, E., Huntley, R., Barrell, D., O'Donovan, C., and Apweiler, R. (2009)

QuickGO: a web-based tool for Gene Ontology searching. *Bioinformatics* **25**: 3045-

3046.

Bobrovskyy, M., Willing, S.E., Schneewind, O., and Missiakas, D. (2018) EssH peptidoglycan
hydrolase enables Staphylococcus aureus type VII secretion across the bacterial cell
wall envelope. *J Bacteriol*.

Burts, M.L., DeDent, A.C., and Missiakas, D.M. (2008) EsaC substrate for the ESAT-6
secretion pathway and its role in persistent infections of Staphylococcus aureus. *Mol*
Microbiol **69**: 736-746.

Burts, M.L., Williams, W.A., DeBord, K., and Missiakas, D.M. (2005) EsxA and EsxB are
secreted by an ESAT-6-like system that is required for the pathogenesis of
Staphylococcus aureus infections. *Proc Natl Acad Sci U S A* **102**: 1169-1174.

Cao, Z., Casabona, M.G., Kneuper, H., Chalmers, J.D., and Palmer, T. (2016) The type VII
secretion system of Staphylococcus aureus secretes a nuclease toxin that targets
competitor bacteria. *Nat Microbiol* **2**: 16183.

Cartron, M.L., England, S.R., Chiriac, A.I., Josten, M., Turner, R., Rauter, Y., Hurd, A., Sahl,
H.G., Jones, S., and Foster, S.J. (2014) Bactericidal activity of the human skin fatty
acid cis-6-hexadecanoic acid on Staphylococcus aureus. *Antimicrob Agents*
Chemother **58**: 3599-3609.

Casabona, M.G., Buchanan, G., Zoltner, M., Harkins, C.P., Holden, M.T.G., and Palmer, T.
(2017a) Functional analysis of the EsaB component of the Staphylococcus aureus
Type VII secretion system. *Microbiology*.

Casabona, M.G., Kneuper, H., Alferes de Lima, D., Harkins, C.P., Zoltner, M., Hjerde, E.,
Holden, M.T.G., and Palmer, T. (2017b) Haem-iron plays a key role in the regulation
of the Ess/type VII secretion system of Staphylococcus aureus RN6390. *Microbiology*
163: 1839-1850.

Cassat, J., Dunman, P.M., Murphy, E., Projan, S.J., Beenken, K.E., Palm, K.J., Yang, S.J.,
Rice, K.C., Bayles, K.W., and Smeltzer, M.S. (2006) Transcriptional profiling of a
Staphylococcus aureus clinical isolate and its isogenic agr and sarA mutants reveals
global differences in comparison to the laboratory strain RN6390. *Microbiology* **152**:
3075-3090.

Chamberlain, N.R., Mehrtens, B.G., Xiong, Z., Kapral, F.A., Boardman, J.L., and Rearick, J.I.

(1991) Correlation of carotenoid production, decreased membrane fluidity, and

resistance to oleic acid killing in Staphylococcus aureus 18Z. *Infect Immun* **59**: 4332-

4337.

Clarke, S.R., Mohamed, R., Bian, L., Routh, A.F., Kokai-Kun, J.F., Mond, J.J., Tarkowski, A.,

and Foster, S.J. (2007) The Staphylococcus aureus surface protein IsdA mediates

resistance to innate defenses of human skin. *Cell Host Microbe* **1**: 199-212.

Conrad, W.H., Osman, M.M., Shanahan, J.K., Chu, F., Takaki, K.K., Cameron, J., Hopkinson-

Woolley, D., Brosch, R., and Ramakrishnan, L. (2017) Mycobacterial ESX-1 secretion

system mediates host cell lysis through bacterium contact-dependent gross membrane

disruptions. *Proc Natl Acad Sci U S A* **114**: 1371-1376.

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014) Accurate

proteome-wide label-free quantification by delayed normalization and maximal peptide

ratio extraction, termed MaxLFQ. *Mol Cell Proteomics* **13**: 2513-2526.

Cruciani, M., Etna, M.P., Camilli, R., Giacomini, E., Percario, Z.A., Severa, M., Sandini, S.,

Rizzo, F., Brandi, V., Balsamo, G., Policelli, F., Affabris, E., Pantosti, A., Bagnoli, F.,

and Coccia, E.M. (2017) Staphylococcus aureus Esx Factors Control Human Dendritic

Cell Functions Conditioning Th1/Th17 Response. *Front Cell Infect Microbiol* **7**: 330.

Delekta, P.C., Shook, J.C., Lydic, T.A., Mulks, M.H., and Hammer, N.D. (2018)

Staphylococcus aureus utilizes host-derived lipoprotein particles as sources of

exogenous fatty acids. *J Bacteriol*.

Dreisbach, A., Hempel, K., Buist, G., Hecker, M., Becher, D., and van Dijl, J.M. (2010) Profiling

the surfacome of Staphylococcus aureus. *Proteomics* **10**: 3082-3096.

Fey, P.D., Endres, J.L., Yajjala, V.K., Widhelm, T.J., Boissy, R.J., Bose, J.L., and Bayles, K.W.

(2013) A genetic resource for rapid and comprehensive phenotype screening of

nonessential Staphylococcus aureus genes. *MBio* **4**: e00537-00512.

Garcia-Fernandez, E., Koch, G., Wagner, R.M., Fekete, A., Stengel, S.T., Schneider, J.,

Mielich-Suss, B., Geibel, S., Markert, S.M., Stigloher, C., and Lopez, D. (2017)

Membrane Microdomain Disassembly Inhibits MRSA Antibiotic Resistance. *Cell* **171**:

1354-1367 e1320.

Geiger, T., Francois, P., Liebeke, M., Fraunholz, M., Goerke, C., Krismer, B., Schrenzel, J.,

Lalk, M., and Wolz, C. (2012) The stringent response of Staphylococcus aureus and

its impact on survival after phagocytosis through the induction of intracellular PSMs

expression. *PLoS Pathog* **8**: e1003016.

George, S.E., Hrubesch, J., Breuing, I., Vetter, N., Korn, N., Hennemann, K., Bleul, L.,

Willmann, M., Ebner, P., Gotz, F., and Wolz, C. (2019) Oxidative stress drives the

selection of quorum sensing mutants in the Staphylococcus aureus population. *Proc*

Natl Acad Sci U S A **116**: 19145-19154.

Giachino, P., Engelmann, S., and Bischoff, M. (2001) Sigma(B) activity depends on RsbU in

Staphylococcus aureus. *J Bacteriol* **183**: 1843-1852.

Goodman, J.K., Zampronio, C.G., Jones, A.M.E., and Hernandez-Fernaud, J.R. (2018)

Updates of the In-Gel Digestion Method for Protein Analysis by Mass Spectrometry.

Proteomics **18**: e1800236.

Gordon, R.J., and Lowy, F.D. (2008) Pathogenesis of methicillin-resistant Staphylococcus

aureus infection. *Clin Infect Dis* **46 Suppl 5**: S350-359.

Greenway, D.L., and Dyke, K.G. (1979) Mechanism of the inhibitory action of linoleic acid on

the growth of Staphylococcus aureus. *J Gen Microbiol* **115**: 233-245.

Groicher, K.H., Firek, B.A., Fujimoto, D.F., and Bayles, K.W. (2000) The Staphylococcus

aureus lrgAB operon modulates murein hydrolase activity and penicillin tolerance. *J*

Bacteriol **182**: 1794-1801.

Hempel, K., Herbst, F.A., Moche, M., Hecker, M., and Becher, D. (2011) Quantitative

proteomic view on secreted, cell surface-associated, and cytoplasmic proteins of the

methicillin-resistant human pathogen Staphylococcus aureus under iron-limited

conditions. *J Proteome Res* **10**: 1657-1666.

Horsburgh, M.J., Ingham, E., and Foster, S.J. (2001) In Staphylococcus aureus, fur is an

interactive regulator with PerR, contributes to virulence, and Is necessary for oxidative

stress resistance through positive regulation of catalase and iron homeostasis. *J*
Bacteriol **183**: 468-475.

Ishii, K., Adachi, T., Yasukawa, J., Suzuki, Y., Hamamoto, H., and Sekimizu, K. (2014)
Induction of virulence gene expression in Staphylococcus aureus by pulmonary
surfactant. *Infect Immun* **82**: 1500-1510.

Jager, F., Kneuper, H., and Palmer, T. (2018) EssC is a specificity determinant for
Staphylococcus aureus type VII secretion. *Microbiology* **164**: 816-820.

Johnson, M., Sengupta, M., Purves, J., Tarrant, E., Williams, P.H., Cockayne, A., Muthaiyan,
A., Stephenson, R., Ledala, N., Wilkinson, B.J., Jayaswal, R.K., and Morrissey, J.A.
(2011) Fur is required for the activation of virulence gene expression through the
induction of the sae regulatory system in Staphylococcus aureus. *Int J Med Microbiol*
301: 44-52.

Kelsey, J.A., Bayles, K.W., Shafii, B., and McGuire, M.A. (2006) Fatty acids and
monoacylglycerols inhibit growth of Staphylococcus aureus. *Lipids* **41**: 951-961.

Kenny, J.G., Ward, D., Josefsson, E., Jonsson, I.M., Hinds, J., Rees, H.H., Lindsay, J.A.,
Tarkowski, A., and Horsburgh, M.J. (2009) The Staphylococcus aureus response to
unsaturated long chain free fatty acids: survival mechanisms and virulence
implications. *PLoS One* **4**: e4344.

Kneuper, H., Cao, Z.P., Twomey, K.B., Zoltner, M., Jager, F., Cargill, J.S., Chalmers, J., van
der Kooi-Pol, M.M., van Dijl, J.M., Ryan, R.P., Hunter, W.N., and Palmer, T. (2014)
Heterogeneity in ess transcriptional organization and variable contribution of the
Ess/Type VII protein secretion system to virulence across closely related
Staphyloccus aureus strains. *Mol Microbiol* **93**: 928-943.

Kohler, T., Weidenmaier, C., and Peschel, A. (2009) Wall teichoic acid protects
Staphylococcus aureus against antimicrobial fatty acids from human skin. *J Bacteriol*
191: 4482-4484.

Korea, C.G., Balsamo, G., Pezzicoli, A., Merakou, C., Tavarini, S., Bagnoli, F., Serruto, D.,
and Unnikrishnan, M. (2014) Staphylococcal Esx proteins modulate apoptosis and

751 release of intracellular Staphylococcus aureus during infection in epithelial cells. *Infect*

752  *Immun* **82**: 4144-4153.

753 Lee, A.S., de Lencastre, H., Garau, J., Kluytmans, J., Malhotra-Kumar, S., Peschel, A., and

754  Harbarth, S. (2018) Methicillin-resistant Staphylococcus aureus. *Nat Rev Dis Primers*

755  **4**: 18033.

756 Liang, X., Zheng, L., Landwehr, C., Lunsford, D., Holmes, D., and Ji, Y. (2005) Global

757  regulation of gene expression by ArlRS, a two-component signal transduction

758  regulatory system of Staphylococcus aureus. *J Bacteriol* **187**: 5486-5492.

759 Lopez, M.S., Tan, I.S., Yan, D., Kang, J., McCreary, M., Modrusan, Z., Austin, C.D., Xu, M.,

760  and Brown, E.J. (2017) Host-derived fatty acids activate type VII secretion in

761  Staphylococcus aureus. *Proc Natl Acad Sci U S A* **114**: 11223-11228.

762 Mashruwala, A.A., Guchte, A.V., and Boyd, J.M. (2017) Impaired respiration elicits SrrAB-

763  dependent programmed cell lysis and biofilm formation in Staphylococcus aureus. *Elife*

764  **6**.

765 Mielich-Suss, B., Wagner, R.M., Mietrach, N., Hertlein, T., Marincola, G., Ohlsen, K., Geibel,

766  S., and Lopez, D. (2017) Flotillin scaffold activity contributes to type VII secretion

767  system assembly in Staphylococcus aureus. *PLoS Pathog* **13**: e1006728.

768 Moran, J.C., Alorabi, J.A., and Horsburgh, M.J. (2017) Comparative Transcriptomics Reveals

769  Discrete Survival Responses of S. aureus and S. epidermidis to Sapienic Acid. *Front*

770  *Microbiol* **8**: 33.

771 Neumann, Y., Ohlsen, K., Donat, S., Engelmann, S., Kusch, H., Albrecht, D., Cartron, M.,

772  Hurd, A., and Foster, S.J. (2015) The effect of skin fatty acids on Staphylococcus

773  aureus. *Arch Microbiol* **197**: 245-267.

774 Nguyen, M.T., Hanzelmann, D., Hartner, T., Peschel, A., and Gotz, F. (2016) Skin-Specific

775  Unsaturated Fatty Acids Boost the Staphylococcus aureus Innate Immune Response.

776  *Infect Immun* **84**: 205-215.

777 Ohr, R.J., Anderson, M., Shi, M., Schneewind, O., and Missiakas, D. (2017) EssD, a Nuclease

778  Effector of the Staphylococcus aureus ESS Pathway. *J Bacteriol* **199**.

Parsons, J.B., Broussard, T.C., Bose, J.L., Rosch, J.W., Jackson, P., Subramanian, C., and

Rock, C.O. (2014) Identification of a two-component fatty acid kinase responsible for

host fatty acid incorporation by Staphylococcus aureus. *Proc Natl Acad Sci U S A* **111**:

10532-10537.

Parsons, J.B., Frank, M.W., Subramanian, C., Saenkham, P., and Rock, C.O. (2011)

Metabolic basis for the differential susceptibility of Gram-positive pathogens to fatty

acid synthesis inhibitors. *Proc Natl Acad Sci U S A* **108**: 15378-15383.

Parsons, J.B., Yao, J., Frank, M.W., Jackson, P., and Rock, C.O. (2012) Membrane disruption

by antimicrobial fatty acids releases low-molecular-weight proteins from

Staphylococcus aureus. *J Bacteriol* **194**: 5294-5304.

Pasztor, L., Ziebandt, A.K., Nega, M., Schlag, M., Haase, S., Franz-Wachtel, M., Madlung, J.,

Nordheim, A., Heinrichs, D.E., and Gotz, F. (2010) Staphylococcal major autolysin (Atl)

is involved in excretion of cytoplasmic proteins. *J Biol Chem* **285**: 36794-36803.

Patton, T.G., Yang, S.J., and Bayles, K.W. (2006) The role of proton motive force in expression

of the Staphylococcus aureus cid and lrg operons. *Mol Microbiol* **59**: 1395-1404.

Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J.,

Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., Perez, E., Uszkoreit, J., Pfeuffer, J.,

Sachsenberg, T., Yilmaz, S., Tiwary, S., Cox, J., Audain, E., Walzer, M., Jarnuczak,

A.F., Ternent, T., Brazma, A., and Vizcaino, J.A. (2019) The PRIDE database and

related tools and resources in 2019: improving support for quantification data. *Nucleic*

Acids Res **47**: D442-D450.

Pulschen, A.A., Sastre, D.E., Machinandiarena, F., Crotta Asis, A., Albanesi, D., de Mendoza,

D., and Gueiros-Filho, F.J. (2017) The stringent response plays a key role in Bacillus

subtilis survival of fatty acid starvation. *Mol Microbiol* **103**: 698-712.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., and Smyth, G.K. (2015) limma

powers differential expression analyses for RNA-sequencing and microarray studies.

Nucleic Acids Res **43**: e47.

Saeloh, D., Tipmanee, V., Jim, K.K., Dekker, M.P., Bitter, W., Voravuthikunchai, S.P., Wenzel,

M., and Hamoen, L.W. (2018) The novel antibiotic rhodomyrtone traps membrane

proteins in vesicles with increased fluidity. *PLoS Pathog* **14**: e1006876.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,

S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V.,

Eliceiri, K., Tomancak, P., and Cardona, A. (2012) Fiji: an open-source platform for

biological-image analysis. *Nat Methods* **9**: 676-682.

Sen, S., Sirobhushanam, S., Johnson, S.R., Song, Y., Tefft, R., Gatto, C., and Wilkinson, B.J.

(2016) Growth-Environment Dependent Modulation of Staphylococcus aureus

Branched-Chain to Straight-Chain Fatty Acid Ratio and Incorporation of Unsaturated

Fatty Acids. *PLoS One* **11**: e0165300.

Smith, A.F., Rihtman, B., Stirrup, R., Silvano, E., Mausz, M.A., Scanlan, D.J., and Chen, Y.

(2019) Elucidation of glutamine lipid biosynthesis in marine bacteria reveals its

importance under phosphorus deplete growth in Rhodobacteraceae. *ISME J* **13**: 39-

49.

Solis, N., Parker, B.L., Kwong, S.M., Robinson, G., Firth, N., and Cordwell, S.J. (2014)

Staphylococcus aureus surface proteins involved in adaptation to oxacillin identified

using a novel cell shaving approach. *J Proteome Res* **13**: 2954-2972.

Tiwari, K.B., Gatto, C., and Wilkinson, B.J. (2018) Interrelationships between Fatty Acid

Composition, Staphyloxanthin Content, Fluidity, and Carbon Flow in the

Staphylococcus aureus Membrane. *Molecules* **23**.

Tong, S.Y., Davis, J.S., Eichenberger, E., Holland, T.L., and Fowler, V.G., Jr. (2015)

Staphylococcus aureus infections: epidemiology, pathophysiology, clinical

manifestations, and management. *Clin Microbiol Rev* **28**: 603-661.

Truong-Bolduc, Q.C., Villet, R.A., Estabrooks, Z.A., and Hooper, D.C. (2014) Native efflux

pumps contribute resistance to antimicrobials of skin and the ability of Staphylococcus

aureus to colonize skin. *J Infect Dis* **209**: 1485-1493.

Unnikrishnan, M., Constantinidou, C., Palmer, T., and Pallen, M.J. (2017) The Enigmatic Esx
Proteins: Looking Beyond Mycobacteria. *Trends Microbiol* **25**: 192-204.

Varemo, L., Nielsen, J., and Nookaew, I. (2013) Enriching the gene set analysis of genome-
wide data by incorporating directionality of gene expression and combining statistical
hypotheses and methods. *Nucleic Acids Res* **41**: 4378-4391.

Ventura, C.L., Malachowa, N., Hammer, C.H., Nardone, G.A., Robinson, M.A., Kobayashi,
S.D., and DeLeo, F.R. (2010) Identification of a novel Staphylococcus aureus two-
component leukotoxin using cell surface proteomics. *PLoS One* **5**: e11634.

von Eiff, C., Becker, K., Machka, K., Stammer, H., and Peters, G. (2001) Nasal carriage as a
source of Staphylococcus aureus bacteremia. Study Group. *N Engl J Med* **344**: 11-16.

Wang, Y., Hu, M., Liu, Q., Qin, J., Dai, Y., He, L., Li, T., Zheng, B., Zhou, F., Yu, K., Fang, J.,
Liu, X., Otto, M., and Li, M. (2016) Role of the ESAT-6 secretion system in virulence
of the emerging community-associated Staphylococcus aureus lineage ST398. *Sci*
Rep **6**: 25163.

Warne, B., Harkins, C.P., Harris, S.R., Vatsiou, A., Stanley-Wall, N., Parkhill, J., Peacock,
S.J., Palmer, T., and Holden, M.T. (2016) The Ess/Type VII secretion system of
Staphylococcus aureus shows unexpected genetic diversity. *BMC Genomics* **17**: 222.

Wood, T.E., Howard, S.A., Forster, A., Nolan, L.M., Manoli, E., Bullen, N.P., Yau, H.C.L.,
Hachani, A., Hayward, R.D., Whitney, J.C., Vollmer, W., Freemont, P.S., and Filloux,
A. (2019) The Pseudomonas aeruginosa T6SS Delivers a Periplasmic Toxin that
Disrupts Bacterial Cell Morphology. *Cell Rep* **29**: 187-201 e187.

Zoltner, M., Ng, W.M., Money, J.J., Fyfe, P.K., Kneuper, H., Palmer, T., and Hunter, W.N.
(2016) EssC: domain structures inform on the elusive translocation channel in the Type
VII secretion system. *Biochem J* **473**: 1941-1952.

Figure legends

Fig. 1. Membrane defects in the absence of EsxC.

A. FM1-43 fluorescence of *S. aureus* WT, Δ*essC*, and Δ*esxC* as measured with a

plate reader. Means ± standard deviation (SD) are shown, *n* = 6; * indicates *P* < 0.05

using a Kruskal-Wallis test with Dunn's test.

B. Survival of Δ*esxC* or Δ*essC* after a 2h-treatment with 0.1% Triton X-100 relative to

the WT. Means are shown; error bars represent standard deviation (SD). *n* = 4, *

indicates *P* < 0.05 using a Kruskal-Wallis test with Dunn's test. The panel on the right

shows a representative picture of bacteria before and after Triton X-100-treatment.

C. The membrane fluidity of WT and Δ*esxC* as measured with a pyrene decanoic acid

staining-based assay. Mean values are shown; error bars represent standard error of

the mean (SEM). *n* = 6, * indicates *P* < 0.05 using a two-tailed t-test.

D. Widefield micrographs of *S. aureus* WT and Δ*esxC* after growth in TSB to OD_{600} of

1.0 in the presence of the lipophilic dye DiIC12. Images are representative of 4

independent experiments.

E. The DiIC12 fluorescence of 80 bacterial clusters from different fields per strain was

quantitated with ImageJ. Means ± standard deviation (SD) are shown, *n* = 4; indicates

**** *P* < 0.0001, using a Kruskal-Wallis rank test

Fig. 2. Enhanced *S. aureus* growth inhibition by linoleic acid in T7SS mutants.

A. *S. aureus* WT USA300, Δ*essC*, and Δ*esxC* were grown in TSB or TSB

supplemented with 80 µM of either linoleic (LA) or stearic acid (SA). Means ± standard

error of the mean (SEM) are shown.

B. After 14h growth as described in (A) bacteria were serially diluted, and CFU were

determined. Mean values are presented, and the error bars represent SD. n = 3, **

indicates $P < 0.01$ using one-way ANOVA with Dunnett's test.

C. USA300 WT with the empty pOS1 plasmid (WT pOS1) and USA300 JE2 *esxC*

mutant with either pOS1 (Δ*esxC* pOS1) or pOS1-*esxC* (Δ*esxC* pOS1-*esxC*) were

grown in TSB or TSB + 80 µM LA as described in (A) followed by CFU estimation.

Mean values are shown; error bars represent SD. n = 5, ** indicates $P < 0.01$ using

one-way ANOVA with Dunnett's test.

D. Newman WT with the empty pOS1 plasmid (WT pOS1) and Newman *esxA* mutant

with either pOS1 (Δ*esxA* pOS1) or pOS1-*esxA* (Δ*esxA* p*esxA*) were grown in TSB or

TSB + 40 µM LA or SA. Means ± standard error of the mean (SEM) are shown, n = 4.

Fig. 3. T7SS mutants display increased membrane permeability upon LA

binding.

A. Chemical structure of azide functionalised linoleic acid (azide-LA; N^6-diazo-N^2-

((9Z,12Z)-octadeca-9,12-dienoyl)lysine, N_3-LA). Highlighted in green is the azido

lysine.

B. *S. aureus* USA300 WT, Δ*essC*, and Δ*esxC* were grown with shaking in TSB to

OD_{600} of 1.0. Bacteria were then stained for 15 min with 10 µM azide-LA prior to

labelling for 1 h with alkyne Alexa Fluor 488. Mean percentage of fluorescence values

relative to WT (100%) are presented; error bars represent SD, n = 5.

C. Micrographs of bacteria grown in TSB and treated as described in (B) and

additionally stained with propidium iodide (PI).

D. ImageJ was used to quantitate PI fluorescence of bacterial clusters from 12 different

fields per strain. Each box-and-whisker plot depicts the minimal and maximal PI

intensities, the median is the vertical bar inside the box, which is delimited by the lower

and upper quartiles. ** indicates $P < 0.01$ using one-way ANOVA with Dunnett's test.

Fig. 4. T7SS mutants display increased PI staining when treated with LA.

Live/Dead staining of *S. aureus* USA300 WT, Δ*essC* or Δ*esxC* mutants after growth

to OD_{600} of 1.0, without (**A**) or with treatment with 80µM (**B**) linoleic acid. Images are

representative of 3 independent experiments.

C. The ratio of SYTO 9: PI fluorescence (live:dead cells) of 25 different fields per strain

was quantitated with ImageJ. Means ± SD are shown, $n = 3$; *** indicates $P < 0.001$,

** indicates $P < 0.01$ using a one-way ANOVA with Tukey's multiple-comparison test

Fig. 5. T7SS mutants are less able to incorporate LA into their phospholipids.

Representative HPLC chromatograms of native phosphatidylglycerol (PG) species of

S. aureus USA300 JE2 WT (**A**) or Δ*esxC* (**B**) grown in TSB (top panel) or in TSB

supplemented with 10 µM LA (bottom panel), in negative ionisation mode. Relative

quantification of the indicated PG species containing an unsaturated FA in WT, Δ*essC*

and Δ*esxC*. C18:2-containing PG species (**C**) and total unsaturated exogenous PG

species (**D**) are presented as ratios of total PG species. Mean values are shown; error

bars represent SD. $n = 3$, * indicates $P < 0.05$ using one-way ANOVA with Dunnett's

test.

Fig. 6. Quantitative proteomics shows altered cellular content and bacterial

response to LA in T7SS mutants. *S. aureus* USA300 WT and mutants (Δ*essC* and

Δ*esxC*) were grown in TSB or TSB supplemented with LA.

A. Venn diagram showing the number of proteins with altered abundance compared

to WT specific to Δ*essC* (23) or Δ*esxC* (10), and common to Δ*essC* and Δ*esxC* (14).

B. The fourteen proteins that are similarly changed in Δ*essC* and Δ*esxC* mutants are

highlighted on a volcano plot.

C. Volcano plot showing the extensive change in the LA-treated WT compared to WT.

D. Venn diagram displaying the numbers of proteins with altered relative abundance

upon LA challenge of WT (LA.WT), Δ*essC* (LA.dEssC) or Δ*esxC* (LA.dEsxC)

compared to the respective untreated samples.

Fig. 7. An altered oxidoreductive response in T7SS mutants in response to LA.

Heatmaps depicting the *P* values of enriched (**A**) or diminished (**B**) molecular functions

following a gene set analysis based on GO (gene ontology) annotations. Molecular

functions that are changed in at least one strain ($P < 0.05$) following growth in presence

of LA are shown. The shades of blue (**A**) or red (**B**) correspond to $-\log_{10}$ (*P* value).

ROS levels were measured in cultures of *S. aureus* USA300 JE2 WT, Δ*essC* or Δ*esxC*

grown to OD_{600} of 1.0 treated (**C**) with or without LA (**D**) using DCF reagent. Means ±

SD are shown N=5. * indicates $P < 0.05$, ** indicates $P < 0.01$, *** indicates $P < 0.001$

using the Kruskal-Wallis rank test.

Fig. S1. USA300 JE2 WT and Δ*esxC* strains display similar growth rates. WT and

Δ*esxC* were grown in TSB, and OD_{600} monitored with a Novaspec® Pro

spectrophotometer. Data shown are means of three independent experiments, and

the error bars indicate the standard errors of the mean.

Fig. S2. EsxC associates with *S. aureus* membrane. Immunoblot analysis of cell

membrane (CM) or cell wall (CW) fractions of WT (USA300), Δ*essC*, and Δ*esxC* with

anti-EsxC sera or anti-PBP2a antibodies (loading control).

Fig. S3. Volcano plot of the quantitative proteomic analysis of surface proteins

in Δ*esxC* compared to WT. The relative abundance of each protein ($\log_2$ fold change,

X-axis) and its statistical significance (P value, Y-axis) are shown in the graph.

Proteins decreased by more than half in Δ$esxC$ ($\log_2$ fold change < -1 and P value <

0.05) are shown in green.

Fig. S4. *S. aureus* growth inhibition by arachidonic acid is increased in T7SS

mutants. *S. aureus* WT USA300, Δ$essC$, and Δ$esxC$ were grown in TSB or TSB

supplemented with 80 µM arachidonic acid (AA). Means ± standard error of the mean

(SEM) are shown, n = 3.

Fig. S5. T7SS substrates contribute to resistance to linoleic acid toxicity.

A. *S. aureus* USA300 wild-type (WT) and USA300 $esxA$ (Δ$esxA$) or $esxB$ (Δ$esxB$)

deletion mutants were grown in TSB or TSB supplemented with 80 µM linoleic (LA) or

stearic acid (SA).

B. *S. aureus* Newman WT and Newman $esxA$ (Δ$esxA$) or $esxB$ (Δ$esxB$) deletion

mutants were grown similarly in TSB or TSB + 40 µM LA or SA.

C. Growth curves as described in (A) were done with RN6390 wild-type (WT) and

RN6390 $essC$ (Δ$essC$) or $esxC$ (Δ$esxC$) deletion mutants. Data shown in (A), (B), and

(C) are representative of at least three independent experiments.

Fig. S6. USA300 JE2 WT and T7SS mutants display similar lipids.

Representative HPLC chromatograms of the indicated bacteria grown in TSB (A) or in

TSB supplemented with LA (B), in negative ionisation mode. Phosphatidylglycerol

(PG) is highlighted.

C. Representative HPLC chromatograms of native PG species of Δ$essC$ grown in TSB

(top panel) or in TSB supplemented with LA (bottom panel), in negative ionisation

mode.

Fig. S7. LA (C18:2) is elongated and incorporated into _S. aureus_ phosphatidylglycerol (PG) species. A-C. Representative mass spectrometry fragmentation spectra for PG species containing unsaturated fatty acids, in negative ionisation mode.

A. PG species with mass 731 m/z, containing C18:2 fatty acid (279 m/z).

B. PG species 759 m/z, containing C20:2 fatty acid (307 m/z)

C. PG species 787 m/z, containing C22:2 fatty acid (335 m/z).

D. and E. Relative quantification of the indicated PG species containing an unsaturated fatty acid in WT, Δ_essC_ and Δ_esxC_. C20:2- (D) and C22:2-containing PG species (E) are presented as ratios of total PG species. Data shown are the means and error bars represent SD of three independent experiments.

Fig. S8. USA300 JE2 WT and Δ_esxC_ growth similarly in presence of sub-inhibitory amounts of linoleic acid. _S. aureus_ WT USA300 and Δ_esxC_ were grown in TSB or TSB supplemented with 40 μM linoleic acid (LA). Means ± SEM are shown, _n_ = 3.

Fig. S9. Principal component analysis (PCA) of the _S. aureus_ cellular proteomic profiles. PCA was performed on all the identified proteins of USA300 JE2 WT and T7SS mutants grown in TSB (untreated) or TSB + LA (LA-treated). Each dot represents a biological replicate.

Fig. S10. ^{1}H NMR spectra of NHS-LA (A) and azide-LA (B) in CDCl$_3$. Both spectra were recorded on a Bruker Advance 300 spectrometer (300 MHz) at 27 °C. The letters indicate the chemical shift δ (in parts per million, ppm) of the protons in each molecule.

Dataset S1. Differentially abundant proteins in WT USA300 JE2 and T7SS mutants in response to linoleic acid.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7