The ER membrane protein complex is important for the biogenesis of flavivirus polyproteins

Ashley M Ngo[1], Matthew J Shurtleff[2], Katerina D Popova[2], Jessie Kulsuptrakul[1], Jonathan S Weissman[2,3], Andreas S Puschnik[1,4]

[1] Chan Zuckerberg Biohub, San Francisco, United States

[2] Department of Cellular and Molecular Pharmacology, University of California, San Francisco, San Francisco, United States

[3] Howard Hughes Medical Institute, University of California, San Francisco, San Francisco, United States

[4] Correspondence to: andreas.puschnik@czbiohub.org

Abstract

Flaviviruses such as dengue (DENV), Zika (ZIKV) or West Nile virus (WNV) translate their genome as a single multi-pass transmembrane (TM) protein at the endoplasmic reticulum (ER) membrane. Several genetic knockout (KO) screens identified the ER membrane protein complex (EMC) as a critical host component for flavivirus infection. The EMC facilitates accurate insertion, topology and/or stabilization of specific cellular TM proteins including a subset of tail-anchored proteins and G protein-coupled receptors. Here, we show that deletion of EMC in human cell lines decreased infection with DENV, ZIKV and WNV by 100 to 10,000-fold. Using replicon and immunoblotting studies in EMC KO cells, we demonstrated that the EMC was essential for viral protein expression. Ribosome Profiling of DENV-infected wild-type and EMC KO cells revealed no drastic differences in translation efficiency. Instead, absence of EMC led to a large fraction of expressed viral proteins being targeted to the proteasome post-translationally. We detected a decrease in stability in NS4A-NS4B of the viral polyprotein, a region rich in transmembrane domains. Additionally, we identified non-synonymous point mutations in NS4A and NS4B by

performing iterative passaging of DENV on EMC KO cells. Adaptive mutations rescued both viral

replication and stable expression of the NS4A-NS4B polyprotein segment in EMC KO cells.

Lastly, we showed a physical interaction between the EMC and DENV NS4B protein post-

cleavage and rapid degradation of processed NS4B in the absence of EMC. Together, our results

suggest that the EMC engages with DENV polyproteins to ensure proper biogenesis of the NS4A-

NS4B region, and provide further evidence for the cellular function of the EMC in the stable

expression of TM proteins.

Introduction

Flaviviruses such as dengue (DENV), Zika (ZIKV) or West Nile virus (WNV) express their genome

as a single multi-pass transmembrane (TM) protein at the endoplasmic reticulum (ER) membrane.

Subsequently, the polyprotein is cleaved by the viral NS2B-NS3 protease and the cellular signal

peptidase into the individual structural (C, prM, E) and non-structural (NS1-5) proteins required

for replication and assembly into new virions (Neufeldt et al., 2018). Similar to cellular TM proteins,

the viral polyprotein relies on the host cell machinery for targeting to the ER, translocation across

the membrane and insertion of transmembrane domains (TMDs) into the lipid bilayer during its

translation. These processes are facilitated by the signal recognition particle (SRP), the

translocon-associated protein (TRAP) complex (also known as signal-sequence receptor

complex), the Sec61 translocon and the signal peptidase complex. These cell components were

all identified as essential host factors for flavivirus infection in genetic screens underscoring their

importance for virus replication (Krishnan et al., 2008; Marceau et al., 2016; Sessions et al., 2009;

Zhang et al., 2016). For most TM proteins, recognition of a hydrophobic TMD or signal sequence

by the SRP followed by transfer to the Sec61 translocation channel are sufficient for membrane

targeting and accurate topogenesis (Shao and Hegde, 2011). Recently, it was shown that the ER

membrane protein complex (EMC) plays a role in the insertion and/or stabilization of certain TM

proteins (Chitwood et al., 2018; Guna et al., 2018; Shurtleff et al., 2018). Intriguingly, several

genome-wide CRISPR knockout (KO) screens for flavivirus dependency factors showed strong

enrichment of the EMC (Ma et al., 2015; Marceau et al., 2016; Savidis et al., 2016; Zhang et al.,

2016). However, its role in the virus life cycle is not yet understood.

The EMC was originally discovered in a genetic screen for yeast mutants modulating the unfolded

protein response (UPR) (Jonikas et al., 2009). Genetic interaction mapping and biochemical

purification revealed a six-subunit core complex (EMC1-6) conserved in yeast and mammals

(Christianson et al., 2011; Jonikas et al., 2009). Other than its more general association with UPR

and ER-associated degradation (ERAD), the EMC was shown to interact co-translationally and

enable biogenesis of a subset of multi-pass TM proteins (Shurtleff et al., 2018). Biochemical

characterization of select EMC-dependent TM proteins, such as tail-anchored proteins and G

protein-coupled receptors (GPCRs), revealed that the EMC is capable of inserting TMDs into the

ER membrane (Chitwood et al., 2018; Guna et al., 2018).

Here, we investigated how the mosquito-borne flaviviruses, DENV, ZIKV and WNV, depend on

the EMC during infection. We showed that the EMC was required for DENV polyprotein

biogenesis and identified a segment spanning NS4A-4B as the region dependent on the EMC for

stable expression. Absence of the EMC resulted in post-translational degradation of viral proteins.

Together, our data highlights how flaviviruses hijack important machinery for TM protein

biogenesis to achieve optimal expression of their polyproteins, which reinforces a role for the EMC

in stabilizing challenging TM proteins during synthesis.

Results

To validate the enrichment of EMC components in the flavivirus CRISPR screens, we generated

isogenic EMC subunit KO cell lines in Huh7.5.1 and HEK293FT, two cell types that were used for

the genetic screens (Figure S1A). Upon infection with DENV expressing Renilla luciferase

(DENV-Luc), we measured a ~10,000-fold reduction in viral replication in Huh7.5.1 EMC1-6 KO

cells (Figure 1A) and a 10-1,000-fold reduction in HEK293FT EMC1-5 KO cells (Figure 1B). The

differences in the extent of the phenotypes may be attributed to the presence of compensatory

pathways or the degree of disruption of the entire protein complex between the two cell types

upon knockout of a single subunit. The strong deficit in DENV replication in the absence of the

EMC persisted over a 30-72h period (Figure S1B). Complementation of EMC2 and EMC4 KO

cells with the respective cDNA completely restored levels of DENV infection (Figures 1C and

S1C). Lastly, we tested the effect of EMC4 KO on several flaviviruses and saw a significant

decrease of viral RNA for DENV, WNV and ZIKV but not hepatitis C virus (HCV), which is a more

distantly related member of the *Flaviviridae* family (Figures 1D and S1D). This is congruent with

results of an HCV CRISPR screen that did not identify the EMC as a critical host factor (Marceau

et al., 2016). This finding highlights that there is a specific, evolutionarily conserved dependency

on the EMC among mosquito-borne flaviviruses and not a universal disruption of trafficking to

and/or translation or replication at the ER membrane, which would affect a wider range of viruses.

Next, we sought to determine at which step of the viral life cycle the EMC is required. As the EMC

is important for the expression of a large set of TM proteins, its disruption leads to an altered cell

surface proteome, which may affect viral entry. We measured effects on viral uptake in WT and

EMC4 KO cells by quantifying the amount of internalized viral RNA at 2 and 6 hours post-infection

(hpi), where we did not observe a difference (Figure 2A). Only at 24hpi, a timepoint when viral

replication had occurred, was a difference in RNA load evident between WT and EMC4 KO cells.

Furthermore, we used a DENV replicon expressing luciferase, which bypasses entry and does

not produce viral particles, thus allowing the analysis of the effect of EMC KO specifically on

translation and replication. In this assay, we saw decreased luminescence in EMC4 KO cells

starting at 12 hours post-electroporation suggesting that the EMC is important during or prior to

viral replication (Figure 2B). We additionally performed the replicon assay in the presence of MK-

0608, a nucleoside analogue that inhibits the viral polymerase (Chen et al., 2015), and observed

that the luminescence signal in EMC4 KO cells was lower than in replication-inhibited WT cells

suggesting that deletion of EMC has an effect on viral translation, which happens prior to

replication (Figure S2A). To further dissect effects on translation or genome replication of DENV,

we conducted an immunoblot timecourse series, where WT and EMC4 KO cells were infected at

a high multiplicity of infection (moi=32) and lysates were harvested at early timepoints post-

infection (6-12hpi), when no or only little DENV replication had occurred. Viral protein expression

was lower in EMC4 KO compared to WT cells as early as 6hpi further supporting a translation

defect (Figures 2C and S2B). To rule out any contribution of low levels of replication to the

observed phenotype, we also performed the immunoblot experiment in the presence of MK-0608

and still saw different amounts of DENV protein accumulation between WT and EMC4 KO cells

indicating that the EMC is indeed important for viral protein expression (Figures S2C and S2D).

Interestingly, while the EMC has been implicated in the stable biogenesis of TM proteins, we

found reduced expression of both TMD and non-TMD containing viral proteins. This suggests that

the defect likely occurs at the polyprotein stage thus affecting all individual, cleaved proteins.

The expression of flaviviral proteins is a complex process that requires several steps. DENV

translation is initiated in a cap-dependent manner, whereupon the single-stranded positive-sense

RNA genome is translated into a single ~3,400 amino acid long multi-pass TM protein. This

polyprotein is co- and post-translationally processed by host and viral proteases producing the

individual functionally active structural and non-structural proteins. We first tested whether the

defect occurs at the stage of polyprotein translation. We used ribosome profiling (Ribo-seq),

which measures the density of ribosomes on a given mRNA species with high positional

resolution, to compare translation efficiencies (TE) and successful full-length synthesis of both

cellular and viral proteins in WT and EMC4 KO cells (Ingolia et al., 2009). Any changes in TE or

premature ribosome stalling in EMC KO vs WT cells would be reflected by changes in ribosomal

footprints (as measured by Ribosome profiling) normalized to changes in mRNA abundance (as

measured by RNA-seq). In the uninfected condition, we did not observe dramatic changes in TE

for cellular mRNAs in EMC4 KO vs WT cells (Figure 3A and Supplementary Table 1). This is in

line with previously reported results using CRISPRi knockdown against EMC subunits suggesting

that deletion of EMC minimally impacts the synthesis rates of its client proteins (Shurtleff et al.,

2018). Similarly, we did not find any substantial differences in viral TE in EMC4 KO cells relative

to WT cells at 18 (0.43x) and 44hpi (2.17x) (Figures 3B and 3C and Supplementary Table 1).

Instead, we observed that DENV RNA was among the highest translated (top 0.1%) mRNAs in

both WT and EMC4 KO cells (Figure S3A). The lower number of ribosomal footprints aligning to

the viral genome in EMC4 KO cells was largely due to the reduced number of viral RNA copies

present in the cells. Moreover, footprints were found to map throughout the viral genome in EMC4

KO cells, indicating that synthesis of full-length polyprotein still occurred (Figure S3B). Therefore,

we concluded that translation of DENV polyproteins is not strongly impacted by loss of EMC,

which led us to explore potential post-translational defects.

Proper insertion of TMDs and correct folding are required for stable expression of TM proteins,

while misfolded proteins are generally targeted to the proteasome for degradation via the ERAD

pathway (Olzmann et al., 2013). To probe whether an increased fraction of viral proteins is

degraded by the proteasome in EMC deficient cells, we infected WT and EMC4 KO cells with

DENV in the presence of Bortezomib (BZ), a proteasome inhibitor, thus preventing removal of

misfolded proteins from the cells. Indeed, BZ treatment led to an increase of detectable viral

proteins in EMC4 KO cells but not WT cells (Figure 3D). Addition of BZ, however, did not restore

DENV replication arguing that the proteasome-targeted proteins are non-functional (Figure S3C).

Together, this suggests that in the absence of EMC viral proteins are still translated but are unable

to achieve a stable conformation resulting in their degradation and inability to establish efficient

replication.

For cellular multi-pass client proteins, such as transporters and GPCRs, the EMC co-

translationally engages with specific TMDs in order to support insertion into the membrane and

allow stable expression of the entire TM protein (Chitwood et al., 2018; Shurtleff et al., 2018). We

used two orthogonal methods, a fluorescence-based protein stability assay and viral evolution, to

examine which region of the viral polyprotein requires the EMC for successful expression. First,

we utilized a dual-color fluorescent reporter as previously described (Chitwood et al., 2018). The

C terminus of different segments of the DENV polyprotein was fused to eGFP followed by a viral

T2A sequence and mCherry (G2C) (Figure 4A). Translation of the fusion constructs generates

two protein products at a 1:1 ratio due to peptide bond skipping by the ribosome at the T2A site:

1) a polyprotein segment fused to eGFP and 2) mCherry. Any post-translational instability of the

eGFP fusion protein due to misfolding results in an eGFP:mCherry ratio of less than 1. As

expected, the cytosolic G2C-only construct displayed a 1:1 ratio in both WT and EMC4 KO cells,

while the β_1-adrenergic receptor (ADRB1) fused to G2C exhibited drastic destabilization in EMC4

KO compared to WT cells as was previously reported (Figure 4B) (Chitwood et al., 2018). Applying

this approach to different DENV polyprotein G2C fusion constructs, we observed that the

segments containing only the structural proteins (capsid, prM and E) or NS1-3 were not

destabilized in EMC4 KO Huh7.5.1 cells (Figure 4B). By contrast, constructs encoding both NS4A

and NS4B (but not either of them alone) showed reduced stability in EMC4 KO cells (Figure 4B).

Similar results were obtained in HEK293FT (Figure S4). NS4A has 3 TMDs with the third (termed

2K peptide) being highly hydrophobic and serving as a signal sequence for the translocation of

NS4B into the ER membrane (Lin et al., 1993). Subsequently, the NS4A-2K junction is cleaved

by the viral protease and the 2K-NS4B junction by the signal peptidase. As the 2K-targeted NS4B-

5-G2C construct did not display a stability defect in EMC deficient cells, the results suggest that

EMC mediates stable expression by first engaging within NS4A prior to the 2K peptide and that a

defect in NS4A topogenesis is extended to the NS4B region. Stability of NS4A-4B-G2C was

restored to near WT levels by complementation of EMC4 KO cells with EMC4 cDNA (Figure 4C).

Moreover, treatment of EMC4 KO cells with BZ increased the eGFP:mCherry ratio of NS4A-4B-

G2C in EMC4 KO cells indicating that this region of the polyprotein is misfolded and targeted for

degradation in the absence of EMC (Figure 4D). Therefore, we linked the DENV replication

phenotype in EMC deficient cells to a specific molecular defect in the NS4A-4B region of the

polyprotein.

In a second, orthogonal approach, we performed viral adaptation to evolve EMC-independent

DENV mutants by passaging the virus consecutively on EMC4 KO Huh7.5.1 cells every 3-4 days.

After 20-22 passages we observed cell death in 3 out of 6 independent adaptation experiments.

Sequencing of RNA extracted from supernatant and alignment to the WT16681 DENV genome

revealed two non-synonymous point mutations in NS4A and NS4B, which were conserved across

all three isolates (Figure 5A). The A6665G mutation led to a tyrosine to cysteine and the A7558T

mutation to an asparagine to tyrosine substitution. The majority of other SNPs was silent (Figure

S5). A6665G is predicted to be inside a TMD helix encoding region of NS4A suggesting that this

TMD relies on the EMC for proper ER membrane insertion (Miller et al., 2007). Only one of 1,359

DENV serotype 2 genomes from the NCBI database has a G at this position encoding cysteine,

while other sequences all encode tyrosine. This evolutionary conservation suggests that this

residue indeed has an important function. Similarly, the frequency of A at position 7558 is very

conserved with 99.6%, and there is no T encoding tyrosine in the 1,359 sequenced strains.

Interestingly, A7558T resides in a non-TMD region at the end of NS4B, thus likely supporting

membrane insertion or folding in a different manner (Miller et al., 2006). To demonstrate that the

acquired mutations confer independence from EMC, we infected WT and EMC4 KO cells with

WT16681 or the three evolved DENV isolates and measured replication. WT16681 DENV

displayed a ~100-fold decrease in viral RNA in EMC4 KO relative to WT cells, while the EMC

adapted isolates had equal RNA levels (Figure 5B). To dissect whether both mutations are

required for optimal replication in EMC KO background, we introduced them individually or in

tandem into the DENV-Luc infectious clone and generated virus. While A6665G and A7558T

individually restored DENV replication in EMC4 KO cells to 30 and 86% of WT levels, respectively,

the double mutant displayed complete rescue of viral growth (Figure 5C). We also introduced the

two adapted mutations into NS4A-5-G2C, which showed increased stability in EMC4 KO cells

compared to the WT construct thus reverting the molecular destabilization phenotype (Figure 5D).

Remarkably, the G2C fluorescence assay and viral adaptation independently pinpointed NS4A-

4B of the polyprotein as the region which requires the EMC. The results of the two experiments

suggest that a defect in the stable insertion of the NS4A TMD2 could lead to subsequent failure

in accurate topogenesis and/or stability of NS4B. By contrast, polyprotein segments starting with

2K-NS4B were stably expressed in EMC KO cells as was shown using NS4B-5-G2C construct

(Figure 4A). Therefore, the EMC is required for correct insertion, processing and/or folding of the

NS4A-4B region.

Finally, to probe how absence of EMC causes failure of DENV replication, we transfected a

subgenomic NS2B-4B construct (containing functional NS2B-NS3 for polyprotein cleavage) with

C terminal fusion to eGFP into WT and EMC4 KO cells and immunoblotted against the different

processed products. While there was no notable difference between the higher molecular weight

intermediates (NS2B-NS3-NS4A-NS4B-eGFP or NS4A-NS4B-eGFP), we observed reduced

accumulation of processed NS4B-eGFP in EMC4 KO cells (Figure 6A). BZ treatment indicated

rapid degradation of cleaved NS4B-eGFP in EMC deficient cells. Furthermore, if the EMC acts to

insert or stabilize NS4A-4B, we hypothesized that direct binding of NS4B to EMC post polyprotein-

processing should be detectable in WT cells. We performed a co-immunoprecipitation using

FLAG-tagged EMC4 and detected NS4B but not prM or NS5 in the pulldown (Figure 6B). Overall,

this supports the finding that EMC physically engages with NS4A-4B to facilitate its biogenesis

and stable expression.

Discussion

Based on our results and previous work (reviewed in Chitwood and Hegde, 2019) we propose the

following model for the role of the EMC in flavivirus infection. During translation of the polyprotein

the EMC engages with TMDs in NS4A and NS4B to facilitate their biogenesis, correct folding and

stable expression (Figure 6C). In the absence of the EMC, TMDs in the NS4A-4B region are not

stably or accurately integrated in the ER membrane leading to misfolding and degradation of

processed NS4B (and potentially also NS4A). NS4A and NS4B fulfill crucial functions for viral

replication. NS4A induces ER membrane rearrangements important for the formation of viral

replication complexes (Miller et al., 2007; Roosendaal et al., 2006) while NS4B interacts with the

NS3 helicase and also suppresses the cellular interferon response (Muñoz-Jordán et al., 2005;

Zou et al., 2015). It remains to be studied whether the other viral proteins are destabilized due to

an overall topology defect in the NS4A-4B region of the polyprotein or if lack of functional NS4A

and/or NS4B inside the replication complexes results in instability of all viral proteins.

In addition to characterizing the role of the EMC in the viral life cycle, our data also helps to further

elucidate its cellular function. The emerging role of the EMC to facilitate topogenesis has been

implicated for several cellular TM proteins (Chitwood et al., 2018; Coelho et al., 2019; Guna et

al., 2018; Shurtleff et al., 2018; Volkmar et al., 2019). For a subset of tail-anchored proteins, the

EMC inserts the sole C-terminal TMD after translation is completed in the cytosol (Guna et al.,

2018). Moreover, the EMC co-translationally inserts the first N-terminal TMD of many GPCRs with

an N_{exo} topology (N terminus in the ER lumen) (Chitwood et al., 2018). By contrast, we found

EMC-dependency within DENV NS4A-4B, a region with several TMDs, which are not close to the

termini of the polyprotein. This provides experimental evidence that the EMC is not only capable

of supporting stable expression of TMDs near the N and C termini but also of internal TMDs,

arguing that the EMC may directly facilitate the biogenesis of a wide variety of implicated TM

proteins beyond tail-anchored proteins and GPCRs (Shurtleff et al., 2018). Additionally, it is

noteworthy that the DENV polyprotein is distinct from human TM proteins in that it contains

multiple signal peptidase cleavage sites along the polyprotein, including at the 2K/NS4B site

between NS4A and NS4B, which removes the 2K peptide acting as signal sequence for NS4B

(Lin et al., 1993; Miller et al., 2006). Interestingly, it was shown (e.g. for ADRB1) that addition of

a signal sequence or an N_{cyt} TMD upstream of the first N_{exo} TMD makes the EMC dispensable for

biogenesis (Chitwood et al., 2018). This observation suggests that EMC-mediated

integration/stabilization of TMDs may be context-dependent. It needs to be elucidated whether

the EMC plays an active role in the TMD insertion of the DENV polyprotein into the ER membrane

or whether it functions as a post-insertional chaperone for specific TMDs.

Generally, for both cellular multi-pass TM proteins and the DENV polyprotein, only one or a subset

of TMDs seem to require the EMC for accurate topogenesis, while the other TMDs rely on the

Sec61 translocon. The exact interplay between Sec61 and the EMC during TM protein biogenesis

warrants further investigation. The simultaneous dependency of flaviviruses on both Sec61 and

EMC as highlighted by the CRISPR KO screens could allow an in-depth analysis of the sequence

and TMD features that may determine membrane insertion via Sec61 or EMC.

The A6665G mutation is predicted to be in a TM helix of NS4A but A7558T is not in a TM region

(Miller et al., 2006, 2007). Therefore, further studies are necessary to determine the exact

interaction between the EMC and the viral TMDs. Interestingly, the A6665G mutation also arose

during DENV adaptation to cell lines deficient in the oligosaccharyltransferase (OST) complex,

another critical host factor inside the ER membrane (Marceau et al., 2016; Puschnik et al., 2017).

The OST complex has been shown to be important for RNA genome replication in a catalytically

independent function and was suggested to act as structural scaffold for viral replication

complexes. It is possible that the tyrosine to cysteine substitution generally modulates ER

anchoring of NS4A. Overall, it remains to be elucidated how the resulting amino acid substitutions

rescue stable expression of NS4A and NS4B.

The isolation of EMC-independent DENV mutants additionally enables drug screening for EMC

modulators by testing compounds against WT DENV-Luc and then counter-screening hits with

EMC-adapted virus. EMC inhibitors may be a useful chemical tool to study EMC function and also

provide a potential antiviral therapeutic against flaviviruses. Host-directed antivirals may have the

advantage of a broader activity against multiple viruses and a higher barrier of drug resistance.

Only 50% of the adaptation experiments yielded escape mutants containing two mutations after

>20 passages. For comparison, for direct-acting antivirals (e.g. viral polymerase inhibitors), one

point mutation is often sufficient to overcome the mechanism of action (Beaucourt and Vignuzzi,

2014).

Lastly, the EMC was originally discovered to be associated with the UPR (Jonikas et al., 2009),

and upregulation of UPR during flavivirus infection has been reported (Peña and Harris, 2011;

Reid et al., 2018; Su et al., 2002). It is possible that competition between viral polyproteins and

cellular proteins for the machinery important for proper TM protein expression (such as EMC,

OST complex and ERAD components) is linked to this phenomenon. The induction of ER stress

can further trigger autophagy, which has also been shown to be beneficial for flavivirus infection

(Abernathy et al., 2019; Lee et al., 2018; Senft and Ronai, 2015). Therefore, hijacking the EMC

may have direct and indirect implications for flavivirus replication.

Overall, our study gives detailed insights into how flaviviruses exploit the EMC to support proper

polyprotein expression, and we provide further evidence for the cellular function of the EMC in

TM protein topogenesis by utilizing DENV as a cell biological tool.

Acknowledgments

We like to thank the following people: Jan Carette for discussions, providing reagents and critically

reading the manuscript; Caleb Marceau for technical advice and providing reagents; members of

the Biohub Infectious Disease Initiative and the Weissman lab for discussions and technical

advice; the Biohub Genomics Platform for help with sequencing. ASP was supported by the Chan

Zuckerberg Biohub. MJS is a Howard Hughes Medical Institute Fellow of the Helen Hay Whitney

Foundation and JSW is an Investigator of the Howard Hughes Medical Institute.

Author contributions

ASP and JSW conceptualized study. AMN and ASP conducted the majority of experiments with help from JK. MJS and KDP performed ribosome profiling. ASP wrote the manuscript with input from all authors.

Figure 1

Figure 1: EMC is required for flavivirus infection

(A) Luminescence of WT and EMC subunit KO Huh7.5.1 cells infected with DENV-Luc at 30hpi.

Dotted line is background from uninfected control.

(B) Luminescence of WT and EMC subunit KO HEK293FT cells infected with DENV-Luc at 48hpi.

Dotted line is background from uninfected control.

(C) Replication of DENV-Luc in WT, EMC2- and EMC4-KO, and cDNA complemented KO HEK293FT cells.

(D) Quantitative RT-PCR of DENV, WNV, ZIKV and HCV RNA in WT or EMC4-KO Huh7.5.1 cells.

In all figures, values are shown as mean of three biological replicates with standard deviation in (A)-(C) and mean with standard error of the mean in (D). *t*-tests were performed to determine statistical significance and *p*-values are defined as ns=non-significant, *<0.05, **<0.01, ***<0.001.

Figure 2

Figure 2: EMC is important for optimal viral protein expression

(A) Quantitative RT-PCR of internalized DENV RNA in WT and EMC4-KO Huh7.5.1 cells at different timepoints post-infection. The data is from three biological replicates and shown as mean with standard error of the mean.

(B) Luminescence of WT and EMC4-KO HEK293FT cells, which were electroporated with DENV replicon expressing Renilla luciferase and collected at different times post-electroporation. The data is from three biological replicates for each timepoint and shown as mean with standard deviation.

(C) Immunoblot analysis of DENV infected WT and EMC4-KO HEK293FT cells at different timepoints post-infection (6/8/10/12h). Lysates were blotted for different viral proteins (Capsid, NS2B, NS3 and NS5) and GAPDH was used as loading control. UI=uninfected control; L=ladder. Quantification of bands is shown in Figure S2B.

Figure 3

349 Figure 3: EMC knockout does not largely affect viral translation efficiency but leads to

350 post-translational protein degradation

(A) Ribosome Profiling (Ribo-seq) and RNA-seq were performed in uninfected WT and EMC4-

KO HEK293FT cells to measure changes in translation efficiency (TE). Log$_2$ fold-change (FC) of

reads per kb of transcript, per million mapped reads (RPKMs) between EMC4-KO and WT cells

for Ribosome Profiling and RNA-seq are displayed on the y- and x-axis, respectively. Each dot

represents one RNA transcript. Fold-change in TE for a given transcript is defined as the change

in Ribosome Profiling RPKMs normalized to the change in RNA-seq RPKMs. The red line

represents fold-change of TE=1.

(B) Ribosome Profiling (Ribo-seq) and RNA-seq were performed in DENV-infected WT and

EMC4-KO HEK293FT cells 18h post-infection to measure changes in translation efficiency (TE).

The DENV transcript is highlighted in red. The red line represents fold-change of TE=1.

(C) Ribosome Profiling (Ribo-seq) and RNA-seq were performed in DENV-infected WT and

EMC4-KO HEK293FT cells 44h post-infection to measure changes in translation efficiency (TE).

The DENV transcript is highlighted in red. The red line represents fold-change of TE=1.

(D) Immunoblot analysis of DENV infected WT and EMC4-KO HEK293FT cells with or without

addition of bortezomib (50nM), a proteasome inhibitor at 18hpi. GAPDH was used as loading

control. UI=uninfected.

Figure 4

A

DENV polyprotein

C-E(TMD1)-G2C
NS1-3-G2C
NS3-4A-G2C
NS2B-4B-G2C
NS4A-4B-G2C
NS4A-5-G2C
NS4B-5-G2C

eGFP-2A-mCherry (G2C)

B

Figure 4: EMC is required for proper biogenesis of the NS4A-NS4B region of the DENV

polyprotein

(A) Overview of dual-fluorescence reporter constructs of DENV polyprotein for analysis of protein

stability. Segments of the DENV genome encoding parts of the polyprotein were cloned with a C-

terminal fusion to eGFP-T2A-mCherry (G2C). The T2A peptide leads to peptide bond skipping

during translation resulting in two protein products, DENV protein fused to eGFP and mCherry.

The GFP:mCherry fluorescence ratio reflects changes in DENV protein stability.

(B) Flow cytometry measurements of GFP:mCherry fluorescence ratio for G2C constructs in WT

and EMC4-KO Huh7.5.1 cells. The fluorescence ratios are depicted as histograms. ADRB1=β1-

adrenergic receptor.

(C) Measurement of GFP:mCherry fluorescence ratio of NS4A-4B-G2C construct in WT, EMC4-

KO and EMC4 cDNA complemented EMC4-KO HEK293FT cells.

(D) Measurement of GFP:mCherry fluorescence ratio of NS4A-4B-G2C construct in WT, EMC4-

KO and bortezomib-treated EMC4-KO HEK293FT cells. Note that (C) and (D) use same WT and

EMC4-KO data as experiments were performed in parallel.

Figure 5: Adaptation of DENV to EMC KO cells reveals non-synonymous point mutations in NS4A and NS4B

(A) Identification of SNPs in EMC4-KO adapted DENV relative to DENV 16681 reference genome by next-generation sequencing of supernatant-extracted viral RNA. Mutations common across all three replicates are highlighted by red boxes and their nucleotide and amino acid changes are shown below. A table of all identified SNPs can be found in Figure S5.

(B) Quantitative RT-PCR of DENV RNA from WT DENV 16681 and the three isolated EMC-adapted DENV in WT and EMC4-KO Huh7.5.1. The data is from three biological replicates and shown as mean with standard error of the mean.

(C) Viral infection assay in WT and EMC4 KO Huh7.5.1 using WT DENV-Luc or mutant DENV-Luc containing A6665G and A7558T either individually or in tandem. The data is from three biological replicates and shown as mean with standard deviation.

(D) Measurement of GFP:mCherry fluorescence ratio of WT NS4A-5-G2C and NS4A-5(A6665G,A7558T)-G2C constructs in WT and EMC4-KO Huh7.5.1.

Figure 6.: EMC physically interacts with DENV NS4B post-cleavage and absence of EMC

leads to rapid degradation of misfolded NS4B.

(A) Immunoblot for processed products of transfected NS2B-4B-GFP construct in WT and EMC4

KO cells in presence or absence of BZ. p84 was used as loading control.

(B) Co-immunoprecipitation of EMC4-FLAG from lysates of uninfected (UI) or DENV infected

HEK293FT EMC4-FLAG cells followed by immunoblotting against viral proteins (prM, NS4B,

NS5). p84 was used as loading and non-specific binding control.

(C) Model for the role of EMC in DENV infection. In DENV infected wild-type cells, the EMC co-

translationally engages with NS4A and/or NS4B TMDs to facilitate correct/stable insertion

ensuring a functional topology. In the absence of the EMC, misfolding of the NS4A-NS4B region

occurs during polyprotein synthesis resulting in degradation of viral proteins.

Figure S1

Figure S1: EMC is required for flavivirus infection

(A) Sanger sequencing of clonal EMC subunit KO Huh7.5.1 and HEK293FT cells. sgRNA

targeting site is highlighted in reference sequence (top) and indel mutation is shown below

sequencing trace of KO cells. For heterozygous mutants, TIDE was used to determine indel

mutations.

(B) Quantitative RT-PCR of DENV RNA in WT or EMC2-KO Huh7.5.1 at 30, 48 and 72h post

infection. Values are shown relative to the WT, 30h condition.

(C) Immunoblot analysis of EMC2 or EMC4 expression in WT, EMC2- and EMC4-KO, and cDNA

complemented KO HEK293FT cells. p84 was used as loading control.

(D) Quantitative RT-PCR of DENV, WNV and ZIKV RNA in WT or EMC4-KO HEK293FT cells.

For QPCR experiments, values are shown as mean of three biological replicates with standard

error of the mean.

Figure S2

A

WT (DMSO)
WT (MK0608)
EMC4-KO (DMSO)
EMC4-KO (MK0608)

B

C

D

Figure S2: EMC is important for optimal viral protein expression

(A) Luminescence of WT and EMC4-KO cells electroporated with DENV replicon expressing

Renilla luciferase and treated with MK-0608 (a nucleoside analogue, which inhibits the DENV

polymerase) or DMSO. Lysates were collected at different times post electroporation. The data is

from three biological replicates and shown as mean with standard deviation.

(B) Quantification of viral protein expression in EMC4-KO cells relative to WT cells by

densitometry. Bands of viral proteins from Figure 2C were quantified and normalized to GAPDH

bands at each timepoint.

(C) Immunoblot analysis of DENV infected WT and EMC4-KO HEK293FT cells at different

timepoints post-infection (6/8/10/12h) with treatment of MK-0608 to inhibit viral replication.

Lysates were blotted for different viral proteins (Capsid, NS2B, NS3 and NS5) and GAPDH was

used as loading control. UI=uninfected control; L=ladder. Quantification of bands is shown in

Figure S2D.

(D) Quantification of viral protein expression in EMC4-KO cells relative to WT cells by

densitometry. Bands of viral proteins from Figure S2C were quantified and normalized to GAPDH

bands at each timepoint.

Figure S3

Figure S3: EMC knockout does not largely affect viral translation efficiency but leads to

post-translational protein degradation

(A) Translation efficiencies (ribosome densities per RNA molecule) of cellular and DENV mRNA

in WT and EMC4-KO cells at 18 and 44hpi. DENV is highlighted as red dot.

(B) Footprint distribution of Ribosome Profiling reads mapped to the DENV genome in WT and

EMC4-KO cells at 18 and 44hpi. Nucleotide position of DENV genome is indicated on x-axis.

(C) DENV-Luc infection of WT and EMC4 KO HEK293FT cells treated with DMSO control or

different concentrations of BZ for 24h. Dotted line indicates background signal from uninfected

cells.

Figure S4

Figure S4: EMC is required for proper biogenesis of the NS4A-NS4B region of the DENV polyprotein

Flow cytometry measurements of GFP:mCherry fluorescence ratio for G2C constructs in WT and EMC4-KO HEK293FT cells. The fluorescence ratios are depicted as histograms. ADRB1=β_1-adrenergic receptor.

Figure S5

Virus isolate	Position	nt change	Coverage	Variant frequency	Reference frequency	aa change
EMC adapted DENV #3	226	T -> C	9754	99.80%	0.10%	synonymous
EMC adapted DENV #1	725	T -> C	5653	99.70%	0.10%	Val->Ala
EMC adapted DENV #1	1531	C -> T	3686	98.40%	0.80%	synonymous
EMC adapted DENV #2	1531	C -> T	9399	99.90%	0.10%	synonymous
EMC adapted DENV #3	1531	C -> T	14698	99.50%	0.20%	synonymous
EMC adapted DENV #1	1818	C -> A	3366	99.00%	0.20%	synonymous
EMC adapted DENV #3	4018	C -> T	16149	94.50%	5.50%	synonymous
EMC adapted DENV #2	5493	A -> G	10626	99.00%	0.50%	synonymous
EMC adapted DENV #1	6665	A -> G	3172	99.70%	0.10%	Tyr->Cys
EMC adapted DENV #2	6665	A -> G	8932	99.40%	0.50%	Tyr->Cys
EMC adapted DENV #3	6665	A -> G	13319	99.50%	0.20%	Tyr->Cys
EMC adapted DENV #1	7558	A -> T	4205	98.40%	0.40%	Asn->Tyr
EMC adapted DENV #2	7558	A -> T	10497	99.40%	0.10%	Asn->Tyr
EMC adapted DENV #3	7558	A -> T	16401	99.00%	0.04%	Asn->Tyr
EMC adapted DENV #3	7581	C -> T	16867	94.20%	5.70%	synonymous
EMC adapted DENV #3	8546	G -> A	15538	94.80%	5.10%	Arg->Lys

467 **Figure S5: Adaptation of DENV to EMC-KO cells reveals conserved point mutations in**

468 **NS4A and NS4B**

469 Table containing all identified SNPs from the three isolated EMC-adapted viruses. A minimum

470 variant frequency cutoff of 80% was used.

Material and methods

Cell culture and viruses

Huh7.5.1 (gift from Frank Chisari) and HEK293FT (Thermo Scientific) were cultured in DMEM (Gibco) supplemented with 10% fetal bovine serum (Omega Scientific), penicillin/streptomycin (Gibco), non-essential amino acids (Gibco) and L-glutamine (Gibco) at 37C and 5% CO_2. Parental cell lines were tested negative for mycoplasma. The following virus strains were used: DENV serotype 2 strain 16681 (stocks kindly provided by Caleb Marceau), DENV-Luc (stocks kindly provided by Jan Carette) (Marceau et al., 2016), WNV Kunjin strain CH16532 (stocks kindly provided by Jan Carette), ZIKV PRVABC59 (stocks kindly provided by Jan Carette) and HCV JFH1 (stocks kindly provided by Jan Carette). Viral titers were determined by plaque- or focus-forming assay.

Chemical compounds

MK-0608 (2'-C-Methylcytidine) was purchased from Carbosynth and used at a concentration of 25 µM. Bortezomib was obtained from Selleckchem and used at concentrations 1-50 nM. For all experiments, DMSO controls were included. Cycloheximide was purchased from Sigma and used at 100 µg/ml for Ribosome Profiling experiments.

Generation of knockout cell lines

Oligos containing sgRNA sequence were annealed and ligated into pX458 (gift from Feng Zhang; Addgene plasmid # 48138). Cells were transfected with pX458 using Lipofectamine 3000 (Invitrogen) and two days later GFP positive cells were single-cell sorted into 96-well plates using a Sony SH800 cell sorter. For genotyping, genomic DNA was isolated from obtained clones using QuickExtract (Lucigen), the sgRNA-targeted sites PCR amplified and the products Sanger-sequenced. Obtained sequences were compared to reference sequences for indel mutations. In the case of heterozygous mutations, TIDE was used to deconvolute sequencing traces (Brinkman

et al., 2014). A list of all used sgRNA oligo and genotyping primer sequences can be found in

Supplementary Table 2.

Dengue Luciferase assay

Huh7.5.1 or HEK293FT cells were seeded in 96-well plates and infected with DENV-Luc at an

moi of 0.01. Cells were harvested in Renilla Lysis buffer and luminescence was measured using

Renilla Luciferase Assay system (Promega) on a Spectramax i3x Multi-Mode Microplate Reader

(Molecular Devices).

Quantitative RT-PCR

Huh7.5.1 or HEK293FT cells were plated in 96-well plates (in triplicates for each condition) and

infected with an moi of 0.5 for all viruses. Cells were then harvested at 30hpi for ZIKV, 48hpi for

DENV and WNV, and 72hpi for HCV using the Power SYBR Cells-to-CT kit (Invitrogen). After

reverse transcription, quantitative PCR was performed on a Bio-Rad CFX96 Touch system. All

viral RNA levels were normalized to 18S levels. For viral internalization assay, 96-well plates

containing WT or EMC4 KO Huh7.5.1 cells were put on ice before infection with DENV at an moi

of 30. Cells were incubated for 1h before moving them to a 37C incubator. At each harvest

timepoint, cells were washed three times with PBS and lysed as described above. All qPCR primer

sequences can be found in Supplementary Table 2.

Lentiviral complementation

EMC2 (GenScript; clone OHu30604) and EMC4 cDNA (GenScript; clone OHu00964) were PCR

amplified   using   tgtggtggaattctgcagataccATGGCGAAGGTCTCAGA   /

CGGCCGCCACTGTGCTGGATTTACTTATCGTCGTCATCCTTGTAATCAGACTGGGTGATC

and   tgtggtggaattctgcagataccATGACGGCCCAGGGG   /

CGGCCGCCACTGTGCTGGATTTACTTATCGTCGTCATCCTTGTAATCCAAAAGCAGTCCT,

respectively, and cloned into EcoRV-cut pLenti CMV Puro DEST (w118-1) (gift from Eric

Campeau & Paul Kaufman; Addgene plasmid # 17452) using NEBuilder® HiFi DNA Assembly

Master Mix (New England BioLabs). Lentivirus was produced in HEK293FT, respective KO cells

transduced with filtered, lentiviral containing supernatant and selected using 2-4 µg/ml puromycin

(Gibco) for 3-4 days.

Replicon assay

DENV replicon plasmid (Marceau et al., 2016) was linearized using XbaI restriction enzyme.

Replicon RNA was generated using the MEGAscript T7 Kit (Invitrogen) with the reaction

containing 5mM m7G(5')ppp(5')G RNA Cap Structure Analog (New England BioLabs). Resulting

RNA was purified by lithium chloride precipitation. For electroporation, 2 million WT or EMC4 KO

HEK293FT cells were washed twice in PBS, resuspended in 100 µl SF Nucleofector solution

(Lonza), mixed with 4 µg replicon RNA, transferred to a 100ul nucleocuvette and pulsed using

program CM-130 on an Amaxa 4D-Nucleofector™ X Unit (Lonza). Cells were then resuspended

in antibiotic-free medium, distributed into 96-wells and lysed at different timepoints post-

electroporation using Renilla Lysis buffer. Luminescence was measured using Renilla Luciferase

Assay system (Promega) on a Spectramax i3x Multi-Mode Microplate Reader (Molecular

Devices).

Immunoblotting

Cell were lysed using Laemmli SDS sample buffer containing 5% beta-mercaptoethanol and

boiled for 10min. Lysates were separated by SDS-PAGE on pre-cast Bio-Rad 4-15% poly-

acrylamide gels in Bio-Rad Mini-Protean electrophoresis system. Proteins were transferred onto

PVDF membranes using Bio-Rad Trans-Blot Turbo transfer system. PVDF membranes were

blocked with PBS buffer containing 0.1% Tween-20 and 5% non-fat milk. Blocked membranes

were incubated with primary antibody diluted in blocking buffer and incubated overnight at 4C on

a shaker. Primary antibodies were detected by incubating membranes with 1:10000 dilution of

HRP-conjugated (Southern Biotech) or IRDye-conjugated (LI-COR) secondary anti-mouse and

anti-rabbit antibodies for 1 h at room temperature. Blots were visualized using a ChemiDoc™ MP

Imaging System (Bio-Rad). The following primary antibodies (and their dilutions) were used in

this study: p84 (Genetex, GTX70220) at 1:2000, GAPDH (SCBT, sc-32233) at 1:2000,

EMC2/TTC35 (Proteintech, 25443-1-AP) at 1:750, EMC4 (Bethyl, A305-752A) at 1:500, Dengue

virus prM protein (Genetex, GTX128093) at 1:1000, Dengue virus Capsid protein (Genetex,

GTX103343) at 1:1000, Dengue virus NS2B (Genetex, GTX124246) at 1:1000, Dengue virus NS3

protein (Genetex, GTX629477) at 1:1000, Dengue virus NS4B protein (Genetex, GTX124250) at

1:1000, Dengue virus Type 2 NS5 protein (Genetex, GTX629447) and GFP (Genetex,

GTX113617) at 1:5000.

Ribosome profiling

WT and EMC4 KO HEK293FT were plated in 10cm dishes and infected with DENV at an of 100

for the 18hpi harvest and an moi of 0.5 for the 44hpi harvest. Cells were treated with 100 µg/ml

cycloheximide for 2 min, washed once with polysome gradient buffer (20 mM Tris pH 7.5, 150

mM NaCl, 5 mM MgCl$_2$, 100 µg/mL cycloheximide, 1 mM DTT) and then lysed using 500 µl of

polysome lysis buffer (20 mM Tris pH 7.5, 150 mM NaCl, 5 MgCl$_2$, 1% Triton X-100, 1 mM DTT,

100 µg/mL cycloheximide, 24 U/ml Turbo DNase (Ambion)) per plate. Cells/lysates were collected

using cell scrapers and incubated on ice for 15min. Crude lysates were centrifuged at 20,000 x g

for 2 min at 4C and the cleared polysome-containing lysates flash frozen by immersion in liquid

nitrogen and stored at -80C until further processing. For ribosomal footprinting, 30 µg total RNA

diluted in polysome buffer to a total volume of 200ul were treated with 750 U/ml RNaseI (Ambion)

for 45min at 25C with shaking (300rpm). RNase activity was quenched by adding 10ul

SuperaseIN (Invitrogen) and placing samples on ice. Note that 10% of the lysates were saved for

RNA-seq library preparation (see below). A 1ml sucrose cushion was added to the bottom of a

13mm x 56mm thick-wall polycarbonate tube (Beckman Coulter) for each sample and the footprint

digests were layered on top. Tubes were ultracentrifuged at 100,000 rpm for 4h at 4C in a

TLA100.3 rotor (Beckman Coulter). After spin, supernatant was removed and the monosome-

containing pellet was resuspended in 300 µl Trizol LS (Life Technologies) for RNA extraction

using the Direct-Zol kit (Zymo). Ribosome Profiling NGS libraries were generated as previously

described (Shurtleff et al., 2018).

For corresponding RNA-seq libraries RNA was extracted from the remaining undigested cleared

lysates using Trizol and the Direct-Zol kit. 1 µg purified total RNA was used as input for the Truseq

stranded total RNA Gold library prep (Illumina). Ribosomal RNA was depleted using Ribo-Zero

H/M/R Gold and depleted RNA fragmented for 8min at 94C. Following first and second strand

synthesis products were purified using SPRI beads. KAPA dual-indexed adapters (Kapa

Biosystems, KK8722) were used for adapter ligation. Finally, PCR enriched products were

cleaned up using SPRI. Ribosome Profiling and RNA-seq libraries were analyzed on Bioanalyzer

or TapeStation (Agilent) before sequencing on a NextSeq 500 platform (Illumina) in separate runs

with 400M reads output.

Analysis of Ribosome Profiling data

Reads from Ribosome Profiling or RNA-seq were preprocessed by removing low-quality reads

and trimming of adaptor sequences. Next, ribosomal RNA reads were removed using Bowtie.

Remaining reads were aligned to the human reference genome file also containing the genomic

DENV2 16681 sequence using Tophat to generate read count tables with RPKMs for each

transcript. A cutoff of at least 5 RPKMs was used for further analysis of transcripts/footprint counts,

and $\log_2$ fold-changes (FC) were calculated between corresponding EMC4 KO and WT samples

for both Ribosome Profiling and RNA-seq. To calculate translation efficiency (TE), the Ribosome

Profiling RPKMs were normalized by the RNA-seq RPKMs for each transcript.

To analyze the footprint distribution of Ribosome Profiling reads across the DENV genome, DENV

specific reads were mapped to the DENV2 16681 reference sequence using Geneious Prime and

coverage at each nucleotide position was determined.

Dual fluorescence cytometry assay

eGFP was amplified from pX458 with adding overhang to plenti-CMV-Puro (EcoRV cut site), a

Kozak sequence and ATG start codon for the G2C-only construct or with an additional overhang

to GlySer-linker sequence for fusion with DENV fragments. T2A-mCherry was amplified from

pX458 where eGFP had been replaced with mCherry. Different regions of the DENV genome

were amplified from the DENV2 16681 infectious clone plasmid (gift from Karla Kirkegaard) with

addition of overhang sequences to plenti-CMV backbone, Kozak sequence and ATG start codon

on 5'end and to GlySer-linker-GFP fragment on 3'end. ADRB1 was amplified from pcDNA3 Flag

beta-1-adrenergic-receptor (gift from Robert Lefkowitz; Addgene plasmid # 14698). All fragments

contained ~20bp overhangs and were assembled into EcoRV cut pLenti CMV Puro DEST (w118-

1) using NEBuilder® HiFi DNA Assembly Master Mix (New England BioLabs) or Gibson-like

master mix (Berkeley MacroLabs) to generate G2C reporter expression constructs. All constructs

were verified by Sanger sequencing. Primer sequences can be found in Supplementary Table 2.

For the G2C dual fluorescence assay, cells were seeded in 24- or 12-well plates and transfected

with G2C constructs using Lipofectamine 3000 (Invitrogen). 24-48h later cells were trypsinized,

filtered (70µm), and analyzed by flow cytometry using a Cytoflex S flow cytometer (Beckman

Coulter). At least 5,000 and 10,000 mCherry positive cells were recorded for Huh7.5.1 and

HEK293FT, respectively. For analysis, cells were gated based on FSC/SSC, FSC-H/FSC-A

(singlets) and finally ECD (mCherry) and FITC (eGFP) using FlowJo 10. Raw fluorescence data

for all gated fluorescent cells was exported and the eGFP:mCherry intensity ratios were calculated

for individual cells. All fluorescence ratios were normalized to the mean of the G2C-only construct.

Distributions of fluorescence ratios were plotted as histograms using GraphPad Prism 8.

Viral adaptation

WT DENV serotype 2 strain 16681 was serially passaged every 3-4 days onto newly plated EMC4

KO Huh7.5.1 cells by transferring increasingly diluted supernatant from infected cells (starting at

2-fold dilutions for the first 3 passages, then stepwise increase of dilutions to 100-fold for the final

passages). Six independent passaging experiments were started and cytopathic effects were

observed in three of the replicates after approximately 20-22 passages. Supernatant was then

harvested, filtered through a 0.45 um filter to remove cellular debris, and RNA was extracted using

Trizol LS and Direct-Zol kit (Zymo). Additionally, RNA was extracted from WT DENV stock. To

prepare sequencing libraries, Truseq stranded total RNA Gold library kit (Illumina) was used.

Isolated RNA was fragmented for 4min at 94C without prior Ribo-Zero depletion, followed by first

and second strand synthesis, and adapter ligation. Fragments were amplified in presence of

SYBR Green on Bio-Rad CFX96 and reaction was stopped for each sample before it reached

plateau (~20-24 cycles). Libraries from different virus isolates were pooled and sequenced on

iSeq 100 (Illumina) with 2x75bp reads.

For analysis, Geneious Prime was used: Fastq files were imported as paired reads and Truseq

adapters and low-quality sequences (minimum 20) were trimmed using BBDuk. Next, reads were

mapped to the WT 16681 reference genome using the Geneious Mapper at medium-low

sensitivity and with 5 iterations for fine-tuning. All samples had a minimum coverage of 1000 at

all nucleotide positions. Finally, SNPs between adapted and WT stock viruses were called with a

minimum variance frequency of 80%. To compare acquired non-synonymous point mutations to

a large dataset of sequenced DENV2 strains, 1,359 complete genome sequences from NCBI

were mapped to the DENV2 16681 strain containing adapted mutations.

Generation of EMC adapted DENV-Luciferase

DENV-Luc infectious clone containing adapted mutations were generated by PCR amplification

from WT DENV-Luc infectious clone plasmid (gift from Jan Carette). For DENV-Luc(A6665G)

fragments were amplified using GAAGAGTGATGGTTATGGTAGGC /

CTGTATTTGTGCGCACCATAGGAGGATG and CATCCTCCTATGGTGCGCACAAATACAG /

GTGATCTTCATTTAAGAATCCTAGGGCTTC. For DENV-Luc(A7558T) fragments were

amplified using GAAGAGTGATGGTTATGGTAGGC / CCCCTTCTTGTGTAGGTTGTGTTCTTC

and GAAGAACACAACCTACACAAGAAGGGG / GTGATCTTCATTTAAGAATCCTAGGGCTTC.

For DENV-Luc(A6665G, A7558T) fragments were amplified using

GAAGAGTGATGGTTATGGTAGGC / CTGTATTTGTGCGCACCATAGGAGGATG,

CATCCTCCTATGGTGCGCACAAATACAG / CCCCTTCTTGTGTAGGTTGTGTTCTTC, and

GAAGAACACAACCTACACAAGAAGGGG / GTGATCTTCATTTAAGAATCCTAGGGCTTC.

PCR products were cloned into NarI and AvrII cut WT DENV-Luc infectious clone using

NEBuilder® HiFi DNA Assembly Master Mix (New England BioLabs) and transformed into Stbl3

bacteria (Berkeley MacroLabs). Constructs were verified by Sanger sequencing. Plasmids were

linearized with XbaI and RNA was generated by in-vitro transcription using the MEGAscript T7 Kit

(Invitrogen) with the reaction containing 5mM m7G(5')ppp(5')G RNA Cap Structure Analog (New

England BioLabs). Resulting RNA was purified by lithium chloride precipitation and transfected

into BHK-21 cells using Lipofectamine 3000 for viral production. Supernatants were harvested 6-

9 days post-transfection.

Co-immunoprecipitation

HEK293FT EMC4 KO cells complemented with FLAG-tagged EMC4 were seeded in a 10cm dish

and infected with DENV for 96h. Cells were washed with PBS and then collected in 1ml Pierce

RIPA buffer (Thermo Scientific) containing 1mM PMSF (Thermo Scientific), 5mM EDTA and 1x

Halt Protease Inhibitor Cocktail (Thermo Scientific) with help of cell scrapers. Cells were lysed on

ice for 1h with occasional vortexing and subsequently centrifuged at 15,000 x g for 10min at 4C

to clear lysates. For co-immunoprecipitation, clarified lysates were incubated at 4C overnight with

Anti-FLAG® M2 Magnetic Beads (Sigma) while rotating. Beads were washed three times using

TBS-T and eluted by boiling in 2x Laemmli buffer. Eluates were loaded for SDS-PAGE and stained

for viral proteins.

Statistical analysis

GraphPad Prism 8 was used to analyze data and perform statistical tests. Unpaired *t*-tests were

applied to measure statistical significance of differences in DENV infection between WT and EMC

KO cells.

Supplementary tables

Supplementary Table 1

Ribosome Profiling data including read counts and fold change for Ribosome Profiling and RNA-

seq used to generate Figures 3A-C.

Supplementary Table 2

Table of oligos used in this study: sgRNA oligos, genotyping primers, qPCR primers, primers to

generate G2C constructs

References

Abernathy, E., Mateo, R., Majzoub, K., van Buuren, N., Bird, S.W., Carette, J.E., and
Kirkegaard, K. (2019). Differential and convergent utilization of autophagy components by
positive-strand RNA viruses. PLoS Biol. *17*, e2006926.

Beaucourt, S., and Vignuzzi, M. (2014). Ribavirin: a drug active against many viruses with
multiple effects on virus replication and propagation. Molecular basis of ribavirin resistance.
Curr. Opin. Virol. *8*, 10–15.

Brinkman, E.K., Chen, T., Amendola, M., and van Steensel, B. (2014). Easy quantitative
assessment of genome editing by sequence trace decomposition. Nucleic Acids Res. *42*, e168–
e168.

Chen, Y.-L., Yokokawa, F., and Shi, P.-Y. (2015). The search for nucleoside/nucleotide analog
inhibitors of dengue virus. Antiviral Res. *122*, 12–19.

Chitwood, P.J., and Hegde, R.S. (2019). The Role of EMC during Membrane Protein
Biogenesis. Trends Cell Biol.

Chitwood, P.J., Juszkiewicz, S., Guna, A., Shao, S., and Hegde, R.S. (2018). EMC Is Required
to Initiate Accurate Membrane Protein Topogenesis. Cell *175*, 1507-1519.e16.

Christianson, J.C., Olzmann, J.A., Shaler, T.A., Sowa, M.E., Bennett, E.J., Richter, C.M., Tyler,
R.E., Greenblatt, E.J., Harper, J.W., and Kopito, R.R. (2011). Defining human ERAD networks
through an integrative mapping strategy. Nat. Cell Biol. *14*, 93–105.

Coelho, J.P.L., Stahl, M., Bloemeke, N., Meighen-Berger, K., Alvira, C.P., Zhang, Z.-R., Sieber,
S.A., and Feige, M.J. (2019). A network of chaperones prevents and detects failures in
membrane protein lipid bilayer integration. Nat. Commun. *10*.

Guna, A., Volkmar, N., Christianson, J.C., and Hegde, R.S. (2018). The ER membrane protein
complex is a transmembrane domain insertase. Science *359*, 470–473.

Ingolia, N.T., Ghaemmaghami, S., Newman, J.R.S., and Weissman, J.S. (2009). Genome-Wide
Analysis in Vivo of Translation with Nucleotide Resolution Using Ribosome Profiling. Science
324, 218–223.

Jonikas, M.C., Collins, S.R., Denic, V., Oh, E., Quan, E.M., Schmid, V., Weibezahn, J.,
Schwappach, B., Walter, P., Weissman, J.S., et al. (2009). Comprehensive characterization of
genes required for protein folding in the endoplasmic reticulum. Science *323*, 1693–1697.

Krishnan, M.N., Ng, A., Sukumaran, B., Gilfoy, F.D., Uchil, P.D., Sultana, H., Brass, A.L.,
Adametz, R., Tsui, M., Qian, F., et al. (2008). RNA interference screen for human genes
associated with West Nile virus infection. Nature *455*, 242–245.

Lee, Y.-R., Kuo, S.-H., Lin, C.-Y., Fu, P.-J., Lin, Y.-S., Yeh, T.-M., and Liu, H.-S. (2018). Dengue
virus-induced ER stress is required for autophagy activation, viral replication, and pathogenesis
both in vitro and in vivo. Sci. Rep. *8*, 489.

Lin, C., Amberg, S.M., Chambers, T.J., and Rice, C.M. (1993). Cleavage at a novel site in the
NS4A region by the yellow fever virus NS2B-3 proteinase is a prerequisite for processing at the
downstream 4A/4B signalase site. J. Virol. *67*, 2327–2335.

Ma, H., Dang, Y., Wu, Y., Jia, G., Anaya, E., Zhang, J., Abraham, S., Choi, J.-G., Shi, G., Qi, L.,
et al. (2015). A CRISPR-Based Screen Identifies Genes Essential for West-Nile-Virus-Induced
Cell Death. Cell Rep. *12*, 673–683.

Marceau, C.D., Puschnik, A.S., Majzoub, K., Ooi, Y.S., Brewer, S.M., Fuchs, G., Swaminathan,
K., Mata, M.A., Elias, J.E., Sarnow, P., et al. (2016). Genetic dissection of Flaviviridae host
factors through genome-scale CRISPR screens. Nature *535*, 159–163.

Miller, S., Sparacio, S., and Bartenschlager, R. (2006). Subcellular Localization and Membrane
Topology of the Dengue Virus Type 2 Non-structural Protein 4B. J. Biol. Chem. *281*, 8854–
8863.

Miller, S., Kastner, S., Krijnse-Locker, J., Bühler, S., and Bartenschlager, R. (2007). The Non-
structural Protein 4A of Dengue Virus Is an Integral Membrane Protein Inducing Membrane
Alterations in a 2K-regulated Manner. J. Biol. Chem. *282*, 8873–8882.

Muñoz-Jordán, J.L., Laurent-Rolle, M., Ashour, J., Martínez-Sobrido, L., Ashok, M., Lipkin, W.I.,
and García-Sastre, A. (2005). Inhibition of alpha/beta interferon signaling by the NS4B protein of
flaviviruses. J. Virol. *79*, 8004–8013.

Neufeldt, C.J., Cortese, M., Acosta, E.G., and Bartenschlager, R. (2018). Rewiring cellular
networks by members of the Flaviviridae family. Nat. Rev. Microbiol. *16*, 125–142.

Olzmann, J.A., Kopito, R.R., and Christianson, J.C. (2013). The Mammalian Endoplasmic
Reticulum-Associated Degradation System. Cold Spring Harb. Perspect. Biol. *5*.

Peña, J., and Harris, E. (2011). Dengue virus modulates the unfolded protein response in a
time-dependent manner. J. Biol. Chem. *286*, 14226–14236.

Puschnik, A.S., Marceau, C.D., Ooi, Y.S., Majzoub, K., Rinis, N., Contessa, J.N., and Carette,
J.E. (2017). A Small-Molecule Oligosaccharyltransferase Inhibitor with Pan-flaviviral Activity.
Cell Rep. *21*, 3032–3039.

Reid, D.W., Campos, R.K., Child, J.R., Zheng, T., Chan, K.W.K., Bradrick, S.S., Vasudevan,
S.G., Garcia-Blanco, M.A., and Nicchitta, C.V. (2018). Dengue Virus Selectively Annexes
Endoplasmic Reticulum-Associated Translation Machinery as a Strategy for Co-opting Host Cell
Protein Synthesis. J. Virol. *92*, e01766-17.

Roosendaal, J., Westaway, E.G., Khromykh, A., and Mackenzie, J.M. (2006). Regulated
cleavages at the West Nile virus NS4A-2K-NS4B junctions play a major role in rearranging
cytoplasmic membranes and Golgi trafficking of the NS4A protein. J. Virol. *80*, 4623–4632.

Savidis, G., McDougall, W.M., Meraner, P., Perreira, J.M., Portmann, J.M., Trincucci, G., John,
S.P., Aker, A.M., Renzette, N., Robbins, D.R., et al. (2016). Identification of Zika Virus and
Dengue Virus Dependency Factors using Functional Genomics. Cell Rep. *16*, 232–246.

Senft, D., and Ronai, Z.A. (2015). UPR, autophagy, and mitochondria crosstalk underlies the
ER stress response. Trends Biochem. Sci. *40*, 141–148.

Sessions, O.M., Barrows, N.J., Souza-Neto, J.A., Robinson, T.J., Hershey, C.L., Rodgers, M.A.,
Ramirez, J.L., Dimopoulos, G., Yang, P.L., Pearson, J.L., et al. (2009). Discovery of insect and
human dengue virus host factors. Nature *458*, 1047–1050.

Shao, S., and Hegde, R.S. (2011). Membrane Protein Insertion at the Endoplasmic Reticulum.
Annu. Rev. Cell Dev. Biol. *27*, 25–56.

Shurtleff, M.J., Itzhak, D.N., Hussmann, J.A., Oakdale, N.T.S., Costa, E.A., Jonikas, M.,
Weibezahn, J., Popova, K.D., Jan, C.H., Sinitcyn, P., et al. (2018). The ER membrane protein
complex interacts cotranslationally to enable biogenesis of multipass membrane proteins. ELife
7, e37018.

Su, H.-L., Liao, C.-L., and Lin, Y.-L. (2002). Japanese Encephalitis Virus Infection Initiates
Endoplasmic Reticulum Stress and an Unfolded Protein Response. J. Virol. *76*, 4162–4171.

Volkmar, N., Thezenas, M.-L., Louie, S.M., Juszkiewicz, S., Nomura, D.K., Hegde, R.S.,
Kessler, B.M., and Christianson, J.C. (2019). The ER membrane protein complex promotes
biogenesis of sterol-related enzymes maintaining cholesterol homeostasis. J. Cell Sci. *132*.

Zhang, R., Miner, J.J., Gorman, M.J., Rausch, K., Ramage, H., White, J.P., Zuiani, A., Zhang,
P., Fernandez, E., Zhang, Q., et al. (2016). A CRISPR screen defines a signal peptide
processing pathway required by flaviviruses. Nature *535*, 164–168.

Zou, J., Lee, L.T., Wang, Q.Y., Xie, X., Lu, S., Yau, Y.H., Yuan, Z., Geifman Shochat, S., Kang,
C., Lescar, J., et al. (2015). Mapping the Interactions between the NS4B and NS3 proteins of
dengue virus. J. Virol. *89*, 3471–3483.

794