1   **Title**

2   **Comprehensive Classification of the RNaseH-like domain-containing Proteins in Plants**

4   Shuai Li, Kunpeng Liu and Qianwen Sun[#]

Tsinghua-Peking Joint Center for Life Sciences and Center for Plant Biology, School of Life Sciences,

Tsinghua University, Beijing 100084, China

[#] Corresponding author Qianwen Sun: sunqianwen@mail.tsinghua.edu.cn.

Main conclusion

This study showed that RNaseH-like domain containing proteins can form 27 subclusters and mainly

regulate nucleic acid metabolism in meristem tissues, revealing these proteins are critical in R-loop

regulation.

Abstract

R-loop is a nucleic acid structure containing an RNA-DNA hybrid and a displaced single-stranded

DNA. Recently, accumulated evidences showed that R-loops occur frequently in various organisms'

genomes and had crucial physiological functions, including replication, transcription and DNA repair.

RNaseH-like superfamily (RNHLS) domain-containing proteins, such as RNase H enzymes, RNase H1

and RNase H2, are important in restricting R-loop levels. However, little is known about other RNHLS

proteins and their relationship and function on modulating R-loops, especially in plants. In this study

we characterized 6193 RNHLS proteins from 13 representative plant species, and clustered them into

27 clusters, among which reverse transcriptases and exonucleases are the two largest groups. Moreover,

we found there are 691 RNHLS proteins in Arabidopsis, and the catalytic alpha helix and beta sheet

domain are conserved among them. Interestingly, each of the Arabidopsis RNHLS proteins is

composed of RNHLS domains and other distinctive protein domains. we further found that *RNHLS*

genes are highly expressed in diverse meristems and metabolic tissues. Various *RNHLS* genes

expressed in different tissues grouped together indicate that they work both isolatedly and

cooperatively. In summary, we systematically analyzed RNHL proteins in plant and found that there are

mainly 27 subclusters of them. Most of these proteins are implicated in DNA replication, RNA

transcription and nucleic acid degradation. We characterized and classified the RNHLS proteins in

plant, which affords new insights on investigation of novel regulatory mechanisms and functions in

R-loop biology.

Keywords: R-loop, RNaseH-like protein, transposon, phylogenetic tree

Introduction

R-loop is a nucleotide structure consists of RNA-DNA hybrid with the displaced single-stranded DNA

(Thomas et al. 1976). Formation of R-loops *in vivo* was found in process of DNA replication, retrovirus

or single stranded RNA virus infection or retrotransposon life cycle (Santos-Pereira and Aguilera 2015).

For a long time, R-loops have been considered as transcription by-products since the nascent transcripts

could bind back to their DNA templates (Huertas and Aguilera 2003). However, recent studies in

different organisms revealed that R-loops are widespread on genomes in yeast, plants and mammals

(Ginno et al. 2012; Chan et al. 2014; Wahba et al. 2016; Xu et al. 2017; Dumelie and Jaffrey 2017;

Sanz et al. 2016). These evidences together revealed that R-loops should be accurately modulated in

different cellular processes.

Ribonucleases H (RNase H), a type of endonuclease, specifically degrades RNA moiety of

RNA-DNA hybrid therefore can destroy and remove R-loops (Stein and Hausen 1969; Hausen and

Stein 1970; Smyshlyaev et al. 2013). RNA molecule might be the origin of life and RNase H mainly

disrupts the direct interaction of RNA and DNA (Joyce 2002). Therefore, it is reasonable that RNase H

is one of the most ancient protein and that many enzymes involved in RNA and DNA related processes

employs RNase H domain during evolution (Majorek et al. 2014; Ma et al. 2008). The RNase H-like

superfamily (RNHLS) proteins are groups of proteins with the similar 3D structure of RNase H domain,

including the type 1 and type 2 RNase Hs, retroviral integrases, DNA transposases, Holliday junction

resolvases, Piwi/Argonaute nucleases, exonucleases and key spliceosome component

Pre-mRNA-processing-splicing factor 8 (Prp8) and other RNA nucleases (V. et al. 2008; Zuo and

Deutscher 2001; Song et al. 2004; Rice and Baker 2001; Parker et al. 2004; Ariyoshi et al. 1994;

Nowotny et al. 2005). Hence, RNHLS proteins play important roles in DNA replication and repair,

RNA transcription and interference, homologous and non-homologous recombination, transposon and

retrovirus transposition, and single-stranded RNA virus infection (Majorek et al. 2014). Previous study

using bioinformatics approaches to search database for all RNHLS proteins revealed that the RNHLS

proteins could be grouped into 152 families and divided mainly into exonucleases and endonucleases

(Majorek et al. 2014). Plants are sessile and need to overcome different biotic or abiotic stress over

their life cycle. Therefore, the expression of *RNHLS* genes should be fine-tuned to maintain

development and overcome these stresses. However, little is known about RNHLS proteins and their

relationship in plants especially on regulation of R-loops.

In this study, to systemically analyze the relationship and diversity of RNHLS proteins in plant, we

using bioinformatics tools to search the database and collect the RNHLS proteins of 13 representative

plant species, including *Solanum lycopersicum* (tomato), *Solanum tuberosum* (potato), *Populus*

trichocarpa (western balsam poplar), *Glycine max* (soybean), *Citrus sinensis* (sweet orange),

Arabidopsis thaliana (arabidopsis), *Oryza sativa* (rice), *Physcomitrella patens* (moss), *Marchantia*

polymorpha (liverwort), *Klebsormidium flaccidum* (filamentous charophyte green algae),

Chlamydomonas eustigma (unicellular flagellate green algae), *Cyanidioschyzon merolae* (red alga), and

Chondrus crispus (carrageen Irish moss). We characterized 6193 RNHLS proteins in these

representative plants. They are clustered into 27 clusters, among them reverse transcriptases and

exonucleases are two largest groups. In Arabidopsis, there are 691 RNHLS proteins, consisting of

RNHLS domain with various types of protein domains in these proteins, while the catalytic alpha helix

and beta sheet domain is conserved among them. In addition, *RNHLS* genes are actively expressed in

meristems and other metabolic tissues. Various *RNHLS* genes expressed in different tissues are grouped

together indicating they work both individually and cooperatively. In summary, we characterized and

classified the RNHLS proteins in plants, which will extend our understanding of R-loop regulation *in*

vivo.

Materials and methods

Co-occurrence of RNHLS domain with other protein domains

The co-occurrence matrix of RNHLS domain (3.30.420.10) with other protein domains was

downloaded from Gene3D (Lewis et al. 2018). The heatmap was processed by online software

Morpheus (https://software.broadinstitute.org/morpheus/) using the default settings.

Protein sequence search

To identify the RNaseH-like superfamily proteins in Arabidopsis comprehensively, we used the

proteins sequences from RNHLS (IPR012337) of InterPro database (http://www.ebi.ac.uk/interpro/)

filtered with different species. In addition, we performed sequence BLAST with AtRNH1A

(At3g01410) as a query in the Phytozome and NCBI database (Goodstein et al. 2012; Altschul et al.

1990). We carried out the analysis after combining the sequence from the three methods and removed

the duplicates.

Protein sequence cluster

The identified proteins were first analyzed with CD-HIT and the threshold was set at 60% with word

size of 3 (Fu et al. 2012). The clustered proteins were used the BLAST to compute the similarity of

RNHLS proteins with an average pairwise E-value at least of 1×10^{-18} and redundancy proteins were

removed (Altschul et al. 1990). The network was visualized using Cytoscape3.7.0 and laid out using

the organic layout algorithm and annotated manually (Shannon et al. 2003).

Protein sequence alignment

To analysis the protein sequences systemically, we used Clustal Omega online service with the default

parameters except for all the iterations set as five (Sievers et al. 2011). The alignment results were

visualized with UGENE and MEGA7 (Okonechnikov et al. 2012). The structure-based sequence

alignment was performed to find the conserved secondary structure using the PROMALS3D and

JPred4 and manually adjusted with UGENE to identify the homologous positions with shift insertions

and deletions from conserved secondary structure elements (Pei et al. 2008; Drozdetskiy et al. 2015).

Phylogenetic tree construction

The phylogenetic tree of the subgroups and Arabidopsis RNHLS proteins were generated using Clustal

Omega at the same parameters for protein sequence alignment. For the total RNHLS proteins

phylogenetic tree, we employed the MAFFT software for multiple sequence alignment and constructed

phylogenetic tree using FastTree (Katoh et al. 2002; Price et al. 2009). We further modified the

phylogenetic tree with online software Interactive tree of life (iTOL) (Letunic and Bork 2016).

Protein domain diagram

The   information   of   protein   domains   was   from   Pfam   (http://pfam.xfam.org/),   InterPro

(http://www.ebi.ac.uk/interpro/) and UniProt (https://www.uniprot.org/). The diagram were depicted

with IBS (Liu et al. 2015).

RNA-seq data of Arabidopsis various tissues

The RNA-seq data of Arabidopsis *RNHLS* genes used here were downloaded from TraVA

(Transcriptome Variation Analysis, http://travadb.org/), except for which were not available (Klepikova

et al. 2016). The heatmap was visualized using Morpheus online software with default settings. GO

analysis of highly expressed genes in meristems was performed on the online database agriGO v2.0

(Tian et al. 2017).

Results

The species selected to cover the characters of RNHLS proteins in plants

To acquire representative plant species for analysis the RNHLS proteins, we employed the Gene3D to

investigate co-occurring domain families, with value for RNHLS proteins with other protein domain

families in 33 plant species (Fig. 1A and Fig. S1) (Lewis et al. 2018). The co-occurrence of RNHLS

domain with DNA polymerase domain, nuclease and nucleotide triphosphate hydrolase domain are in

most of the plant species (Fig. S1). Based on these results, we selected 10 representative plant species

and added three other species including *C. sinensis*, *M. polymorpha*, and *K. flaccidum* to further

represent the evolutionary tree of plants for further investigation (Fig. 1B) (Chang et al. 2016).

A global view of RNHLS protein clusters

To understand the function and distribution of RNHLS proteins in the 13 plant species, we acquired the

RNHLS proteins from the selected plants using the protein database InterPro (Hunter et al. 2009). A

total of 6193 proteins were collected through this method. To elucidate the relationship among these

proteins, we conducted a clustering analysis which relied on the sequence similarity of proteins. First,

we chose the representative proteins using CD-HIT with a threshold of 50 percent and achieved 2066

representative RNHLS proteins. Then, we compared the similarity of the these RNHLS proteins with

total RNHLS (6193) using BLAST with an average pairwise E-value at least of 1×10^{-18}. The network

showed that there are 27 clusters (Fig. 2). Most part of them are Ty1/Copia or Ty3/Gypsy reverse

transcriptases and other transposases or integrases. The second largest part of them are different types

of nucleic acid endonucleases and exonucleases (Fig. 2). The third part of them are Argonaute proteins

and DNA polymerases. The separation of these clusters were consistent with their biological function

and confirmed that our SSN map can show the relative information of the investigated RNHLS proteins.

To further investigation the relationships of all RNHLS proteins and within the 27 subclusters, we

constructed phylogenetic trees of total RNHLS proteins and each of the 27 subclusters. Consistent with

the SSN network, the RNHLS proteins are diversified and most of them are transposon polyprotein

(Fig. S3 and Fig. S4).

To further investigate the composition of these clusters at species level, we indicated the species

information with different colors at the SSN map (Fig. S2). The results showed that most of the

proteins comes from *O. sativa* are clustered in the Ty1/Copia and Ty3/Gypsy polyprotein clusters. In

the Ty1/Copia polyprotein group, proteins from *A. thaliana* and *O. sativa* are clustered in different

subclusters (Fig. S2). Moreover, proteins from *C. sinensis*, *C. crispus*, *S. lycopersicum* and *C. eustigma*

also form subclusters in this cluster. Interestingly, transposases from *C. crispus* and *M. polymorpha*

are clustered into two relative isolated groups (Fig. S2). Consistent with this phenomenon, most of

transposon-related polyprotein clusters prefer to clustering together in a species dependent manner.

Otherwise, the other clusters such as AGO, PRP8, DNA polymerase and RNH2 are distributed more

evenly compared with transposon-related polyprotein clusters (Fig. S2). Additionally, the AGO cluster

is separated into two sub-clusters consistent with their biological function (Fig. 2). Together, these

information indicates that RNHLS proteins are very diversified and the evolution of transposon within

each species are exclusive.

Phylogenetic tree of the Arabidopsis RNHLS proteins

To better understand the RNHLS proteins in plants, we used the RNHLS proteins from Arabidopsis as

candidates to conduct alignment and construct a phylogenetic tree (Fig. 3). Consistent with the protein

clusters, the retrotransposon reverse transcriptases occupy a big part of the clades (Fig. 3). There are 5

clades of Ty1/copia reverse transcriptases, which consistent with the cluster results that there are more

than eight sub-clusters (Fig. 3). Meanwhile, the HAT transposon proteins are also clustered together in

the tree (Fig. 3). Furthermore, the non-LTR retrotransposon reverse transcriptases comprise another

three clades of the tree. The CAF1 proteins are separated into two clades compared with MEN or SDN

proteins which shows only one clade on the tree (Fig. 3). Additionally, the exonucleases are clustered

together. Interestingly, we found that the AGO family proteins are separated into four clades of them

with AGO1/5/10, AGO2/3, AGO4/6/8/9 and AGO7 and this result is identical to previous reports (Fang

and Qi 2016). Various of DNA polymerases with RNHLS domain are clustered into one clade (Fig. 3).

Moreover, we found that one type of RNHLS proteins come from Ring-type E3 ubiquitin transferases.

The PRP8 family proteins are clustered in one clade (Fig. 3). Together these results give us a

comprehensive view of RNHLS proteins and the relationship among them in Arabidopsis.

The domain construction of different types of RNHLS proteins

To better characterize the properties of the RNHLS proteins in plant, we chose the representative

Arabidopsis RNHLS proteins in each of the clades to draw a protein domain diagram to indicate their

domain composition (Fig. 4). The RNH1 contains three important domains with HBD, Dis and RNH

domain. HBD is important for RNA:DNA hybrid binding while RNH domain is for catalytic activity

(Cerritelli and Crouch 2009). The Dis domain provides the ability for RNH to catalyze a variety of

hybrids without unloading (Fig. 4) (Cerritelli and Crouch 2009). Based on this, most RNHLS proteins

combine RNH domain with other domains to exercise their destroying RNA:DNA functions during

different biological processes. To analysis the relationship of RNHLS domain among different RNHLS

proteins, we performed a multiple sequence alignment with the truncated sequences of each

representative Arabidopsis RNHLS proteins (Fig. 5). The results indicate that the conserved alpha helix

and beta sheet domain present in these RNHLS proteins which consistent with their annotated function

in InterPro (Fig. 4 and Fig. 5).

RNHLS proteins are abundant in plant meristems

To predict the biological function of RNHLS proteins in plants, we first analysed their expression

pattern in Arabidopsis and rice using RNA-seq data from database (Fig. 6 and Fig. S5). In Arabidopsis

and rice, various *RNHLS* genes expressed in different tissues are grouped together. In Arabidopsis, we

found these genes are mainly expressed at meristem tissues, such as shoots and root meristems and

inflorescences (Fig. 6A). Some of the genes are mainly expressed in anthers and pistils (Fig. 6A). In

addition, in the young and mature anthers different types of *RNHLS* genes show clearly different

expression patterns (Fig. 6A). Meanwhile, the dry seeds accumulate another group expression of

RNHLS genes (Fig. 6A). However, some genes are expressed at senescent tissues due to programmed

cell death accompanied with nucleic acid degradation (Fig. 6A). We perform a GO analyzer of the

meristem expressed genes and it shows that these proteins are involved in nucleic acid metabolism (Fig.

6B). In rice, the result shows that post-emergence inflorescences and pre-emergence inflorescences are

grouped with totally different expression of *RNHLS* genes consistent with Arabidopsis (Fig. S5). In the

embryo-25 dap (days after plant), there is another group of *RNHLS* expressed (Fig. S5). Moreover, in

seedling four-leaf stage, half of the investigated *RNHLS* genes are expressed in this tissue (Fig. S5).

Taken together, the expression of *RNHLS* genes in different groups are throughout Arabidopsis and rice

development and consistent with their biological function. These results indicate the importance of

RNHLS genes in plants.

Discussion

In this study, we have characterized 6193 RNHLS proteins in 13 plants. Among them, 2066

representative RNHLS proteins are clustered into 27 clusters, including reverse transcriptases,

transposases, integrases, nucleic acid endonucleases, exonucleases, key spliceosome component Prp8,

as well as other RNA nucleases. Moreover, RNHLS domain is combined with various types of protein

domains in different RNHLS proteins. And the catalytic alpha helix and beta sheet domain are

conserved among them in Arabidopsis. In addition, *RNHLS* genes are actively expressed in meristems,

flower tissues and other metabolic tissues, and various *RNHLS* genes expressed in different tissues are

grouped together.

The selected 13 plant species represent the evolutionarily process of plants (Fig. 1B) (Chang et al.

2016). The distribution of RNHLS proteins in all species of plants suggests they are a type of the

evolutionarily oldest proteins (Ma et al. 2008). Among 27 clustered RNHLS proteins, DNA polymerase,

GTF2B, CAF1, PRP8, RRP6L, AGO and EXO worked in basic nucleic acid metabolic processes are

conserved in plant kingdom (Majorek et al. 2014). WRN is able to digest the RNA moiety of the

RNA:DNA hybrid (Suzuki et al. 1999). Recently, it was reported that WRN is involved in

R-loop-related genome stability (Marabitti et al. 2019). Moreover, SDN proteins are involved in

degrading small RNAs and could trimming dsRNA therefore raising the possibility of trimming RNA

strand in the RNA:DNA hybrids because of high structure similarity of dsRNA and RNA-DNA hybrid

(Chen et al. 2018). The family numbers of TE-related RNHLS proteins are larger, especially for

Ty1/Copia and Ty3/Gypsy reverse transcriptase family (Fig.2 and Fig. S4). Some of the TE-related

RNHLS proteins are not present in all selected plants, such as RICESLEEPER is only present in rice

(Fig. S2), Ty1/Copia, Ty3/Gypsy and Non-LTR reverse transcriptase are mainly enriched in rice,

Arabidopsis and other higher plants (Fig. 2 and Fig. S2). This is consistent with the size and

composition of their genome (Chang et al. 2016). As retrotransposons are abundant in higher plants, the

TE-related RNHLS proteins may play a major role in driving genetic diversity and evolution of these

organisms (Smyshlyaev et al. 2013; Moelling et al. 2017).

RNHLS protein is a large group of evolutionarily related, but strongly diverged protein family

(Majorek et al. 2014). The first identified RNHLS protein is RNase H from *Escherichia coli* (Leis et al.

1973). There are two types of RNase H enzymes, which differ in structure and substrate specificity.

RNase H1 is monomeric, whereas RNase H2 is monomeric in bacteria but composed of three subunits

in eukaryotes: RNH2A (the catalytic subunit), RNH2B and RNH2C. Both types of RNase H enzyme

have the ability to remove RNA-DNA hybrids in addition to having different specialized roles. The

phylogenetic analysis of RNHLS proteins in Arabidopsis shows that some proteins with similar

function were clustered in different group according to their sequence, such as AGO proteins. While

some proteins from different protein families clustered together, such as some Ty1/Copia and non-LTR

reverse transcriptase proteins. All these indicate that both amino acid construction and the motif

composition of RNHLS proteins are diversity. Domain diagram analysis of representative proteins

shows that all the RNHLS proteins have RNH domains except AGOs, which use a PIWI domain

instead (Fig.4). The Piwi domain have been reported adopts an RNase H-like fold and enables some,

but not all, AGO proteins to cleave target RNAs complementary to the bound sRNAs (Song et al. 2004).

A catalytic triad (Asp-Asp-His/Asp, DDH/D) is generally thought to be responsible for slicer activity is

present in both RNH domains and PIWI domain (Liu et al. 2004). Further study need to test whether

AGOs have ability to cleave the RNA strand in RNA-DNA hybrid or worked as a sensor for

RNA-DNA hybrid. In addition, RNHLS proteins are abundant in meristems, flower tissues and other

metabolic tissues, and various *RNHLS* genes expressed in different tissues grouped together indicate

they worked division and cooperation (Fig. 6). As RNHLS proteins work on nucleic acid metabolism

especially on R-loop homeostasis, this study characterized the RNHLS Proteins in plants and provide

new direction on R-loop regulation and function identification study in the future.

Conclusion

In this study, we employed the bioinformatics approaches to analyze the RNHLS proteins in

representative 13 plant species. A SSN network showed that these proteins can be separated into 27

clusters and consistent with their biological functions. Moreover, the *RNHLS* genes expression pattern

during arabidopsis and rice development further revealed that these proteins are mainly expressed in

meristems and metabolic tissues. These evidences shed light on our understanding of R-loop regulation

in plants.

Supplementary material

Supplementary material 1: Supplementary figure 1 to supplementary figure 5. (PDF 922 kb)

Supplementary material 2: A representative SSN of the RNHLS proteins. (CYS 4687 kb)

Supplementary material 3: A representative SSN of the RNHLS proteins colored with species

information. (CYS 4657 kb)

Supplementary material 4: Supplementary table 1. Gene3D domain family matrix of RNHLS domain

that co-occur with other protein domains. (XLSX 52.4 kb)

Supplementary material 5: Supplementary table 2. RNHLS protein information used in this study.

(XLS 558 kb)

Supplementary material 6: Supplementary table 3. tissue-specific expression data of *RNHLS* genes of

Arabidopsis and rice. (XLSX 187 kb)

Abbreviation

AGO, Argonaute; CAF1, putative CCR4-associated factor 1 homolog; Dis, disorder protein domain; DUF1744, domain of unknown function 1744 domain; Exo, exonuclease; Gag-pol, putative gag-pol polyprotein; GTF2B, general transcription factor 2-related zinc finger protein; HBD, hybrid binding domain; non-LTR, putative non-LTR retroelement reverse transcriptase; Prp8, pre-mRNA-processing-splicing factor 8; RBRE3UT, RBR-type E3 ubiquitin transferase; RNase H, ribonucleases H; RNH2, ribonuclease H2; RNHLS, RNase H-like superfamily; RRP6L, RRP6-like protein; SDN, small RNA degrading nuclease; SSN, sequence similarity network; WRN, werner syndrome ATP-dependent helicase; ZBED1, zinc finger BED domain-containing protein DAYSLEEPER (Transposase-like protein DAYSLEEPER).

Author contributions: Q.S. conceptualized the study. S.L. analyzed the sequences and prepared the figures, and prepared the manuscript with K.L. and Q.S.

Acknowledgements

This work was funded by grants from the Ministry of Science and Technology of China (grant no. 2016YFA0500800 to Q.S.); and the National Natural Science Foundation of China (grant nos. 91740105 and 31822028 to Q.S.). The Sun Lab was supported by Tsinghua-Peking Joint Center for Life Sciences, Tsinghua University Initiative Scientific Research Program, and the 1000 Young Talent Program of China. K.L. was supported by postdoctoral fellowship from Tsinghua-Peking Joint Center for Life Sciences.

Competing interests: The authors declare no competing interests.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. J Mol

Biol 215 (3):403-410. doi:10.1016/S0022-2836(05)80360-2

Ariyoshi M, Vassylyev DG, Iwasaki H, Nakamura H, Shinagawa H, Morikawa K (1994) Atomic

structure of the RuvC resolvase: a holliday junction-specific endonuclease from *E. coli*. Cell 78

(6):1063-1072. doi:10.1016/0092-8674(94)90280-1

Cerritelli SM, Crouch RJ (2009) Ribonuclease H: the enzymes in eukaryotes. FEBS J 276

(6):1494-1505. doi:10.1111/j.1742-4658.2009.06908.x

Chan YA, Aristizabal MJ, Lu PY, Luo Z, Hamza A, Kobor MS, Stirling PC, Hieter P (2014)

Genome-wide profiling of yeast DNA:RNA hybrid prone sites with DRIP-chip. PLoS Genet 10

(4):e1004288. doi:10.1371/journal.pgen.1004288

Chang C, Bowman JL, Meyerowitz EM (2016) Field Guide to Plant Model Systems. Cell 167

(2):325-339. doi:10.1016/j.cell.2016.08.031

Chen J, Liu L, You C, Gu J, Ruan W, Zhang L, Gan J, Cao C, Huang Y, Chen X, Ma J (2018) Structural

and biochemical insights into small RNA 3' end trimming by Arabidopsis SDN1. Nat Commun 9

(1):3585. doi:10.1038/s41467-018-05942-7

Drozdetskiy A, Cole C, Procter J, Barton GJ (2015) JPred4: a protein secondary structure prediction

server. Nucleic Acids Res 43 (W1):W389-394. doi:10.1093/nar/gkv332

Dumelie JG, Jaffrey SR (2017) Defining the location of promoter-associated R-loops at near-nucleotide

resolution using bisDRIP-seq. Elife 6. doi:10.7554/eLife.28306

Fang X, Qi Y (2016) RNAi in Plants: An Argonaute-Centered View. Plant Cell 28 (2):272-285.

doi:10.1105/tpc.15.00920

Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-generation

sequencing data. Bioinformatics 28 (23):3150-3152. doi:10.1093/bioinformatics/bts565

Ginno PA, Lott PL, Christensen HC, Korf I, Chedin F (2012) R-loop formation is a distinctive

characteristic of unmethylated human CpG island promoters. Mol Cell 45 (6):814-825.

doi:10.1016/j.molcel.2012.01.017

Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, Mitros T, Dirks W, Hellsten U,

Putnam N, Rokhsar DS (2012) Phytozome: a comparative platform for green plant genomics. Nucleic

Acids Res (40) :D1178-1186. doi:10.1093/nar/gkr944

Hausen P, Stein H (1970) Ribonuclease H. An enzyme degrading the RNA moiety of DNA-RNA

hybrids. Eur J Biochem 14 (2):278-283

Huertas P, Aguilera A (2003) Cotranscriptionally formed DNA:RNA hybrids mediate transcription

elongation impairment and transcription-associated recombination. Mol Cell 12 (3):711-721.

doi:http://dx.doi.org/10.1016/j.molcel.2003.08.010

Hunter S, Apweiler R, Attwood TK, Bairoch A, Bateman A, Binns D, Bork P, Das U, Daugherty L,

Duquenne L, Finn RD, Gough J, Haft D, Hulo N, Kahn D, Kelly E, Laugraud A, Letunic I, Lonsdale D,

Lopez R, Madera M, Maslen J, McAnulla C, McDowall J, Mistry J, Mitchell A, Mulder N, Natale D,

Orengo C, Quinn AF, Selengut JD, Sigrist CJ, Thimma M, Thomas PD, Valentin F, Wilson D, Wu CH,

Yeats C (2009) InterPro: the integrative protein signature database. Nucleic Acids Res (37) :D211-215.

doi:10.1093/nar/gkn785

Joyce GF (2002) The antiquity of RNA-based evolution. Nature 418 (6894):214-221.

doi:10.1038/418214a

Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence

alignment based on fast Fourier transform. Nucleic Acids Res 30 (14):3059-3066. doi:DOI

10.1093/nar/gkf436

Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA (2016) A high resolution map

of the Arabidopsis thaliana developmental transcriptome based on RNA-seq profiling. Plant J 88

(6):1058-1070. doi:10.1111/tpj.13312

Leis JP, Berkower I, Hurwitz J (1973) Mechanism of action of ribonuclease H isolated from avian

myeloblastosis virus and Escherichia coli. Proc Natl Acad Sci U S A 70 (2):466-470

Letunic I, Bork P (2016) Interactive tree of life (iTOL) v3: an online tool for the display and annotation

of phylogenetic and other trees. Nucleic Acids Res 44 (W1):W242-245. doi:10.1093/nar/gkw290

Lewis TE, Sillitoe I, Dawson N, Lam SD, Clarke T, Lee D, Orengo C, Lees J (2018) Gene3D:

Extensive prediction of globular domains in proteins. Nucleic Acids Res 46 (D1):D435-D439.

doi:10.1093/nar/gkx1069

Liu J, Carmell MA, Rivas FV, Marsden CG, Thomson JM, Song JJ, Hammond SM, Joshua-Tor L,

Hannon GJ (2004) Argonaute2 is the catalytic engine of mammalian RNAi. Science 305

(5689):1437-1441. doi:10.1126/science.1102513

Liu W, Xie Y, Ma J, Luo X, Nie P, Zuo Z, Lahrmann U, Zhao Q, Zheng Y, Zhao Y, Xue Y, Ren J (2015)

IBS: an illustrator for the presentation and visualization of biological sequences. Bioinformatics 31

(20):3359-3361. doi:10.1093/bioinformatics/btv362

Ma BG, Chen L, Ji HF, Chen ZH, Yang FR, Wang L, Qu G, Jiang YY, Ji C, Zhang HY (2008)

Characters of very ancient proteins. Biochem Biophy Res Co 366 (3):607-611.

doi:10.1016/j.bbrc.2007.12.014

Majorek KA, Dunin-Horkawicz S, Steczkiewicz K, Muszewska A, Nowotny M, Ginalski K, Bujnicki

JM (2014) The RNase H-like superfamily: new members, comparative structural analysis and

evolutionary classification. Nucleic Acids Res 42 (7):4160-4179. doi:10.1093/nar/gkt1414

Marabitti V, Lillo G, Malacaria E, Palermo V, Sanchez M, Pichierri P, Franchitto A (2019) ATM

pathway activation limits R-loop-associated genomic instability in Werner syndrome cells. Nucleic

Acids Res. doi:10.1093/nar/gkz025

Moelling K, Broecker F, Russo G, Sunagawa S (2017) RNase H As Gene Modifier, Driver of Evolution

and Antiviral Defense. Front Microbio 8:1745. doi:10.3389/fmicb.2017.01745

Nowotny M, Gaidamakov SA, Crouch RJ, Yang W (2005) Crystal structures of RNase H bound to an

RNA/DNA hybrid: substrate specificity and metal-dependent catalysis. Cell 121 (7):1005-1016.

doi:10.1016/j.cell.2005.04.024

Okonechnikov K, Golosova O, Fursov M, team U (2012) Unipro UGENE: a unified bioinformatics

toolkit. Bioinformatics 28 (8):1166-1167. doi:10.1093/bioinformatics/bts091

Parker JS, Roe SM, Barford D (2004) Crystal structure of a PIWI protein suggests mechanisms for

siRNA recognition and slicer activity. EMBO J 23 (24):4727-4737. doi:10.1038/sj.emboj.7600488

Pei J, Kim BH, Grishin NV (2008) PROMALS3D: a tool for multiple protein sequence and structure

alignments. Nucleic Acids Res 36 (7):2295-2300. doi:10.1093/nar/gkn072

Price MN, Dehal PS, Arkin AP (2009) FastTree: computing large minimum evolution trees with

profiles instead of a distance matrix. Mol Biol Evol 26 (7):1641-1650. doi:10.1093/molbev/msp077

Rice PA, Baker TA (2001) Comparative architecture of transposase and integrase complexes. Nat Struct

Biol 8 (4):302-307. doi:10.1038/86166

Santos-Pereira JM, Aguilera A (2015) R loops: new modulators of genome dynamics and function. Nat

Rev Genet 16 (10):583-597. doi:10.1038/nrg3961

Sanz LA, Hartono SR, Lim YW, Steyaert S, Rajpurkar A, Ginno PA, Xu X, Chedin F (2016) Prevalent,

Dynamic, and Conserved R-Loop Structures Associate with Specific Epigenomic Signatures in

Mammals. Mol Cell 63 (1):167-178. doi:10.1016/j.molcel.2016.05.032

Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski B, Ideker T

(2003) Cytoscape: a software environment for integrated models of biomolecular interaction networks.

Genome Res 13 (11):2498-2504. doi:10.1101/gr.1239303

Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H, Remmert M,

Soding J, Thompson JD, Higgins DG (2011) Fast, scalable generation of high-quality protein multiple

sequence alignments using Clustal Omega. Mol Syst Biol 7 (1):539. doi:10.1038/msb.2011.75

Smyshlyaev G, Voigt F, Blinov A, Barabas O, Novikova O (2013) Acquisition of an Archaea-like

ribonuclease H domain by plant L1 retrotransposons supports modular evolution. Proc Natl Acad Sci U

S A 110 (50):20140-20145. doi:10.1073/pnas.1310958110

Song JJ, Smith SK, Hannon GJ, Joshua-Tor L (2004) Crystal structure of Argonaute and its

implications for RISC slicer activity. Science 305 (5689):1434-1437. doi:10.1126/science.1102514

Stein H, Hausen P (1969) Enzyme from calf thymus degrading the RNA moiety of DNA-RNA Hybrids:

effect on DNA-dependent RNA polymerase. Science 166 (3903):393-395.

doi:10.1126/science.166.3903.393

Suzuki N, Shiratori M, Goto M, Furuichi Y (1999) Werner syndrome helicase contains a 5'-->3'

exonuclease activity that digests DNA and RNA strands in DNA/DNA and RNA/DNA duplexes

dependent on unwinding. Nucleic Acids Res 27 (11):2361-2368

Thomas M, White RL, Davis RW (1976) Hybridization of RNA to double-stranded DNA: formation of

R-loops. Proc Natl Acad Sci U S A 73 (7):2294-2298

Tian T, Liu Y, Yan H, You Q, Yi X, Du Z, Xu W, Su Z (2017) agriGO v2.0: a GO analysis toolkit for

the agricultural community, 2017 update. Nucleic Acids Res 45 (W1):W122-W129.

doi:10.1093/nar/gkx382

Pena V, Rozov A, Fabrizio P, Luhrmann R, Cwahl M (2008) Structure and function of an RNase H

domain at the heart of the spliceosome. EMBO J 27:2929–40.

Wahba L, Costantino L, Tan FJ, Zimmer A, Koshland D (2016) S1-DRIP-seq identifies high expression

and polyA tracts as major contributors to R-loop formation. Genes Dev 30 (11):1327-1338.

doi:10.1101/gad.280834.116

Xu W, Xu H, Li K, Fan Y, Liu Y, Yang X, Sun Q (2017) The R-loop is a common chromatin feature of

the Arabidopsis genome. Nat plants 3 (9):704-714. doi:10.1038/s41477-017-0004-x

Zuo Y, Deutscher MP (2001) Exoribonuclease superfamilies: structural analysis and phylogenetic

distribution. Nucleic Acids Res 29 (5):1017-1026

Figure legends

Figure 1. Selection of candidate species for investigation of plant RNHLS proteins.

(A) The similarity of Gene3D domain family co-occurrence of RNHLS domain with other protein

domains in all available plant species. The species to be investigated in this study were marked in blue

font. The original matrix of other protein domains that co-occurred with RNHLS domain were

downloaded from Gene3D (http://gene3d.biochem.ucl.ac.uk). The similarity heatmap were performed

in Morpheus (https://software.broadinstitute.org/morpheus/) using the default setting. (B) The

cladogram of species for investigation of RNHLS proteins. The blue font is marked the selected species

from the similarity matrix while the black font is marked the species which are supplemented for better

covering the plant evolutionary.

Figure 2. A representative SSN of the RNHLS proteins.

A total of 4,755 protein sequences were depicted by 2,066 nodes (circles), which represent proteins

sharing >50% sequence identity. Edges between nodes indicate an average pairwise BLAST E-value of

at least 1×10^{-18}. Node coloring represents subgroup classification. Names indicate subgroups that

contain at least one protein with literature-documented functional information except for nodes marked

with gray. The network was visualized by Cytoscape3.7.0 using the organic layout algorithm. WRN,

Werner syndrome ATP-dependent helicase; SDN, small RNA degrading nuclease; RNH2, ribonuclease

H2; AGO, Argonaute; RNH1, ribonuclease H1; RRP6L, RRP6-like protein; RBRE3UT, RBR-type E3

ubiquitin transferase; Exo, exonuclease; Ty1, Ty1/copia-element polyprotein; Ty3, Gypsy/Ty3 element

polyprotein; non-LTR, putative non-LTR retroelement reverse transcriptase; CAF1, putative

CCR4-associated factor 1 homolog; HAT, HAT dimerisation domain-containing protein; GTF2B,

general transcription factor 2-related zinc finger protein; ZBED1, zinc finger BED domain-containing

protein DAYSLEEPER (Transposase-like protein DAYSLEEPER); Gag-pol, putative gag-pol

polyprotein; PRP8, Pre-mRNA-processing-splicing factor 8.

Figure 3. Phylogenetic tree of the Arabidopsis RNHLS proteins.

A total of 691 Arabidopsis RNHLS protein sequences were aligned using Clustal Omega

(https://www.ebi.ac.uk/Tools/msa/clustalo/). The phylogenetic tree was constructed using the results

from this alignment and modified using iTOL (https://itol.embl.de/). Different types of nucleases and

polymerases are clearly separated to different clades. The distribution of transposon-related proteins is

more diversified than AGOs, SDNs and MENs.

Figure 4. Protein domain diagram of Arabidopsis representative RNHLS proteins.

The domain annotations of these proteins were combined from Pfam (http://pfam.xfam.org/), InterPro

(http://www.ebi.ac.uk/interpro/) and UniProt (https://www.uniprot.org/). HBD, hybrid binding domain;

Dis, disorder protein domain; RNH, ribonuclease H domain; RNHS, Ribonuclease H-like superfamily

domain; ZFRT, zinc finger ring-type domain; IBR, In Between Ring fingers domain; RNA B, RNA

binding domain; HRDC, helicase and RNase D C-terminal; DNA POL E/B domain, DNA polymerase

alpha/epsilon subunit B domain; POLAc, DNA polymerase A domain; GR, glycine-rich domain; Nt,

N-terminal domain; L1, linker 1 domain;   AGO1 , Argonaute 1, consists of GR, glycine-rich domain,

Nt, N-terminal domain, L1, linker 1 domain; PAZ, PAZ domain; L2, linker 2 domain; Mid, Mid

domain; Piwi, Mid domain; Ty1POLYP, Ty1/copia-element polyprotein; GAGPI, GAG-pre-integrase

domain; RVT, reverse transcriptase, RNA-dependent DNA polymerase domain; ENDON,

Endonuclease superfamily domain; ZF RVT, reverse transcriptase zinc-binding domain; Ty3POLYP,

Ty3/Gypsy-element polyprotein; GAG, retrotransposon gag domain; ZincB, integrase zinc-binding

domain; POL2B, DNA polymerase epsilon catalytic subunit; DNAPOLB, DNA-directed DNA

polymerase family B multifunctional domain; DUF1744, domain of unknown function 1744 domain;

PRP8, Pre-mRNA-processing-splicing factor 8; PRO8NT, N terminus of PRP8 domain; PROCN,

central domain of PRP8 domain; U6RNAB, U6-snRNA-binding of PRP8 domain; MPN (Mpr1, Pad1

N-terminal) domain. D represents for aspartic acid. E represents for glutamic acid. H represents for

histidine. T represents for threonine. The numbers besides the protein indicate the amino acid number

of that protein.

Figure 5. Multiple sequence alignment of Arabidopsis representative RNHLS proteins.

The RNHLS domains of each protein were selected for this alignment. Each of RNHLS domains were

predicted the secondary structure using Jpred4 (http://www.compbio.dundee.ac.uk/jpred/index.html).

The alignment was visualized using UGENE and adjusted manually.

Figure 6. Tissue-specific expression of Arabidopsis *RNHLS* genes.

(A) The transcriptome of *RNHLS* genes were downloaded from TRAVA (http://travadb.org/browse/)

except for some of the genes could not be retrieved from that database. These genes are mainly

expressed at the meristems such as flowers inflorescences and root meristems. The yellow rectangles

indicate the clusters of genes highly expressed in these tissues. (B) GO annotation of the genes

expressed in Arabidopsis meristems. These genes are mainly involved in nucleic acid metabolism.

Figure 1

A

B

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6