Species specificity and intraspecific variation in the chemical profiles of *Heliconius* butterflies across a large geographic range

Kathy Darragh[1,2], Gabriela Montejo-Kovacevich[1], Krzysztof M. Kozak[2], Colin R. Morrison[2,3], Clarisse M. E. Figueiredo[4], Jonathan S. Ready[4], Camilo Salazar[5], Mauricio Linares[5], Kelsey J. R. P. Byers[1,2], Richard M. Merrill[2,6], W. Owen McMillan[2], Stefan Schulz[7], Chris D. Jiggins[1,2]

[1]*Department of Zoology, University of Cambridge, Cambridge, Cambridgeshire, United Kingdom*

[2]*Smithsonian Tropical Research Institute, Panama City, Panama*

[3]*Department of Integrative Biology, The University of Texas at Austin, Austin, Texas, United States*

[4]*Institute for Biological Sciences, Universidade Federal do Pará, Belém, Pará, Brazil*

[5]*Biology Program, Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Bogota, Colombia*

[6]*Division of Evolutionary Biology, Faculty of Biology, Ludwig-Maximilians-Universität München, Munich, Germany*

[7]*Institute of Organic Chemistry, Technische Universität Braunschweig, Braunschweig, Germany*

Abstract

Traits important for mate choice and behavioural isolation are predicted to be under strong stabilising selection within species, however such traits can also exhibit variation at the population level driven by neutral and adaptive evolutionary processes. Here, we describe patterns of divergence among androconial and genital chemical profiles at inter- and intra-specific levels in mimetic *Heliconius* butterflies. Most variation in chemical bouquets was found between species, but there were also quantitative differences at the population level. The strength of the correlation between chemical and the genetic divergence varies between species and is generally stronger for genital than for androconial chemical profiles. We identified 'indicator' compounds characteristic of particular species that included known biologically active compounds, suggesting a promising approach for identification of candidate compounds for future behavioural studies in novel systems. Overall, the strong signal of species identity suggests a role for these compounds in species recognition, but with additional neutral variation at the population level.

Keywords

Mate choice, reproductive isolation, chemical ecology, Lepidoptera, pheromones, signalling

Introduction

Reproductive isolation between lineages is important for the maintenance of species diversity (Coyne & Orr, 2004). In many animals, mate choice provides a strong pre-mating barrier to hybridisation (Friberg et al., 2008; Gray & Cade, 2000; Martin & Mendelson, 2016; Nagel & Schluter, 1998; Ready et al., 2006; Seehausen et al., 2008; Selz, Pierotti, Maan, Schmid, & Seehausen, 2014). Closely related species often differ in traits important for mate choice, with individuals displaying a preference for conspecific phenotypes (Jiggins, Naisbit, Coe, & Mallet, 2001; Mas & Jallon, 2005; Ryan & Guerra, 2014; Saveer et al., 2014;

Yildizhan et al., 2009). These traits can also exhibit variation both within and between populations of the same species, either due to genetic drift and/or varying selective regimes across their ranges (Bolnick & Kirkpatrick, 2012; Ryan & Guerra, 2014; Ryan & Rand, 1993; Ryan, Rand, & Weigt, 1996). Traits involved in behavioural isolation are predicted to show strong species-specific differences (Gerhardt, 1982), and typically be subject to stabilising selection which can act to decrease intraspecific phenotypic variation (Butlin, Hewitt, & Webb, 1985; Pfennig, 1998; Ptacek, 2000). As a consequence, we would expect to find little trait variability, or at least certain features to be invariant, across species geographic ranges (Benedict & Bowie, 2009; Ferreira & Ferguson, 2002; McPeek, Symes, Zong, & McPeek, 2011; Weber, Mitko, Eltz, & Ramírez, 2016).

Signals important for behavioural isolation could arise from the divergence of traits used in intraspecific communication between populations (Johansson & Jones, 2007; Mendelson & Shaw, 2012; Ryan & Rand, 1993; Smadja & Butlin, 2008). Signal divergence can be driven by various factors, both neutral and adaptive, usually involving multiple evolutionary forces (Leonhardt, Rasmussen, & Schmitt, 2013; Sun et al., 2013). A positive correlation between genetic distance, geographic distance, and phenotypic variation is consistent with stochastic processes, such as genetic drift, playing a prominent role (Irwin, Thimgan, & Irwin, 2008). In contrast, a lack of correlation between phenotypic and genetic divergence suggests that selection is shaping the phenotypic variation, perhaps driving divergence in different directions in each population (Campbell et al., 2010; Conrad, Paxton, Assum, & Ayasse, 2018; Hankison & Ptacek, 2008; Mullen, Vignieri, Gore, & Hoekstra, 2009).

Chemical compounds, such as sex pheromones, mediate intraspecific communication in many systems (Wyatt, 2003, 2014). The role of chemical signalling in behavioural isolation is also well established, especially among moth species (Löfstedt, 1993; Smadja &

Butlin, 2008). Due to the coordination between detection and production, these pheromone blends are traditionally regarded as being under stabilising selection towards a species stereotype (Löfstedt 1993). Even when species-specific characteristics are present, chemical compounds can exhibit intraspecific variation, with both qualitative and quantitative differences found across a species range (Conrad et al., 2018; Gemeno, Lutfallah, & Haynes, 2000; Groot et al., 2009; Huang et al., 1998; McElfresh & Millar, 2001; Ryan, Moore, & Walter, 1995; Weber et al., 2016).

Studies of *Heliconius* butterflies have greatly contributed to our understanding of adaptation and speciation (Jiggins, 2008, 2017; Merrill et al., 2015). Despite the reliance of this group on visual cues for mating (Bybee et al., 2012; Finkbeiner, Fishman, Osorio, & Briscoe, 2017; Jiggins et al., 2001; Merrill, Chia, & Nadeau, 2014; Sánchez et al., 2015), it has long been suggested that male pheromones also play a role in pre-mating barriers (Jiggins, 2008; Merrill et al., 2015), but so far only a few species have been studied. Behavioural experiments reveal that chemical signalling in *Heliconius erato, H. melpomene* and *H. timareta* is important for female mate choice (Darragh et al., 2017; Mérot, Frérot, Leppik, & Joron, 2015). Electroantennographic studies have shown that *Heliconius cydno* and *H. melpomene* respond to both con- and heterospecific androconial chemical bouquets (Byers et al., unpublished), and have identified individual compounds that are biologically active. Furthermore, studies of *H. cydno*, *H. doris*, *H. hecale*, *H. ismenius*, *H. melpomene*, *H. pardalinus*, *H. sara* and *H. timareta* from Panama, Colombia and Ecuador found that major compounds differ between species (Mann et al., 2017; Mérot et al., 2015), suggesting a potential role in reproductive isolation.

The role of chemical signalling is likely to be especially important in co-mimics, where visual signals alone are not sufficient to identify conspecifics (Estrada & Jiggins, 2008; Giraldo, Salazar, Jiggins, Bermingham, & Linares, 2008; Mérot et al., 2013; Sánchez

et al., 2015). In addition, chemical compounds could be part of a multimodal aposematic warning signal (Rojas et al., 2018; Rothschild, 1961), with some tentative evidence that co-mimics exhibit similar chemical bouquets to aid recognition by predators (Mann et al., 2017).

Here, we describe the chemical profiles of seven species of *Heliconius* from over 250 individuals collected across the Neotropics. We focus on *H. melpomene*, *H. erato*, and their relatives, due to the fact that they are the best-studied *Heliconius* species to date. We analysed both wing androconial and genital compounds of male butterflies. We hypothesise that biologically relevant signals for species recognition are likely to be under stronger stabilising selection compared to other compounds. Thus, consistency of compounds across the geographic range of a species might provide an approach to detect putatively biologically active compounds important for mate choice. We use *H. melpomene* as a test-species due to the availability of behavioural and electrophysiological data to investigate this approach, by evaluating consistency in compound blends across different localities.

The extensive dataset analysed here allows us to test evolutionary hypotheses, as well as identifying interesting candidate compounds for future behavioural studies. Firstly, we investigated intraspecific variation in chemical profiles of *H. melpomene* and *H. erato*, which are co-mimetic species with a diversity of geographic colour pattern races found across the Neotropics. We correlated chemical profile data with both geographic and genetic distances between different localities to test the hypothesis of selective evolutionary forces driving the observed variation in chemical profiles. Furthermore, to test if the chemical compounds are part of the aposematic co-mimicry signal we used samples from two different mimicry rings in western Ecuador and Panama (one "*H. melpomene*-like" mimicry ring including *H. melpomene* and *H. erato* and one "*H. cydno*-like" mimicry ring including *H. cydno* and *H. sapho*, as well as *H. eleuchia* in Ecuador only).

Materials and Methods

Figure 1. Map indicating species and populations from twelve localities across the Neotropics.

H. erato (1-11) 1) *H. e. demophoon* 2) *H. e. dignus* 3) *H. e. lativitta* 4) *H. e. cyrbia* 5) *H. e. microclea* 6) *H. e. emma* 7) *H. e. amphitrite* 8) *H. e. hydara* 9) *H. e. amazona* 10) *H. e. phyllis* 11) *H. e. venustus*. *H. melpomene* (12-23) 12) *H. m. rosina* 13) *H. m. bellula* 14) *H. m malleti* 15) *H. m cythera* 16) *H. m. xenoclea* 17) *H. m. schunkei* 18) *H. m. melpomene* 19) *H. m. madeira* 20) *H. m. thelxiope* 21) *H. m. intersectus* 22) *H. m. penelope* 23) *H. m. burchelli*. *H. timareta* (24-25) 24) *H. t. tristero* 25) *H. t. nov. spp. H. elevatus* (26-28) 26) *H. e. willmotti* 27) *H. e. schmassmanni* 28) *H. e. perchlora*. *H. cydno* (29-30) 29) *H. c. chioneus* 30) *H. c. alithea*. *H. sapho* (31-32) 31) *H. s. sapho* 32) *H. s. candidus*. *H. eleuchia* 33) *H. e. primularis*. See Table S1 for sample numbers.

Between February 2016 and August 2017 wild males of *Heliconius cydno*, *H. elevatus*, *H. eleuchia*, *H. erato*, *H. melpomene*, *H. sapho* and *H. timareta* were collected from twelve localities. Between two and fifteen males were chemically analysed per population (Fig. 1, Table S1). We follow the latest *Heliconius* taxonomy (Lamas & Jiggins, 2017).

Extraction and chemical analysis of tissues

The androconial region of the wing, previously described as the grey-brown overlapping region of the hindwing (Darragh et al., 2017), as well as the genitalia, were dissected for analysis immediately after collection. To extract pheromones, the tissue was soaked in 200µl dichloromethane containing 200ng 2-tetradecyl acetate (internal standard) in 2ml glass vials with PTFE-coated caps (Agilent, Santa Clara, USA) for one hour. The solvent was then transferred to new vials and, maintained cool in the field and stored at -20°C upon return. Androconial samples were evaporated to a reduced volume at room temperature prior to analysis. Pheromone extracts were analysed by GC/MS using an Agilent model 5977 mass-selective detector connected to an Agilent GC model 7890B and equipped with an Agilent ALS 7693 autosampler. HP-5MS fused silica capillary columns (Agilent, Santa Clara, USA, 30 m × 0.25 mm, 0.25 µm) were used. Injection was performed in splitless mode (250°C injector temperature) with helium as the carrier gas (constant flow of 1.2 ml/min). The temperature programme started at 50°C, was held for 5 minutes, and then rose at a rate of 5°C/minute to 320°C, before being held at 320°C for 5 minutes. Components were identified by comparison of mass spectra and gas chromatographic retention index with those of authentic reference samples and also by analysis of mass spectra. Components were quantified using 2-tetradecyl acetate as an internal standard. Only compounds eluting earlier than hexacosane were analysed in androconial samples, and those earlier than nonacosane

used in genital samples (Darragh et al. 2017). We removed compounds that were not found in at least half of all individuals from a given population.

Calculation of genetic and geographic distance matrices

To explore genetic distance among the studied *H. erato* and *H. melpomene* subspecies, we computed genetic covariance matrices and performed MDS for each species separately. A whole-genome sequence from a representative individual from each subspecies was used (Table S2). Genotypes were inferred from reads mapped to the *H. melpomene* (v2.5) and *H. erato demophoon* genome scaffolds (Challis, Kumar, Dasmahapatra, Jiggins, & Blaxter, 2016; Davey et al., 2017; Heliconius Genome Consortium, 2012; Van Belleghem et al., 2017) with bwa v0.7.15 (Li & Durbin, 2009). We computed a pairwise identical by state (IBS) matrix with a random sampled read from each position in the genome, implemented in ANGSD v0.912 (Korneliussen, Albrechtsen, & Nielsen, 2014). An interspecific genetic distance matrix was constructed using the function "cophenetic.phylo" from the *ape* package (Paradis & Schliep, 2018) with a previously published phylogeny (Kozak et al., 2015). Geographic distance matrices were created by inputting the co-ordinates of collection localities into the function "distm" in the *geosphere* package to calculate the Haversine great-circle-distance between points (Hijmans, 2017).

Statistical analyses

Inter- and intra-specific indicator compounds

We carried out indicator analysis using the *indicspecies* package (Cáceres & Legendre, 2009). Groupings are decided *a priori* (in this case species or sub-species) and components are determined which act as indicators of these groups. The best indicators are those which are only found in a single group and all group members possess the compound; such a compound would have an indicator value of 1. We used the function "indicators" to

investigate both which single compounds and combinations of compounds best predict group membership. We used the function "pruneindicators" to find the single or combinations of compounds which had the highest indicator values.

Variation in chemical profiles

Divergence in chemical profiles across species and populations was estimated with non-metric multidimensional scaling (NMDS) ordination in three dimensions, based on a Bray-Curtis similarity matrix. We used the "metaMDS" function in the *vegan* package version 2.5-1 (Oksanen et al., 2017), and visualised the NMDS using the *ade4* package (Dray & Dufour, 2007).

We assessed the relative importance of relevant factors in driving the variation in chemical profiles with multivariate statistical analyses. These factors included: species identity, geographic region and individual locality. To compare overall variation in chemical composition between groups, we carried out PERMANOVA (permutational multivariate analysis of variance) testing based on a Bray-Curtis distance matrix, using the "adonis2" function in the *vegan* package (Oksanen et al., 2017) with 1000 permutations. We investigated each term in the model sequentially, starting with species identity, the main clustering factor found from visualisation with NMDS, followed by geographic region (Panama *vs* Western Andes *vs* Eastern Andes *vs* Amazon), and finally individual collecting localities. We followed these PERMANOVA tests with *post hoc* pair-wise testing using the function "pairwise.perm.manova" in the *RVAidememoie* package, with Bonferroni correction, to identify which grouping factors were significantly different (Hervé, 2018). We repeated this within species, in *H. erato* and *H. melpomene,* to investigate fine-scale intraspecific geographic patterns. In the within species analysis we included geographic region (Panama *vs* Western Andes *vs* Eastern Andes *vs* Amazon), and individual collecting localities as the two factors.

One issue with distance-based analyses such as PERMANOVA is that differences in dispersion between groups can be confounded with differences in location (Warton, Wright, & Wang, 2012). To confirm these analyses and account for this issue, we implemented multivariate generalised linear models using the function "ManyGLM" from the *mvabund* package (Wang, Naumann, Wright, & Warton, 2012). We modelled the data using a negative binomial distribution, which we found to be appropriate through examination of residual plots. For interspecific analyses we included species, region, and locality nested within region in the model. For intraspecific analyses we included region and locality nested within region. The "ManyGLM" function fits models to each chemical compound, summing the test statistics to give a multivariate test statistic known as Sum-of-LR. This statistic can be tested for significance using resampling methods. We can also determine which compounds are driving between-group differences by looking at the individual contribution of each compound to the Sum-of-LR, with p-values adjusted for multiple testing using the "adjust" option.

Phylogenetic and geographic distance

Shared ancestry can explain part of the variation in a species' chemical profile. We tested for a correlation between phylogenetic distance and chemical profile divergence with Mantel tests using the *vegan* package (Oksanen et al., 2017). To investigate the role of geographic distance in chemical profile divergence, we compared geographic and chemical distances matrices with partial Mantel tests. To visualise the species phylogeny (Kozak et al., 2015) we used the "plot.phylo" function from the ape package (Paradis & Schliep, 2018).

Genomic and chemical distance within species

We calculated intraspecific genetic distances using genome sequences from 11 *H. erato* and 13 *H. melpomene* populations. We visualised genetic distances in two dimensions

using MDS with the function "cmdscale". We tested for a correlation between intraspecific genetic distance and chemical profile divergence with mantel tests using the *vegan* package (Oksanen et al., 2017). Hybrids between populations of the same species were excluded from this analysis (Table S2). We used partial Mantel tests to investigate the role of geographic distance.

Co-mimics and similarity of chemical profiles

We used samples of two mimicry rings from two localities, Panama and western Ecuador. *H. melpomene* and *H. erato* form one mimicry ring, whilst *H. cydno* and *H. sapho* form another, with the addition of *H. eleuchia* in western Ecuador. To test which factors most influenced the variation in chemical profiles we used both PERMANOVA and multivariate GLM. We included species, region, mimicry group, and interactions between these terms in the models.

All statistical analyses were performed with *R* version 3.5.1 (R Core Team, 2018). Figures were made using a palette of colours optimized for colour-blindness (Wong, 2011). Raw sequence data are available through GenBank Short Read Archive (XXX), GCMS data and R code are available through Dryad (XXX).

Results

Chemical compounds in androconia and genitals:

We sampled 252 androconia and 275 genitals across 42 populations of 33 subspecies of seven species, and identified 349 compounds in the genitals and 157 in the androconia. Of the total number of androconial compounds, 38% are fatty acid derivatives, 20% aromatics, 10% terpenoids, 1% macrolides, <1% lactones and 31% unknown or unidentified compounds. Of the genital compounds, 17% are fatty acid derivatives, 7% aromatics, 10% terpenoids, 1% lactones, 12% macrolides and 44% unknown or unidentified compounds. The main difference is that there are more macrolides in the genitals than androconia.

Heliconius species varied considerably in the amount and abundance of compounds (Fig. 2). Between species there was variation in the number of compounds per individual, and the overall amount of compounds detected (Table S3, S4). For the androconia, *H. eleuchia* had the fewest compounds (13±5), and *H. melpomene* the most (32±11). *H. sapho* had the lowest total abundance of compounds at 1,458±1,606 ng, and *H. melpomene* the most at 7,412±16,484 ng. The species with the fewest genital compounds was *H. sapho* with 32±7, and the most *H. cydno* with 102±21. *H. sapho* also had the lowest total amount of compounds at 6,991±7,950 ng, and *H. cydno* the highest at 91,517±134,225 ng. In general, a higher number of compounds and amount of each is found in the genitals than in the androconial patches of *Heliconius*' wings.

Figure 2. Composition of androconial and genital bouquets across seven *Heliconius* species.
Species significantly differ in: (A) number of androconial compounds (anova, $F_{6,245}$=21.54, p<0.001),
(B) number of genital compounds (anova, $F_{6,268}$=36.15, p<0.001), (C) amount of androconial
compounds (anova, $F_{6,245}$=11.55, p<0.001), (D) amount of genital compounds (anova, $F_{6,268}$=11.62,
p<0.001). Four outlier individuals were removed from C.

Are there species-specific chemical compounds?

In order to identify candidate species recognition pheromones, we examined our data for species specific compounds using indicator analysis. In most species that we examined, there were single androconial compounds that were strong indicators of species identity (Table 1). For example, Geranylgeranylacetone was found only in *H. erato* and was consistently present across samples of this species. Similarily, Octadecanal, a compound previously shown to be electrophysiologically active (Byers et al., unpublished data), was found almost exclusively in *H. melpomene* (specificity=0.999). *H. cydno* and *H. eleuchia* had the weakest indicator scores; in *H. cydno* because the best indicator compound was not found in all individuals examined (coverage= 0.667), and in *H. eleuchia* because the best indicator compound was also found in other species (specificity= 0.747). There were similarly species-specific genital compounds in all species except *H. sapho* and *H. timareta*, where a combination of two compounds was the best predictor (Table 2). Similar to the androconia, in *H. melpomene*, the best indicator compound has known biological activity, in this case the anti-aphrodisiac, (*E*)-β-ocimene (Schulz, Estrada, Yildizham, Boppré, & Gilbert, 2008). For *H. erato* we identified a terpene ester which is only found in *H. erato* individuals and no other species. Other terpene esters were also almost perfect indicator compounds of *H. erato*.

What factors affect interspecific variation in chemical profiles?

Our sampling allowed us to investigate how variation in chemical composition is partitioned within and between species, and determine the extent to which chemistry is a species-diagnostic trait. Visualisation of the chemical profiles reveals that individuals mostly group by species for both androconial and genital chemical bouquets (Fig. 3). Species significantly differ in their androconial bouquet, with species identity accounting for 58% of

the overall variation in chemical profiles (PERMANOVA, Species, $F_{6,251}=74.23$, p<0.001). All pairwise comparisons of species are significant (Table S5). A further 4% of variation can be explained by region (Amazon/Eastern Andes/Western Andes/Panama), and 3% by locality nested within region (PERMANOVA, Region, $F_{3,251}=10.24$, p<0.001; (Region/Locality), $F_{8,251}=2.72$, p<0.001). Finally, 5% of variation is explained by an interaction between species and region, and 2% between species and locality nested within region (PERMANOVA, Species*Region $F_{6,251}=4.96$, p<0.001; Species*(Region/Locality) $F_{8,251}=1.82$, p<0.001).

The results were similar for the genital bouquets, with species identity accounting for 51% of the variation in chemical profiles (PERMANOVA, Species, $F_{6,274}=61.29$, p<0.001). All pairwise comparisons are significant apart from *H. elevatus* and *H. melpomene* (Table S6). A further 5% of variation can be explained by region (Amazon/Eastern Andes/Western Andes/Panama), and 3% by locality nested within region (PERMANOVA, Region, $F_{3,274}=12.73$, p<0.001; (Region/Locality), $F_{8,274}=2.99$, p<0.001). Finally, 6% of variation is explained by an interaction between species and region, and 2% between species and locality nested within region (PERMANOVA, Species*Region $F_{6,274}=6.68$, p<0.001; Species*(Region/Locality) $F_{8,274}=1.77$, p<0.001). For both androconial and genital chemical profiles, most variation is explained by species identity, rather than geographic location, as confirmed by ManyGLM (Tables S7, S8).

Figure 3. NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in chemical compounds of male *Heliconius* of different species. A) Androconial compound bouquets differ significantly between species. Stress=0.155. B) Genital bouquets also differ significantly between species. Stress=0.121.

Does phylogenetic distance explain chemical profile divergence?

Using whole-genome sequence data, we explored the degree to which variation between species can be explained by geographic and genetic distance among the samples. We carried out partial Mantel tests to investigate the correlation between two variables whilst controlling for a third variable. When controlling for geographic distance, genetic and androconial chemical divergence are correlated (Mantel test: r=0.1088, p=0.0001), and there is an even stronger correlation between genetic and genital chemical divergence (Mantel test: r=0.6936, p=0.001). When controlling for genetic distance, geographic distance is significantly but weakly correlated with androconial chemical divergence (Mantel test: r=0.043, p=0.002), with similar results for genital chemical divergence (Mantel test: r=0.046, p=0.007).

Do we find subspecies-specific chemical compounds?

We used an indicator analysis to search for compounds unique to specific populations of *H. erato* and *H. melpomene*. Most intraspecific differences are due to quantitative rather than qualitative differences between populations, perhaps explaining why many population indicators were weak as they are also found in other regions at different amounts (Table S9, S10). The only exception is *H. e. cyrbia* (Western Ecuador) that has many genital compounds unique to this region (Table S9).

What factors affect intraspecific variation in chemical profiles of H. erato *and* H. melpomene*?*

We also wanted to determine the sources of variation within species using our broad sampling of populations across the ranges of *H. erato* and *H. melpomene*. For *H. erato* there

was a strong grouping of individuals by region (Fig. 4), with 27% of variation in androconial profiles being explained by region and 11% by locality nested within region (PERMANOVA, Region $F_{3,87}=11.16$, $p<0.001$, Locality $F_{6,87}=2.35$, $p<0.001$). All four regions are significantly different from each other (Pairwise permutation MANOVAs, $p<0.01$). For *H. erato* genital compounds, 37% of variation is explained by region, and 11% by locality nested within region (PERMANOVA, Region $F_{3,91}=19.01$, $p<0.001$, Locality $F_{6,91}=2.83$, $p<0.01$). All four regions are significantly different from each other (Pairwise permutation MANOVAs, $p<0.05$).

These geographic differences in chemical profiles are not as strong in *H. melpomene* (Fig. 5). For *H. melpomene* androconial compounds, 18% of variation is explained by region and 13% by locality nested within region (PERMANOVA, Region $F_{3,86}=6.73$, $p<0.01$, Locality $F_{8,86}=1.82$, $p<0.001$). The West Andes subspecies (*H. m. cythera*) is not significantly different from either East Andes (multiple populations) or Panama (*H. m. rosina*) (Pairwise permutation MANOVAs, $p=0.072$), however, the other comparisons are significantly different (Pairwise permutation MANOVAs, $p<0.05$). For *H. melpomene* genital compounds, 20% of variation is explained by region, 12% by locality nested within region (PERMANOVA, Region $F_{3,103}=8.91$, $p<0.001$, Locality $F_{7,103}=2.34$, $p<0.001$). All regions are significantly different from each other (Pairwise permutation MANOVAs, $p<0.05$), apart from West Andes and Amazon (Pairwise permutation MANOVAs, $p=0.120$). Within region the amount of variance explained by locality is the same in both *H. erato* and *H. melpomene* but at the larger geographic scale of region, *H. erato* is more structured, with more variance explained. These results were confirmed by ManyGLM tests (Tables S11, S12, S13, S14).

Figure 4. Plots of androconial, genital and genetic distance between *H. erato* populations. A) NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in androconial chemical compounds. Stress=0.174. B) NMDS plot illustrating in three dimensions the variation in genital chemical compounds. Stress=0.118. C) MDS plot illustrating in two dimensions the genetic distance between populations of *H. erato*.

Figure 5. Plots of androconial, genital and genetic distance between *H. melpomene* populations. A) NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in androconial chemical compounds. Stress=0.151. B) NMDS plot illustrating in three dimensions the variation in genital chemical compounds. Stress=0.161. C) MDS plot illustrating in two dimensions the genetic distance between populations of *H. melpomene*.

Does genetic distance explain chemical divergence in H. erato *and* H. melpomene*?*

In *H. erato* both androconial and genital chemical distance are positively correlated with genetic distance, even when accounting for geographic distance (Mantel test, androconia, R=0.255, p=0.001; genitals, R=0.352, p=0.001; partial Mantel test, androconia, R=0.169, p=0.001; genitals, R=0.348, p=0.001). When we account for genetic distance, geographic distance is negatively correlated with androconial chemical distance and not correlated with genital chemical distance (partial Mantel test, androconia, R=-0.149, p=0.001; genitals, R=-0.078, p=0.962).

Heliconius melpomene genitals show similar patterns to *H. erato*, but variation in the androconia is explained by geographic but not genetic distance. Both androconial and genital chemical distance are correlated with genetic distance when tested alone (Mantel test, androconia, R=0.0467, p=0.019; genitals, R=0.146, p=0.001), but when accounting for geography, this correlation is present for the genital bouquet, but disappears for the androconia (partial Mantel test, androconia, R=0.0135, p=0.264, genitals, R=0.120, p=0.001). When we first consider genetic distance, geographic distance is not positively correlated with genital chemical distance, however, it is positively correlated with androconial chemical distance (partial Mantel test, androconia, R=0.231, p=0.001; genitals, R=-0.004, p=0.526).

Figure 6. NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in chemical compounds of male *Heliconius* from Panama and Western Ecuador. *H. erato* and *H. melpomene* are co-mimics (circles), whilst *H. cydno*, *H. eleuchia* and *H. sapho* form a second co-mimicry group (triangles). A) Co-mimicry group contributes 1% of variation in androconial chemical bouquets. (Stress=0.098. B) Co-mimicry group contributes 1.8% of variation in genitals chemical bouquets. Stress=0.094.

Is there evidence for similarity between co-mimics in chemical profile?

Individuals collected in Panama and western Ecuador from two mimicry rings group by species (Fig. 6). In a PERMANOVA model including species, region and co-mimicry group as terms, as well as interactions between these terms, we did not find co-mimicry group to be a significant term. The final model including significant terms included species, region, and an interaction between species and region (PERMANOVA, Species $F_{4,85}=42.94$, p<0.001; Region $F_{1,85}=3.39$, p=0.004; Species*Region $F_{3,85}=2.81$, p=0.002). We confirmed these results using ManyGLM (Table S15).

We found similar trends for genital bouquets. Again, we did not find co-mimicry group to be a significant term, with the final model including species, region, and an interaction between these two (PERMANOVA, Species $F_{4,88}=42.19$, p<0.001; region $F_{1,88}=8.29$, p<0.001; Species*Region $F_{3,88}=5.01$, p<0.001). We confirmed these results using ManyGLM (Table S16). In both PERMANOVA and in multivariate GLMs, variation in both androconial and genital bouquets is explained mostly by the species of the butterfly.

Discussion

Heliconius butterflies are an excellent example of a continental-scale adaptive radiation (Kozak et al., 2015). Speciation in this group is often associated with divergence in wing colour pattern and pattern variation plays an important role in speciation and mate preference (Jiggins, 2008; Jiggins et al., 2001; Merrill et al., 2015, 2011; Sánchez et al., 2015). However, one of the surprising findings to emerge from comparative genomic analysis is the wealth of genes involved in chemosensory reception (Heliconius Genome Consortium, 2012), suggesting that chemical signalling may play an important role in the reproductive biology of the system. To begin to understand the role of chemical signalling in this radiation we have extensively surveyed both inter- and intraspecific variation of *Heliconius*'

androconial and genital chemical profiles across the Neotropics. We find that most of the variation in chemical profile across our samples is explained by species, and we identify key chemicals serving as indicators for each species. Nonetheless, there is also abundant intraspecific variation in chemical profiles. This variation is mainly quantitative in nature, with the exception of *H. erato cyrbia* which has compounds not found in other *H. erato* populations. The hypothesis that co-mimics might also converge in their chemical profiles was not supported. Our work sets the stage for further research into the biology and function of chemical profiles, and their role in within and between-species signalling.

When analysing the chemical profiles across a phylogeny it may not be feasible to carry out behavioural tests in every species. We hypothesised that consistent species-specific compounds are likely to be under stabilising selection and therefore to also be biologically important. In *H. melpomene* the compounds identified as indicators for androconia and genitals, Octadecanal and (*E*)-β-Ocimene respectively, are both known to be biologically active (Schulz et al., 2008). Combining broad geographic sampling with indicator analysis therefore provides a promising approach to determine potential biologically active compounds in other species, which could then be tested behaviourally. Our analyses have already identified a number of compounds in other species that could now be tested functionally, such as the androconial compound Geranylgeranylacetone in *H. erato*.

Chemical profiles are predicted to be highly species-specific if they are involved in species recognition during mating. For instance orchid bee chemical blends, important for mating and species recognition, show high species-specificity, as well as within-species variability, which can be partly explained by geography (Brand et al., 2019; Weber et al., 2016; Zimmermann, Roubik, & Eltz, 2006). We see similar patterns in *Heliconius*, with greater interspecific than intraspecific differences in chemical profiles. Species identity is the best predictor of chemical divergence, with geographic location able to explain some

intraspecific differences. One exception to this is *H. elevatus* which does not group separately from its co-mimic *H. melpomene* for genital compounds, despite the fact that these species are not especially closely related in the *Heliconius* phylogeny. Further samples are needed to confirm that this result is not due to the small sample of *H. elevatus* in this study. As in orchid bees, species differences are often consistent across a large geographic range, suggesting that they could be important for reproductive isolation between species (Weber et al., 2016).

Androconial and genital bouquets show different patterns of evolution. Whilst androconial chemical distance was only weakly correlated with genetic distance between species, genital chemical distance was very strongly correlated. This suggests that neutral evolutionary forces are more important in driving genital chemical evolution. The correlation between genital chemical distance and genetic distance is a much stronger correlation than previously reported (Estrada, Schulz, Yildizhan, & Gilbert, 2011), possibly due to the quantitative nature of our data. This stronger signal of neutrality could be due in part to the fact that there are more compounds in the genitals than androconia, many of which are probably neutrally evolving. For example, in *H. melpomene*, one compound, (*E*)-β-ocimene, can alone act as an anti-aphrodisiac, with other components of the bouquet thought to moderate its evaporation rate (Schulz et al., 2008). Looking at the evolutionary patterns of only the compounds we know to be biologically active, rather than the entire bouquet, may allow us to disentangle the processes involved in the evolution of these profiles.

Heliconius erato and *H. melpomene* both exhibit extensive colour pattern variation across their geographic range (Sheppard, Turner, Brown K. S., Benson W. W., & Singer M. C., 1985). These populations also differ in their androconial and genital bouquets. Whilst traditionally predicted to be under stabilising selection, intraspecific variation between populations in chemical profiles has been documented in other Lepidoptera (Bacquet et al.,

2016; Groot et al., 2009; Huang et al., 1998; McElfresh & Millar, 2001; Takanashi et al., 2005). Chemical divergence in putative male sex pheromones between populations of *Bicyclus anynana* is reported to be as large as differences between *Bicyclus* species, and is greater than predicted by genetic divergence (Bacquet et al., 2016). This is in contrast to what we find here, where interspecific differences are much greater than intraspecific. Interestingly, *Heliconius erato cyrbia*, found west of the Andes, produces many unique genital compounds but is also the most genetically divergent *H. erato* population in our study, which suggests that genetic drift could be important for the evolution of chemical profiles in *Heliconius*. Across all *H. erato* populations we find a correlation between chemical distance, both androconial and genital, and genetic distance. These correlations suggest that the geographic variation between populations could be neutral, with stochastic processes important for bouquet evolution in *Heliconius*.

In *H. melpomene*, correlations between chemical and genetic distance are weaker, with androconial variation better explained by geographic distance. This does not match our expectations of neutrality and might imply that other evolutionary forces are important for chemical profile evolution in *H. melpomene*. One factor is larval host plant use, which affects compound production (Darragh et al., 2019) and varies across the geographic range of *H. melpomene* (Benson, 1978; Benson, Brown, & Gilbert, 1975).

Heliconius butterflies are an excellent example of visual mimicry, with different species converging on the same warning colour patterns (Merrill et al., 2015; Sheppard et al., 1985; Sherratt, 2008). It has been suggested that chemical compounds could also contribute to mimicry between species (Dettner & Liepert, 1994; Mann et al., 2017), but we find no evidence that individuals within co-mimicry groups have more similar chemical profiles than expected stochastically. Most known examples of chemical mimicry come from systems of deception, for example, mimicry of ant alarm pheromones by rove beetles to avoid predation,

rather than mimicry of aposematic warning signals (Dettner & Liepert, 1994; Stoeffler, Maier, Tolasch, & Steidle, 2007; Vereecken & McNeil, 2010). The extreme selection pressure on wing colour and pattern mimicry in *Heliconius* might counteract the potential for chemical mimicry, if chemical bouquets are critical for species recognition in this highly mimetic group.

Even if the androconial and genital chemical compounds are not part of the mimicry, this does not rule out a role in predator deterrence. Genital compounds were originally suggested to form part of the anti-predation signal (Eltringham, 1925). More recently, it has been shown that these compounds act as anti-aphrodisiacs, transferred to females during mating (Gilbert, 1976; Schulz et al., 2008); however, they could be important for both functions. We detected 2-sec-butyl-3-methoxypyrazine in the genitals of *H. melpomene*, *H. cydno* and *H. timareta*, and 2-isobutyl-3-methoxypyrazine in the genitals of *H. melpomene* and *H. cydno*, both compounds known to deter predators in the wood tiger moth (Burdfield-Steel, Pakkanen, Rojas, Galarza, & Mappes, 2018; Rojas et al., 2018, 2017; Rojas, Mappes, & Burdfield-Steel, 2019). More generally, methoxypyrazines act as warning odours in other insects (e.g. Lepidoptera, Rothschild, Moore & Brown, 1984; fireflies, Vencl et al., 2016), effective against avian predators (Guilford, Nicol, Rothschild, & Moore, 1987). Further investigation will be required to determine if odours of *Heliconius* butterflies act as anti-predation signals.

Overall, our study reveals strong species differences in bouquets and the presence of species-specific compounds, as well as intraspecific variation. These results, combined with previous behavioural studies suggest that chemical signalling plays a role in mate choice and reproductive isolation in *Heliconius* butterflies (Darragh et al., 2017; Mérot et al., 2015). A pattern of species-specificity alongside intraspecific variation geographically has also been found in bird song (Benedict & Bowie, 2009). This has been proposed to be a result of a

balance between stabilising selection towards a species stereotype, sexual selection promoting diversity, and geographic segregation alongside selection and drift. Chemical bouquet evolution is also likely to be driven by a combination of these evolutionary processes, both neutral and adaptive. A challenge for the field is the feasibility of testing for the biological relevance of hundreds of compounds in many species, but we hope that our innovative analysis will stimulate not only further targeted functional studies of putatively important compounds, but also large chemical profile surveys in other study systems of evolutionary interest.

References

Bacquet, P. M. B., de Jong, M. A., Brattström, O., Wang, H.-L., Molleman, F., Heuskin, S., …
Nieberding, C. M. (2016). Differentiation in putative male sex pheromone components across
and within populations of the African butterfly *Bicyclus anynana* as a potential driver of
reproductive isolation. *Ecology and Evolution*, *6*(17), 6064–6084. doi: 10.1002/ece3.2298

Benedict, L., & Bowie, R. C. K. (2009). Macrogeographical variation in the song of a widely
distributed African warbler. *Biology Letters*, *5*(4), 484–487. doi: 10.1098/rsbl.2009.0244

Benson, W. W. (1978). Resource partitioning in passion vine butterflies. *Evolution; International
Journal of Organic Evolution*, *32*(3), 493–518. doi: 10.1111/j.1558-5646.1978.tb04593.x

Benson, W. W., Brown, K. S., & Gilbert, L. E. (1975). Coevolution of plants and herbivores: passion
flower butterflies. *Evolution; International Journal of Organic Evolution*, *29*(4), 659–680.
doi: 10.1111/j.1558-5646.1975.tb00861.x

Bolnick, D. I., & Kirkpatrick, M. (2012). The relationship between intraspecific assortative mating
and reproductive isolation between divergent populations. *Current Zoology*, *58*(3), 484–492.
doi: 10.1093/czoolo/58.3.484

Brand, P., Hinojosa-Díaz, I. A., Ayala, R., Daigle, M., Obiols, C. L. Y., Eltz, T., & Ramírez, S. R.
(2019). An olfactory receptor gene underlies reproductive isolation in perfume-collecting
orchid bees. *BioRxiv*, 537423. doi: 10.1101/537423

Burdfield-Steel, E., Pakkanen, H., Rojas, B., Galarza, J. A., & Mappes, J. (2018). De novo Synthesis
of Chemical Defenses in an Aposematic Moth. *Journal of Insect Science*, *18*(2), 1–4. doi:
10.1093/jisesa/iey020

Butlin, R. K., Hewitt, G. M., & Webb, S. F. (1985). Sexual selection for intermediate optimum in
Chorthippus brunneus (Orthoptera: Acrididae). *Animal Behaviour*, *33*(4), 1281–1292. doi:
10.1016/S0003-3472(85)80188-3

Bybee, S. M., Yuan, F., Ramstetter, M. D., Llorente-Bousquets, J., Reed, R. D., Osorio, D., &
Briscoe, A. D. (2012). UV Photoreceptors and UV-Yellow Wing Pigments in *Heliconius*

Butterflies Allow a Color Signal to Serve both Mimicry and Intraspecific Communication. *The American Naturalist*, *179*(1), 38–51. doi: 10.1086/663192

Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of sites: indices and statistical inference. *Ecology*, *90*(12), 3566–3574. doi: 10.1890/08-1823.1

Campbell, P., Pasch, B., Pino, J. L., Crino, O. L., Phillips, M., & Phelps, S. M. (2010). Geographic variation in the songs of neotropical singing mice: testing the relative importance of drift and local adaptation. *Evolution*, *64*(7), 1955–1972. doi: 10.1111/j.1558-5646.2010.00962.x

Challis, R. J., Kumar, S., Dasmahapatra, K. K. K., Jiggins, C. D., & Blaxter, M. (2016). Lepbase: the Lepidopteran genome database. *BioRxiv*, 056994. doi: 10.1101/056994

Conrad, T., Paxton, R. J., Assum, G., & Ayasse, M. (2018). Divergence in male sexual odor signal and genetics across populations of the red mason bee, *Osmia bicornis*, in Europe. *PLOS ONE*, *13*(2), e0193153. doi: 10.1371/journal.pone.0193153

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sunderland, MA: Sinauer.

Darragh, K., Byers, K. J. R. P., Merrill, R. M., McMillan, W. O., Schulz, S., & Jiggins, C. D. (2019). Male pheromone composition depends on larval but not adult diet in Heliconius melpomene. *Ecological Entomology*. doi: 10.1111/een.12716

Darragh, K., Vanjari, S., Mann, F., Gonzalez-Rojas, M. F., Morrison, C. R., Salazar, C., … Jiggins, C. D. (2017). Male sex pheromone components in *Heliconius* butterflies released by the androconia affect female choice. *PeerJ*, *5*, e3953. doi: 10.7717/peerj.3953

Davey, J. W., Barker, S. L., Rastas, P. M., Pinharanda, A., Martin, S. H., Durbin, R., … Jiggins, C. D. (2017). No evidence for maintenance of a sympatric *Heliconius* species barrier by chromosomal inversions. *Evolution Letters*, *1*(3), 138–154. doi: 10.1002/evl3.12

Dettner, K., & Liepert, C. (1994). Chemical Mimicry and Camouflage. *Annual Review of Entomology*, *39*(1), 129–154. doi: 10.1146/annurev.en.39.010194.001021

Dray, S., & Dufour, A. B. (2007). The ade4 package: implementing the duality diagram for ecologists. *Journal of Statistical Software*, *22*(4), 1–20. doi: 10.18637/jss.v022.i04

Eltringham, H. (1925). On the Abdominal Glands in *Heliconius* (Lepidoptera). *Transactions of the Royal Entomological Society of London*, *73*(1–2), 269–275. doi: 10.1111/j.1365-2311.1925.tb02862.x

Estrada, C., & Jiggins, C. D. (2008). Interspecific sexual attraction because of convergence in warning colouration: is there a conflict between natural and sexual selection in mimetic species? *Journal of Evolutionary Biology*, *21*(3), 749–760. doi: 10.1111/j.1420-9101.2008.01517.x

Estrada, C., Schulz, S., Yildizhan, S., & Gilbert, L. E. (2011). Sexual selection drives the evolution of antiaphrodisiac pheromones in butterflies. *Evolution; International Journal of Organic Evolution*, *65*(10), 2843–2854. doi: 10.1111/j.1558-5646.2011.01352.x

Ferreira, M., & Ferguson, J. W. H. (2002). Geographic variation in the calling song of the field cricket *Gryllus bimaculatus* (Orthoptera: Gryllidae) and its relevance to mate recognition and mate choice. *Journal of Zoology*, *257*(2), 163–170. doi: 10.1017/S0952836902000766

Finkbeiner, S. D., Fishman, D. A., Osorio, D., & Briscoe, A. D. (2017). Ultraviolet and yellow reflectance but not fluorescence is important for visual discrimination of conspecifics by Heliconius erato. *Journal of Experimental Biology*, *220*(7), 1267–1276. doi: 10.1242/jeb.153593

Friberg, M., Vongvanich, N., Borg-Karlson, A.-K., Kemp, D. J., Merilaita, S., & Wiklund, C. (2008). Female mate choice determines reproductive isolation between sympatric butterflies. *Behavioral Ecology and Sociobiology*, *62*(6), 873–886. doi: 10.1007/s00265-007-0511-2

Gemeno, C., Lutfallah, A. F., & Haynes, K. F. (2000). Pheromone Blend Variation and Cross-Attraction Among Populations of the Black Cutworm Moth (Lepidoptera: Noctuidae). *Annals of the Entomological Society of America*, *93*(6), 1322–1328. doi: 10.1603/0013-8746(2000)093[1322:PBVACA]2.0.CO;2

Gerhardt, H. C. (1982). Sound Pattern Recognition in Some North American Treefrogs (Anura: Hylidae): Implications for Mate Choice. *Integrative and Comparative Biology*, *22*(3), 581–595. doi: 10.1093/icb/22.3.581

Gilbert, L. E. (1976). Postmating female odor in *Heliconius* butterflies: a male-contributed

   antiaphrodisiac? *Science*, *193*(4251), 419–420. doi: 10.1126/science.935877

Giraldo, N., Salazar, C., Jiggins, C. D., Bermingham, E., & Linares, M. (2008). Two sisters in the

   same dress: *Heliconius* cryptic species. *BMC Evolutionary Biology*, *8*, 324. doi:

   10.1186/1471-2148-8-324

Gray, D. A., & Cade, W. H. (2000). Sexual selection and speciation in field crickets. *Proceedings of

   the National Academy of Sciences*, *97*(26), 14449–14454. doi: 10.1073/pnas.97.26.14449

Groot, A. T., Inglis, O., Bowdridge, S., Santangelo, R. G., Blanco, C., Jr, J. D. L., … Schal, C. (2009).

   Geographic and Temporal Variation in Moth Chemical Communication. *Evolution*, *63*(8),

   1987–2003. doi: 10.1111/j.1558-5646.2009.00702.x

Guilford, T., Nicol, C., Rothschild, M., & Moore, B. P. (1987). The biological roles of pyrazines:

   evidence for a warning odour function. *Biological Journal of the Linnean Society*, *31*(2),

   113–128. doi: 10.1111/j.1095-8312.1987.tb01984.x

Hankison, S. J., & Ptacek, M. B. (2008). Geographical variation of genetic and phenotypic traits in the

   Mexican sailfin mollies, *Poecilia velifera* and *P. petenensis*. *Molecular Ecology*, *17*(9), 2219–

   2233. doi: 10.1111/j.1365-294X.2008.03736.x

Heliconius Genome Consortium. (2012). Butterfly genome reveals promiscuous exchange of mimicry

   adaptations among species. *Nature*, *487*(7405), 94–98. doi: 10.1038/nature11041

Hervé, M. (2018). *RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R package

   version 0.9-69-3.* Retrieved from https://CRAN.R-project.org/package=RVAideMemoire

Hijmans, R. J. (2017). *geosphere: Spherical Trigonometry. R package version 1.5-7*. Retrieved from

   https://CRAN.R-project.org/package=geosphere

Huang, Y., Takanashi, T., Hoshizaki, S., Tatsuki, S., Honda, H., Yoshiyasu, Y., & Ishikawa, Y.

   (1998). Geographic Variation in Sex Pheromone of Asian Corn Borer, *Ostrinia furnacalis*, in

   Japan. *Journal of Chemical Ecology*, *24*(12), 2079–2088. doi: 10.1023/A:1020737726636

Irwin, D. E., Thimgan, M. P., & Irwin, J. H. (2008). Call divergence is correlated with geographic and

   genetic distance in greenish warblers (*Phylloscopus trochiloides*): a strong role for

stochasticity in signal evolution? *Journal of Evolutionary Biology*, *21*(2), 435–448. doi: 10.1111/j.1420-9101.2007.01499.x

Jiggins, C. D. (2008). Ecological Speciation in Mimetic Butterflies. *BioScience*, *58*(6), 541–548. doi: 10.1641/B580610

Jiggins, C. D. (2017). *The Ecology and Evolution of* Heliconius *Butterflies*. Oxford: Oxford University Press.

Jiggins, C. D., Naisbit, R. E., Coe, R. L., & Mallet, J. (2001). Reproductive isolation caused by colour pattern mimicry. *Nature*, *411*(6835), 302–305. doi: 10.1038/35077075

Johansson, B. G., & Jones, T. M. (2007). The role of chemical communication in mate choice. *Biological Reviews*, *82*(2), 265–289. doi: 10.1111/j.1469-185X.2007.00009.x

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. doi: 10.1186/s12859-014-0356-4

Kozak, K. M., Wahlberg, N., Neild, A. F. E., Dasmahapatra, K. K., Mallet, J., & Jiggins, C. D. (2015). Multilocus Species Trees Show the Recent Adaptive Radiation of the Mimetic *Heliconius* Butterflies. *Systematic Biology*, *64*(3), 505–524. doi: 10.1093/sysbio/syv007

Lamas, G., & Jiggins, C. D. (2017). Taxonomic list. In *The ecology and evoution of Heliconius butterflies* (pp. 214–244). Oxford: Oxford University Press.

Leonhardt, S., Rasmussen, C., & Schmitt, T. (2013). Genes versus environment: Geography and phylogenetic relationships shape the chemical profiles of stingless bees on a global scale. *Proceedings. Biological Sciences / The Royal Society*, *280*, 20130680. doi: 10.1098/rspb.2013.0680

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics (Oxford, England)*, *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324

Löfstedt, C. (1993). Moth Pheromone Genetics and Evolution. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, *340*(1292), 167–177. doi: 10.1098/rstb.1993.0055

Mann, F., Vanjari, S., Rosser, N., Mann, S., Dasmahapatra, K. K., Corbin, C., … Schulz, S. (2017). The Scent Chemistry of *Heliconius* Wing Androconia. *Journal of Chemical Ecology*, *43*(9), 843–857. doi: 10.1007/s10886-017-0867-3

Martin, M. D., & Mendelson, T. C. (2016). The accumulation of reproductive isolation in early stages of divergence supports a role for sexual selection. *Journal of Evolutionary Biology*, *29*(4), 676–689. doi: 10.1111/jeb.12819

Mas, F., & Jallon, J.-M. (2005). Sexual Isolation and Cuticular Hydrocarbon Differences between *Drosophila santomea* and *Drosophila yakuba*. *Journal of Chemical Ecology*, *31*(11), 2747–2752. doi: 10.1007/s10886-005-7570-5

McElfresh, J. S., & Millar, J. G. (2001). Geographic Variation in the Pheromone System of the Saturniid Moth *Hemileuca eglanterina*. *Ecology*, *82*(12), 3505–3518. doi: 10.1890/0012-9658(2001)082[3505:GVITPS]2.0.CO;2

McPeek, M. A., Symes, L. B., Zong, D. M., & McPeek, C. L. (2011). Species recognition and patterns of population variation in the reproductive structures of a damselfly genus. *Evolution; International Journal of Organic Evolution*, *65*(2), 419–428. doi: 10.1111/j.1558-5646.2010.01138.x

Mendelson, T. C., & Shaw, K. L. (2012). The (mis)concept of species recognition. *Trends in Ecology & Evolution*, *27*(8), 421–427. doi: 10.1016/j.tree.2012.04.001

Mérot, C., Frérot, B., Leppik, E., & Joron, M. (2015). Beyond magic traits: Multimodal mating cues in *Heliconius* butterflies. *Evolution*, *69*(11), 2891–2904. doi: 10.1111/evo.12789

Mérot, C., Mavárez, J., Evin, A., Dasmahapatra, K. K., Mallet, J., Lamas, G., & Joron, M. (2013). Genetic differentiation without mimicry shift in a pair of hybridizing *Heliconius* species (Lepidoptera: Nymphalidae). *Biological Journal of the Linnean Society*, *109*(4), 830–847. doi: 10.1111/bij.12091

Merrill, R. M., Chia, A., & Nadeau, N. J. (2014). Divergent warning patterns contribute to assortative mating between incipient *Heliconius* species. *Ecology and Evolution*, *4*(7), 911–917. doi: 10.1002/ece3.996

Merrill, R. M., Dasmahapatra, K. K., Davey, J. W., Dell′Aglio, D. D., Hanly, J. J., Huber, B., … Yu, Q. (2015). The diversification of *Heliconius* butterflies: what have we learned in 150 years? *Journal of Evolutionary Biology*, *28*(8), 1417–1438. doi: 10.1111/jeb.12672

Merrill, R. M., Gompert, Z., Dembeck, L. M., Kronforst, M. R., McMillan, W. O., & Jiggins, C. D. (2011). Mate Preference Across the Speciation Continuum in a Clade of Mimetic Butterflies. *Evolution*, *65*(5), 1489–1500. doi: 10.1111/j.1558-5646.2010.01216.x

Mullen, L. M., Vignieri, S. N., Gore, J. A., & Hoekstra, H. E. (2009). Adaptive basis of geographic variation: genetic, phenotypic and environmental differences among beach mouse populations. *Proceedings of the Royal Society of London B: Biological Sciences*, *276*(1674), 3809–3818. doi: 10.1098/rspb.2009.1146

Nagel, L., & Schluter, D. (1998). Body Size, Natural Selection, and Speciation in Sticklebacks. *Evolution*, *52*(1), 209–218. doi: 10.2307/2410936

Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner, H. (2017). *vegan: Community Ecology Package. R package version 2.4-2*. Retrieved from https://CRAN.R-project.org/package=vegan

Paradis, E., & Schliep, K. (2018). ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*. doi: 10.1093/bioinformatics/bty633

Pfennig, K. S. (1998). The evolution of mate choice and the potential for conflict between species and mate–quality recognition. *Proceedings of the Royal Society of London B: Biological Sciences*, *265*(1407), 1743–1748. doi: 10.1098/rspb.1998.0497

Ptacek, M. (2000). The role of mating preferences in shaping interspecific divergence in mating signals in vertebrates. *Behavioural Processes*, *51*(1–3), 111–134. doi: 10.1016/S0376-6357(00)00123-6

R Core Team. (2018). R: A language and environment for statistical computing. (Version 3.3.1). Retrieved from http://www.R-project.org

Ready, J. S., Sampaio, I., Schneider, H., Vinson, C., Santos, T. D., & Turner, G. F. (2006). Colour forms of Amazonian cichlid fish represent reproductively isolated species. *Journal of Evolutionary Biology*, *19*(4), 1139–1148. doi: 10.1111/j.1420-9101.2006.01088.x

Rojas, B., Burdfield-Steel, E., De Pasqual, C., Gordon, S., Hernández, L., Mappes, J., … Lindstedt, C. (2018). Multimodal Aposematic Signals and Their Emerging Role in Mate Attraction. *Frontiers in Ecology and Evolution*, *6*, 93. doi: 10.3389/fevo.2018.00093

Rojas, B., Burdfield-Steel, E., Pakkanen, H., Suisto, K., Maczka, M., Schulz, S., & Mappes, J. (2017). How to fight multiple enemies: target-specific chemical defences in an aposematic moth. *Proceedings. Biological Sciences*, *284*(1863), 20171424. doi: 10.1098/rspb.2017.1424

Rojas, B., Mappes, J., & Burdfield-Steel, E. (2019). Multiple modalities in insect warning displays have additive effects against wild avian predators. *Behavioral Ecology and Sociobiology*, *73*(3), 37. doi: 10.1007/s00265-019-2643-6

Rothschild, M. (1961). Defensive Odours and Mullerian Mimicry Among Insects. *Transactions of the Royal Entomological Society of London*, *113*(5), 101–123. doi: 10.1111/j.1365-2311.1961.tb00803.x

Rothschild, M., Moore, B. P., & Brown, W. V. (1984). Pyrazines as warning odour components in the Monarch butterfly, *Danaus plexippus*, and in moths of the genera *Zygaena* and *Amata* (Lepidoptera). *Biological Journal of the Linnean Society*, *23*(4), 375–380. doi: 10.1111/j.1095-8312.1984.tb00153.x

Ryan, M. A., Moore, C. J., & Walter, G. H. (1995). Individual variation in pheromone composition in *Nezara viridula* (Heteroptera: Pentatomidae): how valid is the basis for designating "pheromone strains"? *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, *111*(2), 189–193. doi: 10.1016/0305-0491(94)00257-U

Ryan, M. J., & Guerra, M. A. (2014). The mechanism of sound production in túngara frogs and its role in sexual selection and speciation. *Current Opinion in Neurobiology*, *28*, 54–59. doi: 10.1016/j.conb.2014.06.008

Ryan, M. J., & Rand, A. S. (1993). Species Recognition and Sexual Selection as a Unitary Problem in Animal Communication. *Evolution*, *47*(2), 647–657. doi: 10.2307/2410076

Ryan, M. J., Rand, A. S., & Weigt, L. A. (1996). Allozyme and Advertisement Call Variation in the Túngara Frog, *Physalaemus Pustulosus*. *Evolution*, *50*(6), 2435–2453. doi: 10.1111/j.1558-5646.1996.tb03630.x

Sánchez, A. P., Pardo-Diaz, C., Enciso-Romero, J., Muñoz, A. G., Jiggins, C. D., Salazar, C., & Linares, M. (2015). An introgressed wing pattern acts as a mating cue. *Evolution; International Journal of Organic Evolution*, *69*(6), 1619–1629. doi: 10.1111/evo.12679

Saveer, A. M., Becher, P. G., Birgersson, G., Hansson, B. S., Witzgall, P., & Bengtsson, M. (2014). Mate recognition and reproductive isolation in the sibling species *Spodoptera littoralis* and *Spodoptera litura*. *Chemical Ecology*, *2*, 18. doi: 10.3389/fevo.2014.00018

Schulz, S., Estrada, C., Yildizham, S., Boppré, M., & Gilbert, L. E. (2008). An antiaphrodisiac in *Heliconius melpomene* butterflies. *Journal of Chemical Ecology*, *34*(1), 82–93. doi: 10.1007/s10886-007-9393-z

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., … Okada, N. (2008). Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), 620–626. doi: 10.1038/nature07285

Selz, O. M., Pierotti, M. E. R., Maan, M. E., Schmid, C., & Seehausen, O. (2014). Female preference for male color is necessary and sufficient for assortative mating in 2 cichlid sister species. *Behavioral Ecology*, *25*(3), 612–626. doi: 10.1093/beheco/aru024

Sheppard, P. M., Turner, J. R. G., Brown K. S., Benson W. W., & Singer M. C. (1985). Genetics and the evolution of Muellerian mimicry in *Heliconius* butterflies. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, *308*(1137), 433–610. doi: 10.1098/rstb.1985.0066

Sherratt, T. N. (2008). The evolution of Müllerian mimicry. *Die Naturwissenschaften*, *95*(8), 681–695. doi: 10.1007/s00114-008-0403-y

Smadja, C. M., & Butlin, R. K. (2008). On the scent of speciation: the chemosensory system and its role in premating isolation. *Heredity*, *102*(1), 77–97. doi: 10.1038/hdy.2008.55

Stoeffler, M., Maier, T. S., Tolasch, T., & Steidle, J. L. M. (2007). Foreign-language Skills in Rove-Beetles? Evidence for Chemical Mimicry of Ant Alarm Pheromones in Myrmecophilous *Pella* Beetles (Coleoptera: Staphylinidae). *Journal of Chemical Ecology*, *33*(7), 1382–1392. doi: 10.1007/s10886-007-9315-0

Sun, K., Luo, L., Kimball, R. T., Wei, X., Jin, L., Jiang, T., … Feng, J. (2013). Geographic Variation
in the Acoustic Traits of Greater Horseshoe Bats: Testing the Importance of Drift and
Ecological Selection in Evolutionary Processes. *PLOS ONE*, *8*(8), e70368. doi:
10.1371/journal.pone.0070368

Takanashi, T., Huang, Y., Takahasi, K. R., Hoshizaki, S., Tatsuki, S., & Ishikawa, Y. (2005). Genetic
analysis and population survey of sex pheromone variation in the adzuki bean borer moth,
Ostrinia scapulalis. *Biological Journal of the Linnean Society*, *84*(1), 143−160. doi:
10.1111/j.1095-8312.2005.00421.x

Van Belleghem, S. M., Rastas, P., Papanicolaou, A., Martin, S. H., Arias, C. F., Supple, M. A., …
Papa, R. (2017). Complex modular architecture around a simple toolkit of wing pattern genes.
Nature Ecology & Evolution, *1*(3), 52. doi: 10.1038/s41559-016-0052

Vencl, F. V., Ottens, K., Dixon, M. M., Candler, S., Bernal, X. E., Estrada, C., & Page, R. A. (2016).
Pyrazine emission by a tropical firefly: An example of chemical aposematism? *Biotropica*,
48(5), 645−655. doi: 10.1111/btp.12336

Vereecken, N. J., & McNeil, J. N. (2010). Cheaters and liars: chemical mimicry at its finest.
Canadian Journal of Zoology, *88*(7), 725−752. doi: 10.1139/Z10-040

Wang, Y., Naumann, U., Wright, S. T., & Warton, D. I. (2012). mvabund− an R package for model-
based analysis of multivariate abundance data. *Methods in Ecology and Evolution*, *3*(3), 471−
474. doi: 10.1111/j.2041-210X.2012.00190.x

Warton, D. I., Wright, S. T., & Wang, Y. (2012). Distance-based multivariate analyses confound
location and dispersion effects. *Methods in Ecology and Evolution*, *3*(1), 89−101. doi:
10.1111/j.2041-210X.2011.00127.x

Weber, M. G., Mitko, L., Eltz, T., & Ramírez, S. R. (2016). Macroevolution of perfume signalling in
orchid bees. *Ecology Letters*, *19*(11), 1314−1323. doi: 10.1111/ele.12667

Wong, B. (2011). Points of view: Color blindness. *Nature Methods*, *8*, 441. doi: 10.1038/nmeth.1618

Wyatt, T. D. (2003). *Pheromones and Animal Behaviour: Communication by Smell and Taste*.
Cambridge: Cambridge University Press.

Wyatt, T. D. (2014). *Pheromones and Animal Behavior: Chemical Signals and Signatures.*

    Cambridge: Cambridge University Press.

Yildizhan, S., van Loon, J., Sramkova, A., Ayasse, M., Arsene, C., ten Broeke, C., & Schulz, S.

    (2009). Aphrodisiac pheromones from the wings of the small cabbage white and large

    cabbage white butterflies, *Pieris rapae* and *Pieris brassicae*. *ChemBioChem*, *10*(10), 1666–

    1677. doi: 10.1002/cbic.200900183

Zimmermann, Y., Roubik, D. W., & Eltz, T. (2006). Species-specific attraction to pheromonal

    analogues in orchid bees. *Behavioral Ecology and Sociobiology*, *60*(6), 833. doi:

    10.1007/s00265-006-0227-8

Table 1: Androconial compounds which are the best indicators of species identity. A is a measure of group specificity of the compounds, B is a measure of group coverage, and sqrtIV is the indicator value which considers both A and B and ranges from 0 (compound not present in any individuals of that species) to 1 (compound only present in that species, and present in all individuals).

Species/compound	A: specificity	B: coverage	sqrtIV
H. cydno			
Unknown aromatic (RI=2130)	1	0.667	0.816

H. eleuchia

Hexahydrofarnesylacetone	0.747	1	0.864

H. elevatus

Homovanillylalcohol	0.912	1	0.955

H. erato

Geranylgeranylacetone	1	1	1

H. melpomene

Octadecanal	0.999	1	1

H. sapho

Methyl 4-hydroxy-3-methoxybenzoate	0.866	1	0.931

H. timareta

5-Decanolide	1	0.889	0.943

Table 2: Genital compounds which are the best indicators of species identity. A, B, and sqrtIV as in Table 1.

Species/compound	A: specificity	B: coverage	sqrtIV
H. cydno			
Unknown ester (RI=1390)	0.999	1	0.999

H. eleuchia

Unknown macrolide RI=1878	0.969	1	0.984

H. elevatus

Icosenol	0.908	1	0.953

H. erato

Unknown terpene ester (RI=2494)	1	1	1

H. melpomene

(*E*)-β-ocimene	0.865	1	0.930

H. sapho

(*Z*)-3-hexenyl isobutyrate & Unknown (RI=1691)	0.957	0.923	0.940

H. timareta

Butyl oleate & (*Z*)-9-Octadecen-13-olide	0.915	1	0.956