Species specificity and intraspecific variation in the chemical profiles of *Heliconius* butterflies across a large geographic range

Kathy Darragh[1,2], Gabriela Montejo-Kovacevich[1], Krzysztof M. Kozak[2], Colin R. Morrison[2,3], Clarisse M. E. Figueiredo[4], Jonathan S. Ready[4], Camilo Salazar[5], Mauricio Linares[5], Kelsey J. R. P. Byers[1,2], Richard M. Merrill[2,6], W. Owen McMillan[2], Stefan Schulz[7], Chris D. Jiggins[1,2]

[1]*Department of Zoology, University of Cambridge, Cambridge, Cambridgeshire, United Kingdom*

[2]*Smithsonian Tropical Research Institute, Panama City, Panama*

[3]*Department of Integrative Biology, The University of Texas at Austin, Austin, Texas, United States*

[4]*Institute for Biological Sciences, Universidade Federal do Pará, Belém, Pará, Brazil*

[5]*Biology Program, Faculty of Natural Sciences and Mathematics, Universidad del Rosario, Bogota, Colombia*

[6]*Division of Evolutionary Biology, Faculty of Biology, Ludwig-Maximilians-Universität München, Munich, Germany*

[7]*Institute of Organic Chemistry, Technische Universität Braunschweig, Braunschweig, Germany*

Running title

Geographic variation in butterfly pheromones

Corresponding author

Kathy Darragh

Department of Zoology

Downing Street

Cambridge

CB23EJ

kd350@cam.ac.uk

+44 (0)1223 336600

Abstract

In many animals, mate choice is important for the maintenance of reproductive isolation between species. Traits important for mate choice and behavioural isolation are predicted to be under strong stabilising selection within species, however such traits can also exhibit variation at the population level driven by neutral and adaptive evolutionary processes. Here, we describe patterns of divergence among androconial and genital chemical profiles at inter- and intra-specific levels in mimetic *Heliconius* butterflies. Most variation in chemical bouquets was found between species, but there were also quantitative differences at the population level. We found a strong correlation between interspecific chemical and genetic divergence, however, this correlation varied in intraspecific comparisons. We identified 'indicator' compounds characteristic of particular species that included known biologically active compounds, suggesting an approach for identification of candidate compounds for future behavioural studies in novel systems. Overall, the strong signal of species identity suggests a role for these compounds in species recognition, but with additional potentially neutral variation at the population level.

Keywords

Mate choice, reproductive isolation, chemical ecology, Lepidoptera, pheromones, signalling

Introduction

Reproductive isolation between lineages is important for the maintenance of species

diversity (Coyne & Orr, 2004). In many animals, mate choice provides a strong pre-mating

barrier, maintaining reproductive isolation (Friberg et al., 2008; Gray & Cade, 2000; Martin

& Mendelson, 2016; Nagel & Schluter, 1998; Ready et al., 2006; Seehausen et al., 2008;

Selz, Pierotti, Maan, Schmid, & Seehausen, 2014). Closely related species often differ in

traits important for mate choice, with individuals displaying a preference for conspecific

phenotypes (Jiggins, Naisbit, Coe, & Mallet, 2001; Mas & Jallon, 2005; M. J. Ryan &

Guerra, 2014; Saveer et al., 2014; Yildizhan et al., 2009). These traits are predicted to show

strong species-specific differences (Gerhardt, 1982), and typically should be subject to

stabilising selection which can act to decrease intraspecific phenotypic variation (Butlin,

Hewitt, & Webb, 1985; Pfennig, 1998; Ptacek, 2000). As a consequence, we would expect to

find little trait variability, or at least certain features to be invariant, across species geographic

ranges (Benedict & Bowie, 2009; Ferreira & Ferguson, 2002; McPeek, Symes, Zong, &

McPeek, 2011; Weber, Mitko, Eltz, & Ramírez, 2016). However, these traits can also exhibit

variation both within and between populations of the same species, either due to genetic drift

and/or varying selective regimes across their ranges (Bolnick & Kirkpatrick, 2012; M. J.

Ryan & Guerra, 2014; M. J. Ryan & Rand, 1993; M. J. Ryan, Rand, & Weigt, 1996).

Signals important for behavioural isolation could arise from the divergence of traits

used in intraspecific communication between populations (Johansson & Jones, 2007;

Mendelson & Shaw, 2012; M. J. Ryan & Rand, 1993; Smadja & Butlin, 2008). Signal

divergence can be driven by various factors, both neutral and adaptive, usually involving

multiple evolutionary forces (Leonhardt, Rasmussen, & Schmitt, 2013; Sun et al., 2013). A

positive correlation between genetic distance and phenotypic variation is consistent with

stochastic processes, such as genetic drift, playing a prominent role (Irwin, Thimgan, &

Irwin, 2008). In contrast, a lack of correlation between phenotypic and genetic divergence

suggests that selection is shaping the phenotypic variation, perhaps driving divergence in

different directions in each population (Campbell et al., 2010; Conrad, Paxton, Assum, &

Ayasse, 2018; Hankison & Ptacek, 2008; Mullen, Vignieri, Gore, & Hoekstra, 2009).

Chemical compounds, such as sex pheromones, mediate intraspecific communication

in many systems (Wyatt, 2003, 2014). The role of chemical signalling in behavioural

isolation is also well established, especially among moth species (Löfstedt, 1993; Smadja &

Butlin, 2008). Due to the coordination between detection and production, these pheromone

blends are traditionally regarded as being under stabilising selection towards a species

stereotype (Löfstedt 1993). Even when species-specific characteristics are present, chemical

compounds can exhibit intraspecific variation, with both qualitative and quantitative

differences found across a species range (Conrad et al., 2018; Gemeno, Lutfallah, & Haynes,

2000; Groot et al., 2009; Huang et al., 1998; McElfresh & Millar, 2001; M. A. Ryan, Moore,

& Walter, 1995; Weber et al., 2016).

Studies of *Heliconius* butterflies have contributed to our understanding of adaptation

and speciation (Jiggins, 2008, 2017; Merrill et al., 2015). Despite the reliance of this group

on visual cues for mating (Bybee et al., 2012; Finkbeiner, Fishman, Osorio, & Briscoe, 2017;

Jiggins et al., 2001; Merrill, Chia, & Nadeau, 2014; Sánchez et al., 2015), it has long been

suggested that male pheromones also play a role in pre-mating barriers (Jiggins, 2008;

Merrill et al., 2015), but so far only a few species have been studied. Behavioural

experiments reveal that chemical signalling in *Heliconius erato, H. melpomene* and *H.*

timareta is important for female mate choice (Darragh et al., 2017; Mérot, Frérot, Leppik, &

Joron, 2015). Electroantennographic studies have shown that *Heliconius cydno* and *H.*

melpomene respond to both con- and heterospecific androconial chemical bouquets (Byers et

al., 2019), and have identified individual compounds that are biologically active.

Furthermore, studies of *H. cydno*, *H. doris*, *H. hecale*, *H. ismenius*, *H. melpomene*, *H.*

pardalinus, *H. sara* and *H. timareta* from Panama, Colombia and Ecuador found that major

compounds differ between species (Mann et al., 2017; Mérot et al., 2015), suggesting a

potential role in reproductive isolation.

The role of chemical signalling is likely to be especially important in co-mimics,

where visual signals alone are not sufficient to identify conspecifics (C. Estrada & Jiggins,

2008; Giraldo, Salazar, Jiggins, Bermingham, & Linares, 2008; Mérot et al., 2013; Sánchez

et al., 2015). In addition, chemical compounds could be part of a multimodal aposematic

warning signal (Rojas et al., 2018; Rothschild, 1961), with some tentative evidence that co-

mimics exhibit similar chemical bouquets to aid recognition by predators (Mann et al., 2017).

Here, we describe the chemical profiles of seven species of *Heliconius* from over 250

individuals collected across the Neotropics. We focus on the co-mimetic species *H.*

melpomene and *H. erato* that are distributed widely across the Neotropics and analysed both

wing androconial and genital compounds of male butterflies. It would be challenging to

conduct electrophysiological experiments on large numbers of species and populations, and

therefore identifying the biologically active components in pheromone blends across a

radiation is beyond the scope of a single study. Instead, we hypothesise that biologically

relevant signals for species recognition are likely to be under stronger stabilising selection

compared to other compounds, and therefore the consistency of phenotypic variation across

the geographic range of a species might provide an initial approach to detect putatively

biologically active compounds important for mate choice. We use *H. melpomene* as a test-

species due to the availability of behavioural and electrophysiological data to investigate this

approach, by evaluating consistency in compound blends across different localities.

The extensive dataset analysed here allows us to test evolutionary hypotheses, as well

as identifying interesting candidate compounds for future behavioural studies. Firstly, we

investigated intraspecific variation in chemical profiles of *H. melpomene* and *H. erato* and

correlated chemical profile data with both geographic and genetic distances between different

localities. Furthermore, to test if the chemical compounds are part of the aposematic co-

mimicry signal we sampled two different mimicry rings in western Ecuador and Panama.

Materials and Methods

130   *Sampling*

131   Between February 2016 and August 2017 wild males of *Heliconius cydno*, *H.*

132   *elevatus*, *H. eleuchia*, *H. erato*, *H. melpomene*, *H. sapho* and *H. timareta* were collected from

133   twelve localities. Between two and fifteen males were chemically analysed per population

134   (Fig. 1, Table S1). We follow the latest *Heliconius* taxonomy (Lamas & Jiggins, 2017).

Figure 1. Map indicating species and populations from twelve localities across the Neotropics. *H. erato* (1-11) 1) *H. e. demophoon* 2) *H. e. dignus* 3) *H. e. lativitta* 4) *H. e. cyrbia* 5) *H. e. microclea* 6) *H. e. emma* 7) *H. e. amphitrite* 8) *H. e. hydara* 9) *H. e. amazona* 10) *H. e. phyllis* 11) *H. e. venustus. H. melpomene* (12-23) 12) *H. m. rosina* 13) *H. m. bellula* 14) *H. m malleti* 15) *H. m cythera* 16) *H. m. xenoclea* 17) *H. m. schunkei* 18) *H. m. melpomene* 19) *H. m. madeira* 20) *H. m. thelxiope* 21) *H. m. intersectus* 22) *H. m. penelope* 23) *H. m. burchelli. H. timareta* (24-25) 24) *H. t. tristero* 25) *H. t. nov. spp. H. elevatus* (26-28) 26) *H. e. willmotti* 27) *H. e. schmassmanni* 28) *H. e. perchlora. H. cydno* (29-30) 29) *H. c. chioneus* 30) *H. c. alithea. H. sapho* (31-32) 31) *H. s. sapho* 32) *H. s. candidus. H. eleuchia* 33) *H. e. primularis*. See Table S1 for sample numbers.

 Extraction and chemical analysis of tissues

The androconial region of the wing, previously described as the grey-brown

overlapping region of the hindwing (Darragh et al., 2017), as well as the genitalia, were

dissected for analysis immediately after collection. To extract pheromones, the tissue was

soaked in 200μl dichloromethane containing 200ng 2-tetradecyl acetate (internal standard) in

2ml glass vials with PTFE-coated caps (Agilent, Santa Clara, USA) for one hour. The solvent

was then transferred to new vials, maintained cool in the field and stored at -20°C upon

return. Androconial samples were evaporated to a reduced volume at room temperature prior

to analysis. Pheromone extracts were analysed by GC/MS using an Agilent model 5977

mass-selective detector connected to an Agilent GC model 7890B and equipped with an

Agilent ALS 7693 autosampler. HP-5MS fused silica capillary columns (Agilent, Santa

Clara, USA, 30 m $\times$ 0.25 mm, 0.25 µm) were used. Injection was performed in splitless mode

(250°C injector temperature) with helium as the carrier gas (constant flow of 1.2 ml/min).

The temperature programme started at 50°C, was held for 5 minutes, and then rose at a rate of

5°C/minute to 320°C, before being held at 320°C for 5 minutes. Components were identified

by comparison of mass spectra and gas chromatographic retention index with those of

authentic reference samples and also by analysis of mass spectra. Components were

quantified using 2-tetradecyl acetate as an internal standard. Only compounds eluting earlier

than hexacosane were analysed in androconial samples, and those earlier than nonacosane in

genital samples (Darragh et al. 2017). We removed compounds that were not found in at least

half of all individuals from a given population.

Calculation of genetic and geographic distance matrices

To explore genetic distance among the studied *H. erato* and *H. melpomene*

subspecies, we computed genetic covariance matrices and performed MDS for each species

separately. A whole-genome sequence from a representative individual from each subspecies

was used (Table S2). Genotypes were inferred from reads mapped to the *H. melpomene*

(v2.5) and *H. erato demophoon* genome scaffolds (Challis, Kumar, Dasmahapatra, Jiggins, &

Blaxter, 2016; Davey et al., 2017; Heliconius Genome Consortium, 2012; Van Belleghem et

al., 2017) with bwa v0.7.15 (Li & Durbin, 2009). We computed a pairwise identical by state

(IBS) matrix with a random sampled read from each position in the genome, implemented in

ANGSD v0.912 (Korneliussen, Albrechtsen, & Nielsen, 2014). An interspecific genetic

distance matrix was constructed using the function "cophenetic.phylo" from the *ape* package

(Paradis & Schliep, 2018) with a previously published phylogeny (Kozak et al., 2015).

Geographic distance matrices were created by inputting the co-ordinates of collection

localities into the function "distm" in the *geosphere* package to calculate the Haversine great-

circle-distance between points (Hijmans, 2017).

Statistical analyses

Inter- and intra-specific indicator compounds

We carried out indicator analysis using the *indicspecies* package (Cáceres &

Legendre, 2009). Groupings are decided *a priori* (in this case species or sub-species) and

compounds are determined which act as indicators of these groups. The best indicators are

those which are only found in a single group and all group members possess the compound;

such a compound would have an indicator value of 1. We used the function "indicators" to

investigate both which single compounds and combinations of compounds best predict group

membership. We used the function "pruneindicators" to find the single or combinations of

compounds which had the highest indicator values.

Variation in chemical profiles

Divergence in chemical profiles across species and populations was estimated with

non-metric multidimensional scaling (NMDS) ordination in three dimensions, based on a

Bray-Curtis similarity matrix. We used the "metaMDS" function in the *vegan* package

version 2.5-1 (Oksanen et al., 2017), and visualised the NMDS using the *ade4* package (Dray

& Dufour, 2007).

We assessed the relative importance of relevant factors in driving the variation in

chemical profiles with multivariate statistical analyses. These factors included: species

identity, geographic region and individual locality. To compare overall variation in chemical

composition between groups, we carried out PERMANOVA (permutational multivariate

analysis of variance) testing based on a Bray-Curtis distance matrix, using the "adonis2"

function in the *vegan* package (Oksanen et al., 2017) with 1000 permutations. We

investigated each term in the model sequentially, starting with species identity, the main

clustering factor found from visualisation with NMDS, followed by geographic region

(Panama *vs* Western Andes *vs* Eastern Andes *vs* Amazon), and finally individual collecting

localities. Model goodness of fit was determined was evaluated by Akaike information

criterion (AIC). In general, we chose the model with the lowest AIC value, however, if two

models were within two AIC of each other we chose the simplest model as the best fit (Table

S3). We followed these PERMANOVA tests with *post hoc* pair-wise testing using the

function "pairwise.perm.manova" in the *RVAidememoie* package, with Bonferroni correction,

to identify which grouping factors were significantly different (Hervé, 2018). We repeated

the PERMANOVA analysis within species, in *H. erato* and *H. melpomene,* to investigate

fine-scale intraspecific geographic patterns. In the within species analysis we included

geographic region (Panama *vs* Western Andes *vs* Eastern Andes *vs* Amazon), and individual

collecting localities as the two factors.

One issue with distance-based analyses such as PERMANOVA is that differences in

dispersion between groups can be confounded with differences in location (Warton, Wright,

& Wang, 2012). To confirm these analyses and account for this issue, we implemented

multivariate generalised linear models using the function "ManyGLM" from the *mvabund*

package (Wang, Naumann, Wright, & Warton, 2012). We modelled the data using a negative

binomial distribution, which we found to be appropriate through examination of residual

plots. For interspecific analyses we included species, region, and locality nested within region

in the model. For intraspecific analyses we included region and locality nested within region.

The "ManyGLM" function fits models to each chemical compound, summing the test

statistics to give a multivariate test statistic known as Sum-of-LR. This statistic can be tested

for significance using resampling methods. We carried out backwards elimination and

compared the fit of models by using the "anova.manyglm" function with a likelihood ratio

test (Table S4). We can also determine which compounds are driving between-group

differences by looking at the individual contribution of each compound to the Sum-of-LR,

with p-values adjusted for multiple testing using the "adjust" option.

Phylogenetic and geographic distance

Shared ancestry can explain part of the variation in a species' chemical profile. We

tested for a correlation between phylogenetic distance and chemical profile divergence with

Mantel tests using the *vegan* package (Oksanen et al., 2017). To investigate the role of

geographic distance in chemical profile divergence, we compared geographic and chemical

distances matrices with partial Mantel tests. To visualise the species phylogeny (Kozak et al.,

2015) we used the "plot.phylo" function from the ape package (Paradis & Schliep, 2018).

Genomic and chemical distance within species

We calculated intraspecific genetic distances using genome sequences from 11 *H.*

erato and 13 *H. melpomene* populations. We visualised genetic distances in two dimensions

using MDS with the function "cmdscale". We tested for a correlation between intraspecific

genetic distance and chemical profile divergence with Mantel tests using the *vegan* package

(Oksanen et al., 2017). Hybrids between populations of the same species were excluded from

this analysis (Table S2). We used partial Mantel tests to investigate the role of geographic

distance.

Co-mimics and similarity of chemical profiles

We used samples of two mimicry rings from two localities, Panama and western

Ecuador. *H. melpomene* and *H. erato* form one mimicry ring, whilst *H. cydno* and *H. sapho*

form another, with the addition of *H. eleuchia* in western Ecuador (Fig. 1). To test which

factors most influenced the variation in chemical profiles we used both PERMANOVA and

multivariate GLM. We included species, region, mimicry group, and interactions between

these terms in the models.

All statistical analyses were performed with *R* version 3.5.1 (R Core Team, 2018).

Figures were made using a palette of colours optimized for colour-blindness (Wong, 2011).

Raw sequence data are available through GenBank Short Read Archive (XXX), GCMS data

is available through Dryad (XXX), R code and data are available on Open Science

Framework (https://osf.io/28yfk/?view_only=c1f7e7a925e74dee84fd2229cbf3f511).

Results

Chemical compounds in androconia and genitals:

We sampled 252 androconia and 275 genitals across 42 populations of 33 subspecies

of seven species, and identified 349 compounds in the genitals and 157 in the androconia. Of

the total number of androconial compounds, 38% are fatty acid derivatives, 20% aromatics,

10% terpenoids, 1% macrolides, <1% lactones and 31% unknown or unidentified

compounds. Of the genital compounds, 17% are fatty acid derivatives, 7% aromatics, 10%

terpenoids, 1% lactones, 12% macrolides and 44% unknown or unidentified compounds. The

main difference is that there are more macrolides in the genitals than androconia.

268   *Heliconius* species varied considerably in the amount and abundance of compounds

269 (Fig. 2). Between species there was variation in the number of compounds per individual, and

270 the overall amount of compounds detected (Table S5, S6). For the androconia, *H. eleuchia*

271 had the fewest compounds (13±5), and *H. melpomene* the most (32±11) (mean ± standard

272 deviation). *H. sapho* had the lowest total abundance of compounds at 1,300±803 ng, and *H.*

273 *melpomene* the most at 7,254±8,242 ng. The species with the fewest genital compounds was

274 *H. sapho* with 32±7, and the most *H. cydno* with 102±21. *H. sapho* also had the lowest total

275 amount of compounds at 6,642±3,975 ng, and *H. cydno* the highest at 91,167±67,122 ng.

276 These values are within the same order of magnitude as expected from previous work on

277 male sex pheromones in the butterfly *Bicyclus anyana* (Nieberding et al., 2012; van Bergen,

278 Brakefield, Heuskin, Zwaan, & Nieberding, 2013). Using *H. erato* as an example, the

279 androconial bouquet is 0.00002%, and genital bouquet 0.0007% of total body weight

280 (Montgomery, Merrill, & Ott, 2016). In general, a higher number of compounds and total

281 abundance of compounds is found in the genitals than in the androconial patches of

282 *Heliconius* wings.

Figure 2. Composition of androconial and genital bouquets across seven *Heliconius* species.
Species significantly differ in: (A) number of androconial compounds (ANOVA, $F_{6,245}=21.54$,
p<0.001), (B) number of genital compounds (anova, $F_{6,268}=36.15$, p<0.001), (C) amount of
androconial compounds (ANOVA, $F_{6,245}=11.55$, p<0.001), (D) amount of genital compounds (anova,
$F_{6,268}=11.62$, p<0.001). Four outlier individuals were removed from C.

Are there species-specific chemical compounds?

In order to identify candidate species recognition pheromones, we examined our data

for species specific compounds using indicator analysis. In most species that we examined

there were single androconial compounds that were strong indicators of species identity

(Table 1). For example, geranylgeranylacetone was found only in *H. erato* and was

consistently present across samples of this species. Similarily, octadecanal, a compound

previously shown to be electrophysiologically active (Byers et al., 2019), was found almost

exclusively in *H. melpomene* (specificity=0.999). *H. cydno* and *H. eleuchia* had the weakest

indicator scores; in *H. cydno* because the best indicator compound was not found in all

individuals examined (coverage= 0.667), and in *H. eleuchia* because the best indicator

compound was also found in other species (specificity= 0.747). There were similarly species-

specific genital compounds in all species except *H. sapho* and *H. timareta*, where a

combination of two compounds was the best predictor (Table 2). Similar to the androconia, in

H. melpomene, the best indicator compound for genitalia has known biological activity, in

this case the anti-aphrodisiac, (*E*)-β-ocimene (Schulz, Estrada, Yildizham, Boppré, & Gilbert,

2008). For *H. erato* we identified a terpene ester which is only found in *H. erato* individuals

and no other species. Other terpene esters were also almost perfect indicator compounds of

H. erato.

What factors affect interspecific variation in chemical profiles?

Our sampling allowed us to investigate how variation in chemical composition is

partitioned within and between species, and determine the extent to which chemistry is a

species-diagnostic trait. Visualisation of the chemical profiles reveals that individuals mostly

group by species for both androconial and genital chemical bouquets (Fig. 3). Species

significantly differ in their androconial bouquet, with species identity accounting for 58% of

the overall variation in chemical profiles (PERMANOVA, Species, $F_{6,251}=72.16$, $p<0.001$).

All pairwise comparisons of species are significant (Table S7). A further 4% of variation can

be explained by region (Amazon/Eastern Andes/Western Andes/Panama), and 3% by locality

nested within region (PERMANOVA, Region, $F_{3,251}=9.96$, $p<0.001$; (Region/Locality),

$F_{8,251}=2.65$, $p<0.001$). Finally, 4% of variation is explained by an interaction between species

and region (PERMANOVA, Species*Region $F_{6,251}=4.82$, $p<0.001$;).

The results were similar for the genital bouquets, with species identity accounting for

51% of the variation in chemical profiles (PERMANOVA, Species, $F_{6,274}=59.81$, $p<0.001$).

All pairwise comparisons are significant apart from *H. elevatus* and *H. melpomene* (Table

S8). A further 5% of variation can be explained by region (Amazon/Eastern Andes/Western

Andes/Panama), and 3% by locality nested within region (PERMANOVA, Region,

$F_{3,274}=12.43$, $p<0.001$; (Region/Locality), $F_{8,274}=2.92$, $p<0.001$). Finally, 6% of variation is

explained by an interaction between species and region (PERMANOVA, Species*Region

$F_{6,274}=6.52$, $p<0.001$). For both androconial and genital chemical profiles, most variation is

explained by species identity, rather than geographic location, as confirmed by ManyGLM

(Tables S9, S10).

Figure 3. NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in chemical compounds of male *Heliconius* of different species. A) Androconial compound bouquets differ significantly between species. Stress=0.155. B) Genital bouquets also differ significantly between species. Stress=0.121.

Does phylogenetic distance explain chemical profile divergence?

Using whole-genome sequence data, we explored the degree to which variation

between species can be explained by geographic and genetic distance among the samples. We

carried out partial Mantel tests to investigate the correlation between two variables whilst

controlling for a third variable. When controlling for geographic distance, genetic divergence

is strongly correlated with both androconial and genital chemical divergence (Mantel test,

androconia, r=0.7871, p=0.001; genitals, r=0.6936, p=0.001). When controlling for genetic

distance, geographic distance is significantly but weakly correlated with androconial and

genital chemical divergence (Mantel test, androconia, r=0.072, p=0.001; genitals, r=0.046,

p=0.007).

Do we find subspecies-specific chemical compounds?

We used an indicator analysis to search for compounds unique to specific populations

of *H. erato* and *H. melpomene*. Most intraspecific differences are due to quantitative rather

than qualitative differences between populations, perhaps explaining why many population

indicators were weak as they are also found in other regions at different amounts (Table S11,

S12). The only exception is *H. e. cyrbia* (Western Ecuador) that has many genital compounds

unique to this region (Table S11).

Figure 4. Plots of androconial, genital and genetic distance between *H. erato* populations. A) NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in androconial chemical compounds. Stress=0.174. B) NMDS plot illustrating in three dimensions the variation in genital chemical compounds. Stress=0.118. C) MDS plot illustrating in two dimensions the genetic distance between populations of *H. erato*.

Figure 5. Plots of androconial, genital and genetic distance between *H. melpomene* populations. A) NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in androconial chemical compounds. Stress=0.151. B) NMDS plot illustrating in three dimensions the variation in genital chemical compounds. Stress=0.161. C) MDS plot illustrating in two dimensions the genetic distance between populations of *H. melpomene*.

 What factors affect intraspecific variation in chemical profiles of H. erato *and* H.

melpomene*?*

We also wanted to determine the sources of variation within species using our broad

sampling of populations across the ranges of *H. erato* and *H. melpomene*. For *H. erato* there

was a strong grouping of individuals by region (Fig. 4), with 27% of variation in androconial

profiles being explained by region and 11% by locality nested within region (PERMANOVA,

Region $F_{3,87}=11.16$, $p<0.001$, Locality $F_{6,87}=2.35$, $p<0.001$). All four regions are significantly

different from each other (Pairwise permutation MANOVAs, $p<0.01$). For *H. erato* genital

compounds, 37% of variation is explained by region, and 11% by locality nested within

region (PERMANOVA, Region $F_{3,91}=19.01$, $p<0.001$, Locality $F_{6,91}=2.83$, $p<0.01$). All four

regions are significantly different from each other (Pairwise permutation MANOVAs,

$p<0.05$).

These geographic differences in chemical profiles are not as strong in *H. melpomene*

(Fig. 5). For *H. melpomene* androconial compounds, the best model only includes region, not

locality, with 18% of variation is explained by region (PERMANOVA, Region $F_{3,86}=6.73$,

$p<0.01$). The West Andes subspecies (*H. m. cythera*) is not significantly different from either

East Andes (multiple populations) or Panama (*H. m. rosina*) (Pairwise permutation

MANOVAs, $p=0.072$), however, the other comparisons are significantly different (Pairwise

permutation MANOVAs, $p<0.05$). For *H. melpomene* genital compounds, 20% of variation is

explained by region, and 12% by locality nested within region (PERMANOVA, Region

$F_{3,103}=8.91$, $p<0.001$, Locality $F_{7,103}=2.34$, $p<0.001$). All regions are significantly different

from each other (Pairwise permutation MANOVAs, $p<0.05$), apart from West Andes and

Amazon (Pairwise permutation MANOVAs, $p=0.120$). Both species show variation between

geographic locations, with more variance explained by region in *H. erato* than *H. melpomene*.

These results were confirmed by ManyGLM tests (Tables S13, S14, S15, S16).

 Does genetic distance explain chemical divergence in H. erato *and* H. melpomene*?*

In *H. erato*, chemical distance is positively correlated with genetic distance, when

accounting for geographic distance, although this correlation is weak for androconia (partial

Mantel test, androconia, R=0.169, p=0.001; genitals, R=0.352, p=0.001). When we account

for genetic distance, geographic distance is weakly correlated with androconial chemical

distance and not correlated with genital chemical distance (partial Mantel test, androconia,

R=0.149, p=0.001; genitals, R=-0.04105, p=0.764). These results were consistent when the

analysis was repeated without unknown compounds (supplementary results).

Heliconius melpomene genitals show similar patterns to *H. erato*, but variation in the

androconia is explained by geographic but not genetic distance. When accounting for

geography, genetic divergence is not correlated with androconial chemical divergence, and

only weakly with genital chemical divergence (partial Mantel test, androconia, R=0.02874,

p=0.141, genitals, R=0.1416, p=0.001). When we first consider genetic distance, geographic

distance is weakly positively correlated with androconial chemical distance, but not genital

chemical distance (partial Mantel test, androconia, R=0.1795, p=0.002; genitals, R=-0.01434,

p=0.744).

Is there evidence for similarity between co-mimics in chemical profile?

We tested for the effect of mimicry ring on chemical profile using individuals

collected in Panama and western Ecuador from two mimicry rings (Fig. 6). For androconial

chemical profiles, 45% of variation is explained by species (nested within mimicry ring, as

each species only belongs to one mimicry ring in our samples), with a further 21% explained

by mimicry ring (PERMANOVA, Mimicry ring $F_{1,85}$=54.29, p<0.001; (Mimicry

ring/Species) $F_{3,85}$=38.83, p<0.001). A further 2% is explained by region, another 2% by the

interaction between region and species nested within mimicry ring, and finally 1% by the

interaction between mimicry ring and region (PERMANOVA, Region $F_{1,85}=4.37$, p<0.001;

Mimicry ring*Region $F_{1,85}=2.70$, p<0.01; (Mimicry ring/Species)*Region $F_{2,85}=2.87$,

p<0.001).We confirmed these results using ManyGLM (Table S17).

We found similar trends for genital bouquets (Fig. 6). Again, species (nested within

mimicry ring) explained the most variation, 41% of genital bouquet variation, with a further

21% explained by mimicry ring (PERMANOVA, Mimicry ring $F_{1,88}=56.90$, p<0.001;

(Mimicry ring/Species) $F_{3,88}=37.03$, p<0.001). A further 3% is explained by region, another

4% by the interaction between region and species nested within mimicry ring, and finally 2%

by the interaction between mimicry ring and region (PERMANOVA, Region $F_{1,88}=9.07$,

p<0.001; Mimicry ring*Region $F_{1,88}=4.82$, p<0.01; (Mimicry ring/Species)*Region

$F_{2,88}=5.10$, p<0.001).We confirmed these results using ManyGLM (Table S18). In both

PERMANOVA and in multivariate GLMs, variation in both androconial and genital

bouquets is explained mostly by the species of the butterfly, with mimicry ring explaining

around half as much variation as species.

All the results described above show a consistent pattern when unidentified

compounds were not included in the analysis (supplementary results). Interspecific analyses

were also consistent when repeated without populations with a sample of fewer than 5

individuals (this removed 7 populations from androconial analysis and 5 from genital

analysis) (supplementary results).

Figure 6. NMDS (non-metric multidimensional scaling) plot illustrating in three dimensions the variation in chemical compounds of male *Heliconius* from Panama and Western Ecuador. *H. erato* and *H. melpomene* are co-mimics (circles), whilst *H. cydno*, *H. eleuchia* and *H. sapho* form a second co-mimicry group (triangles). A) Co-mimicry group contributes 21% of variation in androconial chemical bouquets. Stress=0.098. B) Co-mimicry group contributes 21% of variation in genital chemical bouquets. Stress=0.094.

Discussion

Heliconius butterflies represent a continental-scale adaptive radiation (Kozak et al.,

2015). Speciation in this group is often associated with divergence in wing colour pattern and

pattern variation plays an important role in speciation and mate preference (Jiggins, 2008;

Jiggins et al., 2001; Merrill et al., 2015, 2011, 2019; Sánchez et al., 2015). However, one of

the surprising findings to emerge from comparative genomic analysis is the wealth of

chemosensory genes (Heliconius Genome Consortium, 2012), suggesting that chemical

signalling may play an important role in the reproductive biology of the system. To begin to

understand the role of chemical signalling in this radiation we have extensively surveyed both

inter- and intraspecific variation of *Heliconius'* androconial and genital chemical profiles

across the Neotropics. We find that most of the variation in chemical profile across our

samples is explained by species, and we identify key chemicals serving as indicators for each

species. Nonetheless, there is also abundant intraspecific variation in chemical profiles. This

variation is mainly quantitative in nature, with the exception of *H. erato cyrbia* which has

compounds not found in other *H. erato* populations. We also find that co-mimics do exhibit

convergence in their chemical profiles, confirming an earlier hypothesis (Mann et al., 2017).

Our work sets the stage for further research into the biology and function of chemical

profiles, and their role in within and between-species signalling.

When analysing the chemical profiles across a phylogeny it may not be feasible to

carry out behavioural tests in every species. We hypothesised that consistent species-specific

compounds are likely to be under stabilising selection and therefore to also be biologically

important. In *H. melpomene* the compounds identified as indicators for androconia and

genitals, octadecanal and (*E*)-β-ocimene respectively, are both known to be biologically

active (Byers et al., 2019; Schulz et al., 2008). Combining broad geographic sampling with

indicator analysis therefore provides a promising approach to determine potential biologically

active compounds in other species, which could then be tested behaviourally. Our analyses

have already identified a number of compounds that could now be tested functionally, such as

the androconial compound geranylgeranylacetone in *H. erato*.

Chemical profiles are predicted to be highly species-specific if they are involved in

species recognition during mating. For instance orchid bee chemical blends, important for

mating and species recognition, show high species-specificity, as well as within-species

variability, which can be partly explained by geography (Brand et al., 2019; Weber et al.,

2016; Zimmermann, Roubik, & Eltz, 2006). We see similar patterns in *Heliconius*, with

greater interspecific than intraspecific differences in chemical profiles. Species identity is the

best predictor of chemical divergence, with geographic location able to explain some

intraspecific differences. One exception to this is *H. elevatus* which does not group separately

from its co-mimic *H. melpomene* for genital compounds, despite the fact that these species

are not especially closely related in the *Heliconius* phylogeny. Further samples are needed to

confirm that this result is not due to the small sample of *H. elevatus* in this study. As in

orchid bees, species differences are often consistent across a large geographic range,

suggesting that they could be important for reproductive isolation between species (Weber et

al., 2016).

We found a correlation between chemical distance and genetic distance. This suggests

that neutral evolutionary forces are important in driving chemical bouquet evolution. The

correlation between genital chemical distance and genetic distance is a much stronger

correlation than previously reported (Catalina Estrada, Schulz, Yildizhan, & Gilbert, 2011),

possibly due to the quantitative nature of our data. The strong signal of neutrality suggests

that the majority of compounds in the bouquets are neutrally evolving. For example, in the

genital bouquet of *H. melpomene*, one compound, (*E*)-β-ocimene, can act by itself as an anti-

aphrodisiac, with other components of the bouquet thought to moderate its evaporation rate

(Schulz et al., 2008). In the future, focusing on the evolutionary patterns of only biologically

active compounds, rather than the entire bouquet, may disentangle the processes involved in

the evolution of these profiles.

Heliconius erato and *H. melpomene* both exhibit extensive colour pattern variation

across their geographic range (Sheppard, Turner, Brown K. S., Benson W. W., & Singer M.

C., 1985) and these populations also differ in their androconial and genital bouquets. Whilst

traditionally predicted to be under stabilising selection, intraspecific variation between

populations in chemical profiles has been documented in other Lepidoptera (Bacquet et al.,

2016; Groot et al., 2009; Huang et al., 1998; McElfresh & Millar, 2001; Takanashi et al.,

2005). Chemical divergence in putative male sex pheromones between populations of

Bicyclus anynana is reported to be as large as differences between *Bicyclus* species, and is

greater than predicted by genetic divergence (Bacquet et al., 2016). This is in contrast to what

we find here, where interspecific differences are much greater than intraspecific ones.

Interestingly, *Heliconius erato cyrbia* produces many unique genital compounds and

is also the most genetically divergent *H. erato* population in our study, suggesting that

genetic drift is important for the evolution of chemical profiles in *Heliconius*. Across all *H.*

erato populations we find a correlation between chemical distance and genetic distance,

which is weaker for androconial bouquets. In *H. melpomene*, genetic distance is also weakly

correlated with genital chemical divergence. These correlations suggest that some of the

geographic variation between populations could be neutral, with stochastic processes

important for bouquet evolution in *Heliconius*. In contrast, androconial chemical variation in

H. melpomene is better explained by geographic distance. This does not match our

expectations of neutrality and might imply that other evolutionary forces are important for

chemical profile evolution in *H. melpomene*. One such factor might be larval host plant use,

which affects compound production (Darragh et al., 2019) and varies across the geographic

range of *H. melpomene* (Benson, 1978; Benson, Brown, & Gilbert, 1975).

Heliconius butterflies are an excellent example of visual mimicry, with different

species converging on the same warning colour patterns (Merrill et al., 2015; Sheppard et al.,

1985; Sherratt, 2008). It has been suggested that chemical compounds could also contribute

to mimicry between species (Dettner & Liepert, 1994; Mann et al., 2017). In this study, we

find evidence that individuals within co-mimicry groups have more similar chemical profiles

than expected. Most known examples of chemical mimicry come from systems of deception,

for example, mimicry of ant alarm pheromones by rove beetles to avoid predation, rather than

mimicry of aposematic warning signals (Dettner & Liepert, 1994; Stoeffler, Maier, Tolasch,

& Steidle, 2007; Vereecken & McNeil, 2010). We suggest that in *Heliconius* different

components of the bouquet could be important for chemical mimicry and species recognition,

reducing conflict between these selection pressures.

Convergence of genital bouquets between co-mimics could be due to the anti-

aphrodisiac function of these compounds (Gilbert, 1976; Schulz et al., 2008). Anti-

aphrodisiac compounds are transferred from males to females during mating to deter future

matings from other males. Convergence in wing pattern between co-mimics could result in

harassment not only by conspecific but also heterospecific males (C. Estrada & Jiggins,

2008). The use of the same anti-aphrodisiac by co-mimics could combat interspecific

attraction by deterring males of both species, as highlighted by the production of (E)-β-

ocimene by *H. erato* and *H. melpomene*

Even compounds which are not involved in chemical mimicry could play a role in

predator deterrence. Genital compounds were originally suggested to form part of the anti-

predation signal (Eltringham, 1925). We detected 2-sec-butyl-3-methoxypyrazine in the

genitals of *H. melpomene*, *H. cydno* and *H. timareta*, and 2-isobutyl-3-methoxypyrazine in

the genitals of *H. melpomene* and *H. cydno*, both compounds known to deter predators in the

wood tiger moth (Burdfield-Steel, Pakkanen, Rojas, Galarza, & Mappes, 2018; Rojas et al.,

2018, 2017; Rojas, Mappes, & Burdfield-Steel, 2019). More generally, methoxypyrazines act

as warning odours in other insects (e.g. Lepidoptera, Rothschild, Moore & Brown, 1984;

fireflies, Vencl et al., 2016), effective against avian predators (Guilford, Nicol, Rothschild, &

Moore, 1987). Further investigation will be required to determine if odours of *Heliconius*

butterflies act as anti-predation signals.

Overall, our study reveals strong species differences in bouquets and the presence of

species-specific compounds, as well as intraspecific variation. A pattern of species-specificity

alongside intraspecific variation geographically could be the result of a balance between

stabilising selection towards a species stereotype, sexual selection promoting diversity, and

geographic segregation alongside selection and drift. A challenge for the field is the

feasibility of testing for the biological relevance of hundreds of compounds in many species,

but we hope that our innovative analysis will stimulate not only further targeted functional

studies of putatively important compounds, but also large chemical profile surveys in other

study systems of evolutionary interest.

Acknowledgements

KD and GMK were supported by Natural Research Council Doctoral Training Partnership

(NE/L002507/1) and KD was additionally supported by a Smithsonian Tropical Research

Institute Short Term Fellowship KMK was funded by a Smithsonian Institution Fellowship.

CRM was supported by a Smithsonian Tropical Research Institute Short Term Fellowship

and Predoctoral Fellowship. This study was financed in part by the Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001 (MSc

Scholarship for CMEF). CMEF was also funded by a Smithonsian Tropical Research institute

Short Term Fellowship. CS was funded by COLCIENCIAS (FP44842-5-2017). ML was

supported by research funds from Universidad del Rosario. KJRPB and CDJ were supported

by the European Research Council (339873 SpeciationGenetics). RMM was supported by an

DFG Emmy Noether fellowship. WOM was supported by the Smithsonian Tropical Research

Institute and NSF (DEB 1257689). SS thanks the Deutsche Forschungsgemeinschaft (DFG)

for support through grant Schu984/12-1. We thank Michel Cast for photos of *H. melpomene*

madeira and *H. m. intersectus* (https://cliniquevetodax.com/Heliconius/). The authors have

no conflicts of interest to declare.

Author contributions

Kathy Darragh, Gabriela Montejo-Kovacevich, Krzysztof Kozak, Colin Morrison, Owen McMillan, Clarisse Figueiredo, Jonathan Ready, Camilo Salazar, Mauricio Linares and Chris Jiggins collected samples. Kathy Darragh analysed the data. Stefan Schulz assisted with the chemical analysis. Gabriela Montejo-Kovacevich created the intraspecific genetic distance matrices. Richard Merrill, Owen McMillan, Kelsey Byers and Chris Jiggins contributed to project design. All authors contributed to manuscript revisions.

Data availability

Raw sequence data are available through GenBank Short Read Archive (XXX), GCMS data is available through Dryad (XXX), R code and data are available on Open Science Framework (https://osf.io/28yfk/?view_only=c1f7e7a925e74dee84fd2229cbf3f511).

References

Bacquet, P. M. B., de Jong, M. A., Brattström, O., Wang, H.-L., Molleman, F., Heuskin, S., …

Nieberding, C. M. (2016). Differentiation in putative male sex pheromone components across

and within populations of the African butterfly *Bicyclus anynana* as a potential driver of

reproductive isolation. *Ecology and Evolution*, *6*(17), 6064–6084. doi: 10.1002/ece3.2298

Benedict, L., & Bowie, R. C. K. (2009). Macrogeographical variation in the song of a widely

distributed African warbler. *Biology Letters*, *5*(4), 484–487. doi: 10.1098/rsbl.2009.0244

Benson, W. W. (1978). Resource partitioning in passion vine butterflies. *Evolution; International*

Journal of Organic Evolution, *32*(3), 493–518. doi: 10.1111/j.1558-5646.1978.tb04593.x

Benson, W. W., Brown, K. S., & Gilbert, L. E. (1975). Coevolution of plants and herbivores: Passion

flower butterflies. *Evolution; International Journal of Organic Evolution*, *29*(4), 659–680.

doi: 10.1111/j.1558-5646.1975.tb00861.x

Bolnick, D. I., & Kirkpatrick, M. (2012). The relationship between intraspecific assortative mating

and reproductive isolation between divergent populations. *Current Zoology*, *58*(3), 484–492.

doi: 10.1093/czoolo/58.3.484

Brand, P., Hinojosa-Díaz, I. A., Ayala, R., Daigle, M., Obiols, C. L. Y., Eltz, T., & Ramírez, S. R.

(2019). An olfactory receptor gene underlies reproductive isolation in perfume-collecting

orchid bees. *BioRxiv*, 537423. doi: 10.1101/537423

Burdfield-Steel, E., Pakkanen, H., Rojas, B., Galarza, J. A., & Mappes, J. (2018). De novo Synthesis

of Chemical Defenses in an Aposematic Moth. *Journal of Insect Science*, *18*(2), 1–4. doi:

10.1093/jisesa/iey020

Butlin, R. K., Hewitt, G. M., & Webb, S. F. (1985). Sexual selection for intermediate optimum in

Chorthippus brunneus (Orthoptera: Acrididae). *Animal Behaviour*, *33*(4), 1281–1292. doi:

10.1016/S0003-3472(85)80188-3

Bybee, S. M., Yuan, F., Ramstetter, M. D., Llorente-Bousquets, J., Reed, R. D., Osorio, D., &

Briscoe, A. D. (2012). UV Photoreceptors and UV-Yellow Wing Pigments in *Heliconius*

Butterflies Allow a Color Signal to Serve both Mimicry and Intraspecific Communication.

The American Naturalist, *179*(1), 38–51. doi: 10.1086/663192

Byers, K. J. R. P., Darragh, K., Musgrove, J., Almeida, D. A., Garza, S. F., Warren, I. A., … Jiggins,

C. D. (2019). A major locus controls a biologically active pheromone component in

Heliconius melpomene. *BioRxiv*, 739037. doi: 10.1101/739037

Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of sites: Indices and

statistical inference. *Ecology*, *90*(12), 3566–3574. doi: 10.1890/08-1823.1

Campbell, P., Pasch, B., Pino, J. L., Crino, O. L., Phillips, M., & Phelps, S. M. (2010). Geographic

variation in the songs of neotropical singing mice: Testing the relative importance of drift and

local adaptation. *Evolution*, *64*(7), 1955–1972. doi: 10.1111/j.1558-5646.2010.00962.x

Challis, R. J., Kumar, S., Dasmahapatra, K. K. K., Jiggins, C. D., & Blaxter, M. (2016). Lepbase: The

Lepidopteran genome database. *BioRxiv*, 056994. doi: 10.1101/056994

Conrad, T., Paxton, R. J., Assum, G., & Ayasse, M. (2018). Divergence in male sexual odor signal

and genetics across populations of the red mason bee, *Osmia bicornis*, in Europe. *PLOS ONE*,

13(2), e0193153. doi: 10.1371/journal.pone.0193153

Coyne, J. A., & Orr, H. A. (2004). *Speciation*. Sunderland, MA: Sinauer.

Darragh, K., Byers, K. J. R. P., Merrill, R. M., McMillan, W. O., Schulz, S., & Jiggins, C. D. (2019).

Male pheromone composition depends on larval but not adult diet in Heliconius melpomene.

Ecological Entomology. doi: 10.1111/een.12716

Darragh, K., Vanjari, S., Mann, F., Gonzalez-Rojas, M. F., Morrison, C. R., Salazar, C., … Jiggins, C.

D. (2017). Male sex pheromone components in *Heliconius* butterflies released by the

androconia affect female choice. *PeerJ*, *5*, e3953. doi: 10.7717/peerj.3953

Davey, J. W., Barker, S. L., Rastas, P. M., Pinharanda, A., Martin, S. H., Durbin, R., … Jiggins, C. D.

(2017). No evidence for maintenance of a sympatric *Heliconius* species barrier by

chromosomal inversions. *Evolution Letters*, *1*(3), 138–154. doi: 10.1002/evl3.12

Dettner, K., & Liepert, C. (1994). Chemical Mimicry and Camouflage. *Annual Review of*

Entomology, *39*(1), 129–154. doi: 10.1146/annurev.en.39.010194.001021

Dray, S., & Dufour, A. B. (2007). The ade4 package: Implementing the duality diagram for

ecologists. *Journal of Statistical Software*, *22*(4), 1–20. doi: 10.18637/jss.v022.i04

Eltringham, H. (1925). On the Abdominal Glands in *Heliconius* (Lepidoptera). *Transactions of the*

Royal Entomological Society of London, *73*(1–2), 269–275. doi: 10.1111/j.1365-

2311.1925.tb02862.x

Estrada, C., & Jiggins, C. D. (2008). Interspecific sexual attraction because of convergence in

warning colouration: Is there a conflict between natural and sexual selection in mimetic

species? *Journal of Evolutionary Biology*, *21*(3), 749–760. doi: 10.1111/j.1420-

9101.2008.01517.x

Estrada, Catalina, Schulz, S., Yildizhan, S., & Gilbert, L. E. (2011). Sexual selection drives the

evolution of antiaphrodisiac pheromones in butterflies. *Evolution; International Journal of*

Organic Evolution, *65*(10), 2843–2854. doi: 10.1111/j.1558-5646.2011.01352.x

Ferreira, M., & Ferguson, J. W. H. (2002). Geographic variation in the calling song of the field cricket

Gryllus bimaculatus (Orthoptera: Gryllidae) and its relevance to mate recognition and mate

choice. *Journal of Zoology*, *257*(2), 163–170. doi: 10.1017/S0952836902000766

Finkbeiner, S. D., Fishman, D. A., Osorio, D., & Briscoe, A. D. (2017). Ultraviolet and yellow

reflectance but not fluorescence is important for visual discrimination of conspecifics by

Heliconius erato. *Journal of Experimental Biology*, *220*(7), 1267–1276. doi:

10.1242/jeb.153593

Friberg, M., Vongvanich, N., Borg-Karlson, A.-K., Kemp, D. J., Merilaita, S., & Wiklund, C. (2008).

Female mate choice determines reproductive isolation between sympatric butterflies.

Behavioral Ecology and Sociobiology, *62*(6), 873–886. doi: 10.1007/s00265-007-0511-2

Gemeno, C., Lutfallah, A. F., & Haynes, K. F. (2000). Pheromone Blend Variation and Cross-

Attraction Among Populations of the Black Cutworm Moth (Lepidoptera: Noctuidae). *Annals*

of the Entomological Society of America, *93*(6), 1322–1328. doi: 10.1603/0013-

8746(2000)093[1322:PBVACA]2.0.CO;2

Gerhardt, H. C. (1982). Sound Pattern Recognition in Some North American Treefrogs (Anura:

Hylidae): Implications for Mate Choice. *Integrative and Comparative Biology*, *22*(3), 581–

595. doi: 10.1093/icb/22.3.581

Gilbert, L. E. (1976). Postmating female odor in *Heliconius* butterflies: A male-contributed

antiaphrodisiac? *Science*, *193*(4251), 419–420. doi: 10.1126/science.935877

Giraldo, N., Salazar, C., Jiggins, C. D., Bermingham, E., & Linares, M. (2008). Two sisters in the

same dress: *Heliconius* cryptic species. *BMC Evolutionary Biology*, *8*, 324. doi:

10.1186/1471-2148-8-324

Gray, D. A., & Cade, W. H. (2000). Sexual selection and speciation in field crickets. *Proceedings of*

the National Academy of Sciences, *97*(26), 14449–14454. doi: 10.1073/pnas.97.26.14449

Groot, A. T., Inglis, O., Bowdridge, S., Santangelo, R. G., Blanco, C., Jr, J. D. L., … Schal, C. (2009).

Geographic and Temporal Variation in Moth Chemical Communication. *Evolution*, *63*(8),

1987–2003. doi: 10.1111/j.1558-5646.2009.00702.x

Guilford, T., Nicol, C., Rothschild, M., & Moore, B. P. (1987). The biological roles of pyrazines:

Evidence for a warning odour function. *Biological Journal of the Linnean Society*, *31*(2),

113–128. doi: 10.1111/j.1095-8312.1987.tb01984.x

Hankison, S. J., & Ptacek, M. B. (2008). Geographical variation of genetic and phenotypic traits in the

Mexican sailfin mollies, *Poecilia velifera* and *P. petenensis*. *Molecular Ecology*, *17*(9), 2219–

2233. doi: 10.1111/j.1365-294X.2008.03736.x

Heliconius Genome Consortium. (2012). Butterfly genome reveals promiscuous exchange of mimicry

adaptations among species. *Nature*, *487*(7405), 94–98. doi: 10.1038/nature11041

Hervé, M. (2018). *RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R package*

version 0.9-69-3. Retrieved from https://CRAN.R-project.org/package=RVAideMemoire

Hijmans, R. J. (2017). *Geosphere: Spherical Trigonometry. R package version 1.5-7*. Retrieved from

https://CRAN.R-project.org/package=geosphere

Huang, Y., Takanashi, T., Hoshizaki, S., Tatsuki, S., Honda, H., Yoshiyasu, Y., & Ishikawa, Y.

(1998). Geographic Variation in Sex Pheromone of Asian Corn Borer, *Ostrinia furnacalis*, in

Japan. *Journal of Chemical Ecology*, *24*(12), 2079–2088. doi: 10.1023/A:1020737726636

Irwin, D. E., Thimgan, M. P., & Irwin, J. H. (2008). Call divergence is correlated with geographic and

genetic distance in greenish warblers (*Phylloscopus trochiloides*): A strong role for

stochasticity in signal evolution? *Journal of Evolutionary Biology*, *21*(2), 435–448. doi:

10.1111/j.1420-9101.2007.01499.x

Jiggins, C. D. (2008). Ecological Speciation in Mimetic Butterflies. *BioScience*, *58*(6), 541–548. doi:

10.1641/B580610

Jiggins, C. D. (2017). *The Ecology and Evolution of* Heliconius *Butterflies*. Oxford: Oxford

University Press.

Jiggins, C. D., Naisbit, R. E., Coe, R. L., & Mallet, J. (2001). Reproductive isolation caused by colour

pattern mimicry. *Nature*, *411*(6835), 302–305. doi: 10.1038/35077075

Johansson, B. G., & Jones, T. M. (2007). The role of chemical communication in mate choice.

Biological Reviews, *82*(2), 265–289. doi: 10.1111/j.1469-185X.2007.00009.x

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. doi: 10.1186/s12859-014-0356-4

Kozak, K. M., Wahlberg, N., Neild, A. F. E., Dasmahapatra, K. K., Mallet, J., & Jiggins, C. D.

(2015). Multilocus species trees show the recent adaptive radiation of the mimetic *Heliconius*

butterflies. *Systematic Biology*, *64*(3), 505–524. doi: 10.1093/sysbio/syv007

Lamas, G., & Jiggins, C. D. (2017). Taxonomic list. In *The ecology and evoution of Heliconius*

butterflies (pp. 214–244). Oxford: Oxford University Press.

Leonhardt, S., Rasmussen, C., & Schmitt, T. (2013). Genes versus environment: Geography and

phylogenetic relationships shape the chemical profiles of stingless bees on a global scale.

Proceedings. Biological Sciences / The Royal Society, *280*, 20130680. doi:

10.1098/rspb.2013.0680

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics (Oxford, England), *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324

Löfstedt, C. (1993). Moth Pheromone Genetics and Evolution. *Philosophical Transactions of the*

Royal Society of London B: Biological Sciences, *340*(1292), 167–177. doi:

10.1098/rstb.1993.0055

Mann, F., Vanjari, S., Rosser, N., Mann, S., Dasmahapatra, K. K., Corbin, C., … Schulz, S. (2017).

The Scent Chemistry of *Heliconius* Wing Androconia. *Journal of Chemical Ecology*, *43*(9),

843–857. doi: 10.1007/s10886-017-0867-3

Martin, M. D., & Mendelson, T. C. (2016). The accumulation of reproductive isolation in early stages

of divergence supports a role for sexual selection. *Journal of Evolutionary Biology*, *29*(4),

676–689. doi: 10.1111/jeb.12819

Mas, F., & Jallon, J.-M. (2005). Sexual Isolation and Cuticular Hydrocarbon Differences between

Drosophila santomea and *Drosophila yakuba*. *Journal of Chemical Ecology*, *31*(11), 2747–

2752. doi: 10.1007/s10886-005-7570-5

McElfresh, J. S., & Millar, J. G. (2001). Geographic Variation in the Pheromone System of the

Saturniid Moth *Hemileuca eglanterina*. *Ecology*, *82*(12), 3505–3518. doi: 10.1890/0012-

9658(2001)082[3505:GVITPS]2.0.CO;2

McPeek, M. A., Symes, L. B., Zong, D. M., & McPeek, C. L. (2011). Species recognition and

patterns of population variation in the reproductive structures of a damselfly genus.

Evolution; International Journal of Organic Evolution, *65*(2), 419–428. doi: 10.1111/j.1558-

5646.2010.01138.x

Mendelson, T. C., & Shaw, K. L. (2012). The (mis)concept of species recognition. *Trends in Ecology*

& Evolution, *27*(8), 421–427. doi: 10.1016/j.tree.2012.04.001

Mérot, C., Frérot, B., Leppik, E., & Joron, M. (2015). Beyond magic traits: Multimodal mating cues

in *Heliconius* butterflies. *Evolution*, *69*(11), 2891–2904. doi: 10.1111/evo.12789

Mérot, C., Mavárez, J., Evin, A., Dasmahapatra, K. K., Mallet, J., Lamas, G., & Joron, M. (2013).

Genetic differentiation without mimicry shift in a pair of hybridizing *Heliconius* species

(Lepidoptera: Nymphalidae). *Biological Journal of the Linnean Society*, *109*(4), 830–847.

doi: 10.1111/bij.12091

Merrill, R. M., Chia, A., & Nadeau, N. J. (2014). Divergent warning patterns contribute to assortative

mating between incipient *Heliconius* species. *Ecology and Evolution*, *4*(7), 911–917. doi:

10.1002/ece3.996

Merrill, R. M., Dasmahapatra, K. K., Davey, J. W., Dell'Aglio, D. D., Hanly, J. J., Huber, B., … Yu,

Q. (2015). The diversification of *Heliconius* butterflies: What have we learned in 150 years?

Journal of Evolutionary Biology, *28*(8), 1417–1438. doi: 10.1111/jeb.12672

Merrill, R. M., Gompert, Z., Dembeck, L. M., Kronforst, M. R., McMillan, W. O., & Jiggins, C. D.

(2011). Mate Preference Across the Speciation Continuum in a Clade of Mimetic Butterflies.

Evolution, *65*(5), 1489–1500. doi: 10.1111/j.1558-5646.2010.01216.x

Merrill, R. M., Rastas, P., Martin, S. H., Melo, M. C., Barker, S., Davey, J., … Jiggins, C. D. (2019).

Genetic dissection of assortative mating behavior. *PLOS Biology*, *17*(2), e2005902. doi:

10.1371/journal.pbio.2005902

Montgomery, S. H., Merrill, R. M., & Ott, S. R. (2016). Brain composition in *Heliconius* butterflies,

posteclosion growth and experience-dependent neuropil plasticity. *Journal of Comparative*

Neurology, *524*(9), 1747–1769. doi: 10.1002/cne.23993

Mullen, L. M., Vignieri, S. N., Gore, J. A., & Hoekstra, H. E. (2009). Adaptive basis of geographic

variation: Genetic, phenotypic and environmental differences among beach mouse

populations. *Proceedings of the Royal Society of London B: Biological Sciences*, *276*(1674),

3809–3818. doi: 10.1098/rspb.2009.1146

Nagel, L., & Schluter, D. (1998). Body Size, Natural Selection, and Speciation in Sticklebacks.

Evolution, *52*(1), 209–218. doi: 10.2307/2410936

Nieberding, C. M., Fischer, K., Saastamoinen, M., Allen, C. E., Wallin, E. A., Hedenström, E., &

Brakefield, P. M. (2012). Cracking the olfactory code of a butterfly: The scent of ageing.

Ecology Letters, *15*(5), 415–424. doi: 10.1111/j.1461-0248.2012.01748.x

Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., … Wagner,

H. (2017). *Vegan: Community Ecology Package. R package version 2.4-2*. Retrieved from

https://CRAN.R-project.org/package=vegan

Paradis, E., & Schliep, K. (2018). ape 5.0: An environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*. doi: 10.1093/bioinformatics/bty633

Pfennig, K. S. (1998). The evolution of mate choice and the potential for conflict between species and

mate–quality recognition. *Proceedings of the Royal Society of London B: Biological Sciences*,

265(1407), 1743–1748. doi: 10.1098/rspb.1998.0497

Ptacek, M. (2000). The role of mating preferences in shaping interspecific divergence in mating

signals in vertebrates. *Behavioural Processes*, *51*(1–3), 111–134. doi: 10.1016/S0376-

6357(00)00123-6

R Core Team. (2018). R: A language and environment for statistical computing. (Version 3.3.1).

Retrieved from http://www.R-project.org

Ready, J. S., Sampaio, I., Schneider, H., Vinson, C., Santos, T. D., & Turner, G. F. (2006). Colour

forms of Amazonian cichlid fish represent reproductively isolated species. *Journal of*

Evolutionary Biology, *19*(4), 1139–1148. doi: 10.1111/j.1420-9101.2006.01088.x

Rojas, B., Burdfield-Steel, E., De Pasqual, C., Gordon, S., Hernández, L., Mappes, J., … Lindstedt, C.

(2018). Multimodal Aposematic Signals and Their Emerging Role in Mate Attraction.

Frontiers in Ecology and Evolution, *6*, 93. doi: 10.3389/fevo.2018.00093

Rojas, B., Burdfield-Steel, E., Pakkanen, H., Suisto, K., Maczka, M., Schulz, S., & Mappes, J. (2017).

How to fight multiple enemies: Target-specific chemical defences in an aposematic moth.

Proceedings. Biological Sciences, *284*(1863), 20171424. doi: 10.1098/rspb.2017.1424

Rojas, B., Mappes, J., & Burdfield-Steel, E. (2019). Multiple modalities in insect warning displays

have additive effects against wild avian predators. *Behavioral Ecology and Sociobiology*,

73(3), 37. doi: 10.1007/s00265-019-2643-6

Rothschild, M. (1961). Defensive Odours and Mullerian Mimicry Among Insects. *Transactions of the*

Royal Entomological Society of London, *113*(5), 101–123. doi: 10.1111/j.1365-

2311.1961.tb00803.x

Rothschild, M., Moore, B. P., & Brown, W. V. (1984). Pyrazines as warning odour components in the

Monarch butterfly, *Danaus plexippus*, and in moths of the genera *Zygaena* and *Amata*

(Lepidoptera). *Biological Journal of the Linnean Society*, *23*(4), 375–380. doi:

10.1111/j.1095-8312.1984.tb00153.x

Ryan, M. A., Moore, C. J., & Walter, G. H. (1995). Individual variation in pheromone composition in

Nezara viridula (Heteroptera: Pentatomidae): How valid is the basis for designating

"pheromone strains"? *Comparative Biochemistry and Physiology Part B: Biochemistry and*

Molecular Biology, *111*(2), 189–193. doi: 10.1016/0305-0491(94)00257-U

Ryan, M. J., & Guerra, M. A. (2014). The mechanism of sound production in túngara frogs and its

role in sexual selection and speciation. *Current Opinion in Neurobiology*, *28*, 54–59. doi:

10.1016/j.conb.2014.06.008

Ryan, M. J., & Rand, A. S. (1993). Species Recognition and Sexual Selection as a Unitary Problem in

Animal Communication. *Evolution*, *47*(2), 647–657. doi: 10.2307/2410076

Ryan, M. J., Rand, A. S., & Weigt, L. A. (1996). Allozyme and Advertisement Call Variation in the

Túngara Frog, *Physalaemus Pustulosus*. *Evolution*, *50*(6), 2435–2453. doi: 10.1111/j.1558-

5646.1996.tb03630.x

Sánchez, A. P., Pardo-Diaz, C., Enciso-Romero, J., Muñoz, A. G., Jiggins, C. D., Salazar, C., &

Linares, M. (2015). An introgressed wing pattern acts as a mating cue. *Evolution;*

International Journal of Organic Evolution, *69*(6), 1619–1629. doi: 10.1111/evo.12679

Saveer, A. M., Becher, P. G., Birgersson, G., Hansson, B. S., Witzgall, P., & Bengtsson, M. (2014).

Mate recognition and reproductive isolation in the sibling species *Spodoptera littoralis* and

Spodoptera litura. *Chemical Ecology*, *2*, 18. doi: 10.3389/fevo.2014.00018

Schulz, S., Estrada, C., Yildizham, S., Boppré, M., & Gilbert, L. E. (2008). An antiaphrodisiac in

Heliconius melpomene butterflies. *Journal of Chemical Ecology*, *34*(1), 82–93. doi:

10.1007/s10886-007-9393-z

Seehausen, O., Terai, Y., Magalhaes, I. S., Carleton, K. L., Mrosso, H. D. J., Miyagi, R., … Okada, N.

(2008). Speciation through sensory drive in cichlid fish. *Nature*, *455*(7213), 620–626. doi:

10.1038/nature07285

Selz, O. M., Pierotti, M. E. R., Maan, M. E., Schmid, C., & Seehausen, O. (2014). Female preference

for male color is necessary and sufficient for assortative mating in 2 cichlid sister species.

Behavioral Ecology, *25*(3), 612–626. doi: 10.1093/beheco/aru024

Sheppard, P. M., Turner, J. R. G., Brown K. S., Benson W. W., & Singer M. C. (1985). Genetics and

the evolution of Muellerian mimicry in *Heliconius* butterflies. *Philosophical Transactions of*

the Royal Society of London. B, Biological Sciences, *308*(1137), 433–610. doi:

10.1098/rstb.1985.0066

Sherratt, T. N. (2008). The evolution of Müllerian mimicry. *Die Naturwissenschaften*, *95*(8), 681–

695. doi: 10.1007/s00114-008-0403-y

Smadja, C. M., & Butlin, R. K. (2008). On the scent of speciation: The chemosensory system and its

role in premating isolation. *Heredity*, *102*(1), 77–97. doi: 10.1038/hdy.2008.55

Stoeffler, M., Maier, T. S., Tolasch, T., & Steidle, J. L. M. (2007). Foreign-language Skills in Rove-

Beetles? Evidence for Chemical Mimicry of Ant Alarm Pheromones in Myrmecophilous

Pella Beetles (Coleoptera: Staphylinidae). *Journal of Chemical Ecology*, *33*(7), 1382–1392.

doi: 10.1007/s10886-007-9315-0

Sun, K., Luo, L., Kimball, R. T., Wei, X., Jin, L., Jiang, T., … Feng, J. (2013). Geographic Variation

in the Acoustic Traits of Greater Horseshoe Bats: Testing the Importance of Drift and

Ecological Selection in Evolutionary Processes. *PLOS ONE*, *8*(8), e70368. doi:

10.1371/journal.pone.0070368

Takanashi, T., Huang, Y., Takahasi, K. R., Hoshizaki, S., Tatsuki, S., & Ishikawa, Y. (2005). Genetic

analysis and population survey of sex pheromone variation in the adzuki bean borer moth,

Ostrinia scapulalis. Biological Journal of the Linnean Society, *84*(1), 143–160. doi:

10.1111/j.1095-8312.2005.00421.x

Van Belleghem, S. M., Rastas, P., Papanicolaou, A., Martin, S. H., Arias, C. F., Supple, M. A., …

Papa, R. (2017). Complex modular architecture around a simple toolkit of wing pattern genes.

Nature Ecology & Evolution, *1*(3), 52. doi: 10.1038/s41559-016-0052

van Bergen, E., Brakefield, P. M., Heuskin, S., Zwaan, B. J., & Nieberding, C. M. (2013). The scent

of inbreeding: A male sex pheromone betrays inbred males. *Proceedings of the Royal Society*

of London B: Biological Sciences, *280*(1758), 20130102. doi: 10.1098/rspb.2013.0102

Vencl, F. V., Ottens, K., Dixon, M. M., Candler, S., Bernal, X. E., Estrada, C., & Page, R. A. (2016).

Pyrazine emission by a tropical firefly: An example of chemical aposematism? *Biotropica*,

48(5), 645–655. doi: 10.1111/btp.12336

Vereecken, N. J., & McNeil, J. N. (2010). Cheaters and liars: Chemical mimicry at its finest.

Canadian Journal of Zoology, *88*(7), 725–752. doi: 10.1139/Z10-040

Wang, Y., Naumann, U., Wright, S. T., & Warton, D. I. (2012). Mvabund– an R package for model-

based analysis of multivariate abundance data. *Methods in Ecology and Evolution*, *3*(3), 471–

474. doi: 10.1111/j.2041-210X.2012.00190.x

Warton, D. I., Wright, S. T., & Wang, Y. (2012). Distance-based multivariate analyses confound

location and dispersion effects. *Methods in Ecology and Evolution*, *3*(1), 89–101. doi:

10.1111/j.2041-210X.2011.00127.x

Weber, M. G., Mitko, L., Eltz, T., & Ramírez, S. R. (2016). Macroevolution of perfume signalling in

orchid bees. *Ecology Letters*, *19*(11), 1314–1323. doi: 10.1111/ele.12667

Wong, B. (2011). Points of view: Color blindness. *Nature Methods*, *8*, 441. doi: 10.1038/nmeth.1618

Wyatt, T. D. (2003). *Pheromones and Animal Behaviour: Communication by Smell and Taste*.

Cambridge: Cambridge University Press.

Wyatt, T. D. (2014). *Pheromones and Animal Behavior: Chemical Signals and Signatures*.

Cambridge: Cambridge University Press.

Yildizhan, S., van Loon, J., Sramkova, A., Ayasse, M., Arsene, C., ten Broeke, C., & Schulz, S.

(2009). Aphrodisiac pheromones from the wings of the small cabbage white and large

cabbage white butterflies, *Pieris rapae* and *Pieris brassicae*. *ChemBioChem*, *10*(10), 1666–

1677. doi: 10.1002/cbic.200900183

Zimmermann, Y., Roubik, D. W., & Eltz, T. (2006). Species-specific attraction to pheromonal

analogues in orchid bees. *Behavioral Ecology and Sociobiology*, *60*(6), 833. doi:

10.1007/s00265-006-0227-8

Table 1: Androconial compounds which are the best indicators of species identity. A is a measure of group specificity of the compounds, B is a measure of group coverage, and sqrtIV is the indicator value which considers both A and B and ranges from 0 (compound not present in any individuals of that species) to 1 (compound only present in that species, and present in all individuals).

Species/compound	A: specificity	B: coverage	sqrtIV
H. cydno			
Unknown aromatic (RI=2130)	1	0.667	0.816
H. eleuchia			
Hexahydrofarnesylacetone	0.747	1	0.864
H. elevatus			
Homovanillylalcohol	0.912	1	0.955
H. erato			
Geranylgeranylacetone	1	1	1
H. melpomene			
Octadecanal	0.999	1	1
H. sapho			
Methyl 4-hydroxy-3-methoxybenzoate	0.866	1	0.931
H. timareta			
5-Decanolide	1	0.889	0.943

Table 2: Genital compounds which are the best indicators of species identity. A, B, and sqrtIV as in Table 1.

Species/compound	A: specificity	B: coverage	sqrtIV
H. cydno			
Unknown ester (RI=1390)	0.999	1	0.999
H. eleuchia			
Unknown macrolide RI=1878	0.969	1	0.984
H. elevatus			
Icosenol	0.908	1	0.953
H. erato			
Unknown terpene ester (RI=2494)	1	1	1
H. melpomene			
(*E*)-β-Ocimene	0.865	1	0.930
H. sapho			
(*Z*)-3-Hexenyl isobutyrate & unknown (RI=1691)	0.957	0.923	0.940
H. timareta			
Butyl oleate & (*Z*)-9-octadecen-13-olide	0.915	1	0.956