# Cultured pluripotent planarian stem cells retain potency and express proteins from exogenously introduced mRNAs

Kai Lei[1, 4], Sean A. McKinney[1], Eric J. Ross[1, 2], Heng-Chi Lee[3], Alejandro Sánchez Alvarado[1, 2,*]

[1] Stowers Institute for Medical Research, Kansas City, MO 64110, USA

[2] Howard Hughes Medical Institute, Stowers Institute for Medical Research, Kansas City, MO 64110, USA

[3] Department of Molecular Genetics and Cell Biology, University of Chicago, Chicago, IL 60637, USA

[4] Current Address: School of Life Sciences, Westlake University, Hangzhou, Zhejiang 310024, China

* Correspondence: asa@stowers.org

**Key words:** cell culture, planaria, neoblast, transgenesis, SiR-DNA

**Abstract:**

Planarians possess naturally occurring pluripotent adult somatic stem cells (neoblasts) required for homeostasis and whole–body regeneration. However, methods for culturing neoblasts are currently unavailable, hindering both mechanistic studies of potency and the development of transgenic tools. We report the first robust methodologies for culturing and delivering exogenous mRNA into neoblasts. We identified culture media for maintaining neoblasts in vitro, and showed via transplantation that the cultured stem cells retained pluripotency. By modifying standard flow cytometry methods, we developed a new procedure that significantly improved yield and purity of neoblasts. These methods facilitated the successful introduction and expression of exogenous mRNAs in neoblasts, overcoming a key hurdle impeding the application of transgenics in planarians. The tissue culture advances reported here create new opportunities to advance detailed mechanistic studies of adult stem cell pluripotency in planarians, and provide a systematic methodological framework to develop cell culture techniques for other emerging research organisms.

**Introduction**

Stem cell pluripotency remains an important and still unresolved problem in biology. Several systems have been established to study pluripotency regulation in germlines, embryonic, and induced pluripotent stem cells [1-4]. However, no naturally occurring adult pluripotent stem cells have been identified in traditional model systems, including round worms, flies, fishes, and mice. Unlike traditional research organisms, planarians harbor an abundant population of adult stem cells collectively known as neoblasts. These cells are characteristic of flatworms and acoels [5], and in planarians include a subpopulation of pluripotent stem cells termed clonogenic neoblasts [6-8]. Neoblasts confer planarians with remarkable regenerative abilities and a seemingly limitless capacity for tissue homeostasis. Of the many freshwater planarian species known to exist, *Schmidtea mediterranea* has become one of the most widely studied [9]. Planarians thus provide a unique context in which to explore how nature has solved the complex problem of maintaining stem cell pluripotency in a long-lived adult animal.

Expression of conserved genes regulating pluripotency have been identified in planarian neoblasts and functionally studied using RNA interference [10-13]. However, due in part to the lack of methodologies for cell culture, exogenous gene expression, and transgenesis in planarians, the mechanisms regulating the pluripotency of these adult stem cells *in vivo* are poorly understood. Therefore, developing planarian transgenesis is of great significance [14]. A review of the history of cell culture methodologies and attempts to develop transgenics in planarians

indicated that successful neoblast culture may be a critical first step to develop

transgenic methodologies in planarians.

Transgenic approaches typically take advantage of either early stage embryos

or cultured stem cells. Invertebrates, such as *Caenorhabditis elegans*, *Drosophila*

*melanogaster*, Hydra, *Nematostella vectensis*, and the flatworm *Macrostomum*

*lignano,* have large syncytial germ cells or embryos, respectively, that are highly

amenable to genetic manipulation [14-18]. In vertebrates, such as mice, both zygotes

and cultured embryonic stem cells are used to deliver exogenous genetic material

[19]. Unlike these research organisms, planarians do not possess large, easily

accessible germ cells or early-stage blastomeres amenable to manipulation or

transplantation. Instead, in asexually reproducing planarians, neoblasts are the

only known proliferating cells in the animal [20]. Neoblasts from one animal can be

readily transplanted into a host devoid of its own endogenous neoblasts after lethal

irradiation, resulting in neoblast repopulation and host rescue within 1 month [6,8].

Thus, introduction of exogenous DNA into cultured neoblasts prior to

transplantation is a potential strategy to produce transgenic planarians. Cultured

neoblasts would also be ideal for rapidly screening conditions for delivering and

expressing exogenous mRNA or DNA. Therefore, we aimed to establish a robust

method for culturing pluripotent neoblasts that may allow rapid screening of

conditions for delivering and expressing exogenous mRNA or DNA.

Previous efforts to culture planarian cells were conducted at a time when our

mechanistic understanding of neoblast self-renewal and heterogeneity were

limited [21-24]. In some of these studies, cells with gross morphology similar to

neoblasts survived in an isotonic medium for a couple of weeks, yet neither

functional nor molecular tests on the cultured cells were performed, leaving an

open question as to their identity and potency [23]. Since then, the pan-neoblast

marker *smedwi-1* (a homolog of the Argonaute family of proteins) was identified [25],

allowing us to molecularly define and visualize neoblasts using gene expression

profiling or whole mount in situ hybridization (ISH). Techniques that enrich

neoblasts using flow sorting have also been developed. A cell cycle-based flow

sorting method using Hoechst 33342 staining has been used to isolate S and G2/M

cell cycle phase neoblasts (X1 cells; nearly 90% of X1 cells are *smedwi-1+*

neoblasts) [25,26]. However, Hoechst 33342 is cytotoxic, and X1 neoblasts cannot

proliferate *in vivo* after transplantation into lethally irradiated planarians lacking

stem cells. To solve this technical limitation, a DNA dye free back-gating strategy

using forward scatter (size) and side scatter (complexity) was shown to enrich for

a heterogeneous cell population containing neoblasts (X1(FS)) [8]. Unlike X1

neoblasts, X1(FS) neoblasts proliferate and successfully rescue lethally irradiated

planarians upon transplantation making the X1(FS) population suitable for the

development of an *in vitro* neoblast culture protocol [8]. When considered alongside

the formulation of new types of cell culture media [23,27], these advances provide a

groundwork to attempt establishing new, robust methods for *in vitro* culture of

pluripotent and transplantation-competent neoblasts.

In this study, we performed an unbiased screen of 23 different formulations of

cell culture media to identify the best nutrient conditions for flow cytometrically

isolated neoblasts. Cell morphology, viability, percentage of *smedwi-1+* cells,

clonogenic capacity after transplantation, and rescue efficiency were assayed to

identify the optimal conditions for culturing pluripotent neoblasts. Importantly, time-

lapse imaging captured neoblast division for the first time in culture in real-time.

Moreover, a novel neoblast isolation method using the vital dye SiR-DNA was

developed, improving the purification yields for neoblasts relative to X1(FS), while

preserving the clonogenic and rescue capacity of neoblasts following

transplantation. Finally, we developed electroporation conditions that can deliver

exogenous mRNA into cultured neoblasts providing unambiguous evidence that

exogenous mRNAs can be expressed, albeit with low efficiency, in cultured

neoblasts. Cumulatively, our work provides a foundation for developing long-term

neoblast culture methods and, ultimately, transgenic planarians. It also provides a

systematic methodological framework that may be applied to the development of

cell culture techniques in other invertebrate research organisms.

**Results**

**Identification of seven culture conditions that maintain viable dividing**

**neoblasts**

To test different culture conditions for neoblasts, we first used an established

back-gating method to sort X1(FS) cells, which contain approximately 23.4%±2.5%

neoblasts (*smedwi-1*+) (Fig. 1a-c). We then systematically screened 23 different

types of media, representing most commercially available or previously reported

formulations in the prescence (+) or absence (-) of 5% $CO_2$ (Supplementary Table

1) [23,24]. To assess each culture condition, five criteria were assayed: 1) cell

morphology and viability (viability); 2) percentage of *smedwi-1+* cells maintained

in culture (%neoblasts); 3) cell division; 4) clonogenic capacity after transplantation

(colony expansion); and 5) rescue efficiency in lethally irradiated planarians

(pluripotency) (Fig. 1a).

After 1 day of culture, cell morphologies were observed using transmitted light

microscopy. Cells cultured in CMFB +/- 5% $CO_2$ displayed abnormally roughened

cell morphologies accompanied by abundant cellular debris in the plate,

suggesting poor viability (Fig. 1d). In contrast, cells in all other conditions, such as

IPM +/- 5% $CO_2$, had normal morphology, suggesting high viability (Fig. 1d). Cells

in Leibovitz's L-15 medium (L15) without 5% $CO_2$ extended long processes that

were visible even after 6 days of culture (Supplementary Fig. 1), suggesting

neuronal differentiation as previously observed in cultured *Caenorhabditis elegans*

embryonic cells [28].

To measure viability, cells cultured for 1 day were stained with propidium iodide

(PI), which labels the DNA of dying cells, and the percentage of PI negative cells

was determined using flow cytometry. Consistent with the microscopic evaluation,

cells cultured in CMFB displayed poor survival +/- 5% $CO_2$ (>60% dead cells) (Fig.

1e). In fact, of all media conditions tested, seventeen yielded a viability of at least

60% (Fig. 1e), with only seven of the media performing significantly different in the

presence and absence of 5% $CO_2$ (Fig. 1e).

To determine what proportion of viable cells were neoblasts after 24 hours of

culture, we quantified the number of *smedwi-1+* X1(FS) cells by fluorescent *in situ*

hybridization (FISH). Importantly, all cultures without 5% $CO_2$ maintained fewer

*smedwi-1*+ neoblasts compared to those cultured in the presence of 5% $CO_2$,

except for diluted (d) SFX (Fig. 1f). Furthermore, of the 5% $CO_2$ cultures, seven

media maintained significantly more *smedwi-1*+ neoblasts than all other culture

conditions, including dGrace's medium, IPM, KnockOut DMEM, dL15 medium,

dKnockOut DMEM, dSchneider's medium, and dDMEM (Fig. 1f). Because dSFX

without 5% $CO_2$ failed to support neoblast culture as well as the seven media with

5% $CO_2$ we identified above (Fig. 1e), we did not explore it further in this study.

This result was supported by co-staining cells cultured in IPM + 5% $CO_2$ with

*smedwi-1* and the apoptotic/dead-cell marker, Annexin V (Supplementary Fig. 2);

no    co-labeling    was    observed,    indicating    that    neoblasts    were    viable

(Supplementary Fig. 2). Consistently, cell viability in these seven media + 5% $CO_2$

was consistently greater than 60% (Fig. 1e). We next examined whether *smedwi-*

*1*+ neoblasts were maintained after 3 days in culture using these seven media +

5% $CO_2$, and observed persistent *smedwi-1*+ cells in all culture conditions tested

(Fig. 1g). Thus, neoblasts can be maintained for at least 3 days *in vitro*. Therefore,

we focused on testing dGrace's, IPM, KnockOut DMEM, dL15, dKnockOut DMEM,

dSchneider's, and dDMEM media in subsequent optimization experiments.

Next, we assessed whether cultured neoblasts were capable of dividing *in vitro.*

Although an obvious increase in cell number was not noticed, low levels of both

symmetric and asymmetric neoblast divisions were observed in 1 day cultured

cells, as judged by cell pair size and distribution of *smedwi-1* transcripts (Fig. 1h)

[13]. Confirmation that neoblasts can divide *in vitro* was obtained using time-lapse

imaging microscopy to record the behavior of X1(FS) cells in culture. Both

symmetric and asymmetric cell divisions were observed within the first 24 hours

after culture (Fig. 1i and Supplementary Movies 1 and 2) in IPM, KnockOut DMEM,

and dL15 medium, but not in the other four media tested (Fig. 1i). Consistently, the

percentage of *PCNA+* cells in the cultures of IPM, KnockOut DMEM, and dL15

medium were significantly higher than those in CMFB, Schneider's, and DMEM

medium (Supplementary Fig. 3). Even though we cannot exclude the possibility

that these conditions only allow neoblasts in M phase to complete the cell cycle,

to our knowledge, this is the first time that neoblast divisions have been observed

and recorded *in vitro*. These results suggest that a fraction of X1(FS), *smedwi-1+*

cells can execute cell division within 24 hours after isolation in culture.

**Cultured neoblasts maintain clonogenic capacity**

To determine if X1(FS) neoblasts could divide *in vivo* following *in vitro* culture*,*

we next examined their clonogenic capacity following transplantation into lethally

irradiated planarians, an experimental manipulation that normally leads to robust

neoblast expansion (Fig. 2a). Serial cell dilution experiments indicated that 1,000

freshly collected X1(FS) cells undergo consistent colony expansion in ≥ 80% hosts

upon bulk cell transplantation (Supplementary Fig. 4). Considering the rate of cell

death in culture, 5,000 X1(FS) cells were cultured for each test condition to ensure

that enough cells were viable at the time of transplant. We transplanted X1(FS)

cells cultured in the seven different media + 5% $CO_2$ that showed higher than 15%

*smedwi-1+* cells (Figure 1f) for 1, 2, or 3 days. At 8 days post-transplantation (dpt),

the presence or absence of *smedwi-1+* neoblast colonies and the number of

*smedwi-1+* neoblasts in each colony were determined. All X1(FS) neoblasts

cultured for 1 or 2 days efficiently proliferated *in vivo*, except for those cultured in

dGrace's medium + 5% $CO_2$ (Fig. 2b–d). By comparing the number of *smedwi-1+*

neoblasts in each transplant, X1(FS) cells cultured for 1 day in either IPM or

KnockOut DMEM formed the largest colonies *in vivo* (Fig. 2b, d). X1(FS) cells

cultured for 2 days displayed decreased expansion potential in all conditions, but

all were still capable of forming colonies *in vivo* with the exception of those cultured

in dGrace's medium + 5% $CO_2$. X1(FS) cells cultured for 3 days were largely

incapable of forming colonies following transplantation, though small colonies

formed from cells cultured in dSchneider and dL15 media (Fig. 2c, d). In summary,

IPM and KnockOut DMEM performed best following the first day in culture, but

performed similarly to dKnockOut DMEM, dSchneider's, dL15, and dDMEM after

two days of culture. In addition, we observed that clonogenic capacity of X1(FS)

neoblasts diminished greatly following three days in culture, regardless of the

media used. These results suggest that IPM, KnockOut DMEM, dL15, dKnockOut

DMEM, dSchneider's, and dDMEM are capable of maintaining the proliferation

potential of neoblasts for up to two days in culture in the presence of 5% $CO_2$.

**Cultured neoblasts can rescue stem cell-depleted planarian hosts**

To evaluate the functional pluripotency of neoblasts cultured in these six media

(IPM, KnockOut DMEM, dKnockOut DMEM, dL15, dSchneider's, and dDMEM),

rescue was assessed following bulk-cell transplantation. Genotyping PCR and

restriction fragment length polymorphism (RFLP) assays were performed to test

whether sexual hosts had been transformed into the asexual biotype following

transplantation of the asexual neoblasts (Supplementary Fig. 5a) [8]. For non-

cultured, freshly collected X1(FS) cells, 30–50% of the lethally irradiated (6,000

rads) sexual *S. mediterranea* hosts were rescued (Fig. 3b, c, and Supplementary

Fig. 5b, e). Next, X1(FS) cells cultured in the indicated media for 1, 2, or 3 days

were transplanted into lethally irradiated hosts using the same method. X1(FS)

cells cultured in IPM, dL15, and KnockOut DMEM for 1 and 2 days were capable

of rescuing hosts devoid of stem cells (Fig. 3c and Supplementary Fig. 5c-e), of

which X1(FS) cells cultured in KnockOut DMEM displayed the highest and most

robust rescue efficiency (Fig. 3c and Supplementary Fig. 5e). Consistent with the

clonogenic assay results, none of the X1(FS) neoblasts cultured for 3 days rescued

lethally irradiated hosts. These data indicate that of all conditions tested, KnockOut

DMEM +5% $CO_2$ is the best one for maintaining pluripotent neoblasts in culture for

up to 2 days. IPM and dL15 medium were also capable of maintaining pluripotent

neoblasts in culture for up to 2 days albeit with reduced rates of irradiate animal

rescue after transplantation (Fig. 3c and Supplementary Fig. 5e).

In summary, we found that after screening 23 media followed by assaying 5

criteria (*i.e.*, viability, *smedwi-1* expression, cell division, clonogenic capacity and

rescue efficiency of irradiated animals), three types of media (KnockOut DMEM,

IPM, and dL15) were capable of maintaining pluripotent neoblasts *in vitro*. Of these

three different media, KnockOut DMEM produced cultured neoblasts with the

strongest performance across the multiple measured criteria, with IPM and dL15

medium performing slightly less well (Fig. 3d).

**Electroporation delivers fluorescent dextran into neoblasts**

Following the successful optimization of *in vitro* culture conditions for the

maintenance of pluripotent neoblasts, we next attempted to test conditions for the

delivery of exogenous molecules into neoblasts, the next step required for

developing transgenic methods for planarians. We first used dextran-FITC as a

fluorescent indicator to screen suitable electroporation conditions for neoblasts

labeled by Hoechst 33342 staining (Fig. 4a). We tested 52 electroporation

programs and 10 different buffers using X1 cells [25,26], and found that dextran-FITC

was delivered into neoblasts most efficiently in IPM buffer with electroporation at

100-120V (Supplementary Table 2 and Fig. 4b-d). When similarly applying the

electroporation method to X1(FS) cells, rather than Hoescht 33342 sorted X1 cells,

dextran-FITC+ populations could only be detected with electroporation values of

110V and 120V. However, less than 6% of dextran-FITC+ X1(FS) cells were

*smedwi-1+* neoblasts and virtually no *smedwi-1+* cells could be detected after 1

day culture in KnockOut DMEM +5% $CO_2$ (Fig. 4e). Consistent with the drastic

reduction in *smedwi-1+* cell viability post-electroporation, none of the donor X1(FS)

cell populations subjected to more than 100V formed colonies following

transplantation into lethally irradiated donors (Fig. 4f). We reasoned that the failure

was likely due to the low purity of *smedwi-1+* neoblasts in X1(FS), which was even

further reduced after electroporation. Therefore, it was necessary to develop a new

strategy for neoblast isolation that would result in both higher clonogenic and

pluripotent *smedwi-1+* cell enrichment than the X1(FS) sorting protocol.

We also tested whether X1(FS) can express exogenously delivered mRNA in

current culture conditions. We cloned a planarian endogenous gene, *Smed-*

*histone3.3* and fused with two copies of flag tag (2×flag). After electroporation and

one day of culture, cells electroporated with *Smed-histone3.3-2×flag* mRNA had

more anti-FLAG staining positive cells (9.7 $\pm$ 1.4%) than electroporated cells

without mRNA (1.2$\pm$0.7%) (Fig. 4g). Even though the anti-FLAG antibody stained

enucleated cells, nuclear localization signal in nucleated cells suggested

successful expression of *Smed-histone3.3-2×flag* mRNA (Fig. 4g and 4h). Even

though we did not detect the signal of Smed-HISTONE3.3-2×FLAG in *smedwi-1+*

cells (Supplementary Fig. 6), these data encouraged us to further enrich for

neoblasts to optimize cell culture conditions.

**A new flow cytometry protocol using SiR-DNA and Cell Tracker improves**

**yield of clonogenic, pluripotent, transplantable neoblasts**

To enrich for neoblasts, we tested three major types of cell-permeable DNA

stains  to enrich cycling neoblasts at G2/M cell cycle phases (DRAQ5, Vybrant

DyeCycle, and SiR-DNA). DRAQ5 staining remained cytotoxic similarly to Hoechst

33342. Vybrant DyeCycle staining failed to unambiguously discriminate among

distinct neoblast cell cycle phases by flow cytometry. However, the recently

developed DNA stain, SiR-DNA [29] proved to have low toxicity and enriched

*smedwi-1+* neoblasts to a ratio ~60% (Fig. 5a, b, f and Supplementary Fig. 7).

Comparison of *smedwi-1+* and *smedwi-1-* cell morphology in the isolated

populations showed that *smedwi-1-* cells were generally smaller than *smedwi-1+*

cells (Fig. 5b). To discriminate between small and large cells in the SiR-DNA+

population, the cytoplasmic dyes Cell Tracker Green (CT) and Calcein AM (CAM)

were tested in combination with SiR-DNA in neoblast isolation (Fig. 5c, d). Using

a dual dye staining strategy resulted in a significant increase in neoblast

enrichment, as judged by *smedwi-1+* ISH (Fig. 5e, f); SiR-DNA/Cell Tracker Green

costaining performed comparably to Hoechst 33342 staining for enriching *smedwi-*

*1+* neoblasts (Fig. 5e, f). We termed this new sorted cell population SirNeoblasts.

Unlike neoblasts derived from Hoechst 33342 sorts, SirNeoblasts proliferated *in*

*vitro* and underwent colony expansion *in vivo* after transplantation into lethally

irradiated planarians (Fig. 5g). Facilitated by SiR-DNA staining of DNA, the

separation dynamics of chromosomes in dividing SirNeoblasts were observed *in*

*vitro* (Supplementary Movies 3-5), confirming the occurrence of *bona fide* cell

division in the tested culture condition. Importantly, no noticeable difference in

colony sizes was observed at 7 dpt between X1(FS), single (SiR-DNA), and double

dye (SiR-DNA/CT) stained populations (Fig. 5g). Finally, both freshly isolated

SirNeoblasts and those cultured in KnockOut DMEM +5% $CO_2$ for one day were

capable of rescuing lethally irradiated planarians at frequencies comparable to

those seen with X1(FS) cells (Fig. 3c and Fig. 5h). Based on these results, we

conclude that SiR-DNA/CT dual labeling-based cell sorting can be used to isolate

clonogenic, pluripotent neoblasts that can be maintained in primary culture and

serve as donor cells in transplantation assays. To further characterize the

SirNeoblasts, we stained SirNeoblasts with Hoechst 33342 to analyze their cell

cycle. However, co-staining of SiR-DNA and Hoeschst 33342 resulted in a failure

to detect SiR-DNA staining. We then tested whether Hoeschst 33342 can stain

SiR-DNA stained cells, and found that Hoechst 33342 can replace the staining of

SiR-DNA, which showed the cell cycle distribution of SirNeoblasts consisted of

~17.89% at G1, 13.02% at S, and ~69.09% at G2/M cell cycle phases

(Supplementary Fig. 7). This reversible staining of SiR-DNA may also explain the

reason why SirNeoblasts can proliferate after staining unlike Hoechst 33342

stained X1 cells.

Next, we determined conditions to optimize electroporation efficiency and

viability for SirNeoblasts (Fig. 6a). Consistent with previous studies,

electroporation at 110V-120V was required for dextan-TMR entry into SirNeoblasts

(Fig. 6b, c). As expected, *smedwi-1+* cells were more abundant in the 110 V and

120V electroporated SirNeoblasts compared to X1(FS) cells, and some

electroporated SirNeoblasts persisted for one day in culture (Fig. 6d). Importantly,

110V – 120V electroporated SirNeoblasts were capable of forming colonies and

rescuing lethally irradiated hosts upon transplantation (Fig. 6e, f). However, 120V

electroporations resulted in comparably fewer irradiated hosts being rescued after

SirNeoblast transplantations (Fig. 6e, f), indicating that high voltages may have a

negative impact on pluripotency.

**Exogenous mRNA delivered by electroporation can be successfully**

**expressed in SirNeoblasts**

To assess whether exogenous mRNA could be delivered into SirNeoblasts

using the described electroporation conditions, *tdTomato* mRNA was added to the

electroporation reaction along with Dextran. Dextran positive SirNeoblasts were

sorted and cultured in KnockOut DMEM + 5% $CO_2$. To determine whether mRNA

was successfully delivered, we probed cells via FISH 20 hours after

electroporation. *tdTomato* mRNA signal was detected in both 110V and 120V

electroporated cells, suggesting a successful delivery of exogenous mRNA into

SirNeoblasts (Fig. 6g, h). However, costaining with *smedwi-1+* revealed that not

all *tdTomato* mRNA+ cells retained neoblast identity in culture. The number of

sorted SirNeoblasts responded similarly to X1 and X1(FS) cells with respect to

electroporation in that the number of cells positive for both tdTomato mRNA and

*smedwi-1* expression was significantly higher after 110V electroporation than after

120V (Fig. 6h). This result indicates that under the conditions tested, 110V

electroporation may be the most suitable to both introduce exogenous, charged

molecules such as RNA into neoblasts, while maintaining their viability and

potency.

Unfortunately, expression of tdTomato was not detected by either microscopy

or antibody staining. Two possibilities were suspected: 1) The culure condition is

not good enough to support the translation of the delivered mRNA; 2) There is an

unknown mechanism that prevents the translation of the delivered mRNA. A recent

discovery in *C. elegans* indicated that endogenous piRNAs can target on the

exogenous transgene sequences and prevent their translation [30]. Similarly,

planarian neoblasts contain abundant PIWI and piRNAs. We thus hypothesize that

a similar piRNA targeting mechanism may exist in planarian neoblassts, which may

prevent the translation of the delivered mRNAs. We tested this hypothesis by

synthesizing multiple mRNAs encoding the fluorescent protein mCherry in which

conservative nucleotide substitutions were introduced in order to minimize

potential pairing of the exogenous mRNA with endogenous piRNAs, as was

recently described in *C. elegans* [30]. The synthetic mCherry mRNAs were tested

via electroporation into SirNeoblasts (Fig. 7a). Significantly, we found one mCherry

mRNA construct that resulted in robust mCherry+ cultured SirNeoblasts (Fig. 7b-

e, twice with high expression, five times with medium/low expression, ten times

without expression). Even though we have yet to fully comprehend the

mechanisms that may be underpinning piRNA targeting in neoblasts, the

successful expression results indicated that the culture and electroporation

conditions defined in our study are capable of maintaining neoblasts in culture

capable of retaining both pluripotency (Figures 5g, h and 6e, f) and translational

activity (Figure 7b). Although the efficiency by electroporation is low for mRNA

delivery, our current study is focused on developing a reliable method for culturing

neoblasts. Increasing the efficiency of delivery for mRNA and testing Cas9 and

guide RNAs is clearly necessary and will require further studies.

In summary, we defined a novel FACS isolation strategy and primary cell

culture conditions capable of maintaining clonogenic, pluripotent neoblasts *in vitro*

that are compatible with transplantation, repopulation and rescue of lethally

irradiated hosts. In addition, we optimized electroporation conditions that

successfully introduced fluorescent dextran and exogenous mRNA into

clonogenic, pluripotent neoblasts. These technical milestones are prerequisites for

the successful generation of transgenic planarians.

**Discussion**

Past efforts to culture planarian cells have been unable to convincingly

demonstrate that pluripotent neoblasts could be maintained in culture [23,24,31,32].

Here, we provide definitive molecular and functional evidence that pluripotent

neoblasts can be maintained *in vitro*. This technical advance facilitated the first

real-time observation of neoblast cell division within the first 24 hours after cell

culture *in vitro* (Supplementary Movies 1 and 2) and the first demonstration that

exogenous molecules, including fluorescent conjugated dextrans and mRNA, can

be delivered into planarian cells. This method establishes the required foundation

for future transgenic and genome editing technique development in planarians,

and opens exciting new avenues for a systematic investigation of the biology of a

naturally occurring population of pluripotent adult stem cells.

**The vital fluorescent dye SiR-DNA improves purification of pluripotent**

**neoblasts**

Prior to this study, The use of Hoechst 33342 staining has been broadly

adopted for isolating cycling neoblasts (X1 cells) by FACS. However, X1 cells

labeled with this nuclear dye cannot proliferate *in vivo* following their

transplantation into irradiated hosts. We sought to overcome this limitation by

testing alternative DNA dyes, such as DRAQ5 and Vybrant Dye Cycle, yet they

resulted in cytotoxicity and failed to unambiguously discriminate between distinct

neoblast cell cycle phases by flow cytometry. However, we found that unlike other

vital DNA dyes tested, the recently developed SiR-DNA dye [29] was not cytotoxic,

and when combined with Cell Tracker Green staining, significantly improved

pluripotent neoblast yields by flow cytometry. Together with cell subtype-specific

antibodies, SiR-DNA may allow for more specific dissection of the pluripotent

neoblast population by facilitating the isolation and functional characterization of

different neoblast subpopulations. Furthermore, this reagent may prove useful for

the isolation of viable proliferating cells in other organisms.

**Neoblast cell culture paves the way for transgenesis in planarians**

Transgenesis in planarians has been lacking for decades. Without a planarian-

specific positive control, it has been difficult to determine why exogenous nucleic

acids fail to be translated when introduced into neoblasts. Isolated neoblasts

provide an obvious proving ground for delivery of exogenous materials. While

neoblast transplantation can be performed immediately after delivering exogenous

molecules, the uncertain viability of neoblasts during and after transplantation

made this strategy ineffective. The cell culture system we have developed makes

it possible to trace and study each cell following delivery of exogenous materials

*in vitro*. First, it allows for ease of screening of constructs using a small number of

cells under conditions where neoblast purity and viability are well-established.

Second, when introducing transformed cells into lethally irradiated hosts to monitor

behavior *in vivo*, we can enrich for positive cells via FACS prior to transplantation,

minimizing the effects of cell-cell competition in a heterogeneous donor cell

population. Hence, neoblasts cultured using the methods described here lend

themselves accessible for testing a diversity of delivery methods. For instance,

custom-engineered liposomes were shown to facilitate the transfection of double-

stranded RNA and anti-miRNAs into planarian cells *in vivo* [33]. As such, it should

be possible to use liposomes to deliver larger molecules and genome-editing tools

in an effort to obtain higher neoblast transfection efficiency and further improve the

likelihood of producing transgenic animals. Thus, our methodology not only stands

to facilitate cell transformation, but may also play a key role in efforts to establish

long-term culture systems and/or cell lines.

**piRNA silencing mechanism for transgenes may be of broad occurrence**

**across metazoans**

Efforts to generate transgenic planarians span several decades with little to no

success reported thus far. The reasons for this state of affairs have been generally

associated with technical limitations of both culture conditions and delivery

methods of exogenous nucleic acids into neoblasts. Little consideration, however,

has been given to the possibility that such prolonged failure may be underscored

by unknown aspects of neoblast biology. Given that neoblasts are the *de facto*

units of selection in planarians and that the viability of these animals heavily

depends on the proper function and health of neoblasts, strong positive selection

for evolving robust mechanisms to protect the genome of these cells should be

expected.

piRNAs are small non-coding RNAs that have been shown to be essential for

safeguarding genome integrity by silencing transposable elements [34]. However, it

is also known that many piRNAs do not map to transposable element sequences

in various animals, including mice, *C. elegans* and planarians [35-37]. In fact, the

function of these piRNAs remains largely unknown. Additionally, it has also been

known for decades that transgenes with foreign sequences can be frequently

silenced in the germline of *C. elegans* [38]. Recent studies have begun to shed light

on piRNA function in both *Drosophila* [39] and *C. elegans* [40]. It was recently reported

that the repression of transgenes in the germline of fruitflies could be lifted by using

a UAS-promoter free of interference by Hsp70 piRNAs as silenced [39]. Also, it is known

that the PIWI protein PRG-1 is required for the silencing phenomenon observed in

the germline of *C. elegans*, suggesting a function of piRNAs in this process [40].

More recently, it was discovered that a mechanism targeting transgene sequences

introduced into the syncytial ovary of *C. elegans* involves piRNAs, and that a

sequence-based strategy to bypass transgene silencing by these small non-coding

RNAs allowed expression of exogenously added genes in the germline of this

animal [30].

Given the ancestral origin of PIWI proteins and piRNAs, we hypothesized that

similar mechanisms may be operating in planarian neoblasts. Our current study

showed that exogenous mRNAs in which predicted piRNA targeting sequences

were changed overcame siencing and allowed the translation of the reporter

protein (Figure 7b). However, we do not yet fully understand the piRNA recognition

rules in *S. mediterranea*. The size of planarian piRNAs are ~32nt, in contrast to

~22 nt in *C. elegans* [37,41], so the models prediciting targeting of piRNA in

nematodes do not fully transpose to planarians. Additionally, planarians have at

least three PIWI proteins [25,37], raising another question as to which PIWI proteins

may or may not be required for producing piRNAs that may potentially target

exogenous nucleic acids. Definitive experiments to test this hypothesis are

necessary and future and ongoing research will help in resolving these issues and

testing and refining our predictive piRNA targeting models in the hopes of

producing the most stable exogenous nucleic acid molecules for introduction into

neoblasts.

**A method for mechanistic studies of neoblast proliferation and**
**differentiation** *in vitro*

The paucity of cell culture conditions for invertebrates in general, and
planarians in particular, has hampered our ability to test and identify factors directly
regulating the functions of neoblasts, a remarkably abundant and pluripotent adult
stem cell population. For example, our understanding of how extracellular growth
factors modulate neoblast proliferation is still in its infancy. In planarians, several
of these factors have been shown to have important functions in neoblast
proliferation or homeostasis. For instance, knockdown of *smed-neuregulin (nrg)-7*
or *smed-insulin-like peptide-1* impairs neoblast proliferation *in vivo* [13,42]. We
hypothesize that addition of these factors, or potentially other purified extracellular
growth factors, may boost neoblast proliferation *in vitro*. However, no *in vitro*
culture system had been developed to test this hypothesis. With the methods and
results presented here open the door to test the effects of planarian extracellular
extracts or purified extracellular growth factors from planarian species on the
proliferation and maintenance of neoblasts. Additionally, our protocols lend
themselves to initiate a systematic comparison of the metabolomics of cultured
neoblasts with those found *in vivo*. Such studies will aid in further optimization of
culture conditions and may ultimately lead to the controlled manipulation of cell
metabolism to predictably regulate neoblast proliferation and differentiation *in vitro*.

**Defining the neoblast niche in planarians**

The existence of a niche that supports the proliferation and differentiation of
neoblasts has been previously proposed [43]. This hypothesis has been supported

by indirect evidence [13,44,45]. However, the molecular and cellular nature of the niche

is largely unknown. Transplant experiments carried out in this study showed that

a limited number of neoblasts can be maintained in the transplanted location and

may continue their proliferation and differentiation. Because of the limited number

of cells surviving after transplantation, dissecting the cellular microenvironment of

transplanted neoblasts is likely to be a promising context for a mechanisitic

characterization of the proposed neoblast niche. Together with sublethal irradiation

assays, the cell culture tools reported here should afford us the opportunity to

understand how pluripotency and cell fate may be cell- and non-cell autonomously

regulated in a highly regenerative context.

**A framework for establishing cell culture in new research organisms**

Since the development of cell lines in the 1950s [46], cell culture has enabled

scientists to study fundamental aspects of cell biology. In recent years, the number

of research organisms being employed to address and discover new biology has

steadily increased. However, a comparatively small number of cell types have

been successfully cultured *in vitro*, particularly for invertebrates. The current study

systematically screened the majority of published cell culture media and optimized

culture conditions for planarian neoblasts. Thus, the systematic development of

cell culture methods reported here not only advances the study of cell bbiology in

the highly regenerative planarian *S. mediterranea*, but should also facilitate the

establishment of culture methods for other species, particularly underrepresented

invertebrate research organisms.

## Experimental Procedures

### Planarian care and irradiation treatment

Asexual (Clone CIW4) and sexual (Clone S2F1L3F2) strains of *Schmidtea mediterranea* were maintained in Montjuïc water at 20ºC as previously described [8,20]. Animals were starved for 7–14 days before each experiment. Animals exposed to 6,000 rads of γ rays were used as transplant hosts [8]. After transplantation, hosts were maintained in Montjuïc water with 50 µg/ml Gentamicin (GEMINI, 400-100P). For transplant rescue experiments, host animals were maintained in 3.5 cm Petri dishes (1 worm/dish), and Montjuïc water was changed every 2–3 days.

### Cell collection and culture

X1(FS) cells were collected as previously described with minor modifications [8,25]. Tails from planarians (>8 mm in length) were macerated in Calcium Magnesium free buffer with 1% Bovine Serum Albumin (CMFB) (see Recipe in Table S1) for 20–30 min with vigorous pipetting every 3–5 min. After maceration, dissociated cells were centrifuged at 290g for 10min. Cells were then resuspended in IPM with 10% Fetal Bovine Serum for either Hoechst 33342 or SiR-DNA staining. To gate the X1(FS) cells, the X1 population from a control sample stained with 0.4 mg/ml Hoechst 33342 (ThermoFisher Technologies, H3570) was used to define the forward scatter/side scatter gate. To obtain SirNeoblasts, dissociated cells were stained with SiR-DNA (1µM, Cytoskeleton Inc., CY-SC007) and CellTracker Green CMFDA Dye (2.5µg/ml, Thermo Fisher Technologies, C7025) for 1 hour and 10 min sequentially. Cells were sorted with an Influx sorter using a 100-µm tip. For time-lapse imaging experiments, X1(FS) cells were incubated in

either 5 mL of the indicated culture medium per well in 6-cm dishes (MatTek,

P35G-1.5-14-C) or in 1 mL of the indicated culture medium per well in a 24-well

plate (MatTek, P24G-1.5-13-F). For other experiments, X1(FS) were incubated in

50 µL of the indicated culture medium per well in 384-well plates (Greiner bio-one,

781090). Cells were cultured in indicated media containing 5% Fetal Bovine Serum

(Sigma-Aldrich, F4135) at 22ºC, +/- 5% $CO_2$. Dishes and multi-well plates were

pre-coated with poly-D-lysine (50µg/ml, BD Biosciences).

***In situ* hybridization and antibody staining**

Whole-mount *in situ* hybridization was carried out as previously described [13,47-

49]. For ISH on cultured cells, cell culture plates were centrifuged in an Eppendorf

horizontal centrifuge (Centrifuge 5810 R) at 300 *g* x 3 min. Cells were fixed with

3.7% formaldehyde (Sigma-Aldrich, F8775) or 4% paraformaldehyde (Electron

Microscopy Sciences, 15710) for 20 min. After washing with $1\times$ PBS, cells were

hybridized with riboprobes at 56ºC for at least 15 h. After washing with $2\times$ SSC

and $0.2\times$ SSC, cells were incubated with anti-digoxigenin-POD (Roche

Diagnostics, 11207733910) or anti-fluorescein-POD (Roche Diagnostics,

11426346910) at room temperature for 2 h. After washing with $1\times$ PBS/0.3%

TritonX-100, the signal was developed using tyramide-conjugated Cy3 (Sigma-

Aldrich, PA13101) or Cy5 (Sigma-Aldrich, PA15101).

Anti-phospho-Histone H3 (Ser10) (H3P) antibody (1:1,000, Abcam, ab32107)

and Alexa 555-conjugated goat anti-rabbit secondary antibodies (1:1,000, Abcam,

ab150086) were used to stain proliferating cells at the G2/M phase of the cell cycle.

**Annexin V staining**

Fifty microliters of cultured cells were re-suspended and stained with 2.5 µl of

Annexin V FITC Conjugate (BioLegend, 640905) at room temperature for 15 min.

After washing twice with IPM + 10%FBS, cells were subjected to *smedwi-1* ISH.

Thereafter, anti-fluorescein-POD (Roche Diagnostics, 11426346910) was used to

stain Annexin V for apoptotic and dead cells detection.

**Cell transplantation**

X1(FS) cells collected by flow cytometry were transplanted into irradiated hosts

(6,000 rads) as previously described with minor modifications [8]. Approximately 1

µL of an X1(FS) cell suspension (5,000 cells/µL) was injected into either the post-

pharyngeal midline (of asexual CIW4 hosts) or the post-gonopore midline (of

sexual S2F1L3F2 hosts) at 0.75–1.0 psi (Eppendorf FemtoJet) using a borasilicate

glass microcapillary (Sutter Instrument Co., B100-75-15).

**mRNA synthesis and electroporation**

mRNAs used for electroporation were prepared following the standard

protocols in the mMESSAGE mMACHINE T7 ultra Transcription Kit (ThermoFisher

Technologies, AM1345) and the Ambion RNA Purification Kit (ThermoFisher

Technologies, AM1908). tdTomato mRNA was transcribed from the linearized

plasmid pcDNA3.1(+)-tdTomato. mCherry and T7 promoter sequences were

cloned into the pIDT vector and synthesized by IDT Inc. Primers used to amplify

the template were 5′-CAGATTAATACGACTCACTATAGG-3′ and 5′-

ACTGATAATTAACCCTCACTAAAG-3′.

To screen electroporation conditions, cells from four tail fragments were

suspended in 20 µL electroporation buffers following Heochast 33342 staining. 20

μg Dextran-FITC (ThermoFisher Technologies, D3306) were mixed with cells and

loaded into a 1mm electroporation cuvette for BTX ECM830 electroporator or a

12-well electroporation strip for Lonza 4D electroporator. The buffer SE, SG, SF,

P1-5 were electroporation buffers in Lonza Cell Line and Primary Cell 4D-

Nucleofector Optimization kits (V4XC-9064 and V4XP-9096). Cell viability and

electroporation efficiency were assessed using an Influx sorter.

For exogenous mRNA electroporation, ~1x10$^8$ cells were suspended in 50 μL

IPM following SiR-DNA staining. 50 μg Dextran-FITC and ~5   μg mRNA were

mixed with cells and loaded into a 1mm electroporation cuvette. BTX ECM830

electroporator was used to apply a 110 V and 1 millisecond square wave pulse to

deliver dextran-FITC and mRNA into planarian cells. Dextran-FITC+ SirNeoblasts

were purified using an Influx sorter and cultured in KnockOut DMEM + 5%FBS.

**Microscopy and time-lapse imaging**

The Celigo imaging cell cytometer (Celigo, Inc.) and the Falcon 700 confocal

microscope were used to take pictures of X1(FS) and SirNeoblasts following ISH.

Celigo or ImageJ software was used for quantitative analyses. Time-lapse imaging

of cultured cells was performed using a Nikon Eclipse TE2000-E equipped with

Perfect Focus and a Plan Fluor ELWD 20X/0.45 NA Ph1 objective. Micro-manager

was used to control the microscope and Hamamatsu Orca R2 CCD [50]. Multiple

positions were acquired at 5-min intervals for 24–48 h. *In situ* hybridization

samples were imaged with a Nikon Eclipse Ti equipped with a Yokogawa W1

spinning disk head and a Prior PLW20 Well Plate loader. Several slides were

prepared at once and then loaded and processed automatically using a

combination of Nikon Elements Jobs for all robot and microscope control and Fiji

for object-finding and segmentation. Slides were imaged at low magnification and

objects identified before re-imaging tiled z-stacks using a Plan Apo 10X 0.5NA air

objective. Tiled images were stitched, projected, and *smedwi-1+* puncta were

counted using custom macros and plugins in Fiji.

**Generation of optimized mCherry sequence**

mCherry candidate sequences were generated by means of a custom python

script. Amino acid sequences were back translated to 21 nucleotide sequences

from 7 amino acid words at a time. Each potential nucleotide sequence was

screened against a list of known piRNAs to generate the sequence with the fewest

piRNA matches. A piRNA match consists of no more than a single G/T mismatch

in the 6 nucleotide seed region (positions 2-7 of a piRNA) [30]. Additional G/T

mismatches were scored as .5 and other mismatches as 1. Only the first 21

basepairs of the piRNAs were aligned. The highest scoring piRNA determined the

score for that potential nucleotide sequence. The 21 nucleotide sequence with the

lowest score was retained. The script was run with four alternate coding

tables. The "all" coding table contained all possible codons for each amino acid.

The "smed" coding table contained only those codons known to be most used in

*S. mediterraea* [51]. "lowgc" contained only those codons with the fewest G or C

nucleotides. "highgc" contained only those codons with the most G or C

nucleotides. The "highgc" sequence is shown in Figure 7. The other three

sequences as well as 5 additional sequences generated by shuffling the four

generated sequences and one sequence generated by backtranslating the amino

acid sequence with sms failed to show fluorescence [52].

**Data availability**

637    All codes used for plugins in Fiji are available at:https://github.com/jouyun. All

638    original data underlying this manuscript can be accessed from the Stowers Original

639    Data Repository at: http://www.stowers.org/research/publications/libpb-1281. All

640    reagents are available from the corresponding author upon reasonable request.

**Statistical analyses**

642    Microsoft Excel and Prism 6 were used for statistical analysis. Mean ± s.e.m.

643    is shown in all graphs. Unpaired two-tailed Student's *t*-test was used to determine

644    the significant differences between two conditions. $p < 0.05$ was considered a

645    significant difference.

**Acknowledgments**

We thank I. Wang and P. Reddien for assistance with the transplantation

technique. We thank all members of Sánchez Lab, especially J. Jenkin and C.

Guerrero for animal maintenance and irradiation assistance, L. C. Cheng and E.

Duncan for technical help, and B. Benham-Pyle, E. Davies, and S. Elliot for

comments on the manuscript. We acknowledge all members of the Reptile &

Aquatics, Molecular Biology, Cytometry, and Microscopy Core Facilities at the

Stowers Institute for Medical Research for technical support. This work was

supported by NIH R37GM057260 to A.S.A. A.S.A. is a Howard Hughes Medical

Institute and a Stowers Institute for Medical Research Investigator.

**Author contributions**

K.L. and A.S.A. conceived the project, designed experiments, analyzed data,

and wrote the manuscript. K.L. performed all experiments and data acquisition.

S.A.M. performed the time-lapse imaging experiments and analyzed raw spinning-

disk imaging data. E.J.R. and H.-C.L. designed the variant sequences for mCherry.

**Competing interests**

The authors declare no conflicts of interest.

**Figure Legends**

**Figure 1. Systematic screen identifies cell culture conditions for maintaining**

**X1(FS) neoblasts *in vitro*. (a)** Flowchart illustrating steps of X1(FS) cell culture and

criteria used to identify best culture condition for neoblasts: cell viability, percentage of

*smedwi-1+* neoblasts (%*smedwi-1+*), cell division *in vitro*, colony expansion after

transplantation, and rescue efficiency of irradiated hosts after transplantation

(pluripotency). **(b)** Plots showing the FACS gating to sort X1(FS) cells. **(c)** Representative

images showing *smedwi-1+* neoblasts among the sorted X1(FS) cells. Scale bar, 20 µm.

X1(FS) cells consistently contains 23.4%±2.5% neoblasts in total DAPI+ cells scored.

Three replicates were assayed, n=100 to 150. **(d)** Representative images of cell

morphologies observed after 1 day of culture +5% $CO_2$, including poor cell morphology in

CMFB and healthy cell morphology in IPM (arrowheads). Scale bar, 20 µm. **(e)**

Percentages of live cells (Propidium Iodide-negative) among 23 media, +/- 5% $CO_2$, after

1 day of culture. Knockout DMEM + 5% $CO_2$ yielded best overall cell viability. Three

replicates were assayed, n=500 to 1200. **(f)** Percentage of *smedwi-1+* neoblasts after 1

day of culture under indicated conditions. Significantly more *smedwi-1+* neoblasts were

maintained in seven media + 5% $CO_2$ compared to all other conditions. Three replicates

were assayed, n>500. **(g)** Percentage of *smedwi-1+* neoblasts after 3 days of culture in

indicated media + 5% $CO_2$. **(h)** Representative images of dividing cells undergoing either

symmetric cell division (SCD) or asymmetric cell division (ASCD). Scale bar, 10 µm. **(i)**

Time-lapse images of dividing cells undergoing either SCD or ASCD in IPM + 5% $CO_2$.

Scale bar, 10 µm. Both SCD and ASCD can be observed in ~300 X1(FS) cells cultured

in IPM, KnockOut DMEM, and dL15 + 5% $CO_2$.

Neoblast culture and transformation

**Figure 2. Cultured X1(FS) neoblasts expand after transplantation**. **(a)** Flowchart

showing steps of X1(FS) cell transplantation following *in vitro* culture. **(b)** Representative

images showing colonies of *smedwi-1+* neoblasts at 8 days post-transplantation (dpt)

cultured in the indicated conditions + 5% $CO_2$. Only X1(FS) cells cultured in dGrace's

medium + 5% $CO_2$ did not efficiently form colonies *in vivo*. Scale bar, 200 µm. **(c)**

Percentage of hosts receiving X1(FS) cells cultured in indicated media + 5% $CO_2$ for 1, 2,

or 3 days that possessed *smedwi-1+* colonies (green bars) or H3P+ colonies (red bars)

at 8 dpt. **(d)** Number of *smedwi-1+* neoblasts in colonies formed by X1(FS) cells at 8 dpt

following culturing in indicated media + 5% $CO_2$ for 1, 2, or 3 days. Ten to twelve animals

assayed per condition.

**Figure 3. Cultured X1(FS) cells rescue neoblast-depleted planarians. (a)** Flowchart

illustrates steps of rescue assay using cultured X1(FS) cells. **(b)** Representative images

showing rescue of lethally irradiated hosts following transplantation of freshly isolated

X1(FS) cells, culminating in fission at 95 dpt. Scale bar, 200 µm. **(c)** Rescue rates for

lethally irradiated hosts following the transplantation of X1(FS) cells cultured in the

indicated media + 5% $CO_2$ for 1, 2, or 3 days. Histogram indicates averaged percent from

replicate experiments. Ten to twelve animals assayed per condition in each replicate

experiment. **(d)** Summary of 23 cell culture media screen using the following criteria: cell

morphology, cell viability, %*smedwi-1+* neoblasts, ability of transplanted cells to form

colonies and expand *in vivo* (clonogenesis)*,* and ability to rescue lethally irradiated

Neoblast culture and transformation

animals (pluripotency). Overall, KnockOut DMEM was the most effective medium for

maintaining pluripotent neoblasts in culture for 2 days.

**Figure 4. Electroporation can deliver Dextran-FITC into neoblasts. (a)** Flowchart

describing electroporation assay steps to screen for best conditions for cell viability and

Dextran-FITC deilivery efficiency. **(b)** Plots of X1 viability (upper) and electroporation

efficiency (lower) by using IPM as the electroporation buffer to deliver Dextran-FITC at

120V compared to 0 V controls. **(c)** Representative images of sorted Dextran-FITC$^{low}$ and

Dextran_FITC$^{high}$ cells indicate successful delivery of Dextran-FITC at 120V. **(d)** Viability

(blue) and electroporation efficiency (red) on X1 cells after electroporation using IPM as

electroporation buffer. **(e)** %*smedwi-1+* neoblasts in X1(FS) cells after 100V, 110V, and

120V electroporation immediately (black column) and after 1 day of culture in KnockOut

DMEM + 5% $CO_2$ (white column). Four random fields assayed per condition. $p < 0.05$ for

120 V. N>40. **(f)** Electroporated X1(FS) cells receiving greater than 100 V failed to form

colonies following transplantation. Ten animals assayed per condition. **(g)** Representative

images of electroporated X1(FS) with (upper panel) or without (lower panel) *Smed-*

*histone3.3-2×flag* mRNA. Arrowheads: anti-FLAG+ nucleated cells. Stars: anti-FLAG+

enucleated cells. Scale bar, 20 μm. **(h)** Z-stack images of an nucleated anti-FLAG+ cell.

Scale bar, 10 μm.

**Figure 5. SiR-DNA plus Cell Tracker staining and cell sorting protocol enriches for**

**clonogenic, pluripotent *smedwi-1+* neoblasts. (a)** Plots showing the gate used to

isolate SiR-DNA+ cells for *smedwi-1* ISH. **(b)** *smedwi-1* ISH on isolated cells from the

SiR-DNA+ gate shown in Supplementary Fig. 7a. *smedwi-1-* cells (arrows) were generally

smaller than *smedwi-1+* cells (stars). Scale bar, 20 µm. **(c-d)** Plots showing the gates

used to isolate SiR-DNA+, calcein-AM+ cells **(c)** and SiR-DNA+, Cell Tracker Green+

cells **(d)** for *smedwi-1* ISH. **(e)** *smedwi-1* ISH for SIR-DNA+ neoblasts populations

indicated in **(c)**. Scale bar, 20 µm. **(f)** %*smedwi-1+* neoblasts in indicated FACS isolated

populations. SiR-DNA and Cell Tracker Green dual staining enriches for *smedwi-1+*

neoblasts (SirNeoblasts) comparably to the Hoechst 33342 stained X1 population. *,

p<0.05. **, p<0.01. n.s., no significance. Four random fields assayed per condition. N>70.

**(g)** Representative images showing the clonogenic capacity of transplanted neoblasts

obtained using different FACS isolation protocols. No noticeable difference in the colony

expansion was observed among single and double dye staining populations at 7dpt.

Scale bar, 200 µm. Ten animals assayed per condition. **(h)** Rescue efficiency of fresh and

1-day cultured SirNeoblasts. CT: cell tracker green.

**Figure 6. SirNeoblasts can be used for exogenous mRNA electroporation. (a)**

Flowchart presenting the steps of neoblast electroporation using SirNeoblasts. **(b)** Plots

showing electroporation efficiency of SirNeoblasts at 100V, 110V and 120V compared to

0V. **(c)** Neoblasts after electroporation of Dextran-FITC showing 100% isolation of

positive cells after electroporation at 110V and 120V. All SirNeoblasts were free of

Dextran-FITC without electroporation treatment. Scale bar, 20 µm. **(d)** Percentage of

*smedwi-1+* cells after electroporation, suggesting a relative high ratio of neoblasts after

electrporation by using SirNeoblasts compared to X1(FS) in **Fig. 4e**. Four random fields

assayed per condition. *, p< 0.05 (120V SirNeoblasts vs. 120V X1(FS) at 1 day) . **,

Neoblast culture and transformation

p<0.005 (110V SirNeoblasts vs. 120V SirNeoblasts at 1 day, 110V SirNeoblasts vs. 110V

X1(FS) at 0 day, and 120V SirNeoblasts vs. 120V X1(FS) at 0 day). ***, p<0.001 (110V

SirNeoblasts vs. 110V X1(FS) at 1 day). **(e)** Representative images showing the colony

expansion of electroporated SirNeoblasts after transplanation Scale bar, 200 µm. N=14

for 110V and =10 for 120 V. **(f)** Rescue efficiency of electroporated SirNeoblasts. Scale

bar, 200 µm. **(g)** Representative images showing the mRNA signals (white dots) in cells

1 day after 110V and 120V electroporation. Scale bar, 20 µm. **(h)** Percentage of total cells

and *smedwi-1*+ cells containing mRNA 1 day after 110V and 120V electroporation. n.s.:

not significant. ** < 0.01.s

**Figure 7. mCherry mRNA is expressed in SirNeoblasts. (a)** A flowchart describes

steps of SirNeoblast electroporation using mCherry mRNA. **(b)** Representative images of

mCherry mRNA electroporated SirNeoblasts cultured in KnockOut DMEM for 1 day.

Upper: electroporated SirNeoblast without mRNA in culture. Lower: mCherry mRNA

electroporated SirNeoblasts in culture. Scale bar, 20 µm. **(c)** Plot showing no mCherry

expression after 110V electroporation without mCherry mRNA. **(d)** Plot showing ~5%

mCherry+ cells after 110V electroporation with mCherry mRNA. **(e)** Representative

images of cells from mCherry- population in (upper row) and mCherry+ population in

(lower row). Cells from mCherry+ population showed obvious mCherry localization in

cytoplasm. Scale bar, 20 µm.

Neoblast culture and transformation

**Supplementary Figure 1. X1(FS) cells cultured in L15 extend long cellular**

**processes**. **(a–d)** Four representative images showing long cellular processes from cells

after 6 days of culture in L15 without 5% $CO_2$. Scale bar, 20 µm.

**Supplementary Figure 2. s*medwi-1*+ X1(FS) neoblasts are viable**. X1(FS) cells were

cultured in IPM + 5% $CO_2$ for 2 days. Representative images of apoptotic cells (Annexin

V, green, arrowheads) co-stained with the pan-neoblast marker *smedwi-1* (magenta),

n=37. Two independent replicate experiments were performed. No co-labeling was

observed, suggesting neoblasts examined in study were viable. Scale bar, 20 µm.

**Supplementary Figure 3. IPM, Knockout DMEM, and dL15 maintain more *PCNA+***

**cells.** Percentage of *smedwi-1+* neoblasts after 1 day of culture in indicated media + 5%

$CO_2$.

**Supplementary Figure 4. Determining the number of X1(FS) cells needed for**

**efficient colony expansion**. **(a)** Percentage of lethally irradiated hosts displaying robust

neoblast colony expansion following transplantation with the indicated numbers of sorted

X1(FS) cells. At 7 days post-transplantation (dpt), > 80% of all hosts displayed colony

expansion when 1,000 X1(FS) were transplanted. **(b)** Representative images of hosts

transplanted with X1(FS) cells at 7 dpt. *smedwi-1+* neoblasts: green. DAPI: blue. Scale

bar, 200 µm. Ten animals assayed per condition.

Neoblast culture and transformation

**Supplementary Figure 5. Sexual hosts are rescued and reconstituted by**

**transplantation of cultured asexual X1(FS) cells. (a)** Sequence showing the HpaI

enzyme restriction site, which was used to distinguish between the asexual (donor) and

sexual (host) biotypes by RFLP analyses. **(b)** RFLP data showing rescue of lethally

irradiated sexual worms transplanted with freshly collected, non-cutured X1(FS) cells. **(c-**

**d)** RFLP data showing rescue of lethally irradiated sexual worms transplanted with 1 and

2 day cultured X1(FS) cells. Data from two independent experiments shown replicate 1

(panel c); replicate 2 (panel d). **(e)** Rescue rates for lethally irradiated hosts following

transplantation of X1(FS) cells cultured in the indicated media + 5% $CO_2$ for 1 or 2 days.

None of the conditions rescued lethally irradiated hosts after 3 days. Blue and orange

dots show value of rescue rate from replicate experiments, respectively.

**Supplementary Figure 6. No expression of exogenously delivered *Smed-***

***histone3.3-2✕flag* mRNA in *smedwi-1+* cells.** Representative images of electroporated

X1(FS) without (upper panel) or with (lower panel) *Smed-histone3.3-2✕flag* mRNA. Cells

cultured in Knockout DMEM + 5% $CO_2$ for 1 day were stained with *smedwi-1* riboprobe

and anti-FLAG antibody. Arrowhead: anti-FLAG+ nucleated cells. Scale bar, 20 μm.

**Supplementary Figure 7. Compare SiR-DNA sorted cells. (a)** A plot showing how SiR-

DNA-stained cells are displayed without gates in the flow cytometry analysis using SiR-

DNA versus side scater. **(b)** A plot showing how gates were defined to isolate two SiR-

DNA staining cell populations based on DNA content (SiR-DNA 4n and 2n). **(c)** *smedwi-*

*1* in situ staining for neoblasts in two isolated cell populations based on DNA content as

indicated in (b). SiR-DNA 4n population contains 56.4%±2.6% *smedwi-1*+ neoblasts (also

see Fig. 4f) compared to 26.8%±3.2% in SiR-DNA 2n population, p value = 0.0017. Scale

bar, 20 µm. **(d-g)** Plots showing the cell cycle distribution of SirNeoblasts (SiR-DNA 4n +

CT) (d), cells between SiR-DNA 4n and 2n (e), SiR-DNA 2n (f), and all SiR-DNA+ cells

(g). Sorted cells were stained with Hoechst 33342. Hoechst 33342+ (square gate) cells

were analyzed for cell cycle distribution.

**References**

1      Evans, M. J. & Kaufman, M. H. Establishment in culture of pluripotential cells from mouse

embryos. *Nature* **292**, 154-156 (1981).

2      Martin, G. R. Isolation of a pluripotent cell line from early mouse embryos cultured in

medium conditioned by teratocarcinoma stem cells. *Proc. Natl. Acad. Sci. U. S. A.* **78**,

7634-7638 (1981).

3      Boiani, M. & Scholer, H. R. Regulatory networks in embryo-derived pluripotent stem cells.

*Nat. Rev. Mol. Cell Biol.* **6**, 872-884 (2005).

4      Takahashi, K. & Yamanaka, S. Induction of pluripotent stem cells from mouse embryonic

and adult fibroblast cultures by defined factors. *Cell* **126**, 663-676 (2006).

5      Gehrke, A. R. & Srivastava, M. Neoblasts and the evolution of whole-body regeneration.

*Curr. Opin. Genet. Dev.* **40**, 131-137 (2016).

6      Baguñà, J., Saló, E. & Auladell, C. Regeneration and pattern formation in planarians. III.

Evidence that neoblasts are totipotent stem cells and the source of blastema cells.

*Development* **107**, 77- 86 (1989).

Neoblast culture and transformation

Reddien, P. W. & Sánchez Alvarado, A. Fundamentals of planarian regeneration. *Annu. Rev. Cell Dev. Biol.* **20**, 725-757 (2004).

Wagner, D. E., Wang, I. E. & Reddien, P. W. Clonogenic neoblasts are pluripotent adult stem cells that underlie planarian regeneration. *Science* **332**, 811-816 (2011).

Elliott, S. A. & Sánchez Alvarado, A. The history and enduring contributions of planarians to the study of animal regeneration. *Wiley Interdiscip. Rev. Dev. Biol.* **2**, 301-326 (2013).

Stornaiuolo, A., Bayascas, J. R., Saló, E. & Boncinelli, E. A homeobox gene of the orthodenticle family is involved in antero-posterior patterning of regenerating planarians. *Int. J. Dev. Biol.* **42**, 1153-1158 (1998).

Labbé, R. M. *et al.* A comparative transcriptomic analysis reveals conserved features of stem cell pluripotency in planarians and mammals. *Stem Cells* **30**, 1734-1745 (2012).

Wagner, D. E., Ho, J. J. & Reddien, P. W. Genetic regulators of a pluripotent adult stem cell system in planarians identified by RNAi and clonal analysis. *Cell Stem Cell* **10**, 299-311 (2012).

Lei, K. *et al.* Egf Signaling Directs Neoblast Repopulation by Regulating Asymmetric Cell Division in Planarians. *Dev. Cell* **38**, 413-429 (2016).

Wudarski, J. *et al.* Efficient transgenesis and annotated genome sequence of the regenerative flatworm model Macrostomum lignano. *Nat. Commun.* **8**, 2120 (2017).

Rubin, G. M. & Spradling, A. C. Genetic transformation of Drosophila with transposable element vectors. *Science* **218**, 348-353 (1982).

Fire, A. Integrative transformation of Caenorhabditis elegans. *EMBO J.* **5**, 2673-2680 (1986).

Bottger, A. *et al.* GFP expression in Hydra: lessons from the particle gun. *Dev. Genes Evol.* **212**, 302-305 (2002).

Neoblast culture and transformation

Renfer, E., Amon-Hassenzahl, A., Steinmetz, P. R. & Technau, U. A muscle-specific transgenic reporter line of the sea anemone, Nematostella vectensis. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 104-108 (2010).

Jaenisch, R. & Mintz, B. Simian virus 40 DNA sequences in DNA of healthy adult mice derived from preimplantation blastocysts injected with viral DNA. *Proc. Natl. Acad. Sci. U. S. A.* **71**, 1250-1254 (1974).

Newmark, P. A. & Sánchez Alvarado, A. Bromodeoxyuridine specifically labels the regenerative stem cells of planarians. *Dev. Biol.* **220**, 142-153 (2000).

Betchaku, T. Isolation of planarian neoblasts and their behavior in Vitro with some aspects of the mechanism of the formation of regeneration blastema. *J. Exp. Zool.* **164**, 407–433 (1967).

Schürmann, W., Betz, S. & Peter, R. Separation and subtyping of planarian neoblasts by density-gradient centrifugation and staining. *Hydrobiologia* 383, 117-124 (1998).

Schürmann, W. & Peter, R. Planarian cell culture: a comparative review of methods and an improved protocol for primary cultures of neoblasts. *Belg. J. Zool*, 123-130 (2001).

Asami, M. *et al.* Cultivation and characterization of planarian neuronal cells isolated by fluorescence activated cell sorting (FACS). *Zoolog Sci.* **19**, 1257-1265 (2002).

Reddien, P. W., Oviedo, N. J., Jennings, J. R., Jenkin, J. C. & Sánchez Alvarado, A. SMEDWI-2 is a PIWI-like protein that regulates planarian stem cells. *Science* **310**, 1327-1330 (2005).

Hayashi, T., Asami, M., Higuchi, S., Shibata, N. & Agata, K. Isolation of planarian X-ray-sensitive stem cells by fluorescence-activated cell sorting. *Dev. Growth Differ.* **48**, 371-380 (2006).

Arora, M. Cell Culture Media: A Review. *Mater Methods* **3**, 175 (2013).

Neoblast culture and transformation

Christensen, M. *et al.* A primary culture system for functional analysis of C. elegans neurons and muscle cells. *Neuron* **33**, 503-514 (2002).

Lukinavicius, G. *et al.* SiR-Hoechst is a far-red DNA stain for live-cell nanoscopy. *Nat Commun* **6**, 8497 (2015).

Zhang, D. *et al.* The piRNA targeting rules and the resistance to piRNA silencing in endogenous genes. *Science* **359**, 587-592 (2018).

Teshirogi, W. & Tohya, K. Primary tissue culture of freshwater planarian in a newly devised medium. *Fortschr. Zool.* **36**, 91-96 (1988).

Fukushima, T. & Matsuda, R. Experiments with culture media for planarian cells. *Hydrobiologia* **227**, 187-192 (1991).

Sasidharan, V. *et al.* The miR-124 family of microRNAs is crucial for regeneration of the brain and visual system in the planarian Schmidtea mediterranea. *Development* **144**, 3211-3223 (2017).

Iwasaki, Y. W., Siomi, M. C. & Siomi, H. PIWI-Interacting RNA: Its Biogenesis and Functions. *Annu Rev Biochem* **84**, 405-433 (2015).

Ruby, J. G. *et al.* Large-scale sequencing reveals 21U-RNAs and additional microRNAs and endogenous siRNAs in C. elegans. *Cell* **127**, 1193-1207 (2006).

Aravin, A. A., Sachidanandam, R., Girard, A., Fejes-Toth, K. & Hannon, G. J. Developmentally regulated piRNA clusters implicate MILI in transposon control. *Science* **316**, 744-747 (2007).

Palakodeti, D., Smielewska, M., Lu, Y. C., Yeo, G. W. & Graveley, B. R. The PIWI proteins SMEDWI-2 and SMEDWI-3 are required for stem cell function and piRNA expression in planarians. *RNA* **14**, 1174-1186 (2008).

Neoblast culture and transformation

Kelly, W. G. & Fire, A. Chromatin silencing and the maintenance of a functional germline in Caenorhabditis elegans. *Development* **125**, 2451-2456 (1998).

DeLuca, S. Z. & Spradling, A. C. Efficient expression of genes in the Drosophila germline using a UAS-promoter free of interference by Hsp70 piRNAs. *bioRxiv*, (2018).

Shirayama, M. *et al.* piRNAs initiate an epigenetic memory of nonself RNA in the C. elegans germline. *Cell* **150**, 65-77 (2012).

Friedlander, M. R. *et al.* High-resolution profiling and discovery of planarian small RNAs. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 11546-11551 (2009).

Miller, C. M. & Newmark, P. A. An insulin-like peptide regulates size and adult stem cells in planarians. *The International journal of developmental biology* **56**, 75-82 (2012).

Zhu, S. J. & Pearson, B. J. (Neo)blast from the past: new insights into planarian stem cell lineages. *Curr. Opin. Genet. Dev.* **40**, 74-80 (2016).

Zhu, S. J., Hallows, S. E., Currie, K. W., Xu, C. & Pearson, B. J. A mex3 homolog is required for differentiation during planarian stem cell lineage development. *Elife* **4**, (2015).

Dingwall, C. B. & King, R. S. Muscle-derived matrix metalloproteinase regulates stem cell proliferation in planarians. *Dev. Dyn.* **245**, 963-970 (2016).

Swim, H. E. & Parker, R. F. Method for establishing human cells in culture: susceptibility to poliomyelitis after prolonged cultivation. *Proc. Soc. Exp. Biol. Med.* **83**, 577-579 (1953).

Pearson, B. J. *et al.* Formaldehyde-based whole-mount in situ hybridization method for planarians. *Dev. Dyn.* **238**, 443-450 (2009).

King, R. S. & Newmark, P. A. In situ hybridization protocol for enhanced detection of gene expression in the planarian Schmidtea mediterranea. *BMC Dev. Biol.* **13**, 8 (2013).

Thi-Kim Vu, H. *et al.* Stem cells and fluid flow drive cyst formation in an invertebrate excretory organ. *Elife* **4**, (2015).

Neoblast culture and transformation

50    Edelstein, A. D. *et al.* Advanced methods of microscope control using muManager software. *J. Biol. Methods* **1**, (2014).

51    Nakamura, Y., Gojobori, T. & Ikemura, T. Codon usage tabulated from international DNA sequence databases: status for the year 2000. *Nucleic Acids Res.* **28**, 292 (2000).

52    Stothard, P. The sequence manipulation suite: JavaScript programs for analyzing and formatting protein and DNA sequences. *Biotechniques* **28**, 1102-1104 (2000).

Neoblast culture and transformation

# Lei et al. Figure 1

# Lei et al. Figure 2

# Lei et al. Figure 3

| # | Culture Medium | Normal Cell Morphology | Viability > 50% | %*smedwi-1* + high | Clonogenesis | Rescue (2-day culture) |
|---|---|---|---|---|---|---|
| 1 | IPM | YES | YES | YES | YES | YES |
| 2 | KnockOut DMEM | YES | YES | YES | YES | YES and Best |
| 3 | dKnockOut DMEM | YES | YES | YES | YES | NO |
| 4 | dL15 | YES | YES | YES | YES | YES |
| 5 | dSchneider's | YES | YES | YES | YES | NO |
| 6 | dDMEM | YES | YES | YES | YES | NO |
| 7 | dGrace's | YES | YES | YES | NO | Not tested |
| 8 | Holtfreter's | YES | YES | NO | Not tested | Not tested |
| 9 | Bge | YES | YES | NO | Not tested | Not tested |
| 10 | dRPMI1640 | YES | YES | NO | Not tested | Not tested |
| 11 | RPMI1640 | YES | YES | NO | Not tested | Not tested |
| 12 | DMEM | YES | YES | NO | Not tested | Not tested |
| 13 | L15 | YES | YES | NO | Not tested | Not tested |
| 14 | Essential 8 | YES | NO | NO | Not tested | Not tested |
| 15 | TPP | YES | YES | NO | Not tested | Not tested |
| 16 | M199 | YES | NO | NO | Not tested | Not tested |
| 17 | Schneider's | YES | YES | NO | Not tested | Not tested |
| 18 | CMFB | NO | NO | NO | Not tested | Not tested |
| 19 | dEssential 8 | YES | YES | NO | Not tested | Not tested |
| 20 | dM199 | YES | YES | NO | Not tested | Not tested |
| 21 | SFx | YES | YES | NO | Not tested | Not tested |
| 22 | dSFx | YES | YES | NO | Not tested | Not tested |
| 23 | Grace's | YES | YES | NO | Not tested | Not tested |

# Lei et al. Figure 4

# Lei et al. Figure 5

# Lei et al. Figure 6

# Lei et al. Figure 7

# Lei et al. Supplementary Figure 1

**6 days**

**L15_atmosphere**

# Lei et al. Supplementary Figure 2

**X1(FS) in IPM + 5% $CO_2$ for 2 days**

# Lei et al. Supplementary Figure 3

# Lei et al. Supplementary Figure 4

# Lei et al. Supplementary Figure 5

# Lei et al. Supplementary Figure 6

# Lei et al. Supplementary Figure 7

**a**

**b**

SiR-DNA 4n
SiR-DNA 2n

**c** *smedwi-1* | DAPI | *smedwi-1*/DAPI

SiR-DNA 4n

SiR-DNA 2n

**d** SirNeoblasts (SiR-DNA 4n + CT)

G1%=17.89
S%=13.02
G2/M%=69.09

**e** Between SiR-DNA 4n and 2n

G1%=53.31
S%=40.19
G2/M%=6.49

**f** SiR-DNA 2n

G1%=94.98
S%=5.02
G2/M%=0.00

**g** all SiR-DNA+

G1%=53.23
S%=20.93
G2/M%=15.84