Frequent sugar feeding behavior by *Aedes aegypti* in Bamako, Mali makes

them ideal candidates for control with attractive toxic sugar baits (ATSB)

Fatoumata Sissoko[1*], Amy Junnila[2], Mohamad M. Traore[1], Sekou F. Traore[1], Seydou Doumbia[1],

Seydou Mamadou Dembele[1], Yosef Schlein[2], Petrányi Gergely[6], Rui-De Xue[7], Kristopher L.

Arheart[3], Edita E. Revay[4], Vasiliy D. Kravchenko[5], John C. Beier[3], Gunter C. Müller[1,2]

[*]In memory of Fatoumata Sissoko.

[1] Malaria Research and Training Center, Faculty of Medicine, Pharmacy and Odonto-

Stomatology, University of Sciences, Techniques and Technology of Bamako, Bamako, Mali

[2] Department of Microbiology and Molecular Genetics, IMRIC, Kuvin Centre for the Study of

Infectious and Tropical Diseases, Faculty of Medicine, Hebrew University, Jerusalem, Israel

[3] Department of Public Health Sciences, Miller School of Medicine, University of Miami,

Miami, Florida, USA

[4]Department of Anatomy and Cell Biology, Bruce Rappaport Faculty of Medicine, Technion,

Haifa, Israel

[5] Department of Zoology, Tel Aviv University, Tel Aviv, Israel

[6] Zoologische Staatssammlung München, Germany

[7] Anastasia Mosquito Control District, 500 Old Beach Road, St. Augustine, FL, USA

Abstract

Background

Current tools and strategies are not sufficient to reliably address threats and outbreaks of

arboviruses including Zika, dengue, chikungunya, and yellow fever. Hence there is a growing

public health challenge to identify the best new control tools to use against the vector *Aedes*

aegypti. In this study, we investigated *Ae. aegypti* sugar feeding strategies in Bamako, Mali, to

determine if this species can be controlled effectively using attractive toxic sugar baits (ATSB).

Methodology/Principal findings

We determined the relative attraction of *Ae. aegypti* males and females to a variety of sugar

sources including flowers, fruits, seedpods, and honeydew in the laboratory and using plant-

baited traps in the field. Next, we observed the rhythm of blood feeding versus sugar feeding

activity of *Ae. aegypti* in vegetation and in open areas. Finally, we studied the effectiveness of

spraying vegetation with ATSB on *Ae. aegypti* in sugar rich (lush vegetation) and in sugar poor

(sparse vegetation) urban environments.

Male and female laboratory sugar feeding rates within 24 h, on 8 of 16 plants offered were over

80%. The survival rates of mosquitoes on several plant sources were nearly as long as that of

controls maintained on sucrose solution. In the field, females were highly attracted to 11 of 20

sugar sources, and 8 of these were attractive to males. Peak periods of host attraction for blood-

feeding and sugar feeding in open areas were nearly identical and occurred shortly after sunrise

and around sunset. In shaded areas, the first sugar-seeking peak occurred between 11:30 and

12:30 while the second was from 16:30 to 17:30. In a 50-day field trial, ATSB significantly

reduced mean numbers of landing / biting female *Ae. aegypti* in the two types of vegetation. At

sugar poor sites, the mean pre-treatment catch of 20.51 females on day 14 was reduced 70-fold to

0.29 on day 50. At sugar rich sites, the mean pre-treatment catch of 32.46 females on day 14 was

reduced 10-fold to a mean of 3.20 females on day 50.

Conclusions/Significance

This is the first study to show how the vector *Ae. aegypti* depends on environmental resources of

sugar for feeding and survival. The demonstration that *Ae. aegypti* populations rapidly collapsed

after ATSB treatment, in both sugar rich and sugar poor environments, is strong evidence that

Ae. aegypti is sugar-feeding frequently. Indeed, this study clearly demonstrates that *Ae. aegypti*

mosquitoes depend on natural sugar resources, and a promising new method for vector control,

ATSB, can be highly effective in the fight against *Aedes*-transmitted diseases.

Author summary

Aedes aegypti are notoriously difficult to control since their ubiquitous man-made and

natural breeding sites, in various geographical regions, include almost any receptacle that can

hold water. These diurnal mosquitoes are anthropophilic, a preference that promotes their role as

vectors of many arboviruses including Zika, dengue, chikungunya, and yellow fever. With the

exception of yellow fever, there are no vaccines against any of these arboviruses so that use of

personal protective measures and mosquito vector control are the only means of prevention.

Disease burdens in most endemic areas are not sufficiently reduced by various integrated vector

management (IVM) strategies, hence there is a need for new control tools to complement the

common strategies. Control by Attractive Toxic Sugar Baits (ATSB) appears to be an ideal

candidate for this purpose.

The results of this study support this proposition. They demonstrate that *Ae. aegypti* in

their urban environments in Mali are attracted to and frequently feed on staple diet that includes

a variety of flowers, fruits and seed pods. Therefore, *Ae. aegypti* is a suitable candidate for

control with ATSB. Moreover, the experiments with ATSB, in sparse vegetation or with

competitor plant attractants in rich vegetation, demonstrated that ATSB treatment can cause a

drastic reduction of *Ae. aegypti* populations.

Introduction

Aedes aegypti are vectors for several significant arboviruses including Zika, dengue,

chikungunya, and yellow fever [1-4]. Apart from yellow fever, there are no approved vaccines in

widespread use against these viruses, therefore public health efforts are reliant on effective

vector control methods [5,6]. Common strategies for the control of *Ae. aegypti* include residual

spraying targeted to resting sites [7], space spraying indoors [8], larval control with conventional

pesticides or lethal ovi-traps [9,10], and the use of personal protective measures.

However, the use of these methods is insufficient for achieving significant reductions in

disease burdens thus new control tools are required [11,12]. Methods that are considered

promising for this purpose by the WHO and require further field testing include: the

Incompatible Insect Technique, the sterile insect technique, vector traps, and attractive toxic

sugar baits [13, 14].

For many years, based on published work and experience in rearing laboratory

mosquitoes, it was common knowledge that mosquitoes must feed on sugar to obtain the energy

necessary for survival [15, 16]. However, the observation that *Ae. aegypti* could take blood and

convert it to triglycerides and glycogen that provide energy [17] led to subsequent observations

that female *Ae. aegypti* in the field were taking multiple blood meals within a single gonotrophic

cycle [18-20]. Accordingly, *Ae. aegypti* may be less dependent on sugar for energy than other

mosquito species and presumably rarely feed on sugar [18, 21, 22]. This is not necessarily a

comprehensive rule since there are observations that *Ae. aegypti* are frequent sugar feeders,

given the right environment. It was demonstrated that in Chiapas, Mexico, in an area with

flowering plants, *Ae. aegypti* commonly fed on nectars; 8 to 21% of the mosquitoes tested

positive for sugar using the anthrone test [23]. In the laboratory [24], it was observed that male

and female *Ae. aegypti* displayed two peaks in sugar feeding; a small morning peak with 16 to

18% of mosquitoes sugar-feeding and a larger evening peak with 40 to 42% sugar-feeding

mosquitoes. In another study in Duran, Ecuador, 56.8% of the *Ae. aegypti* females were marked

as sugar positive following feeding on sucrose solution colored with food dye [25]. It has been

concluded that the availability of sugar sources in the local environment affects mosquito

longevity and thus it is a key regulator of mosquito population dynamics and therefore, of their

vectorial capacity [15, 26]. The evidence above indicates that that the success of *Ae. aegypti*, at

least in some regions, also depends on the presence of natural sugar sources. Attractive toxic

sugar bait methods (ATSB) use the staple sugar feeding behavior of mosquitoes in nature for

their control. The potential and efficiency of this method have been demonstrated in several

experiments with various mosquito genera and species [27-34]. The observations on the sugar

feeding behavior of *Ae. aegypti* indicate the potential of also using ATSB methods for the control

of this species. The initial requirement for assessing the possibility of using ATSB against *Ae.*

aegypti is the study of the sugar feeding behavior of the species on natural sugar sources in the

experimental region. This was the basic purpose of this study which was investigated in the

following manner: 1) in the laboratory, we observed whether *Ae. aegypti* groups were feeding on

potentially edible sources of sugar, including flowers, fruits, seedpods, and honeydew, both

available in the mosquito's environment and some known to be highly attractive to other species;

2) We repeatedly offered single plant-sugar sources to mosquitoes as exclusive diets, one species

per series of mosquitoes, and monitored the mortality rate in the mosquitoes; 3) We used plant

baited traps in the field in and around the urban centers of Bamako, Mali and estimated plant

attraction by comparing their catches of mosquitoes; 4) We determined the proportion of sugar

fed mosquitoes and estimated sugar quantities in *Ae. aegypti* males and females from sugar rich

and sugar poor urban neighborhoods; and 5) Finally, we tested the effect of ATSB treatment,

estimated by the number of mosquitoes landing on volunteers, on *Ae. aegypti* population size in

sugar rich and sugar poor areas in Bamako.

Methods

Identification of plants that are potential sugar sources for *Ae. aegypti*

Dominant plants at the study sites were defined with the assistance of the staff of the

department of Traditional Medicine, School of Medicine and Dentistry, University of Sciences

and Technology of Bamako, Mali.

Mosquitoes for laboratory experiments

The *Ae. aegypti* colony used in the laboratory trials were established from the F1

generation of field caught mosquitoes to simulate the field breed as closely as possible. They

were caught close to the campus of the campus of the University of Sciences, Techniques and

Technology of Bamako, Mali, which is inside the city, and kept at their insectary under the

following conditions: 27 ±3ºC, relative humidity 70 ±10%, and photoperiod 12:12 hours

light:dark. Adults for experiments were maintained on 10% sucrose solution and then starved for

24 h before the beginning of experiments. Fresh batches of 5 day old mosquitoes were used for

each experiment.

Testing for sugar

The sugar content in the gut of mosquitoes was determined by a modified cold anthrone

test for fructose [35] and sugar amounts were visually scored by the intensity of the blue colour

reaction under a dissecting microscope. Very light blue colour was defined as class I; darker blue

colour in the reaction fluid was class II; class III was darker, and very intense blue was class IV

(Fig 1).

Fig 1. Microtitre plate with anthrone tested samples. Class I through IV colour intensities as

well as negative control wells are shown.

Laboratory experiments

Sugar feeding of *Ae. aegypti* exposed to potential sugar sources for 24 h. Assays were carried

out with cohorts of 30 female and 30 male *Ae. aegypti* placed together for 24 h in 50×50 cm^2

mosquito cages that contained one of the following: fruits of *Carica papaya* (papaya), *Mangifera*

indica (mango), *Cucumis melo* (honeydew melon), leaves with extra-floral nectar of *Ricinus*

communis, branches of *Lantana camara* soiled with honeydew from the aphid *Aphis gossypii*,

and flowering *Prosopis juliflora*, *Acacia macrostachya*, *Acacia salicina*, *Lantana camara,*

Galphimia gracilis and *Bougainvillia glabra.* Other types of offered sugar source were

undamaged seedpods of *Piliostigma reticulatum*, infested *P. reticulatum* seedpods perforated by

an unknown species of *Microlepidoptera* that were oozing sweet liquid, cut 20 cm segments of

Saccharum officinarum (sugar cane) with the ends tightly sealed with parafilm, and broken /

crushed tissue of similar segments of sugar cane. The general pattern was to offer either 4-5

branches of a target plant with stems in a beaker of distilled water, cut pieces of fruit (100 g), or

complete seed pods (100 g). Additional tests were conducted on *Lantana camara* branches

coated with ATSB solution (active ingredient: microencapsulated garlic oil) and ATSB offered

in bait stations (active ingredient: dinotefuran). The bait stations were experimental prototypes

with a thin protective membrane cover and were supplied from an ongoing Innovative Vector

Control Consortium (IVCC) project carried out by Westham Ltd., Tel Aviv Israel.

A 10% sucrose solution and water soaked-cotton swabs served as the control diet and

were made available in all cages. Each experiment was repeated 6 consecutive times. Mosquitoes

were killed with CO_2 and then either tested immediately for sugar by modified anthrone test or

frozen at -70ºC until anthrone tests could be performed.

Survival of *Ae. aegypti* females with continuous access to different sugar sources. Groups of

100 *Ae. aegypti* females in 50 x 50 cm cages, were allowed to feed on only 1 type of plant and

their survival was monitored for 31 days. Plants were replaced daily, and water-soaked cotton

swabs were generally available. Dead mosquitoes were removed daily. The 11 test diets were

extra-floral nectar on foliage of non-flowering *Ricinus communis*, flowers of *Bougainvillia*

glabra, Prosopis juliflora, or *Galphimia gracilis,* fruits of *Mangifera indica* and *Cucumis melo,*

seed pods of *Piliostigma reticulatum*, either undamaged or perforated by pests and oozing juice,

intact or broken / crushed stems of *Saccharum officinarum*, and honeydew (*Aphis gossypii*)

soiled branches of non-flowering *Lantana camara*. Controls received 10% sucrose solution and

water or water only. Each experiment was repeated 6 consecutive times.

Field experiments

Field sites

All studies took place in Bamako, the capital and largest city of Mali, with ~ 2 million

inhabitants. The city spreads out on both banks of the River Niger and annual flooding limits

building on the banks which are a patchwork of wetlands, parkland, agricultural fields and

forested areas. Some of the richer neighbourhoods and hotel districts include buildings with

parklands or small vegetable fields. Bamako has tropical annual wet and dry periods. The hottest

months are March, April, and May (hottest average temperature $32.4\ ^0C$). The average

temperature in the coldest month (December) is $25.1°C$. Total annual rainfall averages 1098.5

mm; the rainy period is May through September, with peak rain occurring in August/September.

The driest periods are late October through April.

'Sugar rich' and 'sugar poor' sites for field studies were chosen around Bamako and were

identified primarily by land use which was determined by scouting areas by foot with trained

botanists who could estimate flowering vegetation cover. Urban areas with the lush vegetation of

irrigated parkland and gardens containing $\geq 50\%$ flowering vegetation, were defined as 'sugar

rich' and densely populated urban neighborhoods with sparse vegetation containing $< 5\%$

flowering vegetation, were defined as 'sugar poor'.

Attraction of *Ae. aegypti* to plant baited glue net traps (GNT's) in the field. Experiments

were carried out for 10 consecutive days at the end of the dry season in 2016 along the shady

margins of a forest gallery parallel to the River Niger.

Specifically designed glue net traps (GNTs) [30], were used to test attraction to plants.

Briefly, cut bottom halves of 1.5 L plastic bottles were set in the ground, with the margins

protruding to the surface, and filled with water. Dark green, rigid plastic netting, with mesh size

0.8 cm x 0.2 cm, was cut into 70×70 cm squares, rolled into cylinders, and each was put

vertically above one of the cut bottles with the water, fastened to the ground with pegs and fitted

with mesh covers. About 0.5 kg of test plant material was fixed in the center of a cylinder that

was then covered and coated externally with glue (Tangle Foot, Tel Aviv, Israel). The different

plant baits are listed in Table 2A and 2B. Controls were water-soaked sponges, empty traps, or

naked *L. camara* branches, sprayed with either 10% sucrose solution or ATSB solution used in

this study. Mosquitoes caught in the glue were removed, counted, and stored in 70% ethanol for

identification. Each morning of the 10 day monitoring period, the baits were replaced, cylinders

were repainted with glue, and trap locations were rotated to avoid location bias.

Timing of host-seeking and sugar-seeking activities in the field. The rhythm of the search for

blood meals was shown by the number of landing/biting events on human volunteers in the field,

in 30 min. intervals, for 18 hours (05:00 h to 23:00 h). This was done at 6 separate sites: 3 shady,

3 open and mosquitoes on volunteers from each group of sites were pooled and averaged.

Mosquitoes were collected with aspirators near the shady margins of the forest gallery with thick

undergrowth along the River Niger, and in open, sun-exposed grassland 30 m away from the

trees. The United States Environmental Protection Agency guidelines and protocols for the use

of human volunteers in landing catch experiments were carefully followed [36]. Three

volunteers, 2 males and 1 female, all professional entomologists/medics participated in this

study. As part of the consent process, the participants in all human trials were fully advised of

the nature and objectives of the test and the potential health risks from exposure to mosquito

bites. According to EPA regulations, they were required to avoid alcohol, caffeine, and fragrance

products (e.g., perfume, cologne, hairspray, lotion, etc.) during the entire test period. For the

tests, volunteers were wearing long trousers and long-sleeved shirts as protection against

mosquito bites. One leg of the trousers was rolled up to expose the skin used as the test area.

Volunteers were seated motionless in chairs with the exposed leg extended while observation,

counting and recording of mosquitoes was made by assistants. The distance between the

volunteers was 20 m and their locations were rotated through 9 stations every 30 min. to

eliminate positional bias.

The rhythm of activity in the search for sugar meals was observed in both shaded and

open areas by counting the catches of 9 GNTs in each type of area (18 total) baited with highly

attractive flowering *P. juliflora* branches in 30 min. intervals, for 18 hours (05:00 h to 23:00 h).

The baits were replaced by fresh branches at each time interval.

Comparing sugar feeding rates of *Ae. aegypti* in 'sugar rich' and 'sugar poor' habitats. The

sugar-fed status of blood searching mosquitoes was observed. Catches on volunteers were

carried out in the morning (11:00 h – 12:00 h) at 'sugar poor' and 'sugar rich' sites (Fig 2). Three

volunteers, 2 males and 1 female, all professional entomologists/medics participated in this study

and set-up was as described under "Timing of host seeking activity in the field".

Fig 2. Pictures of sugar rich and sugar poor habitats in Bamako neighbourhoods. A and B)

Densely populated urban areas with sparse vegetation (<5% flowering), **C and D**) Lush

vegetation of irrigated gardens and parkland with areas of ≥50% flowering vegetation.

Sweep- net and aspirator (battery powered vacuum aspirators, John W. Hock, Gainesville, FL)

catches for periods of 30 min. were also carried out in nearby vegetation. Captured mosquitoes

were put on ice or in cages immediately, and random subsamples were tested for sugar [35].

Mosquito samples taken were either processed within 1 hour of collection or 6 hours after

collection to assess the rate of sugar digestion from the guts of the mosquitoes. This process is

expressed by the loss of positive anthrone reactions [35] with time. Random samples of 100

female and 100 male mosquitoes were tested from each site and for each processing time point.

Sampling continued on consecutive days until these numbers were obtained, 4 days for the sugar

rich area and 7 for the sugar poor area.

ATSB experiments. The ATSB method was field tested in 50 day experiments (Fig 2) in

parklands and irrigated gardens ('sugar rich') and also in dry residential areas ('sugar poor') of

Bamako from mid-June to the end of July 2016 (Fig 3).

Fig 3. Local weather conditions in Bamako during June and July 2016. Rain occurred during

the study, and two treatments were applied at the sites to avoid a "wash off" effect.

A total of 12 sites, 6 'sugar rich' and 6 'sugar poor' areas of ~ 1 ha^2 were selected. Three of each

type were untreated controls and 3 in their vicinity received ATSB spray treatment. The effect of

ATSB treatment was evaluated by recording biting / landing rates of female *Ae. aegypti* on

human volunteers. The procedure was as described under "timing of host-seeking activity" with

the same consent process [36]. The volunteers were placed in the center of each plot and each of

them was moved among 3 monitoring stations per site that amounted to 9 repetitions per site.

ATSB was sprayed mainly on broad-leafed shrubs, bushes, and small trees up to 1.5 m high.

Plants with flowers or fruit were not treated to minimize the impact on non-target organisms

[37]. At the sugar poor sites, for the lack of suitable vegetation, artificial structures and buildings

were sprayed. Treated areas were sprayed twice, once on day 15 and again on day 32 of the

experiments.

ATSB formulations. For the ATSB spray trials, we used ATSB Mosquito Bait Concentrate,

with the active ingredient of microencapsulated garlic oil (Universal Pest Solutions, Dallas TX,

USA). The material was used according to label instructions, the concentrate was diluted 1:3,

and was applied with a backpack sprayer on vegetation or on suitable artificial structures to

cover 5% of the targeted area (four plots of 1 ha $500m^2$ were treated). Perimeter treatments were

used at sites with continuous lines of vegetation (e.g. hedges, other shrubs) or extensive

landscaping. ATSB was applied to vegetation in a continuous band measuring about 0.5 m wide,

between 0.3 m to 1.5 m above the ground, and to the point of runoff at a rate of 500 to 600 ml of

mixture per 30 m. Spot-treatments were applied to single shrubs, small patches of vegetation or

artificial structures at least 30 cm above the ground in patches of approximately 1 m^2 to the point

of runoff. The spot applications were repeated, if possible, every 4 m.

Statistical analysis

For the statistics reported in Table 1, we used a generalized linear model for a Poisson

distributed outcome, number of female and male mosquitos. Separate models were employed for

females and males. The independent variable was plant. We present model means and standard

deviations along with raw and Dunnett adjusted p-values for comparing means with the control,

ATSB on *L. camara* branches.

Figures 4 A and B present survival plots for a series of female *Ae. aegypti* that were each

exposed for 31 days to two types of plants. We used Kaplan-Meyer statistics to compute the data

for these figures.

Tables 2 A and B report the mean number of *Ae. aegypti* caught overnight by GNTs each with a

different bait of fruit, seedpods, or flowers. We used a generalized linear model for a Poisson

distributed outcome, number of female and male mosquitos. Overdispersion was evident;

therefore, we changed to a negative binomial model that employs a scale parameter to adjust the

model for overdispersion. Separate models were analyzed for females and males. The

independent variable was plant. We present model means and standard deviations along with raw

and Dunnett adjusted p-values for comparing means with the control, an empty trap. The data for

Figures 4 A and B come from the data used in the analysis of Tables 2 A and B.

The percentage of *Ae. aegypti* with different amounts of sugar in the gut is reported in Table 3.

The mosquitos were caught at 'sugar rich' and 'sugar poor' sites on human volunteers or with

sweep-nets in the vegetation. We totaled the number of sugar positive mosquitos in each

category as determined by anthrone testing for each time, sugar status, and catch type and then

divided by the total number of sugar positive mosquitos. The data for Figures 5 A and B come

from the data analyzed for Table 3.

Table 4 presents the mean number of *Ae. aegypti* following ATSB treatment at sugar poor and

sugar rich sites. The data is the number of female mosquitos landing on human volunteers

compared to their frequency at untreated control sites. We applied a generalized linear mixed

model analysis to the mean catch for control and treatment period in each sugar rich and sugar

poor control and experimental groups. Each mean came from nine catches: three volunteers each

providing landing catches in three different monitoring stations. The Shapiro-Wilk test

determined that the mean counts were normally distributed. The model included fixed effects for

group (sugar rich and sugar poor), treatment period (control [days 1-14], ATSB treatment 1 [days

16-31], and ATSB treatment 2 [days 35-50]), condition (control and experimental), plus all two-

way interactions and the three-way interaction. Appropriate error terms for the fixed effects are

random effects. The error term for the group effect was site nested within group. The treatment

and condition effects were repeated measures within the groups. Therefore, the error term for

treatment and group x treatment was group x treatment x site nested within group. Likewise, the

error term for condition and condition x group was condition x group x site nested within group.

The error term for treatment x condition and group x treatment x condition was treatment x

condition x site nested within group. Means and standard errors are reported for the group x

treatment x condition interaction. We report *p*-values and Bonferroni adjusted *p*-values for

comparisons between control and experimental means at each treatment level for sugar poor and

sugar rich groups as well as comparisons between sugar rich and sugar poor means at each

treatment level for control and experimental groups. Finally, we report *p*-values and Dunnett

adjusted *p*-values for comparisons between control and the two ATSB levels for sugar rich and

poor experimental and control groups. The data for this analysis was used to produce Figures 7 A

and B.

Plants in field experiments were also ranked by being assigned an attraction index (AI),

which was calculated by the following equation: average catch with the plant bait (PB) ÷ by the

average catch with the empty trap control (ET); PB/ET = AI.

Results

Sugar feeding of *Ae. aegypti* exposed to potential sugar sources for 24 h

Sugar feeding rates for females on 8 out of 16 of the sugar sources were high, ranging

from 81.67% on *Mangifera indica* (mango), to 93.89% on *Cucumis melo* (honeydew melon).

Males fed avidly on 8 sugar sources. Exposure to fluids oozing from decomposing seedpods of

P. reticulatum resulted in 80% sugar positive mosquitoes and the highest result was the 97.22%

of positive specimens in the series that received *C. papaya* (papaya). Also, 88.89% of the

females and 95.00% of the males in the ATSB on *L. camara* group were sugar positive after the

exposure (Table 1).

Table 1. **Mean numbers ±SE over replicate cages in six experiments, in descending order,**

of sugar positive mosquitoes after 24 h exposure to a single plant-sugar source.

Females					Males				
Species	Mean	±SE	% Positive	*p	Species	Mean	±SE	% Positive	*p
C. melo F	28.17	2.17	93.89%	0.621	*C. papaya* F	29.17	2.21	97.22%	0.830
A. macrostachya FL	28.00	0.91	93.00%	0.601	[§]ATSB on *L. camara* G	28.50	2.18	95.00%	N/A
P. juliflora FL	27.83	0.70	92.78%	0.700	*P. juliflora* FL	28.00	2.16	93.33%	0.871
[§]ATSB on *L. camara* G	26.67	1.33	88.89%	N/A	*A. macrostachya* FL	27.83	2.15	92.78%	0.828
A. salicina FL	26.50	2.10	88.33%	0.956	*G. gracilis* FL	26.67	2.11	88.89%	0.547
G. gracilis FL	25.50	2.06	85.00%	0.693	*A. salicina* FL	26.50	2.10	88.33%	0.511
C. papaya F	25.33	2.06	84.44%	0.652	*M. indica* F	25.33	2.06	84.44%	0.293
P. reticulatum (oozing juice) F	24.67	2.03	82.22%	0.496	*C. melo* F	25.17	2.05	83.89%	0.268
M. indica F	24.50	2.02	81.67%	0.460	[§]Sucrose 10% on cotton	24.16	2.01	85.56%	0.147

§Sucrose 10% on cotton	24.17	2.01	80.56%	0.393	*P. reticulatum* (oozing juice) F	24.00	2.00	80.00%	0.132
§ATSB Bait Station	18.83	1.77	62.78%	0.006	*L. camara* FL	18.65	1.76	62.22%	0.010
L. camara FL	18.81	1.75	61.11%	0.003	*L. camara* with *A. gossypii* G, H	13.96	1.53	46.67%	<0.001
L. camara with *A. gossypii* G, H	14.50	1.56	48.33%	<0.001	*R. communis* G, E	13.31	1.49	45.00%	<0.001
S. officinarum (broken) G	13.17	1.48	43.89%	<0.001	§ATSB Bait Station	12.97	1.47	43.33%	<0.001
R. communis G, E	4.67	0.99	15.88%	<0.001	*S. officinarum* (broken) G	11.29	1.37	37.78%	<0.001
B. glabra FL	4.62	0.88	15.56%	<0.001	*B. glabra* FL	3.23	0.73	11.11%	<0.001
P. reticulatum (intact) F	0.67	0.33	2.22%	<0.001	*P. reticulatum* (intact) F	0.58	0.31	2.22%	<0.001
N. glauca FL	0.50	0.29	1.67%	<0.001	*N. glauca* FL	0.32	0.23	1.11%	<0.001
S. officinarum (intact) G	0.33	0.24	1.11%	<0.001	*S. officinarum* (intact) G	0.25	0.21	0.56%	<0.001

*Compared to positive control ATSB on *L. camara* branches

§Positive control

N/A Not applicable

F: Fruit, seedpod

FL: Blossom

G: Branches

H: Honeydew on branches (*Aphis gossypii*)

E: Extra-floral nectar

Survival of *Ae. aegypti* females with continuous access to different sugar sources

Of the different female groups fed exclusively for 31 days on one diet, the negative control group

of 100 starved and thirsty females died within 4 days. Mosquitoes survived for up to 6 days on

water alone and the provision of 10% sucrose solution allowed 68% survival of up to 31 days.

Among the plant diet series, the survival proportion by day 31 was the highest (85.89%) in the

group fed on *P. juliflora* whereas the lowest survival rate of 5.00% was in the group that

received intact seedpods of *P. reticulatum* (Fig 4A and 4B).

**Fig 4. Survival in a series of female *Ae. aegypti* that were each exposed for 31 days to one
type of plant.** (**A**) Survival after exposure to branches with plant blossoms. *Ricinus communis*
baits were branches with extra-floral nectaries. (**B**) Sugar source types other than blossoms.

Attraction of *Ae. aegypti* to plant baited glue net traps (GNTs) in the field

Females were attracted to 11 of the 23 baits. The highest mean catches of 18.70, 13.70, 11.20, and 9.80 specimens were from traps baited with *P. juliflora, A. macrostachya, A. salicina, M. indica,* and ATSB on branches, respectively. Males were attracted to 8 baits, the 4 most attractive being *P. juliflora* (12.90), *A. macrostachya* (10.20), *A. salicina* (7.60), and the positive control of ATSB coated branches (6.50) (Tables 2A and 2B).

Table 2. Mean number (± SE) of *Ae. aegypti* caught overnight by GNTs each with a different bait of fruit, seedpods, or flowers. (A) Females. (**B**) Males.

Table 2A.

Origin	Condition	Species	Mean	±SE	Attraction Index	P Value	Sig.
Ornamental	Flowering	*P. juliflora*	18.70	8.18	37.40	<0.001	*
Native	Flowering	*A. macrostachya*	13.70	6.02	27.40	<0.001	*
Ornamental	Flowering	*A. salicina*	11.20	5.57	22.40	<0.001	*
Positive control	Non-flowering	ATSB on *L. camara* branches	9.80	4.80	19.60	<0.001	*
Agricultural	Flowering	*M. indica* (Mango)	5.60	2.99	11.20	<0.001	*
Agricultural	Fruit	*C. papaya* (Papaya)	4.90	3.00	9.80	<0.001	*
Agricultural	Fruit	*M. indica*	4.30	2.67	8.60	<0.001	*
Agricultural	Fruit	*C. melo*	3.70	2.41	7.40	0.001	*
Native	Flowering	*R. communis*	3.00	2.05	6.00	0.002	*
Aphids	Non-flowering	*L. camara* with *A. gossypii*	2.60	1.84	5.22	0.011	*
Ornamental	Flowering	*G. gracilis*	2.20	1.69	4.40	0.031	*
Ornamental	Non-flowering	*P. juliflora*	1.61	0.99	3.22	0.154	NS
Negative control	Cotton Towel	Water	1.60	0.97	3.20	0.154	NS
Negative control	Cotton Towel	Sucrose 10%	1.50	1.08	3.00	0.200	NS
Ornamental	Flowering	*B. glabra*	1.40	0.84	2.80	0.258	NS
Agricultural	Non-flowering	*M. indica*	1.30	1.34	2.60	0.332	NS
Ornamental	Flowering	*L. camara*	1.10	0.74	2.20	0.536	NS
Native	Non-flowering	*A. macrostachya*	1.00	1.05	2.00	0.651	NS
Positive control	Non-flowering	Sucrose 30% on *L. camara*	1.00	0.94	2.00	0.669	NS

Origin	Condition	Species	Mean	±SE	Attraction Index	P Value	Sig.
Native	Non-flowering	R. communis	0.90	0.99	1.80	0.332	NS
Agricultural	Crushed	S. officinarum	0.80	0.63	1.60	1.000	NS
Ornamental	Non-flowering	A. salicina	0.70	0.67	1.40	0.812	NS
Aphids	Non-flowering	Aphid infested L. camara	0.60	0.70	1.20	0.622	NS
Agricultural	Intact	S. officinarum	0.50	0.71	1.00	0.442	NS
Negative control	Empty	Empty trap	0.50	0.53	N/A	N/A	N/A

* Significant compared to negative control empty trap

NS - Not significant compared to negative controls

NA - Not applicable

382 **Table 2B.**

Origin	Condition	Species	Mean	±SE	Attraction Index	P Value	Sig.
Ornamental	Flowering	P. juliflora	12.90	5.51	32.25	<0.001	*
Native	Flowering	A. macrostachya	10.20	5.37	25.50	<0.001	*
Ornamental	Flowering	A. salicina	7.60	4.99	19.00	<0.001	*
Positive control	Non-flowering	Sucrose 30% on L. camara	6.60	0.70	16.50	1.000	NS
Positive control	Non-flowering	ATSB on L. camara	6.50	4.45	16.25	<0.001	*
Agricultural	Flowering	M. indica	3.20	1.93	8.00	0.001	*
Agricultural	Fruit	C. melo	2.90	2.47	7.25	0.002	*
Agricultural	Fruit	M. indica	2.80	1.32	7.00	0.002	*
Agricultural	Fruit	C. papaya	2.70	1.77	6.75	0.003	*
Native	Flowering	R. communis	1.90	1.29	4.75	0.459	NS
Aphids	Non-flowering	L. camara with A. gossypii	1.80	1.03	4.50	0.036	*
Ornamental	Non-flowering	P. juliflora	1.30	1.06	3.25	0.154	NS
Ornamental	Flowering	G. gracilis	1.20	0.92	3.00	0.002	*
Agricultural	Non-flowering	M. indica	1.10	0.99	2.75	0.274	NS
Native	Non-flowering	A. macrostachya	1.00	0.94	2.50	0.342	NS
Ornamental	Flowering	B. glabra	0.90	0.74	2.25	0.478	NS
Native	Non-flowering	R. communis	0.60	0.70	1.50	0.459	NS
Aphids	Non-flowering	L. camara with A. gossypii	0.60	0.70	1.50	1.000	NS
Agricultural	Intact	S. officinarum	0.60	0.84	1.50	1.000	NS
Ornamental	Flowering	L. camara	0.60	0.70	1.50	1.000	NS
Agricultural	Crushed	S. officinarum	0.50	0.53	1.25	0.777	NS
Negative control	Cotton Towel	Water	0.50	0.53	1.25	0.777	NS
Ornamental	Non-flowering	A. salicina	0.40	0.52	1.00	0.553	NS

| Negative control | Cotton Towel | Sucrose 10% | 0.40 | 0.52 | 1.00 | 0.553 | NS |
| Negative control | Empty | Empty trap | 0.40 | 0.52 | N/A | N/A | N/A |

* Significant compared to negative control empty trap

NS - Not significant compared to negative controls

NA - Not applicable

Timing of host-seeking and sugar-seeking activities in the field

In shady areas, the three volunteers caught an average 54.5 females and 4.6 males over 18

hours of monitoring. The same volunteers in an open area caught an average of 22.0 females and

0.3 males over 18 hours. In open areas, attraction of females to the volunteers showed a first

peak shortly after sunrise between 07:30 and 08:00 h and a second, larger peak was observed

around sunset between 19:00 and 21:30 h. At shady sites, the first peak was delayed to between

09:30 to 10:30 h and the second appeared earlier, between 18:00 to 19:30 h. The morning peak

of female activity in shaded areas, averaging 15 landings per volunteer, was greater than the

average of 9 landings per volunteer in open areas. In the evening, the average number of female

landings per volunteer was 18 in open areas, while in shade it was nearly 27. Interestingly, in

shaded areas there were also small numbers of male landings, 5 landings per volunteer, roughly

at the same time as the peak of female activity. In open areas, landing of males was negligible

(Fig 5A).

Fig 5. Periodicity of host-seeking and sugar-seeking behavior of *Ae. aegypti* over 18 hours

(**A**) evaluated by average catches of mosquitoes landing on a human volunteer, in 30 min

intervals (±SE) and (**B**) evaluated by catches of GNTs baited with highly attractive flowering

branches of *P. juliflora*. Numbers shown are the average catch per volunteer, per time period and

average catch per trap, per time period (±SE).

The attraction rate to a sugar source (traps baited with flowers of *P. juliflora*) in sunny

and shaded areas also exhibited two peaks as did the search for a host (Fig 5B). In open areas,

the first smaller peak of female landings was just after sunrise and the second followed sunset. In

shaded areas, the first peak occurred between 11:30 h and 12:30 h while the second took place

from around 16:30 h to 17:30 h. Sugar questing activity of males in the shady areas followed the

pattern and duration of the females but the numbers caught were about a half. Only 2 males were

caught in open sunny areas throughout the entire study period (Fig 5B).

Comparing sugar feeding rates of *Ae. aegypti* in 'sugar rich' and 'sugar poor' habitats

Less than 20% of females and nearly 80% of males landing on human volunteers were

sugar positive in both sugar rich and sugar poor environments. From resting sites within

vegetation, 60-65% of 'sugar rich' site females or males were sugar positive and so were 39-40%

of the females or males sampled from the 'sugar poor' site. Leaving mosquitoes alive in cages

for six hours before testing decreased the proportion of sugar positive specimens by about a half

(Fig 6A and 6B).

Fig 6. Percentage of sugar positive mosquitoes from the catch on volunteers and from

resting habitat sweep-net catches at sugar rich (SR) and sugar poor (SP) sites. (**A**) Tested

for sugar within 1 hour after collection and (**B**) 6 hours after collection.

Table 3. The percentage of *Ae. aegypti* with different amounts of sugar in the gut that were

caught at 'sugar rich' and 'sugar poor' sites on human volunteers or with sweep-nets in the

vegetation. Sugar quantities were classed by the intensity of the anthrone reaction.

Site	Time*	Catch type	Females				Males			
			I	II	III	IV	I	II	III	IV
Sugar rich	1 hr	Human landing catch	60.32	32.54	6.35	0.79	23.00	40.00	29.00	8.00
Sugar poor	1 hr	Human landing catch	68.37	24.49	7.14	0.00	28.57	48.57	18.57	4.29
Sugar rich	1 hr	Resting site catch	13.18	21.47	40.49	24.86	11.60	20.63	42.54	25.23
Sugar poor	1 hr	Resting site catch	46.31	28.19	19.80	5.71	53.94	37.54	7.26	1.26
Sugar rich	6 hr	Human landing catch	90.39	9.62	0.00	0.00	81.81	15.91	2.27	0.00
Sugar poor	6 hr	Human landing catch	92.31	7.69	0.00	0.00	56.94	30.56	12.50	0.00
Sugar rich	6 hr	Resting site catch	47.15	36.94	12.61	3.30	56.69	29.94	11.78	1.59
Sugar poor	6 hr	Resting site catch	80.00	16.92	3.08	0.00	75.00	23.53	1.47	0.00

*Time between collection and testing

Classification of *Ae. aegypti* females or males with different quantities of sugar in the gut

showed that in 'sugar rich' environments, 60 to 90% of the sugar positive females caught on

volunteers had mostly small (class I) sugar quantities in the gut (Table 3). Samples from 'sugar

rich' resting sites, contained various sugar quantities in similar proportions (classes I to IV). A 6

hour delay in testing for sugar caused the sugar contents in the gut to degrade, and thus the

reactions in the sugar testing to fall below classes III and IV. The sugar feeding status of

mosquitoes caught at the 'sugar poor' sites, on volunteers or in resting habitats, followed the

pattern observed for mosquitoes from the 'sugar rich' sites.

ATSB experiments

The baseline for evaluating the impact of ATSB treatment was the density of female

mosquitoes landing on volunteers at the control site, and the pre-treatment period at the

experimental site. The mean daily catches on the 3 volunteers increased gradually at both 'sugar

rich' and 'sugar poor' control sites. At the 'sugar poor' control site, the mean began at 5.46

±0.43 females and was 32.93±2.89 on the last experimental day. At the 'sugar rich' control site,

the initial mean increased from 12.27 ±1.20 to 65.89 ±5.55 females on the last day. ATSB

treatment at both experimental sites on day 15 caused a drastic reduction in the numbers of *Ae.*

aegypti females from about 5 days post-application until the end of the experiments ($P < 0.001$

days 16-50; Table 4). At the treated 'sugar poor' site, the mean of captured females on the last

day was 0.29 ±0.10 and at the 'sugar rich' site it was 3.20 ±0.43 (Fig 7A and 7B, Table 4).

ATSB significantly reduced mean numbers of landing / biting female *Ae. aegypti* at both 'sugar

rich' and 'sugar poor' sites. At the sugar poor site, the mean catch pre-treatment on day 14 was

20.51 females and was reduced 70-fold to 0.29 on day 50. At the sugar rich site, the mean pre-

treatment catch on day 14 was 32.46 and was reduced 10-fold to 3.20 on day 50 (Fig 7A and 7B,

Table 4).

Table 4. Reduction of the *Ae. aegypti* population following ATSB treatment at 'sugar poor' and

'sugar rich' sites as indicated by the decrease in landings of females on human volunteers

compared to their frequency at untreated control sites.

	Sugar Poor					Sugar Rich					SP vs. SR	
	Control		Experimental			Control		Experimental			Cont	Exp
Day	Mean	SE	Mean	SE	P	Mean	SE	Mean	SE	P	P	P
1	5.46	0.63	9.10	0.94	0.001	12.27	1.20	19.41	1.78	0.001	<0.001	<0.001
3	8.51	0.89	8.98	0.93	0.716	13.69	1.32	17.53	1.63	0.065	0.001	<0.001
6	6.66	0.73	10.17	1.03	0.005	19.68	1.80	20.39	1.86	0.785	<0.001	<0.001
9	7.99	0.84	12.96	1.26	0.001	16.00	1.50	22.94	2.07	0.006	<0.001	<0.001
12	14.05	1.35	16.67	1.56	0.203	25.70	2.29	23.62	2.12	0.505	<0.001	0.007
14	16.46	1.55	20.51	1.88	0.096	22.67	2.05	32.46	2.84	0.004	0.014	0.000
16	19.19	1.77	8.58	0.90	<0.001	24.59	2.21	22.77	2.06	0.548	0.054	<0.001
19	21.04	1.93	4.81	0.58	<0.001	28.21	2.50	17.46	1.62	0.000	0.022	<0.001
22	17.06	1.59	2.49	0.36	<0.001	25.17	2.25	10.31	1.04	<0.001	0.003	<0.001

25	23.28	2.10	1.56	0.27	<0.001	30.99	2.73	7.60	0.81	<0.001	0.024	<0.001
28	18.76	1.73	0.70	0.17	<0.001	38.10	3.31	3.95	0.50	<0.001	<0.001	<0.001
31	23.57	2.12	0.47	0.14	<0.001	44.29	3.82	6.62	0.73	<0.001	<0.001	<0.001
35	26.04	2.32	0.22	0.09	<0.001	40.65	3.53	2.44	0.36	<0.001	0.000	<0.001
38	24.34	2.19	0.41	0.13	<0.001	48.95	4.18	1.14	0.22	<0.001	<0.001	0.005
41	28.89	2.55	0.26	0.10	<0.001	55.37	4.69	0.92	0.20	<0.001	<0.001	0.004
44	24.23	2.18	0.63	0.16	<0.001	62.17	5.24	1.00	0.21	<0.001	<0.001	0.156
47	30.14	2.66	0.47	0.14	<0.001	56.52	4.78	0.59	0.16	<0.001	<0.001	0.561
50	32.93	2.89	0.29	0.10	<0.001	65.89	5.55	3.20	0.43	<0.001	<0.001	<0.001

Fig 7. Reduction of *Ae. aegypti* female population following ATSB treatment. Results are shown as the reduction in mean number of landing/biting attempts per 30-minute interval for the treatment and control sites. (A) Sugar rich sites. (B) Sugar poor sites.

Discussion

In this study, we investigated the feeding of *Ae. aegypti* in the laboratory and in the field on some natural sugar sources. We compared sugar feeding of the mosquitoes in urban 'sugar poor' and 'sugar rich' habitats, and then tested the potential of mosquito control by ATSB in the two different types of environments.

The 24 hour exposure to single, potential sugar sources demonstrated that the proportion of sugar positive *Ae. aegypti* was high in those offered flowers or fruit: 93.00% of females and 92.78% of males fed on *Acacia macrostachia*, 84.44% of females and 97.22% of males fed on *C. papaya*, 93.89% of females and 83.89% of males fed on *C. melo*, 81.67% of females and 84.44% of males fed on on *M. indica*, and 92.78% of females and 93.33% of males fed on *P. juliflora.* It is interesting to note that feeding rates for females (and for males) on some flowers was low: 15.56% of females fed on *B. glabra* and 1.67% of females fed on *N. glauca.* Seedpods *(P.*

reticulatum) and sugarcane *(S. officinarum)* have tough outer coverings and therefore their sugar

is apparently inaccessible to *Ae. aegypti* (Table 1). It is not surprising that mosquitoes died in

less than a week on the sole diet of fairly inedible *B. glabra* while ~ 90% survived in a series

maintained on the highly attractive nectar of *P. juliflora.* A high rate of feeding does not

necessarily mean high quality of meals. In the survival experiment, there was no link between

the 85.00% of females feeding on *G. gracilis* flowers within 24 hrs (Table 1) and the high (~

50%) mortality in mosquitoes exposed to these flowers for 30 days (Fig 4A and 4B). In this

context, it should be noted that the mosquitoes in the above experiments may have fed on a given

source of sugar because there was no alternative. It should also be noted that in the tight space of

50 X 50 cm cages, contact between the flying mosquitoes and the offered diet is presumably

inevitable, hence feeding is not necessarily the outcome of attraction.

The preference of sugar sources in nature is exhibited in the size of catches by baited

traps which show the relative attraction compared to other tested baits and in competition with

other environmental olfactory cues. An example of the difference between direct contact with a

sugar source in cages and the performance of a sugar source as an attractant is *L. camara.*

Overnight exposure to it in a cage resulted in 60.0 to 63.3% feeding (Table 1) but used as a bait

in the field, the low catch amounted to a mean of 1.3% to 3.6% females per trap (Table 2A).

Such differences in attraction in laboratory versus field settings were noted and discussed in

early studies [15,38,39]. Some of these attractive baits were also highly attractive to *An. gambiae*

s.l., *An. sergentii, Ae. albopictus* and *Culex pipiens* [27,28,30,39] showing that mosquitoes of

different genera and of both sexes are commonly attracted to specific emanations mostly to

flowers of some plants. It would be interesting to see whether it is a developed adaptation that

guides different mosquitoes to the richest sources of sugar in their environment.

The observation of peak times for searching for sugar meals, which are at dawn and dusk

(Fig 5A and 5B), are useful for obtaining maximal information on the sugar-feeding status of *Ae.*

aegypti populations. This is particularly important in view of the rapid digestion of sugar meals

which may be misleading when mosquitoes are caught at the wrong time of the day or kept alive

before being tested for sugar. For example, in catches on volunteers in sugar rich area only

60.32% blood fed mosquitoes had stage 1 (minimal quantities) sugar quantities in the gut.

Examination of a similar series of mosquitoes caught in sugar rich areas, after 6 hrs delay, the

proportion of stage 1 blood meals increased to 90.39%. In previous studies that estimated the

digestion rates of mosquito species in the field, the half-life of fructose was approximately 7 to

10 hours [15,40,41].

To a certain degree, sugar meals inhibit the taking of blood and vice versa, both

competing for space in the gut system [42, 43] and as a result, host seeking behaviour is

terminated [44]. In Israel [34], it was shown that *Ae. albopictus* females that have fed on natural

sugar sources or ATSB did not come to humans to blood feed. An opposite extreme case is *Ae.*

aegypti that are adapted to domestic environments where sugar may be rare but there is an easy

and unlimited access to blood [45]. This mosquito uses supplementary blood meals during the

gonotrophic cycle as energy reserves [19] that convert blood into survival time in a ratio reported

to be higher than that of nondomestic species [17]. In the domestic environment, females of this

species seldom feed on sugar, but feral populations often do [45, 46]. In the laboratory, all sugar

feeding ceases when blood-host stimuli are present, but without such stimuli, sugar feeding is

frequent [47]. Thus, the option to take sugar is retained for the competitive advantage it affords

under some circumstances. A similar strategy may be used by some anthropophilic *Anopheles*

spp. [48]. Whether *Ae. aegypti* commonly, rarely, or never take sugar in nature remains

controversial [48-51].

Theoretically, such variations can be interpreted as results of metabolic differences

between mosquito subpopulations. Otherwise, assuming that producing energy from blood meals

is general in *Ae. aegypti,* differences in the rates of blood and sugar feeding could be a response

dictated by the environment that is, according to the relative abundance of sources.

It was also concluded that the enhanced blood-feeding capability among older sugar-deprived

An. gambiae demonstrated the close association between sugar-feeding and blood-feeding

behavior [52]. Our results similarly portray dependence of the blood feeding drive on sugar

feeding. In catches on volunteers, the proportion of sugar positive females was similar in sugar

poor (68.37% class I, Table 3) and in sugar rich areas (60.32% class I, Table 3). On the other

hand, in sugar rich resting habitats, the quantity of class I sugar meals was 13.18% and 65.35%

were of classes III and IV. Moreover, their proportion was about 40% greater than in the sugar

poor resting habitats where 46.31% were of class I and 25.51% were of classes III and IV. These

results indicate that following feeding on larger sugar quantities in the relatively lush vegetation

of urban sugar rich habitats, mosquitoes were less interested in blood meals. Also, in the sugar

poor and sugar rich areas, the effect of ATSB treatments were manifested at similarly rapid rates

and the reduction of the mosquito population was to similar levels. In other words, mosquitoes

responded equally when the sugar bait (ATSB) was uniformly offered in both habitats. In both

experiments, the catch on volunteers and the results of ATSB treatment, it appears that the

intensity of the search for a host blood meal depends on the prevalence or scarcity of sugar meals

in different types of *Ae. aegypti* urban habitats.

Attractive Toxic Sugar Bait (ATSB) treatment applied in the sugar poor and sugar rich

areas caused a drastic reduction in *Ae. aegypti* approaches to volunteers in both environments.

The initial effect of ATSB was apparently somewhat delayed since there was a massive supply

of newly emerged mosquitoes. Later, high female mortality reduced the oviposition and the *Ae.*

aegypti populations collapsed almost completely. At the sugar rich sites, the daily approaches of

Ae. aegypti to the volunteers decreased from an average of 28.0 before treatment to 1.7 landings /

bites a week later (Fig 7A). At the sugar poor sites, an average of 19.0 landings was reduced to

0.44 in the first week and these levels remained similarly low until the end of the experiment

(Fig 7B). The drastic effect of ATSB is comparable to the results obtained in Israel, Florida,

Morocco and Mali [28, 30, 33, 53-55].

Since sugar is the only food source for male mosquitoes [15], ATSB should be highly

effective against male *Ae. aegypti* but should also affect survival and fecundity of females as

well. Our data demonstrates that ATSB treatment can be highly effective against *Ae. aegypti* in

both sugar rich and sugar poor environments. ATSB was also highly effective against *Ae.*

albopictus in Mali [34] and it may be a significant treatment against both species, particularly if

their distribution overlaps. Generally, it appears from this study that ATSB is a new promising

tool for the control of *Ae. aegypti.*

Acknowledgements

We dedicate this work to Fatoumata Sissoko who passed away suddenly after the completion of

this work. We also wish to thank the participating communities of Bamako in Mali. This work,

titled "ICEMR-Mali: Multidisciplinary research for malaria control and prevention in West

Africa", was supported by NIH NIAID grant number U19 AI129387 awarded to JB. The funders

had no role in study design, data collection and analysis, decision to publish, or preparation of

the manuscript. The authors declare that no competing interests exist in relation to this project.

References

571 1. Gubler DJ. Epidemic dengue/dengue hemorrhagic fever as a public health, social and
572 economic problem in the 21st century. Trends in Microbiology. 2002;10(2):100-3.
573
574 2. Barrett AD, Higgs S. Yellow fever: a disease that has yet to be conquered. Annual Review of
575 Entomology 2007;52:209-29.
576
577 3. Caglioti C, Lalle E, Castilletti C, Carletti F, Capobianchi MR, Bordi L. Chikungunya virus
578 infection: an overview. New Microbiology 2013;36(3):211-7.
579
580 4. Chouin-Carneiro T, Vega-Rua A, Vazeille M, Yebakima A, Girod R, Goindin D, Dupont-
581 Rouzeyrol M, Lourenço-de-Oliveira R, Failloux AB. Differential susceptibilities of *Aedes*
582 *aegypti* and *Aedes albopictus* from the Americas to Zika virus. PLoS Neglected Tropical
583 Diseases. 2016;10(3):e0004543.
584
585 5. WHO. Yellow fever in 1998. Weekly Epidemiological Records. 1990;65:213-219.
586
587 6. Garske T, Van Kerkhove MD, Yactayo S, Ronveaux O, Lewis RF, Staples JE, Perea W,
588 Ferguson NM, Yellow Fever Expert Committee. Yellow Fever in Africa: estimating the burden
589 of disease and impact of mass vaccination from outbreak and serological data. PLoS Medicine.
590 2014; 11(5):e1001638.
591
592 7. WHO. Dengue hemorrhagic fever, diagnosis, treatment and control. WHO Geneva. 1986;7-15.
593
594 8. Chadee DD. Resting behaviour of *Aedes aegypti* in Trinidad: with evidence for the re-
595 introduction of indoor residual spraying (IRS) for dengue control. Parasites and Vectors.
596 2013;6(1):255.
597
598 9. Rapley LP, Johnson PH, Williams CR, Silcock RM, Larkman M, Long SA, Russell RC,
599 Ritchie SA. A lethal ovitrap□based mass trapping scheme for dengue control in Australia: II.
600 Impact on populations of the mosquito *Aedes aegypti*. Medical and Veterinary Entomology.
601 2009;23(4):303-16.
602

603 10. Tsuzuki A, Huynh T, Tsunoda T, Luu L, Kawada H, Takagi M. Effect of existing practices
604 on reducing *Aedes aegypti* pre-adults in key breeding containers in Ho Chi Minh City, Vietnam.
605 The American Journal of Tropical Medicine and Hygiene. 2009;80(5):752-7.
606
607 11. Benelli G, Mehlhorn H. Declining malaria, rising of dengue and Zika virus: insights for
608 mosquito vector control. Parasitology Research. 2016;115(5):1747-54.
609
610 12. Von Seidlein L, Kekulé AS, Strickman D. Novel vector control approaches: The future for
611 prevention of Zika virus transmission. PLoS Medicine. 2017; 17;14(1):e1002219.
612
613 13. WHO. Draft global vector control response 2017-2030 (Version 4.3)
614 www.who.int/malaria/areas/vector_control/Draft-WHO-GVCR-2017-2030.pdf
615 2016. Last accessed Oct. 10, 2017.
616
617 14. WHO. Mosquito (vector) control emergency response and preparedness for Zika virus
618 http://www.who.int/neglecteddiseases/news/mosquitovectorcontrolresponse/en/ 2017
619   6;11(5):e1001638.
620
621 15. Foster WA. Mosquito sugar feeding and reproductive energetics. Annual Review of
622 Entomology. 1995;40(1):443-74.
623
624 16. Yuval B. The other habit- sugar feeding by mosquitoes. Bulletin of the Society for Vector
625 Ecology. 1992; 17:150-156.
626
627 17. Nayar JK, Sauerman DM. The effects of nutrition on survival and fecundity in Florida
628 mosquitoes: Part 2. Utilization of a blood meal for survival. Journal of Medical Entomology.
629 1975;12(1):99-103.
630
631 18. Edman JD, Strickman D, Kittayapong P, Scott TW. Female *Aedes aegypti* (Diptera:
632 Culicidae) in Thailand rarely feed on sugar. Journal of Medical Entomology. 1992;29(6):1035-8.
633
634 19. Scott TW, Clark GG, Lorenz LH, Amerasinghe PH, Reiter P, Edman JD. Detection of
635 multiple blood feeding in *Aedes aegypti* (Diptera: Culicidae) during a single gonotrophic cycle
636 using a histologic technique. Journal of Medical Entomology. 1993;30(1):94-9.
637
638 20. Scott JM, Seeger KE, Gibson-Corrado J, Muller GC, Xue RD. Attractive Toxic Sugar Bait
639 (ATSB) mixed with pyriproxyfen for control of larval *Aedes albopictus* (Diptera: Culicidae)
640 through fecal deposits of adult mosquitoes. Journal of Medical Entomology. 2017;54(1):236-8.
641
642 21. France KR, Judson CL. Nitrogen partitioning and blood meal utilization by *Aedes aegypti*
643 (Diptera Culicidae). Journal of Insect Physiology. 1979;25(11):841-6.
644
645 22. Briegel H. Protein catabolism and nitrogen partitioning during oögenesis in the mosquito
646 *Aedes aegypti*. Journal of insect physiology. 1986;32(5):455-62.
647

23. Martinez-Ibarra JA, Rodriguez MH, Arredondo-Jimenez JI, Yuval B. Influence of plant abundance on nectar feeding by *Aedes aegypti* (Diptera: Culicidae) in southern Mexico. Journal of medical entomology. 1997;34(6):589-93.

24. Chadee DD, Sutherland JM, Gilles JR. Diel sugar feeding and reproductive behaviours of *Aedes aegypti* mosquitoes in Trinidad: With implications for mass release of sterile mosquitoes. Acta Tropica. 2014;132:S86-90.

25. Qualls WA, Naranjo DP, Subía MA, Ramon G, Cevallos V, Grijalva I, Gómez E, Arheart KL, Fuller DO, Beier JC. Movement of *Aedes aegypti* following a sugar meal and its implication in the development of control strategies in Durán, Ecuador. Journal of Vector Ecology. 2016;41(2):224-31.

26. Gu W, Müller G, Schlein Y, Novak RJ, Beier JC. Natural plant sugar sources of *Anopheles* mosquitoes strongly impact malaria transmission potential. PLoS One. 2011 Jan 20;6(1):e15996.

27. Müller G, Schlein Y. Sugar questing mosquitoes in arid areas gather on scarce blossoms that can be used for control. International Journal for Parasitology. 2006;36(10):1077-80.

28. Müller GC, Schlein Y. Efficacy of toxic sugar baits against adult cistern-dwelling *Anopheles claviger*. Transactions of The Royal Society of Tropical Medicine and Hygiene. 2008;102(5):480-4.

29. Müller GC, Junnila A, Schlein Y. Effective control of adult *Culex pipiens* by spraying an attractive toxic sugar bait solution in the vegetation near larval habitats. Journal of Medical Entomology. 2010a;47(1):63-6.

30. Müller GC, Beier JC, Traore SF, Toure MB, Traore MM, Bah S, Doumbia S, Schlein Y. Field experiments of *Anopheles gambiae* attraction to local fruits/seedpods and flowering plants in Mali to optimize strategies for malaria vector control in Africa using attractive toxic sugar bait methods. Malaria Journal. 2010b;9(1):262.

31. Müller GC, Beier JC, Traore SF, Toure MB, Traore MM, Bah S, Doumbia S, Schlein Y. Successful field trial of attractive toxic sugar bait (ATSB) plant-spraying methods against malaria vectors in the *Anopheles gambiae* complex in Mali, West Africa. Malaria Journal. 2010c;9(1):210.

32. Beier JC, Müller GC, Gu W, Arheart KL, Schlein Y. Attractive toxic sugar bait (ATSB) methods decimate populations of *Anopheles* malaria vectors in arid environments regardless of the local availability of favoured sugar-source blossoms. Malaria journal. 2012;11(1):31.

33. Revay EE, Müller GC, Qualls WA, Kline DL, Naranjo DP, Arheart KL, Kravchenko VD, Yefremova Z, Hausmann A, Beier JC, Schlein Y. Control of *Aedes albopictus* with Attractive Toxic Sugar Baits (ATSB) and potential impact on non-target organisms in St. Augustine, Florida. Parasitology Research. 2014;113(1):73-9.

694   34. Junnila A, Revay EE, Müller GC, Kravchenko V, Qualls WA, Allen SA, Beier JC, Schlein
695   Y. Efficacy of attractive toxic sugar baits (ATSB) against *Aedes albopictus* with garlic oil
696   encapsulated in beta-cyclodextrin as the active ingredient. Acta Tropica. 2015;152:195-200.
697
698   35. Schlein Y, Jacobson RL. Mortality of *Leishmania major* in *Phlebotomus papatasi* caused by
699   plant feeding of the sand flies. American Journal of Tropical Medicine and Hygiene.
700   1994;50(1):20-7.
701
702   36. United States Environmental Protection Agency (EPA). Product performance test guidelines
703   OPPTS 810.3700: Insect repellents to be applied to human skin. 2010 July EPA 712-C-10-00.
704
705   37. Qualls WA, Müller GC, Revay EE, Allan SA, Arheart KL, Beier JC, Smith ML, Scott JM,
706   Kravchenko VD, Hausmann A, Yefremova ZA. Evaluation of attractive toxic sugar bait
707   (ATSB)—barrier for control of vector and nuisance mosquitoes and its effect on non-target
708   organisms in sub-tropical environments in Florida. Acta Tropica. 2014;131:104-10.
709
710   38. Healy TP, Jepson PC. The location of floral nectar sources by mosquitoes: the long-range
711   responses of *Anopheles arabiensis* Patton (Diptera: Culicidae) to *Achillea millefolium* flowers
712   and isolated floral odour. Bulletin of Entomological Research. 1988;78(4):651-7.
713
714   39. Muller, GC, Junnila, A, Traore, MM, Traore, SF, Doumbia, S, Sissoko, F, Dembele, SM,
715   Schlein, Y, Arheart, KL, Revay, EE and Kravchenko, VD. The invasive shrub *Prosopis juliflora*
716   enhances the malaria parasite transmission capacity of *Anopheles* mosquitoes: a habitat
717   manipulation experiment. Malaria Journal. 2017;16(1):237.
718
719   40. Reisen WK, Meyer RP, Milby MM. Patterns of fructose feeding by *Culex tarsalis* (Diptera:
720   Culicidae). Journal of Medical Entomology. 1986;23(4):366-73.
721
722   41. Andersson IH, Jaenson TG. Nectar feeding by mosquitoes in Sweden, with special reference
723   to *Culex pipiens* and *Cx torrentium*. Medical and Veterinary Entomology. 1987;1(1):59-64.
724
725   42. Vargo AM, Foster WA. Responsiveness of female *Aedes aegypti* (Diptera: Culicidae) to
726   flower extracts. Journal of Medical Entomology. 1982;19(6):710-8.
727
728   43. Hancock RG, Foster WA. Effects of preblood-meal sugar on sugar seeking and upwind flight
729   by gravid and parous *Aedes aegypti* (Diptera: Culicidae). Journal of Medical Entomology.
730   1993;30(2):353-9.
731
732   44. Klowden MJ, Lea AO. Abdominal distension terminates subsequent host-seeking behaviour
733   of *Aedes aegypti* following a blood meal. Journal of Insect Physiology. 1979;25(7):583-5.
734
735   45. Nayar JK. *Aedes aegypti* (L.)(Diptera: Culicidae): Observations on dispersal, survival,
736   insemination, ovarian development and oviposition characteristics of a Florida population.
737   Journal of the Florida Anti-Mosquito Association. 1981;52:24-40.
738

46. Van Handel E, Edman JD, Day JF, Scott TW, Clark GG, Reiter P, Lynn HC. Plant-sugar,
glycogen, and lipid assay of *Aedes aegypti* collected in urban Puerto Rico and rural Florida.
Journal of the American Mosquito Control Association. 1994;10(2):149-53.

47. Yee WL, Foster WA. Diel sugar-feeding and host-seeking rhythms in mosquitoes (Diptera:
Culicidae) under laboratory conditions. Journal of Medical Entomology. 1992 Sep 1;29(5):784-
91.

48. Briegel H. Metabolic relationship between female body size, reserves, and fecundity of
Aedes aegypti. Journal of Insect Physiology. 1990a;36(3):165-72.

49. Holstein M. Biology of *Anopheles gambiae*: Research in Western Africa. Geneva: WHO.
1954.

50. Laarman JJ. The intake of sugars by females of wild *Anopheles gambiae* and *Anopheles
funestas*. Acta Leidensia. 1968;36.

51. Muirhead-Thomson RC. Mosquito behaviour in relation to malaria transmission and control
in the Tropics. London: Edward Arnold and Co.; 1951.

52. Straif SC, Beier JC. Effects of sugar availability on the blood-feeding behavior of *Anopheles
gambiae* (Diptera: Culicidae). Journal of Medical Entomology. 1996;33(4):608-12.

53. Müller GC, Xue RD, Schlein Y. Seed pods of the carob tree *Ceratonia siliqua* are a favored
sugar source for the mosquito *Aedes albopictus* in coastal Israel. Acta Tropica.
2010d;116(3):235-9.

54. Müller GC, Xue RD, Schlein Y. Differential attraction of *Aedes albopictus* in the field to
flowers, fruits and honeydew. Acta Tropica. 2011;118(1):45-9.

55. Khallaayoune K, Qualls WA, Revay EE, Allan SA, Arheart KL, Kravchenko VD, Xue RD,
Schlein Y, Beier JC, Müller GC. Attractive toxic sugar baits: control of mosquitoes with the low-
risk active ingredient dinotefuran and potential impacts on nontarget organisms in Morocco.
Environmental Entomology. 2013;42(5):1040-5.

Fig 1. Microtitre plate with anthrone tested samples. Class I through IV colour intensities as well as negative control wells are shown.

Fig 2. Pictures of sugar rich and sugar poor habitats. A and B) Densely populated urban areas with sparse vegetation (<5% flowering), **C and D)** Lush vegetation of irrigated gardens and parkland with areas of ≥50% flowering vegetation.

Fig 3. Local weather conditions during June and July 2016. Weather conditions in Bamako at the time of the ATSB study. Rain occurred during the study, and two treatments were applied at the sites to avoid a "wash off" effect.

Fig 4. Survival in a series of female *Ae. aegypti* that were each exposed for 31 days to one type of plant. (**A**) Survival after exposure to branches with plant blossoms. *Ricinus communis* baits were branches with extra-floral nectaries. (**B**) Sugar source types other than blossoms.

Fig 5A. Periodicity of host-seeking and sugar-seeking behavior of *Ae. aegypti* over 18 hours

(**A**) evaluated by average catches of mosquitoes landing on a human volunteer, in 30 min intervals (±SE) and (**B**) evaluated by catches of GNTs baited with highly attractive flowering branches of *P. juliflora*. Numbers shown are the average catch per volunteer, per time period and average catch per trap, per time period (±SE).

Fig 6. Percentage of sugar positive mosquitoes from the catch on volunteers and from resting habitat sweep-net catches at sugar rich (SR) and sugar poor (SP) sites. (**A**) Tested for sugar within 1 hour after collection and (**B**) 6 hours after collection.

Fig 7. Reduction of *Ae. aegypti* female population following ATSB treatment. Results are shown as the reduction in mean number of landing/biting attempts per 30-minute interval for the treatment and control sites. (A) Sugar rich sites. (B) Sugar poor sites.