Propagation of cortical activity via open-loop intrathalamic architectures: a computational analysis

Jeffrey W. Brown[a], Aynaz Taheri[b], Robert V. Kenyon[b], Tanya Berger-Wolf[b], and Daniel A. Llano[a,c,d,e]

    a. University of Illinois College of Medicine at Urbana-Champaign, Urbana, IL 61801
    b. Department of Computer Science, University of Illinois at Chicago, Chicago, IL 60607
    c. Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana-Champaign, Urbana, IL 61801
    d. Neuroscience Program, University of Illinois at Urbana-Champaign, Urbana, IL 61801
    e. Department of Molecular and Integrative Physiology, University of Illinois at Urbana-Champaign, Urbana, IL 61801

Corresponding Author:
Daniel Llano, M.D., Ph.D.
2355 Beckman Institute
405 North Mathews Avenue
Urbana, IL 61801
d-llano@illinois.edu
(217) 244-0740

Keywords: Thalamus; thalamic reticular nucleus; intrathalamic signaling; cortical signaling; open-loop; propagation; computational model

Abstract:

Propagation of signals across the cerebral cortex is a core component of many cognitive processes and is generally thought to be mediated by direct intracortical connectivity. The thalamus, by contrast, is considered to be devoid of internal connections and organized as a collection of parallel inputs to the cortex. Here, we provide evidence that "open-loop" intrathalamic connections involving the thalamic reticular nucleus (TRN) can support propagation of oscillatory activity across the cortex. Recent studies support the existence of open-loop thalamo-reticulo-thalamic (TC-TRN-TC) synaptic motifs in addition to traditional closed-loop architectures. We hypothesized that open-loop structural modules, when connected in series, might underlie thalamic and, therefore cortical, signal propagation. Using a supercomputing platform to simulate thousands of permutations of a thalamo-reticular-cortical network and allowing select synapses to vary both by class and individually, we evaluated the relative capacities of closed- and open-loop TC-TRN-TC synaptic configurations to support both propagation and oscillation. We observed that 1) signal propagation was best supported in networks possessing strong open-loop TC-TRN-TC connectivity; 2) intrareticular synapses were neither primary substrates of propagation nor oscillation; and 3) heterogeneous synaptic networks supported more robust propagation of oscillation than their homogeneous counterparts. These findings suggest that open-loop heterogeneous intrathalamic architectures complement direct intracortical connectivity to facilitate cortical signal propagation.

Significance Statement:

Interactions between the dorsal thalamus and thalamic reticular nucleus (TRN) are speculated to contribute to phenomena such as arousal, attention, sleep, and seizures. Despite the importance of the TRN, the synaptic microarchitectures forming the basis for dorsal thalamus-TRN interactions are not fully understood. The computational neural model we present incorporates "open-loop" thalamo-reticular-thalamic (TC-TRN-TC) synaptic motifs, which have been experimentally observed. We elucidate how open-loop motifs possess the capacity to shape the propagative properties of signals intrinsic to the thalamus and evaluate the wave dynamics they support relative to closed-loop TC-TRN-TC pathways and intrareticular synaptic connections. Our model also generates predictions regarding how different spatial distributions of reticulothalamic and intrareticular synapses affect these signaling properties.

Introduction:

Propagation of activity across the cerebral cortex is thought to underlie multiple cognitive processes, as well as pathological processes such as epilepsy and migraine (1-4). Cortical regions are highly interconnected via direct axonal projections as well as via polysynaptic pathways involving the basal ganglia and thalamus (5, 6). Cortical signal propagation is generally thought to be mediated via direct cortical connections (7, 8), but recent evidence suggests that the thalamus serves as a control point to modify cortical activity during cognitive processes such as attentional shifting (9). An advantage of a thalamic mode of signal propagation is the efficiency by which modulatory influences may control thalamic, and therefore cortical, propagation. The thalamus, however, is generally thought to have limited internal connectivity and therefore limited capacity to serve as a substrate for signal propagation.

A major intermediary allowing for communication between thalamocortical neurons, the thalamic reticular nucleus (TRN), is a sheet of GABAergic neurons that partially envelops the dorsal thalamus (10). It has been speculated to participate in phenomena ranging from selective attention (11-13) to sleep and arousal (12-15) and fear responses (16), and may play a role in generating absence seizures (17-21), symptoms of neurodevelopmental disorders (22, 23), and schizophrenia (24). The TRN projects exclusively to TC neurons, while receiving reciprocal, glutamatergic thalamoreticular (TC-TRN) connections (25).

The structural microarchitecture of bidirectional pathways connecting the dorsal thalamus and TRN has been the subject of ongoing debate. It was originally assumed that thalamo-reticulo-thalamic (TC-TRN-TC) pathways were reciprocal, forming "closed loops" of recurrent inhibition delivered to TC neurons (Fig. 1A, left) (10, 15, 26-28). While closed disynaptic loops have indeed been confirmed, they were only identified in a minority of examined TC-TRN pairs (10, 29-33). Another connectional scheme between the dorsal thalamus and TRN is the so-called "open-loop" TC-TRN-TC pathway, wherein a TC neuron is not reciprocally inhibited by the TRN neuron it excites (Fig. 1A, right). Open-loop configurations have been inferred from recordings in rodent thalamic slice preparations (34-38) and confirmed in anatomical studies (32, 39, 40). Furthermore, open-loop pathway variants in the form of X-TRN-TC are also known to exist, with X representing indirect sources of modulation to the sensory thalamus via the TRN, such as monoaminergic and cholinergic brainstem nuclei, nuclei of the basal forebrain, amygdala, and prefrontal cortex (9, 41-45).

We previously observed through a computational model that the open-loop TC-TRN-TC pathway, rather than uniformly depressing thalamic (and consequently cortical) activity, paradoxically enhanced thalamocortical output over a range of TC and TRN input frequencies (46). This finding demonstrated the capacity of an open-loop system to function as a tunable filter of thalamocortical transmission, subject to the temporal dynamics of input to the TRN, whether from other, non-reciprocally connected TC neurons or extrinsic sources. In both our previous model and earlier models built on closed-loop TC-TRN-TC synaptic motifs, the post-inhibitory rebound exhibited by TC neurons, as mediated by T-type Ca^{2+} channels and driven by inhibition from the TRN, served as a catalyst of signal propagation within the networks (9, 46-55).

Based on previous studies of open-loop TC-TRN-TC synaptic organization, we hypothesized that these synaptic modules might underlie intrathalamic and therefore intracortical signal propagation. Accordingly, we sought here to evaluate the efficacy of open-loop pathways relative to other potential synaptic configurations in mediating signal transmission across the thalamus and cortex. To this end, we constructed a model network based on that of (46) by connecting in series three thalamo-reticulo-layer-4-cortical (TC-TRN-L4) pathways, potentially featuring both closed- and/or open-loop TC-TRN-TC motifs, with the latter constituting one mode of connectivity between parallel TC-TRN-L4 pathways. Intrareticular synapses represented the other structural connections between pathways, based on the identification of both GABAergic (56-62) and electrical synapses (61-65) between TRN neurons. Thus,

we included three different polysynaptic configurations between vertical pathways in our network (Fig.
1B, from left to right): 1) those with a chemical intrareticular synapse; 2) those with anelectrical
intrareticular synapse; and 3) open-loop TC-TRN-TC pathways. To analyze how each variety of inter-
pathway connection contributed to network dynamics, permutations of the baseline network were
generated by varying three properties associated with each of the inter-pathway synaptic motifs. We
quantified network dynamics as a function of variable TC-TRN-TC and intrareticular synaptic
architectures by defining and measuring two properties inherent to stimulus-evoked responses in each
network variant: propagation and oscillation, with the latter included in light of the fact that many
characterized thalamic waveforms both oscillate and propagate through the thalamus and cortex (25).
113
114 **Network architecture and simulations:**
115 We constructed a neuronal network comprising three interconnected thalamo-reticulo-cortical pathways
116 (Fig. 1C). Thalamic, reticular, and cortical cell layers were aligned topographically, such that TC_A
117 projected to both TRN_A and $L4_A$ (10, 25, 52, 66, 67).
118
In the case of homogeneously varied synaptic network permutations, the synaptic parameters associated
with three inter-pathway motifs varied as a class, with all external, TC-TRN, and TC-L4 synaptic
conductances held constant: 1) GABAergic intrareticular ($TRN-TRN_{GABA}$) synapses ranged in
conductance between 0 and 450 nS; 2) electrical intrareticular ($TRN-TRN_{Elec}$) synapses ranged in
coupling coefficient between 0 and 0.36; and 3) a TC-TRN-TC "openness" coefficient, defined as the
weight distribution of lateral (open-loop, comprising 2 synapses of the form $TRN_i \rightarrow TC_{i+1}$) vs. recurrent
(closed-loop, comprising 3 synapses of the form $TRN_i \rightarrow TC_i$) reticulothalamic connectivity, varied
between 0 (completely closed-loop) and 1.0 (completely open-loop) and with a baseline TRN-TC
conductance of 80 nS.
128
129 For the heterogeneously varied synaptic network variants, all TRN-TRN and TRN-TC synapses were
130 allowed to vary independently. Domains for each of the synaptic variables were selected to include the
131 range of conductance or coupling strengths reported in physiological measurements and/or used in similar
132 neural models (19, 49, 52, 54, 63, 64, 67).
133
Ongoing afferent synaptic input was delivered to every TC neuron in the model as Poisson-modulated
spike trains centered at 40 Hz. An additional 200-Hz pulse train was applied to neuron TC_A between
$t=0.400$ and $t=1.500$ s during every network simulation run. This high-frequency stimulus was modeled
on those used to elicit spindle-like waves in a ferret thalamoreticular slice preparation (18, 68). A given
network's output was compiled by assembling spike histograms (10-ms bins) averaging 1,000 simulations
for every L4 neuron (Fig. 1D). Network properties were quantified in the most downstream element of the
cortical output layer, $L4_C$. Propagation across a network was quantified as the amplitude of the initial
stimulus-evoked response in the detrended $L4_C$ histogram. The degree of oscillation supported by each
network permutation was defined as the amplitude of the first off-center peak in the normalized
autocorrelogram of post-stimulation activity (Fig. 1D). Both propagation and oscillation scores are
reported as normalized to the maximum scores tabulated for each property. Given the high prevalence of
propagating oscillatory waves in the cerebral cortex [reviewed in (69)], we furthermore defined a
composite "optimization" (*Op*) metric to measure the capacity of networks to simultaneously support and
balance between propagation (*Pr*) and oscillation (*Os*):

$$Op = \sqrt{Pr^2 + Os^2} - |Pr - Os| \qquad (1)$$

149
Results:
Homogeneously varied synaptic models
Stimulus-evoked responses propagated linearly across the length of homogeneous synaptic networks,
occurring at average fixed intervals of 93.31 ± 0.35 ms (mean ± standard error of the mean; range, 60-110

ms) between adjacent TC-TRN-L4 pathways, across all model permutations and with a mean velocity of
0.54 mm/s, assuming a 50 μm separation between adjacent neurons in each network layer. All 770
homogeneous network variants were ranked according to their cortical propagation scores (Fig. 2A, top).
Linear regression analysis (R^2=0.793, root-mean-square-error or RMSE=0.047, p<0.0001) demonstrated a
strong positive correlation between the TC-TRN-TC openness coefficient and propagation score
(normalized regression coefficient or NRC=1.000). By contrast, chemical and electrical TRN-TRN
synaptic connectivity tended to modestly diminish propagation (NRC=-0.173 and NRC=-0.136,
respectively; Table S1). Further, other excitatory connectivity, such as cortico-cortical or corticothalamic
connectivity, often postulated as being important for cortical signal propagation (5, 7, 8), was not
necessary. Thus, the homogeneously varied synaptic network permutations that best accommodated
signal propagation were generally ones with weak or absent synapses between TRN neurons and strong
open-loop TC-TRN-TC connections. For example, Network a, which epitomizes this architecture,
exhibited robust signal propagation in response to a fixed stimulus delivered to TC_A; a representative
simulation of this network is shown in Fig. 2B, left, and its position in Fig. 2C is labeled. Stimulus-
evoked activity in this network tended to propagate efficiently from $L4_A$ to $L4_C$: near-synchronous
propagation cascades were elicited in both the TRN and L4 layers of the model, having been stimulated
by propagating activity in upstream TC neurons. Smooth, linear propagation of action potentials across
the network depended on the synchronous induction of inhibitory postsynaptic potentials (IPSPs) and the
ensuing post-inhibitory rebound spikes in TC neurons, which occurred reliably and at fixed intervals in
Network a.

A 2° multiple regression model of propagation as a function of all three synaptic class variables
(R^2=0.842, RMSE=0.041, p<0.0001; Table S1) revealed modestly negative interaction term between
TRN-TRN_{Elec} synapses and TC-TRN-TC openness (NRC=-0.365), indicating that in networks where both
electrical synapses were strong and TC-TRN-TC openness high, the extent of supported propagation
diminished nonlinearly; a smaller negative interaction between TRN-TRN_{GABA} synapses and TC-TRN-TC
openness was also observed (NRC=-0.152). Together, these terms suggested that propagation was more
significantly affected by connections in the TRN layer as a function of increasing open-loop TC-TRN-TC
architecture. This relationship is evident in Fig. 2C, as propagation scores conspicuously decreased in
network variants with an openness coefficient of 1.0 as either chemical or electrical synapses increase in
weight.

Oscillatory responses recurred in $L4_C$ neurons at a mean frequency of 9.07 ± 0.2 Hz (range, 7.14-12.50
Hz) across all homogeneous model permutations. Propagation and oscillation scores across all 770
homogeneous networks were strongly anticorrelated (Pearson's r=-0.671, p<0.0001). Accordingly,
oscillation was best accommodated in network permutations exhibiting strongly closed-loop connectivity
(Fig. 2A, bottom), however the capacity to support oscillation was neither markedly linear nor
monotonically decreasing as a function of increasing openness coefficient (Fig. 2D). Rather, a one-way
analysis of variance (ANOVA) with Tukey's tests revealed that, on average, oscillation scores peaked and
remained statistically indistinguishable from one another across the subset of network permutations with
openness coefficients between 0 and 0.4, with scores then decreasing in a roughly linear fashion with
increasing TC-TRN-TC openness [$F(10,759)$=137.8, p<0.0001]. These data suggest that networks with
mixed open- and closed-loop connectivity (which is likely close to physiological reality) can support the
coexistence of oscillation and propagation (see *Heterogeneously varied synaptic models*, below).

The predominant mechanism by which oscillation arose in $L4_C$ was through post-inhibitory rebound in
TC_C, as engendered by the strong recurrent inhibition found in network permutations exhibiting primarily
closed-loop TC-TRN-TC connectivity. This mode of oscillation was exemplified by Network b, a
strongly closed-loop network variant. In the simulation shown of this network (Fig. 2B, right), oscillatory
activity was enabled by a single epoch of signal propagation. Notably, neither the presence of strong

GABAergic nor electrical intrareticular synapses in Network b exerted much effect on its ability to
support oscillation, as predicted by the regression models.
206
Heterogeneously varied synaptic models
Recent studies have highlighted heterogeneity in TRN neuronal connectivity, synaptic physiology and
chemical identities (70-72). We therefore examined the impact of allowing all synaptic connections
involving the TRN to be independently varied. We constructed circuit-level schematics of linear
regression models for propagation (Fig. 3A, top) and oscillation (Fig. 3A, bottom) as functions of the 14
synaptic variables in heterogeneous networks.
213
Propagation in heterogeneously varied synaptic networks increased chiefly as a function of increasing the
strength of the more downstream of the two laterally inhibitory TRN-TC synapses, $TRN_B \rightarrow TC_C$: the
corresponding term in a linear regression model of propagation (R^2=0.742, RMSE=0.069, p<0.0001;
Table S2) possessed an NRC of 1.000 (Fig. 3A, top). Propagation scores also scaled to a lesser extent
with the more upstream laterally inhibitory reticulothalamic synapse, $TRN_A \rightarrow TC_B$ (NRC=0.608). The
two inhibitory intrareticular synapses originating at the rightmost end of the model network,
$TRN_C \rightarrow TRN_A$ and $TRN_C \rightarrow TRN_B$, both exerted a small negative effect on propagation (NRC=-0.087 and
NRC=-0.084, respectively). Additionally, two TRN-TRN_{Elec} synapses, $TRN_A = TRN_B$ and $TRN_A = TRN_C$
(where the "=" denotes an electrical synapses), marginally decremented propagation in heterogeneous
networks, with NRCs of -0.051 and -0.072, respectively. These findings at an individual synaptic level
comported with the observation that strong TRN-TRN interactions, whether chemical or electrical, tended
to impede signal propagation in homogeneous network variants.
226
A 2° regression model (R^2=0.857, RMSE=0.051, p<0.0001; Table S2) disclosed a large, propagation-
enhancing interaction between the two laterally inhibitory synapses (NRC=0.753), underscoring the same
dependence of propagation on strong open-loop TC-TRN-TC connectivity as seen in homogeneously
synaptic networks, but additionally demonstrating that propagation scores increased nonlinearly as a
function of simultaneously increasing the weights of $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$. Interactions between
TRN-TRN synapses of either variety and TRN-TC synapses tended diminish propagation, as did those
between recurrent and lateral inhibitory TRN-TC synapses. Taken together, the linear and 2° regression
models indicated that heterogeneous network permutations with strong laterally inhibitory TRN-TC
synapses tended to best support propagation. Consistent response propagation across the length of the
network was epitomized by Network a', in which $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$ were both relatively
strong and those synapses impeding propagation relatively weak (Fig. 3B, left).
238
Comparisons between homogeneously and heterogeneously varied synaptic architectures
In contrast to the homogeneous models, there was a very small negative correlation between the
propagation and oscillation scores of these networks (r=-0.0296, p=0.0008), suggesting that propagation
and oscillation more easily coexist in heterogeneous than homogeneous models. This supposition was
confirmed through a 2° regression analysis (R^2=0.388, RMSE=0.118, p<0.0001), which suggested that
interactions between recurrently and laterally inhibitory TRN-TC synapses (NRCs ranging between 0.345
and 0.669) facilitated the propagation of oscillation, a mechanism typified by Network b' (Fig. 3B, right).
Two intrareticular synapses, TRN_A-TRN_C and $TRN_A = TRN_C$, tended to contribute modestly to oscillation
(NRCs of 0.115 and 0.117, respectively, in the linear regression model, R^2=0.253, RMSE=0.131,
p<0.0001; Fig. 3A, bottom), while, in their individual capacities, $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$
diminished oscillation (NRCs of -1.000 and -0.892, respectively).
250
We analyzed the relative capacities of homogeneously and heterogeneously varied synaptic networks to
support propagation, oscillation, and optimization by comparing the 20 highest scores achieved by
homogeneous and heterogeneous network permutations with respect to each performance metric. No
significant differences in mean propagation scores between top-performing homogeneous and

heterogeneous networks were disclosed [unpaired *t*-test, $t(38)=0.46$, $p=0.647$; Fig. 4]. We attributed this
lack of differences to the fact that network permutations in which the synapses TRN$_A$→TC$_B$ and
TRN$_B$→TC$_C$ were both maximally weighted would be equally capable of supporting robust signal
propagation, regardless of whether these synapses were varied homogeneously or heterogeneously. By
contrast, top-scoring heterogeneous network variants better supported both oscillation [$t(38)=13.88$,
$p<0.0001$] and optimization [$t(38)=18.04$, $p<0.0001$] than their homogeneous counterparts. Because
networks supporting the propagation of oscillatory activity would, by definition, score high with respect
to optimization, these results not only confirmed that heterogeneous networks were more likely than
homogenous networks to accommodate this oscillatory mechanism, but furthermore that propagation of
oscillation across the thalamocortical network was associated with higher oscillation scores than post-
inhibitory-driven oscillation in TC$_C$, the predominant form of oscillation observed in homogeneous
networks.
267
Discussion:
The simulations presented here suggest that open-loop TC-TRN-TC synaptic motifs (Fig. 1B, right) could
function as a substrate for signal propagation across cortical networks without the need for direct cortico-
cortical, intra-reticular or corticothalamic connectivity. Post-inhibitory rebound mediated by T-type
calcium channels served as a substrate for both propagation and oscillation in the simulated networks.
TRN-TRN connections, either chemical or electrical (Fig. 1B, left and middle), diminished horizontal
propagation by disrupting the precise timing relationships required to propagate a signal across the
network. Models with heterogeneously varied TRN synapses outperformed those whose synapses varied
as a class with respect to the propagation of oscillatory activity, consistent with the emerging literature
documenting cellular and synaptic heterogeneity in the TRN (70-72). These data suggest that widespread
propagating cortical activity, under both pathological and physiological conditions, may be mediated, at
least in part, by intrathalamic connections. The model makes strong predictions that can be tested
physiologically. Finally, the approach used here, which employed supercomputing applications to search
through a very large parameter space, serves as model for future computational models with large
parameter spaces.
283
Like most of the thalamic (19, 47-50) and thalamocortical models (52, 53, 55) that inspired our model, we
utilized single-compartment, Hodgkin-Huxley neurons. While these model cells contribute to the
computational parsimony and practicality of network models, particularly where the analysis of network
dynamics is prioritized, they neglect the intrinsic cable properties of real neurons and, relatedly, the
spatially disparate nature of synaptic integration and heterogeneous expression of intrinsic and synaptic
conductances (73, 74). Such considerations are particularly relevant here relative to dendritic distributions
of T- and H-currents in TC neurons (52, 54, 75, 76) and TRN neurons (54, 77-79). Although
multicompartment neuronal models incorporating such details could conceivably alter the network
dynamics being studied, they were not necessary to simulate the propagation of oscillatory waves seen
physiologically (18, 19, 48-50).
294
Additionally, the present model omitted explicit corticothalamic and corticoreticular synapses, both of
which have been identified and physiologically characterized to varying degrees (80-86), though the
former were effectively amalgamated with both feedforward sensory and modulatory projections to the
thalamus in the form of the Poisson-modulated external input we delivered to individual TC neurons.
Both forms of feedback have been implicated in the spread of spindle waves and in the maintenance of
their synchronization over large distance scales (on the order of the length of the mammalian forebrain)
and are furthermore known to drive spindle wave formation and propagation *in vivo* by polysynaptically
recruiting TC neurons via TRN-mediated post-inhibitory rebound (80, 83, 86-91). It should be noted,
however, that short-range coherence of spindle waves, which can be elicited in isolated thalamic slice
preparations (18, 68), is preserved following decortication, both *in vivo* and *in silico* (52, 83, 88). By
extension, it is reasonable to assume that the dynamics of the spindle-like waveforms generated in our

small-scale, broadly feedforward model, in which the cortex served solely as an output layer, would not
be qualitatively altered by corticothalamic or corticoreticular feedback.
308
Comparison to related computational models and physiological data
Although the production of spindle waves was not an explicit objective of our study, some of the wave
dynamics arising in our networks were nevertheless consistent with those inherent to spindle or spindle-
like waves. Despite possessing higher degrees of TC→TRN and TRN→TC synaptic divergence and
lacking the exclusively open-loop TC-TRN-TC architecture characterizing a subset of our network
variants, other isolated thalamic models allowing for longitudinal wave propagation similarly
accommodated this propagation along the lattice of interconnected TC and TRN neurons by way of
laterally inhibitory TRN-TC synapses (19, 48, 50, 92); at short ranges, this mechanism of signal
propagation also prevailed in larger-scale thalamo-reticulo-cortical models, while corticothalamic
projections acted to propagate activity to more distal sites [(52); see (93), for a schematic illustrating
short- and long-range thalamocortical wave propagation]. Comparably, recurrently inhibitory TRN-TC
synapses have been documented to play a vital role in the generation of oscillatory behavior in the
thalamus (17, 93). The temporal parameters of propagating and oscillation signals in our model also
matched some of those previously reported: the mean signal propagation velocity and oscillation
frequency measured across homogeneous networks fell within the ranges of spindle wave propagation
velocities and intraspindle spike frequencies reported in both physiological and computational spindle
wave studies (19, 48, 68, 94, 95).
326
Several key structural elements of our set of network models and the range of phenomena they produced
distinguish them from previous thalamic and thalamocortical models. One particularly notable point of
departure relative to similar network models was the extent to which thalamoreticular, reticulothalamic,
and thalamocortical synapses diverged. Although all three classes of synapses are known to diverge
significantly and have been observed to target neuronal somata hundreds of microns from their origins
(25, 32, 58, 96-100), the TC-TRN, TRN-TC, and TC-L4 synapses in our model were constrained to
remain strictly local and minimally divergent (or non-divergent, in the case of TC-TRN and TC-L4
synapses). With respect to the first two classes of synapses, this constraint was imposed to probe the
impact the disynaptic TC-TRN-TC open-loop motifs characterizing a subset of network permutations,
which constituted one of the foci of our study, and analyze the signal propagation they may support. This
neuroanatomical scheme contrasted with previous computational models featuring parallel,
interconnected thalamoreticular pathways, in which both TC and TRN synapsed bidirectionally with
several neighboring TRN and TC cells, respectively, within a radius of several hundred microns (e.g., 19,
48-50, 52-54, 67, 101). It is highly likely that if more divergent synaptic connections were used in the
current model, even greater propagation would have been observed.
342
The functional implications open-loop thalamo-reticulo-thalamic synaptic motifs
The spread of activity from one cortical region to another is a foundational concept at the core of our
understanding of sensory processing, higher order-cognitive functions such as attention and language,
sleep-related oscillatory phenomena, and pathological findings such as propagation of ictal discharges and
migraine. Despite the importance of communication between cortical regions, its underlying substrates
are not well understood. It has long been speculated that the TRN could serve as a control point for large-
scale cortical signal processing given its central location, the high degree of convergence of projections
involved in attention, arousal and emotion onto the TRN and the TRN's particularly strong control over
TC firing properties (11-13, 102-104). Although the anatomical bases of open-loop TC-TRN-TC motifs
have been partially characterized, their functional significance in the brain lingers as a subject of
continued speculation. Here we show that open-loop TC-TRN-TC architectures can support at least short-
range cortical signal propagation. Within the thalamus, these configurations have thus far been observed
both within and across individual thalamic nuclei and are thought to serve as pathways for intra- and
cross-modal modulation, respectively (32, 34-40); as has been previously surmised, these synaptic

pathways could also plausibly lend themselves to sensory enhancement, multisensory integration, and
attentional mechanisms (10, 35, 46, 105). At a minimum, and as inferred from physiological studies,
open-loop pathways should be fully capable of supporting signaling propagation from one thalamic relay
neuron to another through a limited number of intervening synapses (with a disynaptic pathway serving as
the shortest such configuration). Moreover, interference with thalamoreticular transmission should cause
a breakdown in some forms of cortical signal propagation. Recent work has established that stimulation of
the TRN *in vivo* can induce propagating rhythmic activity across the cortex (106-108). These data suggest
that abnormal cortical signal propagation seen in seizures, migraines or hallucinations may be disrupted
by targeted therapeutics applied to the TRN. Both forthcoming physiological investigation and future
modeling studies will be able to evaluate these predictions and help provide a full accounting of the role
of the various modes of connectivity between cortical regions.
368
Methods:
Intrinsic neuronal models
Our network model was directly based on an earlier incarnation published by our research group (46).
Single-compartment TC, TRN, and L4 model neurons obeyed Hodgkin-Huxley kinetics, with membrane
potentials V varying according to the first-order differential equation
$$C \frac{dV}{dt} = -g_L(V - E_L) - \sum_i g_i(V)(V - E_i)$$
                                                                        (2)

where C is the membrane capacitance, g_L and E_L are the leakage conductance and reversal potential,
respectively, and $g_i(V)$ and E_i are the dynamic conductance and reversal potential, respectively, of the ith
voltage-gated, ligand-gated (chemical synaptic), or electrical synaptic conductance (for electrical synaptic
conductances, the effective reversal potential is equal to the presynaptic membrane potential; see
Equation 3a). All three varieties of model neurons expressed both the standard transient sodium (I_{Na}) and
delayed-rectifier potassium (I_K) currents, as reported by (46). TC and TRN neurons additionally included
a T-type calcium conductance (T-current; I_T) and hyperpolarization-activated cation current (H-current;
I_H), following the TC model of (109). Both TRN and L4 cells expressed a slow, non-inactivating
potassium conductance (I_M), following the modeling of (110), which accounts for the spike-frequency
adaptation previously reported in physiological recordings from these neurons (46, 111). A list of intrinsic
model cell parameters, including current conductances, reversal potentials, selected gating kinetics, and
membrane capacitance, can be found in Table S3.
387
Synaptic models
The kinetics of chemical synapses in our model network conformed to the synaptic depression model of
(112), following our previous computational network model (46). This model presupposes a finite
quantity of "resources," akin to synaptic vesicles, capable of being released by the presynaptic neuron;
these resources can exist in an active, inactive, or recovered state. A parameter U_{SE} characterizes the
fraction of recovered resources that can be converted to an active state (i.e., for release by the presynaptic
neuron) following action potential induction in the presynaptic axon terminal(s). Following resource
activation, synapses inactivate according to the time constant τ_{inact}; resources become available again for
activation after a recovery period described by the time constant τ_{recov}. These parameters, along with the
neurotransmitters, postsynaptic conductances, and reversal potentials characterizing all of the chemical
synapses in our model, are given in Table S4.
399
Glutamatergic thalamoreticular and thalamocortical (TC-L4) and baseline GABAergic reticulothalamic
synaptic parameters matched those of our earlier model (46), with the latter synapses allowed to vary in
conductance. TRN-TC signaling was mediated exclusively through GABA$_A$ receptors, mirroring other
thalamic and thalamocortical models in which the slower TRN-TC GABA$_B$ conductance was omitted (51,
54, 55). Both GABAergic (TRN-TRN$_{\text{GABA}}$) and electrical synapses (TRN-TRN$_{\text{Elec}}$) were included
between TRN neurons; as with TRN-TC synapses, both varieties of TRN-TRN synapses were allowed to

vary in strength. Although evidence has been presented challenging the existence of GABAergic
intrareticular synapses in certain mammalian species and age groups (10, 63, 113-115), our model
avoided making assumptions regarding their presence, strength, or spatial distribution by allowing the
associated synaptic conductances to vary over a range of physiological values, including zero, and in
distribution. The reversal potential, conductance, and kinetics of the external synapses projecting to the
TC neurons were directly based on retinogeniculate synapses (116), although the generic nature of the
external inputs in our model allows them to represent not only immediately upstream sensory input but
also brainstem modulation (e.g., serotonergic, adrenergic) known to act on thalamic nuclei (117).

Electrical synapses between TRN neurons were based on the Cx36-dependent intrareticular gap junctions
first identified by (58). For TRN neurons, the sum of electrical synaptic currents (I_{Elec}) entering any
postsynaptic neuron j from presynaptic neurons i was included in the rightmost term from Equation 1 and
calculated as

$$I_{Elec(j)} = \sum_i g_{ij} \left(V_j - V_i \right) \tag{3a}$$

420   where g_{ij} was itself calculated as

$$g_{ij} = D(x) \frac{g_{gap}}{1/_{CC} - 1} \tag{3b}$$

where CC was the electrical coupling coefficient between TRN neurons i and j, g_{gap} is the gap junction
conductance (set at 5 nS), and $D(x)$ was a scaling factor that depended on the physical distance between
the coupled TRN neurons (54, 73, 118). TRN-TRN$_{Elec}$ synapses were symmetrical (non-rectifying), such
that $G_{ij} = G_{ji}$.

We extrapolated the attenuation of intrareticular synaptic strength as a function of intracellular distance
based on mappings of intrinsic connections within the TRN along a horizontal (anteroposterior) plane
assembled by (61). Assuming 1) an intracellular distance of 50 µm between adjacent TRN neurons, 2) a
distance x (in multiples of 50 µm) between non-adjacent neurons, and 3) a Gaussian falloff in synaptic
strength (119), the baseline (adjacent-neuron) conductances of TRN-TRN$_{GABA}$ and TRN-TRN$_{Elec}$
synapses were scaled for non-adjacent synapses using the function

$$D(x) = e^{-\frac{x^2}{2\lambda^2}} \tag{4}$$

where λ_{GABA}=531 µm and λ_{Elec}=130 µm.

Given the small spatial scale of our model, synaptic delays associated with finite axonal conductance
times within the TRN and between the TRN and dorsal thalamus were disregarded, mirroring the
simplification incorporated into previous thalamic and thalamocortical models simulating synaptic
interactions on the order of 100 microns (48, 54). Although small (~1 ms) thalamocortical delays were
inserted into the network model of (54), these were likewise omitted on the basis of the cortex functioning
solely as an output layer in our model.

Computations and statistics
Our model was coded, simulated, and analyzed in MATLAB R2018b (MathWorks), utilizing both a Dell
Inspiron 3847 and Hewlett-Packard Z840 running Windows 10 and nodes on the Illinois Campus Cluster
(National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign).
Simulations employed 0.1-ms time steps, with temporal integration based on the hybrid analytic-numeral
integration method of (120), which optimizes between accurate solutions to Hodgkin-Huxley and synaptic
models and computational efficiency. All simulations commenced with a 200-ms equilibration period,
during which no external stimulation was delivered to TC neurons; this allowed all network elements to
attain steady-state conditions. Statistical analysis was performed in both MATLAB and R (121), with the
glmnet package (122) utilized within the latter platform to perform regression analyses. Multiple linear
regression was employed to establish rudimentary relationships between synaptic classes (homogeneously
synaptic networks) or individual synapses (heterogeneously synaptic networks) and each of the two

studied network properties, even in instances where these relationships deviated from linearity. 2°
regression models with interaction terms elucidated how synaptic interactions and nonlinearities affected
these network properties. Regressions were optimized using elastic net regularization, with the specific
regularization hyperparameter α selected to minimize each regression model's root-mean-square error. To
convey the relative influence of different synaptic classes or individual synapses on dynamic network
properties, all regression coefficients are reported here as normalized to the coefficient with the largest
absolute value; the effects corresponding to NRCs with absolute values of less than 0.05 were disregarded
as negligibly influential on network dynamics. Both unpaired Student t-tests and one-way ANOVA
models were used to compare the mean property scores between different sets of networks, with Tukey's
honestly significant difference tests used to ascertain pairwise difference between groups in the latter.
Kolmogorov-Smirnov and Levene's tests were employed to confirm normality and homogeneity of
variance, respectively, when utilizing parametric mean-comparison tests; data were log-transformed as
needed to conform to these prerequisites.
468

Acknowledgments:
This work made use of the Illinois Campus Cluster, a computing resource that is operated by the Illinois
Campus Cluster Program (ICCP) in conjunction with the National Center for Supercomputing
Applications (NCSA) and which is supported by funds from the University of Illinois at Urbana-
Champaign. We additionally thank Profs. Thomas Anastasio, Ruoqing Zhu, and Rama Ratnam, Drs. Kush
Paul and Baher Ibrahim, and Mr. Weddie Jackson for their valuable analytical, statistical, and technical
insights.

References:
1.   L. Muller, A. Reynaud, F. Chavane, A. Destexhe, The stimulus-evoked population response
in visual cortex of awake monkey is a propagating wave. *Nat. Commun.* **5**, 3675 (2014).
480

481   2.   L. Muller *et al.*, Rotating waves during human sleep spindles organize global patterns of
482   activity that repeat precisely through the night. *eLife* **5**, e17267 (2016).
483

484   3.   V. Kokkinos, A.M. Koupparis, M. Koutroumanidis, G.K. Kostopoulos, Spatiotemporal
485   propagation patterns of generalized ictal spikes in childhood absence epilepsy. *Clin.*
486   *Neurophysiol.* **128**(9), 1553-1562 (2017).
487

488   4.   A.A.P. Leao, Spreading depression of activity in the cerebral cortex. *J. Neurophysiol.* **7**, 359–
489   390 (1944).
490

491   5.   B.B. Theyel, D.A. Llano, S.M. Sherman, The corticothalamocortical circuit drives higher-
492   order cortex in the mouse. *Nat. Neurosci.* **13**(1), 84-88 (2010).
493

494   6.   A. Parent, L.N. Hazrati, Functional anatomy of the basal ganglia. I. The cortico-basal
495   ganglia-thalamo-cortical loop. *Brain Res. Brain Res. Rev.* **20**(1), 91-127 (1995).
496

497   7.   R. Kötter, F.T. Sommer, Global relationship between anatomical connectivity and activity
498   propagation in the cerebral cortex. *Philos, Trans. R. Soc. Lond. B. Biol. Sci.* **355**(1393), 127-
499   134 (2000).
500

501   8.   D.J. Felleman, D.C. Van Essen, Distributed hierarchical processing in the primate cerebral
502   cortex. *Cereb. Cortex.* **1**(1), 1-47 (1991).
503

504   9.   R.D. Wimmer *et al.*, Thalamic control of sensory selection in divided attention. *Nature*
505   **526**(7575), 705-709 (2015).

506
507   10.   D. Pinault, The thalamic reticular nucleus: structure, function and concept. *Brain Res. Brain*
508   *Res. Rev.* **46**(1), 1-31 (2004).
509
510   11.   F. Crick, Function of the thalamic reticular complex: the searchlight hypothesis. *Proc. Natl.*
511   *Acad. Sci. U.S.A.* **81**(14), 4586-4590 (1984).
512
513   12.   R.W. Guillery, S.L. Feig, D.A. Lozsádi, Paying attention to the thalamic reticular nucleus.
514   *Trends Neurosci.* **21**(1), 28-32 (1998).
515
516   13.   K. McAlonan, J. Cavanaugh, R.H. Wurtz, Attentional modulation of thalamic reticular
517   neurons. *J. Neurosci.* **26**(16), 4444-4450 (2006).
518
519   14.   R.R. Llinás RR, D. Paré, Of dreaming and wakefulness. *Neuroscience* **44**(3), 521-535 (1991).
520
521   15.   M. Steriade, D. McCormick, T. Sejnowski, Thalamocortical oscillations in the sleeping and
522   aroused brain. *Science* **262**(5134), 679-685 (1993).
523
524   16.   P. Dong *et al.*, A novel cortico-intrathalamic circuit for flight behavior. *Nat. Neurosci.* doi:
525   10.1038/s41593-019-0391-6 (29 April 2019).
526
527   17.   M. von Krosigk, T. Bal, D.A. McCormick, Cellular mechanisms of a synchronized oscillation
528   in the thalamus. *Science* **261**(5119), 361-364 (1993).
529
530   18.   T. Bal, M. von Krosigk, D.A. McCormick, Role of the ferret perigeniculate nucleus in the
531   generation of synchronized oscillations in vitro. *J. Physiol.* **483**(3), 665-685 (1995).
532
533   19.   A. Destexhe, T. Bal, D.A. McCormick, T.J. Sejnowski, Ionic mechanisms underlying
534   synchronized oscillations and propagating waves in a model of ferret thalamic slices. *J.*
535   *Neurophysiol.* **76**(3), 2049-2070 (1996).
536
537   20.   J.R. Huguenard, Neuronal circuitry of thalamocortical epilepsy and mechanisms of
538   antiabsence drug action. *Adv. Neurol.* **79**, 991-999 (1998).
539
540   21.   D.A. McCormick, D. Contreras, On the cellular and network bases of epileptic seizures.
541   *Annu. Rev. Physiol.* **63**, 815-846 (2001).
542
543   22.   M.F. Wells, R.D. Wimmer, L.I. Schmitt, G. Feng, M.M. Halassa, Thalamic reticular
544   impairment underlies attention deficit in Ptchd1(Y/-) mice. *Nature* **532(7597)**, 58-63 (2016).
545
546   23.   A. Krol, R.D. Wimmer, M.M. Halassa, G. Feng, Thalamic reticular dysfunction as a circuit
547   endophenotype in neurodevelopmental disorders. *Neuron* **98(2)**, 282-295 (2018).
548
549   24.   F. Ferrarelli, G. Tononi, The thalamic reticular nucleus and schizophrenia *Schizophr. Bull.*
550   **37(2)**, 306-315 (2011).
551
552   25.   S.M. Sherman, R.W. Guillery, Exploring the thalamus (Academic Press, 2001).
553
554   26.   P.T. Hale, A.J. Sefton, L.A. Bauer, L.J. Cottee, Interrelations of the rat's thalamic reticular
555   and dorsal lateral geniculate nuclei. *Exp. Brain Res.* **45(1-2)**, 217-229 (1982).
556

557   27.   R.A. Warren, A. Agmon, E.G. Jones, Oscillatory synaptic interactions between
558   ventroposterior and reticular neurons in mouse thalamus in vitro. *J. Neurophysiol.* **72(4)**,
559   1993-2003 (1994).

561   28.   S.M. Sherman, R.W. Guillery, Functional organization of thalamocortical relays. *J.*
562   *Neurophysiol.* **76(3)**, 1367-1395 (1996).

564   29.   A. Shosaku, Cross-correlation analysis of a recurrent inhibitory circuit in the rat thalamus. *J.*
565   *Neurophysiol.* **55(5)**, 1030-1043 (1986).

567   30.   F.S. Lo, S.M. Sherman, Feedback inhibition in the cat's lateral geniculate nucleus. *Exp. Brain*
568   *Res.* **100(2)**, 365-368 (1994).

570   31.   L.J. Gentet, D. Ulrich, Strong, reliable and precise synaptic connections between thalamic
571   relay cells and neurones of the nucleus reticularis in juvenile rats. *J. Physiol,* **546(3)**, 801-811
572   (2003).

574   32.   D. Pinault, M. Deschênes, Anatomical evidence for a mechanism of lateral inhibition in the
575   rat thalamus. *Eur. J. Neurosci.* **10(11)**, 3462-3469 (1998).

577   33.   T. FitzGibbon, S.G. Solomon, A.K. Goodchild, Distribution of calbindin, parvalbumin, and
578   calretinin immunoreactivity in the reticular thalamic nucleus of the marmoset: evidence for a
579   medial leaflet of incertal neurons. *Exp. Neurol.* **164(2)**, 371-383 (2000).

581   34.   J.W. Crabtree, G.L. Collingridge, J.T. Isaac, A new intrathalamic pathway linking modality-
582   related nuclei in the dorsal thalamus. *Nat. Neurosci.* **1(5)**, 389-394 (1998).

584   35.   J.W. Crabtree, J.T. Isaac, New intrathalamic pathways allowing modality-related and cross-
585   modality switching in the dorsal thalamus. *J. Neurosci.* **22(19)**, 8754-8761 (2002).

587   36.   Y.W. Lam, S.M. Sherman, Mapping by laser photostimulation of connections between the
588   thalamic reticular and ventral posterior lateral nuclei in the rat. *J. Neurophysiol.* **94(4)**, 2472–
589   2483 (2005).

591   37.   S.C. Lee, S.J. Cruikshank, B.W. Connors, Electrical and chemical synapses between relay
592   neurons in developing thalamus. *J. Physiol.* **588(13)**, 2403-2415 (2010).

594   38.   Y.W. Lam, S.M. Sherman, Functional topographic organization of the motor reticulothalamic
595   pathway. *J. Neurophysiol.* **113(9)**, 3090-3097 (2015).

597   39.   A. Kimura, H. Imbe, T. Donishi, Y. Tamai, Axonal projections of single auditory neurons in
598   the thalamic reticular nucleus: implications for tonotopy-related gating function and cross-
599   modal modulation. *Eur. J. Neurosci.* **26(12)**, 3524-3535 (2007).

601   40.   A. Kimura, Diverse subthreshold cross-modal sensory interactions in the thalamic reticular
602   nucleus: implications for new pathways of cross-modal attentional gating function. *Eur. J.*
603   *Neurosci.* **39(9)**, 1405-1418 (2014).

605   41.   C. Asanuma, L.L. Porter, Light and electron microscopic evidence for a GABAergic
606   projection from the caudal basal forebrain to the thalamic reticular nucleus in rats. *J. Comp.*
607   *Neurol.* **302(1)**, 159-172 (1990).

42. M.E. Bickford, A.E. Günlük, S.C. Van Horn, S.M. Sherman, GABAergic projection from the basal forebrain to the visual sector of the thalamic reticular nucleus in the cat. *J. Comp. Neurol.* **348(4)**, 481-510 (1994).

43. B. Zikopoulos, H. Barbas, Prefrontal projections to the thalamic reticular nucleus form a unique circuit for attentional mechanisms. *J. Neurosci.* **26(28)**, 7348-7361 (2006).

44. Y.G. Sun YG *et al.*, Biphasic cholinergic synaptic transmission controls action potential activity in thalamic reticular nucleus neurons. *J. Neurosci.* **33(5)**, 2048–2059 (2013).

45. J.D. Pita-Almenar, D. Yu, H.C. Lu, M. Beierlein, Mechanisms underlying desynchronization of cholinergic-evoked thalamic network activity. *J. Neurosci.* **34(43)**, 14463–14474 (2014).

46. A.M. Willis, B.J. Slater, E.D. Gribkova, D.A. Llano, Open-loop organization of thalamic reticular nucleus and dorsal thalamus: a computational model. *J. Neurophysiol.* **114(4)**, 2353-2367 (2015).

47. A. Destexhe, D.A. McCormick, T.J. Sejnowski, A model for 8–10 Hz spindling in interconnected thalamic relay and reticularis neurons. *Biophys. J.* **65(6)**, 2473–2477 (1993).

48. D. Golomb, X.J. Wang, J. Rinzel, Propagation of spindle waves in a thalamic slice model. *J. Neurophysiol.* **75(2)**,750-769 (1996).

49. V.S. Sohal, J.R. Huguenard, Long-Range Connections Synchronize Rather Than Spread Intrathalamic Oscillations: Computational Modeling and In Vitro Electrophysiology. *J. Neurophysiol.* **80(4)**, 1736-1751 (1998).

50. M. Bazhenov, I. Timofeev, M. Steriade, T.J. Sejnowski, Computational models of thalamocortical augmenting responses. *J. Neurosci.* **18(16)**, 6444-6465 (1998).

51. T. Pham, J.S. Haas, Electrical synapses between inhibitory neurons shape the responses of principal neurons to transient inputs in the thalamus: a modeling study. *Sci. Rep.* **8(1)**, 7763 (2018).

52. A. Destexhe, D. Contreras, M. Steriade, Mechanisms underlying the synchronizing action of corticothalamic feedback through inhibition of thalamic relay cells. *J. Neurophysiol.* **79(2)**, 999-1016 (1998).

53. M. Bazhenov, I. Timofeev, M. Steriade, T.J. Sejnowski, Model of thalamocortical slow-wave sleep oscillations and transitions to activated states. *J. Neurosci.* **22(19)**, 8691–8704 (2002).

54. R.D. Traub *et al.*, Single-column thalamocortical network model exhibiting gamma oscillations, sleep spindles, and epileptogenic bursts. *J. Neurophysiol.* **93(4)**, 2194-2232 (2005).

55. J. Rogala, W.J. Waleszczyk, S. Lęski, A. Wróbel, D.K. Wójcik, Reciprocal inhibition and slow calcium decay in perigeniculate interneurons explain changes of spontaneous firing of thalamic cells caused by cortical inactivation. *J. Comput. Neurosci.* **34(3)**, 461-476 (2013).

56.   G. Ahlsén, S. Lindström, Mutual inhibition between perigeniculate neurones. *Brain Res.* **236(2)**, 482-486 (1982).

57.   M. Steriade, E.G. Jones, R.R. Llinás, Thalamic Oscillations and Signaling (Wiley-Interscience, 1990).

58.   C.L. Cox, J.R. Huguenard, D.A. Prince, Heterogeneous axonal arborizations of rat thalamic reticular neurons in the ventrobasal nucleus. *J. Comp. Neuro.* **366(3)**, 416-430 (1996).

59.   M.V. Sanchez-Vives, T. Bal, D.A. McCormick, Inhibitory interactions between perigeniculate GABAergic neurons. *J. Neurosci.* **17(22)**, 8894-8908 (1997).

60.   Y. Shu, D.A. McCormick, Inhibitory interactions between ferret thalamic reticular neurons. *J. Neurophysiol.* **87(5)**, 2571-2576 (2002).

61.   C. Deleuze, J.R. Huguenard, Distinct electrical and chemical connectivity maps in the thalamic reticular nucleus: potential roles in synchronization and sensation. *J. Neurosci.* **26(33)**, 8633-8645 (2006).

62.   Y.W. Lam, C.S. Nelson, S.M. Sherman, Mapping of the functional interconnections between thalamic reticular neurons using photostimulation. *J. Neurophysiol.* **96(5)**, 2593–2600 (2006).

63.   C.E. Landisman *et al.*, Electrical synapses in the thalamic reticular nucleus. *J. Neuro.* **23(3)**, 1002-1009 (2002).

64.   M.A. Long, C.E. Landisman, B.W. Connors, Small clusters of electrically coupled neurons generate synchronous rhythms in the thalamic reticular nucleus. *J. Neurosci.* **24(2)**, 341-349 (2004).

65.   P. Fuentealba *et al.*, Experimental evidence and modeling studies support a synchronizing role for electrical coupling in the cat thalamic reticular neurons in vivo. *Eur. J. Neurosci.* **20(1)**, 111-119 (2004).

66.   E.G. Jones, Some aspects of the organization of the thalamic reticular complex. *J. Comp. Neurol.* **162(3)**, 285-308 (1975).

67.   V.S. Sohal, M.M. Huntsman, J.R. Huguenard, Reciprocal inhibitory connections regulate the spatiotemporal properties of intrathalamic oscillations. *J. Neurosci.* **20(5)**, 1735-1745 (2000).

68.   U. Kim, T. Bal, D.A. McCormick, Spindle waves are propagating synchronized oscillations in the ferret LGNd in vitro. *J. Neurophysiol.* **74(3)**, 1301-1323 (1995).

69.   L. Muller, F. Chavane, J. Reynolds, T.J. Sejnowski, Cortical travelling waves: mechanisms and computational principles. *Nat. Rev. Neurosci.* **19(5)**, 255-268 (2018).

70.   S.H. Lee, G. Govindaiah, C.L. Cox, Heterogeneity of firing properties among rat thalamic reticular nucleus neurons. *J. Physiol. (Lond.)* **582(1)**, 195-208 (2007).

71.   M.M. Halassa *et al.*, State-dependent architecture of thalamic reticular subnetworks. *Cell* **158(4)**, 808-821 (2014).

72. A. Clemente-Perez *et al.*, Distinct Thalamic Reticular Cell Types Differentially Modulate Normal and Pathological Cortical Rhythms. *Cell Rep.* **19(10)**, 2130-2142 (2017).

73. P. Dayan, L.F. Abbott, Theoretical Neuroscience, revised edition (MIT Press, 2005).

74. A.V. Herz, T. Gollisch, C.K. Machens, D. Jaeger, Modeling single-neuron dynamics and computations: a balance of detail and abstraction. *Science* **314(5796)**, 80-85 (2006).

75. D.A. McCormick, H.C. Pape, Properties of a hyperpolarization-activated cation current and its role in rhythmic oscillation in thalamic relay neurones. *J. Physiol. (Lond.)* **431**, 291-318 (1980).

76. S.R. Williams, G.J. Stuart, Action potential backpropagation and somato-dendritic distribution of ion channels in thalamocortical neurons. *J. Neurosci.* **20(4)**, 1307-1317 (2000).

77. D. Contreras, R. Curró Dossi, M. Steriade, Electrophysiological properties of cat reticular thalamic neurones in vivo. *J. Physiol.* **470**, 273-294 (1993).

78. A. Destexhe, D. Contreras, M. Steriade, T.J. Sejnowski, J.R. Huguenard, In vivo, in vitro, and computational analysis of dendritic calcium currents in thalamic reticular neurons. *J. Neurosci.* **16(1)**, 169-185 (1996).

79. S.R. Crandall, G. Govindaiah, C.L. Cox, Low-threshold Ca^{2+} current amplifies distal dendritic signaling in thalamic reticular neurons. *J. Neurosci.* **30(46)**, 15419-15429 (2010).

80. M. Steriade, P. Wyzinski, V. Apostol, "Corticofugal projections governing rhythmic thalamic activity" in Corticothalamic Projections and Sensorimotor Activities, T.L. Frigyesi, E. Rinvik, M.D. Yahr, Eds. (Raven, 1972), pp. 221-272.

81. E.L. White, S.M. Hersch, A quantitative study of thalamocortical and other synapses involving the apical dendrites of corticothalamic projection cells in mouse SmI cortex. *J. Neurocytol.* **11(1)**, 137-157 (1982).

82. M. De Curtis, R. Spreafico, G. Avanzini, Excitatory amino acids mediate responses elicited in vitro by stimulation of cortical afferents to reticularis thalami neurons of the rat. *Neuroscience* **33(2)**, 275-283 (1989).

83. D. Contreras, A. Destexhe, T.J. Sejnowski, M. Steriade, Control of spatiotemporal coherence of a thalamic oscillation by corticothalamic feedback. *Science* **274(5288)**, 771-774 (1996).

84. H. Blumenfeld, D.A. McCormick, Corticothalamic inputs control the pattern of activity generated in thalamocortical networks. *J. Neurosci.* **20(13)**, 5153-5162 (2000).

85. L. Zhang, E.G. Jones, Corticothalamic inhibition in the thalamic reticular nucleus. *J. Neurophysiol.* **91(2)**, 759-766 (2004).

86. S.R. Crandall, S.J. Cruikshank, B.W. Connors, A corticothalamic switch: controlling the thalamus with dynamic synapses. *Neuron* **86(3)**, 768-782 (2015).

87. J.P. Roy, M. Clercq, M. Steriade, M. Deschênes, Electrophysiology of neurons of lateral thalamic nuclei in cat: mechanisms of long-lasting hyperpolarizations. *J. Neurophysiol.* **51(6)**, 1220-35 (1984).

88. D. Contreras, M. Steriade, Spindle oscillation in cats: the role of corticothalamic feedback in a thalamically generated rhythm. *J. Physiol.* **490(1)**, 159-179 (1996).

89. N. Suga, X. Ma, Multiparametric corticofugal modulation and plasticity in the auditory system. *Nat. Rev. Neurosci.* **4(10)**, 783-794 (2003).

90. A.M. Sillito, J. Cudeiro, H.E. Jones, Always returning: feedback and sensory processing in visual cortex and thalamus. *Trends Neurosci.* **29(6)**, 307-316 (2006).

91. J.M. Sorokin *et al.,* Bidirectional Control of Generalized Epilepsy Networks via Rapid Real-Time Switching of Firing Mode. *Neuron* **93(1)**, 194-210 (2017).

92. L. Muller, A. Destexhe, Propagating waves in thalamus, cortex and the thalamocortical system: Experiments and models. *J. Physiol. Paris* **106(5-6)**, 222-238 (2012).

93. A. Destexhe, T.J. Sejnowski, Interactions between membrane conductances underlying thalamocortical slow-wave oscillations. *Physiol. Rev.* **83(4)**, 1401-1453 (2003).

94. M. Steriade, M. Deschênes, The thalamus as a neuronal oscillator. *Brain Res.* **320(1)**, 1-63 (1984).

95. P. Andersen, S.A. Andersson, The physiological basis of the alpha rhythm (Appleton-Century-Crofts, 1968).

96. C.L. Cox, J.R. Huguenard, D.A. Prince, Nucleus reticularis neurons mediate diverse inhibitory effects in thalamus. *Proc. Natl. Acad. Sci. U.S.A.* **94(16)**, 8854-8859 (1997).

97. E.G. Jones, The Thalamus (Plenum Press, 1985).

98. J.W. Crabtree, Organization in the somatosensory sector of the cat's thalamic reticular nucleus. *J. Comp. Neurol.* **366(2)**, 207-222 (1996).

99. J.M. Alonso, W.M. Usrey, R.C. Reid, Rules of connectivity between geniculate cells and simple cells in cat primary visual cortex. *J. Neurosci.* **21(11)**, 4002-4015 (2001).

100. L.M. Miller, M.A. Escabí, H.L. Read, C.E. Schreiner, Functional convergence of response properties in the auditory thalamocortical system. *Neuron* **32(1)**, 151-160 (2001).

101. E.M. Izhikevich, G.M. Edelman, Large-scale model of mammalian thalamocortical systems. *Proc. Natl. Acad. Sci. U.S.A.* **105(9)**, 3593-3598 (2008).

102. C.H. Brunia, G.J. Van Boxtel, Wait and see. *Int. J. Psychophysiol.* **43(1)**, 59-75 (2001).

103. C.D. Yingling, J.E. Skinner, Selective regulation of thalamic sensory relay nuclei by nucleus reticularis thalami. *Electroencephalogr. Clin. Neurophysiol.* **41(5)**, 476-482 (1976).

808 104. Y.J. John, D. Bullock, B. Zikopoulos, H. Barbas, Anatomy and computational modeling of
809    networks underlying cognitive-emotional interaction. *Front. Hum. Neurosci.* **7**, 101 (2013).
810

811 105. J.W. Crabtree, Functional Diversity of Thalamic Reticular Subnetworks. *Front. Syst.*
812    *Neurosci.* **12**, 41 (2018). doi:10.3389/fnsys.2018.00041
813

814 106. M.M. Halassa *et al.*, Selective optical drive of thalamic reticular nucleus generates thalamic
815    bursts and cortical spindles. *Nat. Neurosci.* **14(9)**, 1118–1120 (2011).
816

817 107. L.D. Lewis *et al.*, Thalamic reticular nucleus induces fast and local modulation of arousal
818    state. *eLife* **4**, e08760 (2015).
819

820 108. A. Kim *et al.*, Optogenetically induced sleep spindle rhythms alter sleep architectures in
821    mice. *Proc. Natl. Acad. Sci. U.S.A.* **109(50)**, 20673-20678 (2012).
822

823 109. C. Deleuze *et al.*, T-type calcium channels consolidate tonic action potential output of
824    thalamic neurons to neocortex. *J. Neurosci.* **32(35)**, 12228–12236 (2012).
825

826 110. M. Pospischil *et al.*, Minimal Hodgkin-Huxley type models for different classes of cortical
827    and thalamic neurons. *Biol. Cybern.* **99(4-5)**, 427–441 (2008).
828

829 111. W.M. Yamada, C. Koch, P.R. Adams, "Multiple channels and calcium dynamics" in Methods
830    in Neuronal Modeling, C. Koch, I. Segev, Eds. (MIT Press, 1989), pp. 97–133.
831

832 112. M.V. Tsodyks, H. Markram, The neural code between neocortical pyramidal neurons
833    depends on neurotransmitter release probability. *Proc. Natl. Acad. Sci. U.S.A.* **94(2)**, 719–723
834    (1997).
835

836 113. D. Pinault, Y. Smith, M. Deschênes, Dendrodendritic and axoaxonic synapses in the thalamic
837    reticular nucleus of the adult rat. *J. Neurosci.* **17(9)**, 3215-3233 (1997).
838

839 114. S.J. Cruikshank, H. Urabe, A.V. Nurmikko, B.W. Connors, Pathway-specific feedforward
840    circuits between thalamus and neocortex revealed by selective optical stimulation of axons.
841    *Neuron* **65(2)**, 230-245 (2010).
842

843 115. G. Hou, A.G. Smith, Z.W. Zhang, Lack of Intrinsic GABAergic Connections in the Thalamic
844    Reticular Nucleus of the Mouse. *J. Neurosci.* **36(27)**, 7246-7252 (2016).
845

846 116. C. Chen, W.G. Regehr, Presynaptic modulation of the retinogeniculate synapse. *J. Neurosci.*
847    **23(8)**, 3130–3135 (2003).
848

849 117. A. Siegel, H.N. Sapru, Essential Neuroscience, Third Edition (Lippincott Williams &
850    Wilkins, 2015).
851

852 118. K. Shimizu, M. Stopfer, Gap junction. *Curr. Biol.* **23(23)**, R1026-1031 (2013).
853

854 119. V.S. Sohal, J.R. Huguenard, Inhibitory interconnections control burst pattern and emergent
855    network synchrony in reticular thalamus. *J. Neurosci.* **23(26)**, 8978-8988 (2003).
856

857 120. J.W. Moore, F. Ramon, On numerical integration of the Hodgkin and Huxley equations for a
858    membrane action potential. *J. Theor. Biol.* **45(1)**, 249–273 (1974).

859
860   121.   R Core Team, R: A language and environment for statistical computing (R Foundation for
861   Statistical Computing, 2013). URL http://www.R-project.org/.
862
863   122.   J. Friedman, J. Hastie, R. Tibshirani, Regularization paths for generalized linear models via
864   coordinate descent. *J. Stat. Softw.* **33(1)**, 1-22 (2010).
865

Figure Legends:
Figure 1. Pathways and properties of thalamocortical signaling. **A:** Closed- vs. open-loop thalamo-
reticulo-thalamic configurations. **B:** Three possible pathways through which a signal might propagate
from one thalamocortical (TC) neuron to another via the thalamic reticular nucleus (TRN). **C:** Baseline
thalamo-reticulo-cortical model network. Broken-line synapses were allowed to vary either as a class
(homogeneously) or independently of one another (heterogeneously). **D:** Sample L4 spike histograms
(detrended) in a network permutation responding to a fixed, sustained stimulus delivered to TC_A (yellow
arrow). The propagation score assigned to any network permutation was quantified as the amplitude of
the initial stimulus-evoked response in the detrended $L4_c$ histogram; response propagation across the L4
subnetwork (orange arrow) was consistently linear, and thus the initial response in $L4_C$ was observed at a
fixed interval relative to the onset of stimulation. Oscillation intrinsic to any network variant was
quantified as the amplitude of the first off-center peak in the normalized autocorrelogram (right) of post-
stimulation activity in the detrended $L4_C$ histogram (within broken black box). The initial 400 ms of
activity preceding the fixed stimulus (in grey) is shown here for each histogram but was not included in
the calculations of either propagation or oscillation. Note that the bin heights in the $L4_A$ histogram shown
here were truncated in order to maintain identical vertical scaling across all three L4 histograms.
882
Figure 2. Propagation and oscillation in homogeneously varied synaptic networks (N=770). **A:** Ordinal
heat maps ranking homogeneously varied synaptic network permutations according to the extent of
supported signal propagation and oscillation. The network property ranks and synaptic makeups of two
selected networks, Networks a and b, are indicated. **B:** Representative simulations and circuit diagrams
depicting the normalized synaptic makeups for the two selected networks. The yellow arrow indicates
when the fixed stimulus was delivered to TC_A in each simulation. Orange highlighting indicates epochs of
linear propagation, while circles are placed above spikes occurring during periods of oscillatory activity.
C: A heat map displaying propagation scores in TRN-TRN synaptic parameter space for the 70 fully
open-loop networks (openness coefficient=1.0), with Network a highlighted. **D:** Mean oscillation scores
for networks varied nonlinearly as a function of their openness coefficients, with networks possessing
openness coefficients of 0 and 0.4 supporting oscillation to equal extents (one-way ANOVA with Tukey
post-hoc tests, $F(10,759)=137.8$, $p<0.0001$). Error bars indicate standard errors of the mean; N.S.=not
significant.
896
Figure 3. Propagation and oscillation in heterogeneously varied synaptic networks (N=12,681). **A:**
Network regression models illustrating how propagation (top) and oscillation (bottom) varied as a
function of individual synaptic weights across simulated heterogeneously synaptic network permutations.
Gray synapses are either non-variable or associated with normalized regression coefficients with absolute
values under 0.05. Synapses with positive and negative coefficients in the regression models are depicted
separately in the left- and right-sided circuit diagrams, respectively. **B:** Representative simulations for two
selected heterogeneous networks, whose normalized synaptic weights are depicted in the circuit diagrams.
Networks a' and b' respectively illustrate propagation and propagation of oscillation from Column A to
Column C.
906
Figure 4. Propagation, as measured in those network permutations scoring highest with respect to the
property, was equally supported in networks where synaptic weights varied independently of one another
(heterogeneously; red) as in networks where synaptic strength varied homogeneously (blue) by class

[unpaired *t*-test, $t(38)=0.46$, $p=0.647$]. By contrast, oscillation and optimization scores were significantly
higher in top-performing heterogeneous networks than their homogeneous counterparts [oscillation:
$t(38)=13.88$, $p<0.0001$; optimization: $t(38)=18.04$, $p<0.0001$]. Each bar corresponds to a mean of the top
20 network propagation or oscillation scores within each synaptic architecture group; error bars indicate
standard errors of the mean. ****=$p<0.0001$; N.S.=not significant.

A

Propagation

Oscillation

TRN-TRN
GABA_A Conductance

TRN-TRN
Electrical Coupling

TC-TRN-TC Openness

B

Network a

Network b

L4

TRN

TC

200 mV

300 ms

Synaptic Weight
(Normalized)

0   1

C

Propagation Scores
(Openness Coeff. = 1.0)

GABA_A Cond. (nS)

Electrical Coupling Coeff.

0   0.36

0.95

0.75

D

N.S.

Mean Oscillation Score

Openness Coefficient

Table S1. Normalized linear and second-order regression coefficients for propagation and oscillation in homogeneously varied synaptic networks.

Normalized Regression Coefficients (NRCs)				
Synaptic Variable	Propagation Linear	Propagation 2°	Oscillation Linear	Oscillation 2°
TRN-TRN$_{GABA}$	-0.173	-0.670	-	0.060
TRN-TRN$_{Elec}$	-0.136	-0.347	-	-
TRN-TC	1.000	1.000	-1.000	-0.052
(TRN-TRN$_{GABA}$)^2	-	0.332	-	-
(TRN-TRN$_{Elec}$)^2	-	0.164	-	-
(TRN-TC)^2	-	0.594	-	-1.000
TRN-TRN$_{GABA}$ **x** TRN-TRN$_{Elec}$	-	0.262	-	-
TRN-TRN$_{GABA}$ **x** TRN-TC	-	-0.152	-	-
TRN-TRN$_{Elec}$ **x** TRN-TC	-	-0.365	-	-

• The regressions include 1°, 2°, and interaction terms corresponding to TRN-TRN$_{GABA}$, TRN-TRN$_{Elec}$, and open-loop TC-TRN-TC synapses/pathways. Terms associated with regression coefficients of absolute values < 0.05 are omitted. Positive and negative terms are highlighted in red and blue, respectively. Linear regression for propagation, R^2=0.793, RMSE=0.047, p<0.0001; second-order regression for propagation, R^2=0.842, RMSE=0.041, p<0.0001; linear regression for oscillation, R^2=0.526, RMSE=0.145, p<0.0001; second-order regression for oscillation, R^2=0.630, RMSE=0.128, p<0.0001.

Table S2. Normalized linear and second-order regression coefficients for propagation and oscillation in heterogeneously varied synaptic networks.

Normalized Regression Coefficients (NRCs)				
Synaptic Variable	Propagation Linear	Propagation 2°	Oscillation Linear	Oscillation 2°
$TRN_A\text{-}TRN_C$	-	-	0.115	-
$TRN_C\text{-}TRN_A$	-0.088			
$TRN_C\text{-}TRN_B$	-0.084	-0.073	-	-
$TRN_A\text{=}TRN_B$	-0.051	-0.091	-	-
$TRN_A\text{=}TRN_C$	-0.072	-	-	-
$TRN_B\text{=}TRN_C$	-	-0.113	0.117	-
$TRN_A\text{-}TC_A$	-0.075	-	0.621	0.077
$TRN_A\text{-}TC_B$	0.608	0.571	-0.289	-1.000
$TRN_B\text{-}TC_B$	-0.128	-0.196	0.333	0.417
$TRN_B\text{-}TC_C$	1.000	1.000	-0.379	-0.892
$TRN_C\text{-}TC_C$	-0.207	-0.239	1.000	0.107
$(TRN_C\text{-}TRN_B)^2$	-	0.079	-	-
$(TRN_A\text{-}TC_B)^2$	-	-0.245	-	0.189
$(TRN_B\text{-}TC_B)^2$	-	0.174	-	-0.093
$(TRN_B\text{-}TC_C)^2$	-	-0.472	-	0.278
$(TRN_C\text{-}TC_C)^2$	-	0.187	-	-0.146

Normalized Regression Coefficients (NRCs), continued		
Synaptic Variable	Propagation 2°	Oscillation 2°
$TRN_A\text{-}TRN_B$ x $TRN_A\text{-}TC_B$	0.070	-
$TRN_A\text{-}TRN_C$ x $TRN_C\text{-}TC_C$	-	0.215
$TRN_B\text{-}TRN_A$ x $TRN_A\text{=}TRN_B$	-	0.111
$TRN_B\text{-}TRN_A$ x $TRN_A\text{-}TC_A$	-	-0.186
$TRN_C\text{-}TRN_A$ x $TRN_A\text{-}TC_A$	-	-0.172
$TRN_C\text{-}TRN_A$ x $TRN_A\text{-}TC_B$	-0.119	-
$TRN_C\text{-}TRN_A$ x $TRN_B\text{-}TC_C$	-0.096	-
$TRN_C\text{-}TRN_B$ x $TRN_B\text{-}TC_C$	-0.153	-
$TRN_A\text{=}TRN_B$ x $TRN_B\text{-}TC_C$	-	-0.129
$TRN_A\text{=}TRN_C$ x $TRN_A\text{-}TC_A$	-	-0.114
$TRN_A\text{=}TRN_C$ x $TRN_C\text{-}TC_C$	-0.079	-
$TRN_A\text{-}TC_A$ x $TRN_A\text{-}TC_B$	-	0.634
$TRN_A\text{-}TC_A$ x $TRN_B\text{-}TC_C$	-	0.449
$TRN_A\text{-}TC_B$ x $TRN_B\text{-}TC_B$	-0.166	0.361
$TRN_A\text{-}TC_B$ x $TRN_B\text{-}TC_C$	0.753	-0.274
$TRN_A\text{-}TC_B$ x $TRN_C\text{-}TC_C$	-0.106	0.669
$TRN_B\text{-}TC_B$ x $TRN_B\text{-}TC_C$	-	0.345
$TRN_B\text{-}TC_B$ x $TRN_C\text{-}TC_C$	-	-0.192
$TRN_B\text{-}TC_C$ x $TRN_C\text{-}TC_C$	-0.124	0.399

• The regressions include 1°, 2°, and interaction terms corresponding to the 14 variable synapses in the networks. Equal signs denote gap junctions. Linear regression for propagation, R^2=0.742, RMSE=0.069, p<0.0001; second-order regression for propagation, R^2=0.857,

RMSE=0.051, p<0.0001; linear regression for oscillation, R^2=0.253, RMSE=0.131, p<0.0001; second-order regression for oscillation, R^2=0.388, RMSE=0.118, p<0.0001.

Table S3. Intrinsic model cellular parameters.

Model Cellular Parameters			
Parameter	TC cell	TRN cell	L4 cell
Leak conductance, g_L (nS)	3.263	3.7928	4.8128
Leak reversal potential, E_L (mV)	-60.03	-57	-60.2354
Transient sodium conductance, g_{Na} (nS)	1,500	3,000	3,000
Sodium equilibrium potential, E_{Na} (mV)	50		
Delayed-rectifier potassium conductance, g_K (nS)	520	400	140
M-type potassium conductance, g_M (nS)	-	3.5	1.5
M-type potassium time constant, τ_M (ms)	-	200	180
Potassium equilibrium potential, E_K (mV)	-100		-90
T-type calcium conductance, g_T (nS)	45	21	-
Calcium equilibrium potential, E_T (mV)	120		
H-current conductance, g_H (nS)	0.608	0.0192	-
H-current reversal potential, E_H (mV)	-33		-
Membrane capacitance, C_m (pF)	100.4	75.0	109.3865

Table S4. Model synaptic parameters.

Model Synaptic Parameters						
Synapse	Neurotransmitter	Conductance (nS)	τ_{recov} (ms)	τ_{inact} (ms)	Reversal Potential (mV)	U_{SE}
External synapse to TC cell	(Glutamate)	32	125	2.64	0	0.76
TC-to-TRN cell synapse (TC-TRN)	Glutamate	150	500	2.64	0	0.76
TC-to-L4 cell synapse (TC-L4)	Glutamate	50	160	11.52	0	0.8113
TRN-to-TC cell synapse (TRN-TC)	GABA$_A$	Variable (0-80)	167.29	16.62	-80	0.62
Chemical TRN-to-TRN cell synapse (TRN-TRN$_{GABA}$)	GABA$_A$	Variable (0-450)	225	15	-75	0.62