A computational model of intrathalamic signaling via open-loop thalamo-reticular-thalamic architectures

Jeffrey W. Brown[1], Aynaz Taheri[2], Robert V. Kenyon[2], Tanya Berger-Wolf[2], and Daniel A. Llano[1,3,4,5]

1.   University of Illinois College of Medicine at Urbana-Champaign, Urbana, IL 61801

2.   Department of Computer Science, University of Illinois at Chicago, Chicago, IL 60607

3.   Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana-

Champaign, Urbana, IL 61801

4.   Neuroscience Program, University of Illinois at Urbana-Champaign, Urbana, IL 61801

5.   Department of Molecular and Integrative Physiology, University of Illinois at Urbana-

Champaign, Urbana, IL 61801

Corresponding Author:

Daniel Llano, M.D., Ph.D.

2355 Beckman Institute

405 North Mathews Avenue

Urbana, IL 61801

d-llano@illinois.edu

(217) 244-0740

**Abstract**:

The thalamic reticular nucleus (TRN), a sheet of GABAergic neurons that partially envelops, receives

excitatory input from, and projects inhibitory output to the dorsal thalamus, is known to form part of the

thalamus' intrinsic connectivity. In this capacity, the TRN has been shown to play a critical role in

shaping physiological phenomena such as spindle-wave and absence-seizure activity, while it is also

speculated to contribute to integrative neural functions such as arousal and attention. It was long supposed

that pairs of thalamic relay (TC) and TRN neurons formed "closed" disynaptic loops, in which a TC

neuron was inhibited by the TRN neuron it excited. Recent experimental observations and modeling

studies, however, support both the existence and potential functional significance of "open-loop"

thalamo-reticulo-thalamic (TC-TRN-TC) synaptic motifs, in which neurons from the TRN are not

reciprocally excited by the TC neurons they inhibit. We hypothesized that these structural modules, when

connected in series, might underlie certain modes of signal propagation from one part of the thalamus to

another. In the present study, we sought to evaluate the relative capacities of closed- and open-loop TC-

TRN-TC synaptic configurations to support both stimulus-evoked propagation and oscillation, both of

which characterize a variety thalamic and thalamocortical waveforms, while simultaneously exploring the

possibility that synaptic connections exclusive to the TRN, of which both chemical or electrical varieties

have been identified, might cooperatively or separately underlie these wave properties. To this end, we

generated and simulated permutations of a small thalamo-reticular-cortical network, allowing select

synapses to vary both by class (homogeneously) and independently (heterogeneously), and examined how

synaptic variations altered the propagative and oscillatory properties of the stimulus-driven responses

arising in the networks. Our analysis revealed that 1) stimulus-evoked signal propagation was best

supported in networks possessing strong open-loop TC-TRN-TC connectivity; 2) oscillation arose most

commonly though one of two mechanisms, one of which involved periodically occurring post-inhibitory

rebound induced by the TRN in the thalamus and required strongly closed-loop TC-TRN-TC motifs and

the other of which was characterized by the propagation of oscillatory activity across a network and was

dependent on uniformly strong reticulothalamic synapses; 3) intrareticular synapses were neither primary

substrates of propagation nor oscillation, tending to interfere with the former and either attenuating or

facilitating a weak, nondominant form of the latter; 4) neither the average propagative nor oscillatory

efficiency of those network permutations best accommodating these properties significantly changed as a

function of altering the duration of a fixed, external stimulus applied to them; and 5) heterogeneously

synaptic networks tended to support more robust oscillation than their homogeneous counterparts, while

the capacity to support propagation did not depend on the spatial uniformity synaptic weights. We relate

these findings to those elucidated by related modeling studies constructed around exclusively closed-loop

TC-TRN-TC connectivity and discuss the functional implications of both thalamic architectures relative

to experimental data concerning normal and pathological processes in the thalamus.

**Significance Statement:**

Interactions between the dorsal thalamus, which functions as a relay for sensory, motor, and integrative

information from subcortical brain structures to the cerebral cortex, and thalamic reticular nucleus (TRN)

are known to underlie various neurophysiological waveforms and are speculated to contribute to

phenomena such as arousal, attention, sleep, and epileptic processes. Despite this, the synaptic

microarchitectures forming the basis for dorsal thalamus-TRN interactions are not fully understood. The

computational neural model we present in this manuscript is among the first to incorporate so-called

"open-loop" thalamo-reticular-thalamic (TC-TRN-TC) synaptic motifs, which have been experimentally

verified in both anatomical and physiological studies. We elucidate how open-loop motifs possess the

capacity to shape the propagative and oscillatory properties of signals intrinsic to the thalamus and

evaluate the wave dynamics they support relative to closed-loop TC-TRN-TC pathways and intrareticular

synaptic connections. Our model also generates predictions regarding how different spatial distributions

of reticulothalamic and intrareticular synapses affect these signaling properties.

## Introduction:

The thalamus has long been regarded as the "relay station" of the brain, through which most sensory information travels to reach dedicated areas of the cerebral cortex. While pathways between the thalamus and cortex have been extensively studied, both structurally and functionally, the nature and functional significance of intrathalamic pathways and signal propagation represent outstanding questions. Monosynaptic connections between thalamic relay (thalamocortical; TC) neurons are not believed to exist abundantly within the mature mammalian brain (Smith et al., 2006; Lee et al., 2010), and while inhibitory interneurons found in thalamic nuclei of certain mammalian species project to TC neurons (Guillery and Sherman, 2002; Sherman, 2004), there is no evidence that these interneurons participate in disynaptic pathways between TC neurons.

In fact, the major intermediary allowing for communication between thalamic relay neurons, the thalamic reticular nucleus (TRN), is extrinsic to the thalamic nuclei in which these neurons reside. The TRN is a sheet of GABAergic neurons in the ventral thalamus that partially envelops the dorsal thalamus (typically considered the "thalamus proper," in which all first- and higher-order thalamic nuclei are found; Pinault, 2004). Although the TRN was identified in the late 19[th] Century and has been recorded from extensively in slice preparations, a dearth of *in vivo* recordings owing to the TRN's deep position within the brain has left a lack of definitive information on the structure's integrative functional significance; the structure has, however, been speculated to participate in phenomena ranging from selective attention (Crick, 1984; Guillery et al., 1998; McAlonan et al., 2006) to sleep and arousal (Llinás and Paré, 1991; Steriade et al., 1993), and may also play a role in generating certain kinds of seizures (von Krosigk et al., 1993; Bal et al., 1995; Destexhe et al., 1996a; Huguenard, 1998; McCormick and Contreras, 2001) and neurodevelopmental disorders (Wells et al., 2016; Krol et al., 2018). The TRN projects exclusively to TC neurons of the dorsal thalamus, with the former functioning as the latter's primary source of inhibition and, consequently, as a putative filter of thalamocortical signaling (Sherman and Guillery, 2001). Reciprocal, glutamatergic thalamoreticular (TC-TRN) connections are also known to exist, with the bidirectional communication between the dorsal thalamus and TRN thought to form the minimal network

necessary for the generation and propagation of sleep spindles, a waveform endemic to stage 2 NREM

sleep (Steriade and Deschênes, 1984; Steriade et al., 1987; von Krosigk et al., 1993; Bal et al., 1995; Kim

et al., 1995; Ulrich and Huguenard, 1997).

The structural constitution of bidirectional pathways connecting the dorsal thalamus and TRN has been

the subject of ongoing debate. It was originally assumed that thalamo-reticulo-thalamic (TC-TRN-TC)

pathways were reciprocal in nature, forming "closed loops" of recurrent inhibition delivered to TC

neurons (Fig. 1A, left; Hale et al., 1982; Steriade et al., 1993; Warren et al., 1994; Sherman and Guillery,

1996; Pinault, 2004). While closed disynaptic loops have indeed been confirmed with *in vivo* and *in vitro*

electrophysiological studies in both the rat and cat thalamus, they were only identified in a minority of

examined TC-TRN pairs (Shosaku, 1986; Lo and Sherman, 1994; Gentet and Ulrich, 2003; Pinault,

2004); these observations were consistent with anatomical tracing studies in both species (Pinault and

Deschênes, 1998; FitzGibbon et al., 2000). Another connectional scheme between the dorsal thalamus

and TRN is the so-called "open-loop" TC-TRN-TC pathway, wherein a TC neuron is not reciprocally

inhibited by the TRN neuron it excites (Fig. 1A, right); within the thalamus, this disynaptic motif

functions as a substrate for lateral inhibition. Open-loop configurations have been inferred from

recordings in rodent thalamic slice preparations (Crabtree et al., 1998; Crabtree and Isaac, 2002; Lam and

Sherman, 2005; Lee et al., 2010; Lam and Sherman, 2015) and identified in anatomical labeling studies

(Pinault and Deschênes, 1998; Kimura et al., 2007; Kimura, 2014). Furthermore, open-loop pathway

variants in the form of X-TRN-TC are also known to exist, with X representing potentially indirect

sources of modulation to the sensory thalamus via the TRN, such as monoaminergic and cholinergic

brainstem nuclei, nuclei of the basal forebrain, amygdala, and multimodal association cortex (Asanuma

and Porter, 1990; Bickford et al., 1994; Zikopoulos and Barbas, 2006; Sun et al., 2013; Pita-Almenar et

al., 2014).

Several computational models simulating either thalamic (Destexhe et al., 1993; Destexhe et al., 1994;

Golomb et al., 1996; Destexhe et al., 1996a; Sohal and Huguenard, 1998; Bazhenov et al., 1998;

Wasylenko et al., 2010; Pham and Haas, 2018) or thalamocortical networks (Destexhe et al., 1998;

Bazhenov et al., 2002; Traub et al., 2005; Izhikevich and Edelman, 2008; Rogala et al., 2013) and

incorporating closed-loop TC-TRN-TC pathways have reproduced propagative and/or oscillatory

waveforms intrinsic to the thalamus with high fidelity, including sleep spindles, epileptiform activity, and

gamma oscillations, and have engendered important dynamical clarifications or experimentally verifiable

predictions in relation to these phenomena. In a departure from these computational studies and in an

effort to explore the effects of thalamocortical transmission within an open-loop thalamoreticular system,

our research group previously simulated a one-dimensional, three-neuron thalamo-reticulo-cortical

computational network, in which a TRN neuron projected to a TC neuron non-reciprocally (i.e., in an

open-loop manner), with both the TC and TRN neurons receiving stochastic external input (Willis et al.,

2015). We reported that the open-loop TC-TRN-TC pathway, rather than uniformly depressing thalamic

(and consequently cortical) activity, paradoxically enhanced thalamocortical output over a domain of TC

and TRN input frequencies. This demonstrated the capacity of an open-loop system to function as a

tunable filter of thalamocortical transmission, subject to the temporal dynamics of input to the TRN,

whether from other, non-reciprocally connected TC neurons or extrinsic sources. In both our previous

model and earlier models built on closed-loop TC-TRN-TC synaptic motifs, the post-inhibitory rebound

(PIR) exhibited by TC neurons, as mediated by T-type $Ca^{2+}$ channels and driven by inhibition from the

TRN, served as a catalyst of signal propagation within the networks: in the case of the open-loop network,

PIR could enhance vertical (thalamocortical) transmission, while in closed-loop networks, it could drive

both horizontal (intrathalamic) and vertical signal propagation (Destexhe et al., 1993; Golomb et al.,

1996; Destexhe et al., 1996a; Destexhe et al., 1998; Sohal and Huguenard, 1998; Bazhenov et al., 1998;

Bazhenov et al., 2002; Traub et al., 2005; Rogala et al., 2013; Willis et al., 2015; Pham and Haas, 2018).

In physiological studies, TRN-driven PIR has been observed to promote thalamic bursting behavior (von

Krosigk et al., 1993; Sherman, 2001; Halassa et al., 2011).

Based on previous physiological characterizations and computational modeling of open-loop TC-TRN-

TC synaptic organization, we hypothesized that these synaptic modules might underlie intrathalamic

signal propagation. Accordingly, we sought in the present study to test this hypothesis, specifically

evaluating the efficacy of open-loop pathways relative to other potential synaptic configurations in

mediating signal transmission across the thalamus. To this end, we constructed a baseline model network

based on that of Willis et al. (2015) by connecting in series three thalamo-reticulo-cortical (layer 4)

pathways, or TC-TRN-L4 "columns," potentially featuring either or both closed- and open-loop TC-TRN-

TC motifs, with the latter constituting one mode of connectivity between parallel TC-TRN-L4 columns.

Intrareticular synapses represented the other structural connections between columns, based on the

identification of both GABAergic (Ahlsén and Lindström, 1982; Steriade et al., 1990; Cox et al., 1996;

Sanchez-Vives et al., 1997; Shu and McCormick, 2002; Deleuze and Huguenard, 2006; Lam et al., 2006)

and electrical synapses (Landisman et al., 2002; Long et al., 2004; Fuentealba et al., 2004; Deleuze and

Huguenard, 2006; Lam et al., 2006) between TRN neurons. Thus, we included three different

polysynaptic, intercolumnar pathway configurations in our network (Fig. 1B, from left to right): 1) those

with a central chemical intrareticular synapse; 2) those with a central electrical intrareticular synapse; and

3) open-loop TC-TRN-TC pathways.

To analyze how each variety of intercolumnar pathway contributed to network dynamics, permutations of

the baseline network were generated by varying three properties associated with each of the intercolumnar

synaptic motifs: 1) the conductance of GABAergic TRN-TRN synapses; 2) the electrical coupling

coefficient between TRN neurons; and 3) a TC-TRN-TC "openness" coefficient that corresponded to the

ratio of lateral to recurrent reticulothalamic connections within the network. In an effort to account for the

varying degrees of spatial uniformity with which intrareticular and TC-TRN-TC synaptic architectures

might be distributed within the thalamus, the synaptic parameters were varied in one of two manners: 1)

homogeneously, whereby parameters were varied systematically within each variable synaptic class and

across a range of discrete physiological values, or 2) heterogeneously, where parameters varied discretely

for every individual intracolumnar synapse and in which the TC-TRN-TC openness coefficient was

replaced by the conductances of individual reticulothalamic (TRN-TC) synapses. We quantified network

dynamics as a function of variable TC-TRN-TC and intrareticular synaptic architectures by defining and

measuring two properties inherent to stimulus-evoked responses in each network variant: propagation and

oscillation, with the latter included in light of the fact that many characterized thalamic waveforms both

oscillate and propagate through the thalamus (Sherman and Guillery, 2001). We furthermore compared

the relative capacities of homogeneous and heterogeneous network permutations to support these

properties.

**Computational Model and Methods:**

*Intrinsic neuronal models*

Our network model was directly based on an earlier incarnation published by our research group (Willis

et al., 2015). Single-compartment TC, TRN, and cortical (L4) model neurons obeyed Hodgkin-Huxley

kinetics, with membrane potentials $V$ varying according to the first-order differential equation

$$C\frac{dV}{dt} = -g_L(V - E_L) - \sum_i g_i(V)(V - E_i) \tag{1}$$

where $C$ is the membrane capacitance, $g_L$ and $E_L$ are the leakage conductance and reversal potential,

respectively, and $g_i(V)$ and $E_i$ are the dynamic conductance and reversal potential, respectively, of the $i$th

voltage-gated, ligand-gated (chemical synaptic), or electrical synaptic conductance (for electrical synaptic

conductances, the effective reversal potential is equal to the presynaptic membrane potential; see

Equation 2a). All three varieties of model neurons expressed both the standard transient sodium ($I_{Na}$) and

delayed-rectifier potassium ($I_K$) currents, as reported by Willis et al. (2015). TC and TRN neurons

additionally included a T-type calcium conductance (t-current; $I_T$) and hyperpolarization-activated cation

current (h-current; $I_H$), following the TC model of Deleuze et al. (2012). Both TRN and L4 cells

expressed a slow, non-inactivating potassium conductance ($I_M$), following the modeling of Pospichil et al.

(2008), which accounts for the spike-frequency adaptation previously reported in physiological recordings

from these neurons (Yamada et al., 1989; Willis et al., 2015). A list of intrinsic model cell parameters,

including current conductances, reversal potentials, selected gating kinetics, and membrane capacitance,

can be found in Table 1.

*Synaptic models*

The kinetics of chemical synapses in our model network conformed to the synaptic depression model of

Tsodyks and Markram (1997; Tsodyks et al., 1998), following our previous computational network model

(Willis et al., 2015). The Tsodyks and Markram model presupposes a finite quantity of "resources," akin

to synaptic vesicles, capable of being released by the presynaptic neuron; these resources can exist in an

active, inactive, or recovered state. A parameter $U_{SE}$ characterizes the fraction of recovered resources that

can be converted to an active state (i.e., for release by the presynaptic neuron) following action potential

induction in the presynaptic axon terminal(s). Following resource activation, synapses inactivate

according to the time constant $\tau_{inact}$; resources become available again for activation after a recovery

period described by the time constant $\tau_{recov}$. These parameters, along with the neurotransmitters,

postsynaptic conductances, and reversal potentials characterizing all of the chemical synapses in our

model, are given in Table 2.

Glutamatergic thalamoreticular and thalamocortical (TC-L4) and baseline GABAergic reticulothalamic

synaptic parameters matched those of our earlier model (Willis et al., 2015), with the latter synapses

allowed to vary in conductance (see the *Network architecture and permutations* subsection for additional

details). TRN-TC signaling was mediated exclusively through GABA$_A$ receptors, mirroring other

thalamic and thalamocortical models in which the slower TRN-TC GABA$_B$ conductance was omitted

(Traub et al., 2005; Rogala et al., 2013; Willis et al., 2015; Pham and Haas, 2018). Both GABAergic

(TRN-TRN$_{GABA}$) and electrical synapses (TRN-TRN$_{Elec}$) were included between TRN neurons; as with

TRN-TC synapses, both varieties of TRN-TRN synapses were allowed to vary in strength. Although

evidence has been presented challenging the existence of GABAergic intrareticular synapses in certain

mammalian species and age groups (Pinault et al., 1997; Landisman et al., 2002; Pinault, 2004;

Cruikshank et al., 2010; Hou et al., 2016), our model avoided making assumptions regarding their

presence, strength, or spatial distribution by allowing the associated synaptic conductances to vary over a

range of physiological values, including zero, and in distribution.

Electrical synapses between TRN neurons were based on the Cx36-dependent intrareticular gap junctions

first identified by Landisman et al. (2002). For TRN neurons, the sum of electrical synaptic currents ($I_{Elec}$)

entering any postsynaptic neuron $j$ from presynaptic neurons $i$ was included in the rightmost term from

Equation 1 and calculated as

$$I_{Elec(j)} = \sum_i g_{ij} \left( V_j - V_i \right) \tag{2a}$$

248 where $g_{ij}$ was itself calculated as

$$g_{ij} = D(x) \frac{g_{gap}}{1/_{CC} - 1} \tag{2b}$$

where $CC$ was the electrical coupling coefficient between TRN neurons $i$ and $j$, $g_{gap}$ is the gap junction

conductance (set at 5 nS), and $D(x)$ was a scaling factor that depended on the physical distance between

the coupled TRN neurons (see the *Network architecture and permutations* subsection for additional

details; Dayan and Abbott, 2005; Traub et al., 2005; Shimizu and Stopfer, 2013). TRN-TRN$_{Elec}$ were

symmetrical (non-rectifying), such that $G_{ij}=G_{ji}$.

A generalized afferent synaptic input was delivered to every TC neuron in the model. Given that the

temporal profiles of both spontaneous and stimulus-evoked inputs impinging on the dorsal thalamus can

vary drastically from moment to moment, these external inputs were delivered as Poisson-modulated

spike trains centered at 40 Hz; the central frequency was chosen to maximize thalamocortical output

relative to the saturation of the external synapse and concurrent inhibitory input being received from the

TRN (Willis et al., 2015). Individual pulses lasted 0.1 ms. An additional high-frequency pulse train was

inserted into neuron TC$_A$ over a fixed time interval during every network simulation run (see the

*Computational methods and calculation of network dynamics* subsection for additional details). The

reversal potential, conductance, and kinetics of these synapses were directly based on retinogeniculate

synapses (Chen and Regehr, 2003), although the generic nature of the external inputs in our model allows

them to represent not only immediately upstream sensory input but also brainstem modulation (e.g.,

serotonergic, adrenergic) known to act on thalamic nuclei (Siegel and Sapru, 2015).

Given the small spatial scale of our model (see the *Network architecture and permutations* subsection for

additional details), synaptic delays associated with finite axonal conductance times within the TRN and

between the TRN and dorsal thalamus were disregarded, mirroring the simplification incorporated into

previous thalamic and thalamocortical models simulating synaptic interactions on the order of 100

microns (Golomb et al. 1996; Traub et al., 2005). Although small (~1 ms) thalamocortical delays were

inserted into the network model of Traub et al. (2005), these were likewise omitted on the basis of the

cortex functioning solely as an output layer in our model. A table containing the conductances, reversal

potentials, and gating kinetics for synaptic currents and further details thereof may be found in SI:

Materials and Methods.

*Network architecture and permutations*

We constructed a 3 x 3-neuronal network comprising three interconnected thalamo-reticulo-cortical

columns for use in this study (Fig. 2A). Thalamic, reticular, and cortical cell layers were aligned with one

another topographically, such that $TC_A$ projected to both $TRN_A$ and $L4_A$ (Jones, 1975; Destexhe et al.,

1998; Sohal et al., 2000; Sherman and Guillery, 2001; Pinault, 2004). The TC-TRN and TRN-TC

synapses in our model were strictly local and minimally divergent (or non-divergent, in the case of TC-

TRN synapses) in order to preserve disynaptic TC-TRN-TC open-loop motifs and analyze the signal

propagation they may support; see the Discussion section for an elaboration of this point. Both varieties

of intrareticular synapse, by contrast, were allowed to diverge, targeting both adjacent and non-adjacent

TRN neurons. We extrapolated the attenuation of intrareticular synaptic strength as a function of

intracellular distance based on mappings of intrinsic connections within the TRN along a horizontal

(anteroposterior) plane assembled by Deleuze and Huguenard (2006). Assuming 1) an intracellular

distance of 50 µm between adjacent TRN neurons, 2) a distance $x$ (in multiples of 50 µm) between non-

adjacent neurons, and 3) a Gaussian falloff in synaptic strength (Sohal and Hueguenard, 2000), the

baseline (adjacent-neuron) conductances of $TRN\text{-}TRN_{GABA}$ and $TRN\text{-}TRN_{Elec}$ synapses were scaled for

non-adjacent synapses using the function

295
$$D(x) = e^{-\frac{x^2}{2\lambda^2}}$$
(3)

where $\lambda_{GABA}$=531 μm and $\lambda_{Elec}$=130 μm.

In the case of so-called homogeneously synaptic network permutations, the conductances or coupling

coefficients associated with three classes of synapses were allowed to vary discretely and uniformly

within their class, with all external, TC-TRN, and TC-L4 synaptic conductances held constant: 1) TRN-

$TRN_{GABA}$ synapses ranged in conductance between 0 and 450 nS in steps of 50 nS; 2) TRN-$TRN_{Elec}$

synapses ranged in coupling coefficient between 0 and 0.36 in steps of 0.06; and 3) a TC-TRN-TC

"openness" coefficient, defined as the weight distribution of lateral (open-loop, comprising 2 synapses of

the form $TRN_i \rightarrow TC_{i+1}$) vs. recurrent (closed-loop, comprising 3 synapses of the form $TRN_i \rightarrow TC_i$)

reticulothalamic connectivity, varied between 0 (completely closed-loop) and 1.0 (completely open-loop)

in steps of 0.1 (e.g, with a baseline TRN-TC conductance of 80 nS, a network with a TC-TRN-TC

openness coefficient of 0.7 would set the conductance of all laterally inhibitory TRN-TC synapses at 56

nS and all recurrently inhibitory TRN-TC synapses at 24 nS). Accordingly, 10 (TRN-$TRN_{GABA}$) x 7

(TRN-$TRN_{Elec}$) x 11 (TC-TRN-TC) or 770 homogeneous network variants were generated.

For the heterogeneously synaptic network variants, all TRN-TRN and TRN-TC synapses (6 TRN-

$TRN_{GABA}$ + 3 TRN-$TRN_{Elec}$ + 5 TRN-TC) were allowed to vary discretely and independently of one

another, with all other synaptic weights held constant. Here, TRN-$TRN_{GABA}$ synapses ranged in

conductance between 0 and 400 nS in steps of 100 nS, TRN-$TRN_{Elec}$ synapses ranged in coupling

coefficient between 0 and 0.36 in steps of 0.12, and TRN-TC synapses (both laterally and recurrently

inhibitory) ranged in conductance between 0 and 80 nS in steps of 16 nS. Due to both computational and

time constraints, approximately 12,600 heterogeneous network permutations out of a theoretical >7

billion were generated at random. For both homogeneous and heterogeneous network variants, domains

for each of the three synaptic variables were selected to include the range of conductance or coupling

strengths reported in physiological measurements and/or used in similar neural models (Destexhe et al.,

1996a; Destexhe et al., 1998; Sohal and Hueguenard, 1998; Sohal et al., 2000; Landisman et al., 2002;

Long et al., 2004; Traub et al., 2005). For both varieties of TRN-TRN synapse, we modestly extended the

upper bounds of the synaptic parameter domains to capture a broader range of dynamics potentially

generated by intrareticular synapses in the thalamoreticular system.

*Computational methods and calculation of network dynamics*

Our model was coded, simulated, and analyzed in MATLAB R2018b (MathWorks), utilizing both a Dell

Inspiron 3847 and Hewlett-Packard Z840 running Windows 10 and nodes on the Illinois Campus Cluster

(National Center for Supercomputing Applications, University of Illinois at Urbana-Champaign).

Simulations employed 0.1-ms time steps, with temporal integration based on the hybrid analytic-numeral

integration method of Moore and Ramon (1974), which optimizes between accurate solutions to Hodgkin-

Huxley and synaptic models and computational efficiency. Statistical analysis was performed in both

MATLAB and R (R Core Team, 2013), with the *glmnet* package (Friedman et al., 2010) utilized within

the latter platform to perform regression analyses. Multiple linear regression was employed to establish

rudimentary relationships between synaptic classes (homogeneously synaptic networks) or individual

synapses (heterogeneously synaptic networks) and each of the two studied network properties, even in

instances where these relationships deviated from linearity. Second-order regression models with

interaction terms elucidated how synaptic interactions and nonlinearities affected these network

properties. Regressions were optimized using elastic net regularization, with the specific regularization

hyperparameter $\alpha$ selected to minimize each regression model's root-mean-square error (RMSE). To

convey the relative influence of different synaptic classes or individual synapses on dynamic network

properties, all regression coefficients are reported here as normalized to the coefficient with the largest

absolute value; the effects corresponding to normalized regression coefficients (NRCs) with absolute

values of less than 0.05 were disregarded as negligibly influential on network dynamics. Both paired

Student *t*-tests and one-way analysis of variance (ANOVA) models were used to compare the mean

property scores between different sets of networks, with Tukey's honestly significant difference tests used

to ascertain pairwise difference between groups in the latter. Kolmogorov-Smirnov and Levene's tests

were employed to confirm normality and homogeneity of variance, respectively, when utilizing

parametric mean-comparison tests; data were log-transformed as needed to conform to these

prerequisites.

Individual simulations of network activity lasted either 1.000 or 1.500 s. During every run of the 1.000-s

simulations, a 200-Hz spike train lasting 50 ms and beginning at the 400-ms time index was inserted at

the external synapse to $TC_A$ (fixed, "punctate" stimulation; Fig. 2B). During 1.500-s simulation runs, an

extended version of the same spike train lasting 1.100 s was delivered externally to $TC_A$ (fixed,

"sustained" stimulation; Fig. 2C). These high-frequency, time-locked stimuli were modeled on those used

to elicit spindle-like waves in a ferret thalamoreticular slice preparation (Kim et al., 1995; Bal et al.,

1995) and designed to allow for the averaging of responses within a given network permutation across

simulation runs, with variations in neuronal responses from one run to the next arising due to the

stochastic (Poisson-modulated) input delivered to every TC neuron. Homogeneously synaptic networks

were simulated for both 1.000 s and 1.500 s, while heterogeneously synaptic networks were simulated for

1.500 s. All simulations commenced with a 200-ms equilibration period, during which no external

stimulation was delivered to TC neurons; this allowed all network elements to attain steady-state

conditions.

To analyze the properties of propagation and oscillation inherent to a network's stimulus-driven response,

every network permutation was simulated 1,000 times. A given network's output across these

simulations, as represented by action potentials in the L4 layer of the model, was compiled by assembling

spike histograms consisting of 10-ms bins for every L4 neuron (Fig. 3). Both dynamic network properties

were defined relative to the most downstream element of the cortical output layer, $L4_C$. Propagation

across a network was quantified as the amplitude of the initial stimulus-evoked response in the detrended

$L4_C$ histogram. Because response propagation across the L4 subnetwork was consistently linear, the

initial response in $L4_C$ could be predictably observed within a small, fixed interval relative to the onset of

stimulation (see the Results section for additional information on signal propagation velocities). The

degree of oscillation supported by each network permutation was defined as the amplitude of the first off-

center peak in the normalized autocorrelogram of post-stimulation activity (i.e., activity measured

between 400 ms and 1500 ms) in the detrended $L4_C$ histogram; this corresponded to the degree to which

activity in $L4_C$ was able to periodically recur. (Note that although other simulated neurons exhibited

oscillatory activity as well, the term "oscillation" henceforth refers explicitly to oscillation in $L4_C$ as

defined above, unless otherwise stated.) Both propagation and oscillation scores are reported as

normalized to the maximum scores tabulated for each property across all the homogeneously and

heterogeneously synaptic networks simulated in the study.

**Results:**

*Response of homogeneously synaptic models to a fixed, punctate stimulus*

Delivering a fixed, punctate external stimulus to $TC_A$ was adequate to elicit strong propagative responses

within a subset of homogeneously synaptic network permutations. Signals propagated linearly across the

length of networks, with stimulus-evoked responses occurring at average fixed intervals of $102.29 \pm 0.51$

ms (mean $\pm$ standard error of the mean) between adjacent TC-TRN-L4 columns and with a mean velocity

of 0.49 mm/s, assuming a 50 µm separation between adjacent neurons in each network layer.

All 770 homogeneous network variants were ranked according to their propagation scores (Fig. 4A, top).

Linear regression analysis ($R^2$=0.829, RMSE=0.042, $p$<0.0001) demonstrated a strong positive correlation

between the TC-TRN-TC openness coefficient and propagation scores (normalized regression coefficient

or NRC=1.000), with propagation better supported in networks with stronger open-loop TC-TRN-TC

pathways. By contrast, propagation tended to be modestly diminished as a function of increasing both

chemical and electrical TRN-TRN synaptic connectivity (NRC=-0.135 and NRC=-0.193, respectively).

Mechanistically, we interpreted this result to indicate that intrareticular synapses tended to interfere with

the spike timing necessary for linear signal propagation via the open-loop TC-TRN lattice through which

they primarily traveled. Thus, the homogeneously synaptic network permutations that best accommodated

signal propagation were generally ones with weak or absent synapses between TRN neurons and strong

open-loop TC-TRN-TC connections. A second-order multiple regression model of propagation as a

function of all three synaptic class variables ($R^2$=0.863, RMSE=0.039, $p$<0.0001) revealed a significant

but small negative interaction term between TRN-TRN$_{Elec}$ synapses and TC-TRN-TC openness (NRC=-

0.239), indicating that in networks where both electrical synapses were strong and TC-TRN-TC openness

high, the extent of supported propagation diminished nonlinearly. A positive interaction of comparable

magnitude between chemical and electrical intrareticular synapses (NRC=0.226) tempered the negative

linear effects that these two synaptic classes exerted on propagation in networks where TRN-TRN

interaction was strong. No other non-negligible interaction terms with NRCs were revealed by the

second-order regression model, and all the terms therein were qualitatively consistent with those in the

linear regression model (Table 3). Moreover, the relative magnitude of the coefficients in the second-

order regression model indicated that linear synaptic effects tended to dominate over nonlinear effects

across the domains of studied synaptic weights. The mean propagation score for all networks was 0.648 ±

0.006, with scores ranging from 0.435 to 0.923.

Two network permutations exhibiting fully open-loop TC-TRN-TC connectivity (openness coefficient or

OC=1.0), in which propagation was generally robust, were selected for further analysis (these networks

are labeled in Figs. 4A, top, and 4B, left). Each network was indexed by its synaptic variables ([TRN-

TRN$_{GABA}$ conductance (in nS), TRN-TRN$_{Elec}$ coupling coefficient, TC-TRN openness coefficient]):

Network a ([0,0,1.0]) and Network b ([450,0.36,1.0]). The propagation score of Network a, 0.913, was the

2$^{nd}$ highest of the 770 ranked networks, with Network b scoring 0.757 (89$^{th}$). Consistent with regression

analysis, the extent of response propagation in fully open-loop networks (i.e., OC=1.0) was maximal

when TRN-TRN synapses, both chemical and electrical, were absent, with increasing electrical coupling

between TRN neurons attenuating propagation more rapidly than increasing GABAergic projection

strength (Fig. 4B, left). Voltage traces of all network neurons depict activity, both preceding and

following fixed, punctate external stimulation of $TC_A$ (yellow arrows), during representative 1.0-s

simulations of Networks a and b (Fig. 4C, left and center). Both isolated spikes and bursts, defined as at

least three spikes occurring in rapid succession, were observed intermittently in neurons found in all three

network layers. In Network a, stimulus-evoked activity tended to propagate smoothly from $TC_A$ to $TC_C$;

near-synchronous propagation cascades were elicited in both the TRN and L4 layers of the model, having

been stimulated by propagating activity in upstream TC neurons. Smooth, linear propagation of action

potentials across the network depended on the synchronous induction of inhibitory postsynaptic potentials

(IPSPs) and the ensuing post-inhibitory rebound spikes in TC neurons, which occurred reliably and at

fixed intervals in Network a due to both its fully open-loop TC-TRN-TC connectivity and the absence of

any synapses within the network's TRN layer. Relative to Network a, Network b was not capable of

supporting activity propagation with the same fidelity, as predicted by the regression analyses.

Mechanistically, this was attributable to both the presence of strong TRN-$TRN_{GABA}$ synaptic connections,

which reduced the incidence of IPSPs in TC neurons required for signal propagation across the network,

and Network b's strong electrical coupling between TRN neurons, which destructively shunted a

propagating signal away from the thalamoreticular lattice through which it predominantly traversed the

network.

Propagation and oscillation scores across all 770 homogeneous networks, as driven by a fixed, punctate

stimulus, were strongly anticorrelated (Pearson's $r$=-0.739, $p$<0.0001). Accordingly, oscillation was best

accommodated in network permutations exhibiting strongly closed-loop connectivity (Fig. 4A, bottom),

as confirmed quantitatively through a linear regression model in which TC-TRN-TC openness was

associated with an NRC of -1.000 ($R^2$=0.661, RMSE=0.137, $p$<0.0001). This analysis furthermore

demonstrated that both varieties of TRN-TRN synapse played nearly negligible roles in mediating or

otherwise modulating oscillation, yielding NRCs of 0.058 and 0.064 for TRN-$TRN_{GABA}$ and TRN-

$TRN_{Elec}$ synapses, respectively. The insignificant relationship between oscillation and the weights of

intrareticular synapses was further reflected by the largely unpatterned distribution of oscillation scores in

TRN-$TRN_{GABA}$ x TRN-$TRN_{Elec}$ phase space for the 70 fully closed-loop network variants (Fig. 4B, right).

No significant interactions or nonlinearities were disclosed through a second-order regression model

relating oscillation to the three variable synaptic classes; indeed, this model was fully consistent with and

only marginally more explanatory than its linear counterpart ($R^2$=0.687, RMSE=0.137, $p$<0.0001; Table

3). The mean oscillation score across network variants in this set of simulations was $0.300 \pm 0.006$, with

scores ranging from 0.012 to 0.758.

Although oscillation was observed in homogeneously synaptic networks driven by a fixed, punctate

stimulus to $TC_A$, any oscillation in $L4_C$ was judged to be generally unrelated to the fixed stimulus;

equivalently stated, stimulus-induced oscillatory activity induced in $L4_A$ did not propagate well through

the network to $L4_C$. This deduction followed from the facts that 1) networks that supported robust

oscillation (strongly closed-loop networks) did not facilitate strong signal propagation, and in particular,

the propagation of oscillatory activity from one end of the network to the other and 2) the remaining

synaptic motifs connecting adjacent columns in the network and potentially serving as substrates for

signal propagation, namely those featuring a central intrareticular synapse, were demonstrated to exert

generally negligible or weakly attenuating effects on network propagation and oscillation. In absence of

stimulus-driven oscillatory activity propagating through the network, the default and indeed predominant

mechanism by which oscillation arose in $L4_C$ was through post-inhibitory rebound in $TC_C$, as engendered

by the strong recurrent inhibition found in network permutations exhibiting primarily closed-loop TC-

TRN-TC connectivity. This mechanism of oscillation was exemplified by Network c ([50, 0.18, 0]),

which received an oscillation score of 0.690 (7th highest of 770 network permutations): as illustrated in a

representative simulation of this network (Fig. 4C, right), an initial action potential in $TC_C$, elicited

through a stochastic external input and notably in the absence of upstream oscillatory activity propagating

through the network, drove recurrent inhibition in this neuron via $TRN_C$, causing oscillation in $TC_C$ over

several cycles and consequently driving concurrent oscillation in $L4_C$.

Based on the nature of oscillation in homogeneously synaptic network variants responding to a fixed,

punctate stimulus, we speculate that the rate of network oscillation corresponded to the kinetics of the t-

current, which underlies PIR. Across these simulations, oscillatory responses recurred linearly at mean

intervals of $109.27 \pm 0.31$ ms, or equivalently, at a mean frequency of $9.15 \pm 0.02$ Hz.

*Response of homogeneously synaptic models to a fixed, sustained stimulus*

As was the case when homogenously synaptic networks responded to a punctate external stimulus,

homogeneous network responses to a fixed, sustained stimulus propagated in a linear manner, recurring in

adjacent TC/L4 neurons at averaged fixed intervals of $99.96 \pm 0.69$ ms and propagating at a velocity of

0.50 mm/s. All networks variants driven by a fixed, sustained external stimulus to $TC_A$ were ranked

separately according to their scores for each network property (Fig. 5A, top). Network permutations that

allowed for robust propagative responses when $TC_A$ was stimulated in a punctate manner similarly

supported such responses when external stimulation to $TC_A$ was sustained (linear regression, $R^2$=0.793,

RMSE=0.047, $p$<0.0001). Propagation scores rose with increasing TC-TRN-TC openness (NRC=1.000),

while GABAergic and electrical TRN-TRN synapses both tended to exert moderate attenuating effects on

network propagation (NRC=-0.173 and NRC=-0.136, respectively). A second-order regression model

($R^2$=0.842, RMSE=0.041, $p$<0.0001) disclosed qualitatively similar synaptic interactions when

homogeneous networks were stimulated in a sustained manner as when they were driven by a punctate

stimulus (Table 4). The one exception to this was an additional though small negative interaction between

TRN-TRN$_{GABA}$ synapses and TC-TRN-TC openness (NRC=-0.152). Both this negative interaction and

the one between TRN-TRN$_{Elec}$ and TC-TRN-TC openness indicated that propagation was more

significantly affected by connections in the TRN layer as a function of increasing open-loop TC-TRN-TC

architecture; this is evident in Fig. 5B (left), as propagation scores conspicuously decrease in network

variants with an OC of 1.0 as either chemical or electrical synapses increase in weight. The mean

propagation score arising from sustained stimulation to $TC_A$ was $0.633 \pm 0.006$, with scores ranging from

0.414 to 0.961; these scores were not significantly different than those associated with fixed, punctate

stimulation of the same network permutations [paired $t$-test, $t(769)=1.067$, $p=0.287$].

A simulation of Network d ([0,0.06,1]) illustrates typical signal propagation in a network permutation

exhibiting high open-loop TC-TRN-TC connectivity and weak intrareticular synapses responding to a

fixed, sustained stimulus delivered to $TC_A$ (Fig. 5C, left); its position as a function of propagation score is

indicated in Figs. 5A (top) and 5B (left). The capacity of Network d to accommodate response

propagation throughout the length of the network in response to sustained stimulation was

mechanistically comparable to the similarly constituted Network a in the presence of punctate stimulation

(see Fig. 4C), with the former earning a propagation score of 0.961, the highest score among all

homogeneous network permutations exhibiting propagation as a function of fixed, sustained external

stimulation.

Oscillatory responses in networks when subjected to fixed, sustained stimulation recurred linearly in $L4_C$

at mean intervals of $110.26 \pm 0.25$ ms ($9.07 \pm 0.02$ Hz). Propagation and oscillation scores across

networks were similarly anticorrelated ($r$=-0.671, $p<0.0001$), though not as strongly as when $TC_A$ was

stimulated in a fixed, punctate manner. While network permutations with stronger closed-loop TC-TRN-

TC architectures were most permissive of cortical oscillation, this relationship was neither markedly

linear nor monotonically decreasing as a function of increasing openness coefficient (linear regression,

$R^2$=0.526, RMSE=0.145, $p<0.0001$; Table 4). Rather, a one-way ANOVA with Tukey's tests revealed

that, on average, oscillation scores peaked and remained statistically indistinguishable from one another

across the subset of network permutations with OCs between 0 and 0.4, with scores then decreasing in a

roughly linear fashion with increasing TC-TRN-TC openness (Fig. 5D; $F(10,759)=137.8$, $p<0.0001$). This

trend was partially accounted for in a second-order regression model ($R^2$=0.630, RMSE=0.129,

$p<0.0001$), in which the quadratic term in TC-TRN-TC openness had an NRC of -1.000 and was thus the

major determinant of network oscillation (all other synaptic variables in this model possessed NRCs with

absolute values less than 0.06). As with oscillation under the previous fixed stimulation condition, the

inconsequential roles played collectively by TRN-TRN synapses in oscillation here were reflected by the

haphazard distribution of scores in intrareticular synaptic parameter space (Fig. 5B, middle and right).

The mean oscillation score of homogeneously synaptic networks subjected to fixed, sustained stimulation

was $0.369 \pm 0.006$ (scores ranged from 0.006 to 0.827), with networks exhibiting a greater oscillation

score under this fixed stimulation condition than when stimulated in fixed, punctate manner [paired *t*-test,

$t(769)=-13.345, p<0.0001$].

We surmise that the discrepancy in network oscillation scores between homogeneous network variants

excited in punctate and sustained manners owes to a mode of oscillation that, though still infrequently

observed, became more prevalent in the latter set of simulations. Network permutations with somewhat

stronger closed- than open-loop TC-TRN-TC connectivity (OCs=0.3 and 0.4), such as Network e

([100,0.12,0.4]; oscillation score, 0.662; rank, 21$^{st}$ of 770), were intermittently able to support the

propagation of oscillatory activity from $TC_A$ to $TC_C$, and thus between their downstream cortical

counterparts, by virtue of 1) receiving sustained external input through $TC_A$ in a manner sufficient to elicit

linearly recurring action potentials and 2) possessing sufficiently strong laterally inhibitory TRN-TC

synapses to support propagation (Fig. 5C, middle). As previously mentioned, this mechanism for

generating oscillation in $L4_C$ was comparatively rarer when the same subset of networks was stimulated

in a fixed, punctate manner, as oscillation in $TC_A$ could not often be sustained without concurrent, high-

frequency external stimulation. Network f ([450,0.36,0.2]) typified the still-dominant mode of oscillation

exhibited in homogeneous networks stimulated in a punctate manner, namely that oscillation observed in

strongly closed-loop networks deriving from PIR in $TC_C$ (Fig. 5C, right). In the selected simulation of this

network, oscillatory activity was enabled by a single epoch of signal propagation, although Poisson-

mediated external input to $TC_C$ remained the more common catalyst for this mode of oscillation. Notably,

neither the presence of strong GABAergic nor electrical intrareticular synapses in Network f exerted

much effect on its ability to support oscillation (score, 0.726, 4[th] of 770), as predicted by the regression

models.

*Response of heterogeneously synaptic models to a fixed, sustained stimulus*

Given both the number of individually varying synaptic parameters within heterogeneously synaptic

networks (14, vs. 3 in the homogeneous networks) and the large number of heterogeneously synaptic

network permutations simulated in this study (~12,600), we relied solely on regression analysis to

ascertain the ultimate relationship between individual synaptic variables and network properties of

interest; neither ranking network variants in order of performance nor examining specific parameter

subspaces, as was done in the case of homogeneous networks, was practical. Accordingly, we constructed

circuit-level schematics of linear regression models for propagation (Fig. 6A) and oscillation (Fig. 6B) as

functions of the 14 synaptic variables in heterogeneous networks. While second-order regression models

for both performance metrics were also generated, these models revealed only a small number of

significant quadratic or interaction terms that could not more parsimoniously be accounted for by single

synaptic variables in the corresponding linear regressions. We note any nonlinearities of interest here and

report all regression results in Table 5.

The extent of stimulus-evoked response propagation in those heterogeneously synaptic networks

generated increased chiefly as a function of increasing the strength of the more downstream of the two

laterally inhibitory TRN-TC synapses, $TRN_B \rightarrow TC_C$: the corresponding term in a linear regression model

of propagation ($R^2$=0.742, RMSE=0.069, $p$<0.0001) possessed an NRC of 1.000 (Fig. 6A). Propagation

scores, which averaged at $0.601 \pm 0.001$ and ranged from 0.299 to 1.000, also scaled to a lesser extent

with the more upstream laterally inhibitory reticulothalamic synapse, $TRN_A \rightarrow TC_B$ (NRC=0.608). The

two inhibitory intrareticular synapses originating at the rightmost end of the model network,

$TRN_C \rightarrow TRN_A$ and $TRN_C \rightarrow TRN_B$, both exerted a small negative effect on propagation (NRC=-0.087 and

NRC=-0.084, respectively). Additionally, two TRN-TRN$_{Elec}$ synapses, $TRN_A = TRN_B$ and $TRN_A = TRN_C$

(where the "=" denotes an electrical synapses), marginally decremented propagation in heterogeneous

networks, with NRCs of -0.051 and -0.072, respectively. These findings clarified at an individual synaptic

level the observation that strong TRN-TRN interactions, whether chemical or electrical, tended to impede

signal propagation in homogeneous network variants. The three recurrently inhibitory synapses also

decremented propagation efficiency to minor degrees, with $TRN_C \rightarrow TC_C$, which stood to interfere directly

with $TC_C$'s reception of a propagating signal, associated with the largest NRC of these three synapses (-

0.207; Table 5).

A second-order regression model ($R^2$=0.857, RMSE=0.051, $p$<0.0001) disclosed a large, propagation-

enhancing interaction between the two laterally inhibitory synapses (NRC=0.753), underscoring the same

dependence of propagation on strong open-loop TC-TRN-TC connectivity as seen in homogeneously

synaptic networks, but additionally demonstrating that propagation scores increased nonlinearly as a

function of simultaneously increasing the weights of $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$. Quadratic terms

elucidated by this model confirmed that nearly all of the synapses highlighted in the linear regression

analysis retained the same qualitative effects on propagation over the domain of their weights, despite the

magnitudes of those effects changing as a function of synaptic weight (Table 5); one exception to this was

$TRN_C \rightarrow TRN_B$, whose individual influence on propagation became negligibly positive at 450 nS.

Interactions between TRN-TRN synapses of either variety and TRN-TC synapses tended diminish

propagation, as did those between recurrent and lateral inhibitory TRN-TC synapses. Taken together, the

linear and 2° regression models indicated that heterogeneous network permutations with strong laterally

inhibitory TRN-TC synapses tended to best support propagation. We speculate that those synapses

tending to diminish propagation, albeit all to a relatively small degree, shared in common the capacity to

interfere with the precise spike timing required to propagate a signal continuously along the open-loop

TC-TRN-TC subnetwork from $L4_A$ to $L4_C$: in some cases, this interference would arise due to

uncorrelated TRN-mediated inhibition of TC neurons (polysynaptically in the cases of $TRN_A$=$TRN_B$ and

$TRN_A$=$TRN_c$ and monosynaptically for $TRN_A \rightarrow TC_A$, $TRN_B \rightarrow TC_B$, and $TRN_C \rightarrow TC_C$) or in the case of

$TRN_C \rightarrow TRN_A$ and $TRN_C \rightarrow TRN_B$, by disrupting the requisite TRN-driven PIR in the TC neurons.

Consistent response propagation across the length of the network was epitomized by Network a'

(propagation score=0.868), in which $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$ were both relatively strong and those

synapses impeding propagation relatively weak (Fig. 6C, left).

Across the heterogeneously synaptic network permutations generated and analyzed, the mean oscillation

score was $0.438 \pm 0.001$, with scores ranging between $4 \times 10^{-4}$ and 1; there was a very small negative

correlation between the propagation and oscillation scores of these networks ($r$=-0.0296, $p$=0.0008).

Although neither a linear ($R^2$=0.253, RMSE=0.131, $p$<0.0001) nor a second-order ($R^2$=0.388,

RMSE=0.118, $p$<0.0001) regression model accounted for a majority of the variability in oscillation scores

across surveyed heterogeneous networks, each analysis nevertheless illuminated different aspects of this

property's synaptic substrates. Linear regression analysis demonstrated that oscillation scores scaled

positively with all three recurrently inhibitory TRN-TC synapses (Fig. 6B), mirroring the dependence of

this network property on strong, global closed-loop TC-TRN-TC connectivity in homogeneously synaptic

network variants. That the most downstream of these synapses, $TRN_C \rightarrow TC_C$, was strongly correlated with

oscillation, with an NRC=1.000, corroborated the inferred role played by this synapse in generating what

was the dominant mode of oscillation for homogeneous networks. Network b' (oscillation score, 0.508)

was representative of a heterogeneous network variant in which this mechanism of oscillation was

prevalent (Fig. 6C, middle). Two intrareticular synapses, $TRN_A$-$TRN_C$ and $TRN_A$=$TRN_C$, tended to

contribute modestly to oscillation (NRCs of 0.115 and 0.117, respectively), apparently representing

alternate though nondominant pathways through which oscillatory activity might propagate from one end

of the network to the other or be directly induced through successive bouts of PIR-mediated excitation in

$TC_C$ via $TRN_C$ (see Fig. 1B, left and middle pathways). By contrast, the two laterally inhibitory

reticulothalamic synapses, $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$, tended to impede oscillation (NRCs of -0.289

and -0.379, respectively).

Although a 2° regression analysis confirmed that $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$ were, in their individual

capacities, strongly anticorrelated with oscillation in heterogeneous networks (NRCs of -1.000 and -

0.892, respectively), interaction terms in this model involving these two synapses and any of the three

recurrently inhibitory TRN-TC synapses ($TRN_A \rightarrow TC_A$, $TRN_B \rightarrow TC_B$, $TRN_C \rightarrow TC_C$) were moderately to

strongly positive, with NRCs ranging between 0.345 and 0.669 (Table 5). Furthermore, in network

permutations in which there was a continuum of strongly closed- and open-loop TC-TRN-TC motifs (i.e.,

networks in which both varieties of TRN-TC synapse were strongly expressed, allowing oscillatory

activity to propagate reliably towards $TC_C/L4_C$), the cumulative oscillation-facilitating effects of these

synaptic interactions superseded the attenuating, noninteractive effects of $TRN_A \rightarrow TC_B$ and $TRN_B \rightarrow TC_C$.

Indeed, some such heterogeneous networks, as typified by Network c' (oscillation score, 1.000; Fig. 6C,

right), supported degrees of oscillation that exceeded those of top-performing homogeneous networks,

whose synaptic uniformity precluded architectures in which both recurrently and laterally inhibitory

TRN-TC synapses were maximally weighted. Other interactions, such as those between $TRN_A \rightarrow TC_A$ and

several intrareticular synapses, modestly hindered oscillation, apparently by diminishing the probability

of periodically induced PIR in $TC_A$ via $TRN_A$ underlying oscillation in the former cell and the subsequent

downstream propagation of this oscillatory activity; this interference could arise either from aperiodic

inhibition of $TRN_A$ ($TRN_B \rightarrow TRN_A$ x $TRN_A \rightarrow TC_A$, NRC=-0.186; $TRN_C \rightarrow TRN_A$ x $TRN_A \rightarrow TC_A$, NRC=-

0.172) or through delaying/accelerating its spiking threshold via electrical coupling ($TRN_A = TRN_C$ x

$TRN_A \rightarrow TC_A$, NRC=-0.114). Although there were fewer heterogeneous network variants simulated (or

theoretically possible) that possessed a synaptic architecture capable of supporting the propagation of

oscillation than networks in which oscillation was borne of PIR-mediated oscillation via $TRN_C \rightarrow TC_C$, the

former mode of oscillation was more strongly correlated with high oscillation scores among these

networks, the opposite of what was observed in homogeneous networks: this was evidenced by the

relative magnitudes of the second-order regression NRC of $TRN_C \rightarrow TC_C$ (0.107) and those of the synaptic

interactions mediating propagation of oscillation.

*Comparisons of network properties across synaptic architecture and external stimulation groups*

We analyzed the relative capacities of homogeneously synaptic networks, whether stimulated externally

in a fixed, punctate or sustained manner, and heterogeneous synaptic networks (for which $TC_A$ was

stimulated in a fixed, sustained manner) to support propagation and oscillation by comparing the 20

highest scores achieved by homogeneous and heterogeneous network permutations with respect to each

performance metric. We decided against performing a direct comparison of mean performance scores

across the full sets of homogeneous network permutations and surveyed heterogeneous network variants

due to both unequal sample sizes (770 homogeneous vs. ~12,600 heterogeneous network permutations)

and incomplete sampling of the full heterogeneous synaptic parameter space (less than 0.001% of all

possible heterogeneously synaptic network variants were simulated in this study). As these caveats did

not apply to the homogeneous networks alone, we were nevertheless able to employ repeated-measures

comparisons of propagation and oscillation scores across the full set of 770 homogeneous network

variants as a function of fixed stimulation condition (punctate vs. sustained) to more directly gauge the

effect of different external stimulation durations on these network properties (see the *Response of*

*homogeneously synaptic models to a fixed, sustained stimulus* subsection).

A one-way ANOVA disclosed no significant differences in mean propagation scores between top-

performing network permutations across any of the three synaptic/stimulation groups [$F_{(2,57)}=0.84$,

$p=0.437$; Fig. 7]. We attributed this to the fact that synaptic regression analyses of both homogenously

and heterogeneously networks, regardless of the fixed, eternal stimulation condition imposed on the

former, predicted that similarly constituted homogeneous and heterogeneous network variants, namely

those with 1) fully open-loop TC-TRN-TC architectures (equivalently, those in which $TRN_A \rightarrow TC_B$ and

$TRN_B \rightarrow TC_C$ were maximally weighted) and 2) nonexistent intrareticular and recurrently inhibitory TRN-

TC synapses, would rank among the top signal propagators. In contrast to propagation, the top scorers

among heterogeneous network variants better supported oscillation than the top homogeneous performers,

regardless of fixed stimulation condition [one-way ANOVA, $F_{(2,57)}=166.14$, $p<0.0001$]. Pairwise mean

oscillation differences assessed through Tukey's tests between top-performing heterogeneous networks

and both groups of homogeneous networks were highly significant ($p<0.0001$), while there was no

significant difference in top homogeneous oscillation scores as a function of differing fixed, external

stimulation ($p$=0.153). We concluded that heterogeneously synaptic networks' superior capacity to

oscillate was most closely related to their greater relative ability to propagate oscillatory activity across

TC-TRN-L4 columns; as related in the preceding section, this mode of oscillation, when available,

yielded appreciably higher fidelity of oscillation than afforded by the effect of PIR-driven oscillation in

$TC_C$, itself predominant in homogeneous networks.

**Discussion:**

The data generated and analyzed in the present study confirmed our central hypothesis, that open-loop

TC-TRN-TC synaptic motifs (Fig. 1B, right) could function as a substrate for signal propagation within

the thalamus. Propagation scores were more strongly dependent on the more downstream of the two

laterally inhibitory reticulothalamic synapses, $TRN_B \rightarrow TC_C$, in heterogeneously synaptic networks. By

contrast, not only was propagation poorly supported through intrareticular synapses, both chemical or

electrical (Fig. 1B, left and middle, respectively), these pathways in fact generally interfered with the

spike timing underlying propagation across the TC-TRN subnetwork; for the same reason, recurrently

inhibitory TRN-TC synapses, which served no role in mediating signal propagation, also tended to

diminish networks' capacity to support this property. Changing the duration of the generalized external

stimulus delivered to networks from punctate (50 ms) to sustained (1,100 ms) did not significantly affect

propagation scores across the set of homogeneously synaptic network permutations, nor were there

differences observed between homogeneous and heterogeneous networks accommodating high degrees of

propagation.

Oscillation in model networks typically arose through one of two mechanisms: the first, which was both

more prevalent and strongly correlated with oscillation scores in homogeneous networks, relied on $TRN_C$-

mediated post-inhibitory rebound in $TC_C$ that recurred at periodic intervals and could be sustained over

several hundred milliseconds. As such, this mode of oscillation was observed in networks exhibiting a

strongly weighted $TRN_C \rightarrow TC_C$ synapse, whether individually, as in heterogeneous network variants, or as

part of homogeneous networks with a uniformly strong closed-loop TC-TRN-TC architecture. This

mechanism could be elicited either indirectly through stimulation of the network upstream of $TC_C$,

including as a consequence of the fixed, external stimulus delivered to $TC_A$, or directly through

spontaneous external stimulation of the former neuron. Comparatively more robust oscillation, persisting

over a greater number of cycles and consequently associated with higher oscillation scores, was observed

as a result of the other mode of oscillation, which entailed the propagation of oscillatory activity, as

induced through sustained stimulation of $TC_A$, through a network. This oscillation mechanism was most

likely to occur in networks exhibiting simultaneously strong closed- and open-loop TC-TRN-TC motifs

and was therefore more prevalent in heterogeneous network permutations, where it was moreover more

strongly determinative of oscillation scores than the other mode of oscillation. For this reason, the mean

oscillation score of top-performing heterogeneous networks was significantly higher than those of their

homogeneous counterparts; (across the full set of homogeneously synaptic networks, fixed, sustained

stimulation slightly enhanced oscillation scores over those attained through punctate stimulation).

Synaptic interactions that impeded the latter mode of oscillation typically interfered with the induction of

oscillatory activity through PIR in $TRN_A \rightarrow TC_A$ by disrupting the temporal dynamics of this process and

precluding this activity from propagating downstream. A third, relatively weak and infrequently observed

form of oscillation also involved oscillatory activity propagating from one end of the network to the other

by way of $TRN_A$-$TRN_C$ and $TRN_A$=$TRN_C$.

*Methodological considerations of the study*

In an effort to maximize the predictive potential of our neural model relative to computational demands

that can increase exponentially with increasing degrees of freedom, we relied in some instances on

simplifications and/or idealizations, while in other cases, deliberately omitted experimentally verified

biophysical and neurophysiological details of the thalamo-reticulo-cortical networks being simulated and

their individual neuronal constituents. Such details can, in principle, can be selectively added to future

iterations of this model to explore questions and phenomena related to those posed and examined in the

present study.

Two points should be made regarding the constitution of the model neurons used in our study. First, like

most of the thalamic (Destexhe et al., 1993; Destexhe et al., 1994; Golomb et al., 1996; Destexhe et al.,

1996a; Sohal and Huguenard, 1998; Bazhenov et al., 1998) and thalamocortical models (Destexhe et al.,

1998; Bazhenov et al., 2002; Rogala et al., 2013) that inspired our model, we utilized single-

compartment, Hodgkin-Huxley neurons. While these model cells contribute to the computational

parsimony and practicality of network models, particularly where the analysis of network dynamics is

prioritized, they neglect the intrinsic cable properties of real neurons and, relatedly, the spatially disparate

nature of synaptic integration and heterogeneous expression of intrinsic and synaptic conductances

(Dayan and Abbott, 2005; Herz et al., 2006). Such considerations are particularly relevant here relative to

dendritic distributions of t- current and h-current in TC neurons (McCormick and Pape, 1990; Destexhe et

al., 1998; Williams and Stuart, 2000; Traub et al., 2005) and TRN neurons (Contreras et al., 1993;

Destexhe et al., 1996b; Traub et al., 2005; Crandall et al., 2010). Multicompartment neuronal models

incorporating such details could conceivably alter the network dynamics being studied. Second, although

we allowed for heterogeneous connections within a given synaptic class for a subset of simulated network

permutations, all cells within each of the three network layers were modeled with identical intrinsic

parameters. In reality, even anatomically proximal populations of TC, TRN, and layer 4 cortical neurons

each exhibit a wide spectrum of different intrinsic properties (Leresche et al., 1991; Spreafico et al., 1991;

Lee et al., 2007; Landau et al., 2016), with the variability of these neurons' synaptic "footprints" in space

often well exceeding what we capture in our model (Cox et al., 1996; Cox et al., 1997; Pinault et al.,

1997). If larger-scale elaborations of the model are simulated in the future, both the cellular and spatial

synaptic variability observed experimentally could be approximated by allowing given parameters to vary

systematically or at random within physiological ranges, much as we did with TRN-TRN and TRN-TC

synaptic conductances in the present study; this would allow any such model to approach a more

representative degree of anatomical and functional heterogeneity.

In the interest of isolating the feedforward dynamics of thalamocortical transmission intrinsic to our

present model, certain synaptic connections to TC, TRN, and L4 neurons were either excluded or

generalized. First, while Poisson processes are often utilized to represent complex, noisy, and/or

generalized synaptic inputs (Dayan and Abbott, 2005), it is reasonable, given the coherence of activity

across anatomically disparate regions of the brain that prevails during sensory processing, that canonically

non-sensory brain regions projecting to the both the TRN and dorsal thalamus might exhibit activity

correlated with ascending sensory input, making the use of stochastic inputs to TC neurons of our model a

crucially simplifying but imprecise approximation. Similarly, although the high-frequency pulse trains we

employed as generalized external stimuli within the model belie the rich and variable stimulus encoding

schemes inherent to different sensory systems, these rudimentary stimulus approximations, in

combination with the baseline stochastic input, were nevertheless sufficient to elicit propagating and/or

oscillating waves in the model network and analyze their dynamics across network permutations,

reflecting the use of similar pulse trains to generate spindle-like waves in isolated thalamic slices and

analysis of how those waves were perturbed as a function of various pharmacological manipulations (Bal

et al., 1995; Kim et al., 1995). Although we chose not to emphasize the correlation between stimulus and

response in the present study, in keeping with similar thalamo-reticulo(-cortical) modeling studies (e.g.,

Destexhe et al., 1996a, Golomb et al., 1996; Sohal and Huguenard, 1998; Bazhenov et al., 1998, Traub et

al., 2005), future efforts using the present baseline network model or elaborations thereof might

investigate response dynamics as a function of varying spatial and/or temporal stimulus profiles (see, for

example, Pham and Haas, 2018). A more realistic rendering of stimulus representations in this modeling

paradigm might also account for the subset of inhibitory afferents projecting to parts of the thalamus,

including the lateral geniculate nuclei and medial geniculate bodies (Cucchiaro et al., 1991; Winer et al,

1996; Peruzzi et al., 1997; Llano et al., 2014).

Additionally, the present model omitted explicit corticothalamic and corticoreticular synapses, both of

which have been identified and physiologically characterized to varying degrees (Steriade et al., 1972;

White and Hersch, 1982; DeCurtis et al., 1989; Contreras et al., 1996; Blumenfeld and McCormick, 2000;

Zhang and Jones, 2004; Crandall et al., 2015), though the former were effectively amalgamated with both

feedforward sensory and modulatory projections to the thalamus in the form of the generalized, Poisson-

modulated external input we delivered to individual TC neurons. Both forms of feedback have been

implicated in the spread of spindle waves and in the maintenance of their synchronization over large

distance scales (on the order of the length of the mammalian forebrain) and are furthermore known to

drive spindle wave formation *in vivo* by polysynaptically recruiting TC neurons via TRN-mediated PIR

(Steriade et al., 1972; Roy et al., 1984; Contreras and Steriade, 1996; Contreras et al., 1996; Suga and Ma,

2003; Sillito et al., 2006; Crandall et al., 2015). It should be noted, however, that short-range coherence of

spindle waves, which can be elicited in isolated thalamic slice preparations (Bal et al., 1995; Kim et al.,

1995), is preserved following decortication, both *in vivo* and *in silico* (Contreras and Steriade, 1996;

Contreras et al., 1996; Destexhe et al., 1998). By extension, it is reasonable to assume that the dynamics

of the spindle-like waveforms generated in our small-scale, broadly feedforward model, in which the

cortex served solely as an output layer, would not be qualitatively altered by corticothalamic or

corticoreticular feedback. An additional challenge in modeling descending cortical projections lies in

rendering the complex intracolumnar interactions between multiple cortical layers upstream of them:

several thalamocortical models incorporating corticothalamic and corticoreticular feedback limited

cortical representation to layer 6 neurons and/or fast-spiking interneurons (Destexhe et al., 1998;

Bazhenov et al., 2002; Rogala et al., 2013), while Traub et al. (2005) simulated six distinct cortical

populations within layers 2-6 as part of an expansive thalamocortical network model comprising 3,560

multicompartment neurons. Future efforts within our modeling paradigm stand to both expand the spatial

scale of the thalamocortical network and incorporate reciprocal cortical projections involving interactions

between multiple cortical layers.

Finally, external inputs to the TRN were also omitted from the model. Although it is recognized that the

TRN receives a variety of inputs from beyond the thalamus and sensory cortex (Asanuma and Porter,

1990; Bickford et al., 1994; Zikopoulos and Barbas, 2006; Sun et al., 2013; Pita-Almenar et al., 2014), an

early version of our model that included stochastic external inputs to TRN neurons mirroring those

delivered to the TC layer did not significantly alter network dynamics beyond marginally increasing

variability in those network properties studied.

*Comparison to related computational models and physiological data*

Although the production of spindle waves was not an explicit objective of our study, some of the wave

dynamics arising in our networks were nevertheless consistent with those inherent to spindle or spindle-

like waves, as observed empirically or generated in other modeling studies. Despite possessing higher

degrees of TC→TRN and TRN→TC synaptic divergence and lacking the exclusively open-loop TC-

TRN-TC architecture characterizing a subset of our network variants, other isolated thalamic models

allowing for longitudinal wave propagation similarly accommodated this propagation along the lattice of

interconnected TC and TRN neurons by way of laterally inhibitory TRN-TC synapses (Kim et al., 1995;

Destexhe et al., 1996a; Golomb et al., 1996; Bazhenov et al., 1998; Muller and Destexhe, 2012); at short

ranges, this mechanism of signal propagation also prevailed in large-scale thalamo-reticulo-cortical

models, while corticothalamics acted to propagate activity to more distal sites (Destexhe et al., 1998; see

Destexhe and Sejnowski, 2003, for a schematic illustrating short- and long-range thalamocortical wave

propagation). Comparably, recurrently inhibitory TRN-TC synapses have been documented to play a vital

role in the generation of oscillatory behavior in the thalamus (Steriade and Deschênes, 1984; von Krosigk

et al., 1993). The temporal parameters of propagating and oscillation signals in our model also matched

some of those previously reported: among homogeneously synaptic network permutations, the mean

velocity of signal propagation across the length of the network was approximately 0.50 mm/s, which was

consistent with the propagation velocity of spindle waves measured in a ferret thalamic slice and about

half as fast as values reported in computational models of this slice preparation (Kim et al., 1995; Golomb

et al., 1996; Destexhe et al., 1996a). Similarly, the roughly 9.1-Hz mean oscillation frequency measured

across homogeneous networks fell within the 6-12 Hz range of intraspindle spike frequencies reported in

both physiological and computational spindle wave studies (Andersen and Andersson, 1968; Steriade and

Deschênes, 1984; Kim et al., 1995; Golomb et al., 1996; Destexhe et al., 1996a).

Several key structural elements of our set of network models and the range of phenomenology they

produced distinguish them from previous thalamic and thalamocortical models. One particularly notable

point of departure relative to similar network models was the extent to which thalamoreticular,

reticulothalamic, and thalamocortical synapses diverged. Although all three classes of synapses are

known to diverge significantly and have been observed to target neuronal somata hundreds of microns

from their origins (Jones, 1985; Crabtree, 1996; Cox et al., 1996; Cox et al., 1997; Pinault and Deschênes,

1998; Sherman and Guillery, 2001; Alonso et al., 2001; Miller et al., 2001), the TC-TRN, TRN-TC, and

TC-L4 synapses in our model were constrained to remain strictly local and minimally divergent (or non-

divergent, in the case of TC-TRN and TC-L4 synapses). With respect to the first two classes of synapses,

this constraint was imposed in order to preserve the disynaptic TC-TRN-TC open-loop motifs

characterizing a subset of network permutations, which constituted one of the focuses of our study, and

analyze the signal propagation they may support. This neuroanatomical scheme contrasted with previous

computational models featuring parallel, interconnected thalamoreticular pathways, in which both TC and

TRN synapsed bidirectionally with several neighboring TRN and TC cells, respectively, within a radius

of several hundred microns (e.g., Golomb et al., 1996; Destexhe et al., 1996a; Destexhe et al., 1998; Sohal

and Huguenard, 1998; Sohal et al., 2000; Bazhenov et al., 1998; Bazhenov et al., 2002; Traub et al., 2005;

Izhikevich and Edelman, 2008).

The contrasting synaptic architectures between our network model and other isolated thalamic models

accounted for several important differences in the waveforms documented in this study and those

previously observed *in simulo*. First, the asymmetric distribution of the laterally inhibitory TRN-TC

synapses in our model generally restricted the propagation of signals to a single direction, whereas

bidirectional wave propagation was observed in several other network models marked by greater degrees

TC-TRN and TRN-TC synaptic divergence when these networks were externally stimulated in their

anatomical centers (Bazhenov et al., 1998; Sohal et al., 2000). Relatedly, whereas some percentage of the

stimulus-driven responses and spontaneous signals arising in Column A of our network tended to

dissipate before arriving at Column C due to either spike failure or extraneous network activity interfering

with the temporal dynamics required for uninterrupted propagation (see, for example, Fig. 3, in which

spikes associated with a single epoch of propagation that were tallied across simulations of the same

network diminished between $L4_A$ and $L4_C$), the presence of bidirectional, divergent thalamoreticular and

reticulothalamic synapses in other network models were associated with less dampened, longer-duration,

and in larger networks, longer-distance signal propagation, a property ostensibly attributable to the greater

interconnectedness between TC and TRN neurons throughout these networks and a resulting degeneracy

of signaling pathways (Destexhe et al., 1996a; Golomb et al., 1996; Bazhenov et al., 1998; Sohal and

Huguenard, 1998).

We emphasize, however, that the discrepancy in spatiotemporal signal coherence between our network

model and others was also likely a function of disparate network inputs: on the one hand, fixed external

input was applied periodically to certain networks (Destexhe et al., 1996a; Bazhenov et al., 1998; Rinzel

et al., 1998), rather than over a singular interval as in ours, or distributed over multiple neurons in a

spatially dependent manner (Bazhenov et al., 1998; Sohal et al., 2000; Pham and Haas, 2018), in contrast

to the application of fixed stimulation to a single neuron (in our model, $TC_A$). Arguably more integral to

the difference in signal integrity was the stochastic input received by neurons in the TC layer of our

model network: as detailed in our analysis, these inputs elicited sufficient noise to interfere intermittently

with efficient propagation and oscillation. Other thalamic and thalamocortical model networks lacked

these inputs, and indeed some remained fully quiescent in the absence of external stimulation (Destexhe

et al., 1993; Golomb et al., 1996; Rinzel et al., 1998; Bazhenov et al., 1998; Sohal et al., 2000; Traub et

al., 2005). We moreover ascribe the absence of the "waxing and waning" patterns of activation lasting

0.5-2 s in TC, TRN, and cortical cell populations that characterize spindle waves, as measured

experimentally in individual TC and TRN neurons (Steriade and Llinàs, 1988; Leresche et al., 1991;

Soltesz et al., 1991; von Krosigk et al., 1993), observed across neuronal population through

electroencephalography and local field potential recordings (Gibbs and Gibbs, 1950; Andersen and

Andersson, 1968; Steriade and Deschênes, 1984; Steriade et al., 1987), and generated in network models

aimed at reproducing the spindling phenomenon (Destexhe et al., 1993; Destexhe et al., 1994; Destexhe et

al., 1996a; Bazhenov et al. 1999), to the presence of stochastic activity in our model. Although the

thalamic relay neurons in our model expressed the h-current and, in a subset of network permutations, the

reciprocal TC-TRN connections thought to account for spindling, the phenomenon also presupposes a

sustained antiphase relationship between TC and TRN spiking over the course of an individual spindle

wave that was invariably disrupted by the stochastic input to our network (Destexhe et al., 1993; Bal and

McCormick, 1996; Luthi and McCormick, 1998; Timofeev et al., 2001). That spindle waves still occur *in*

*vivo* during sleep despite a myriad of inputs impinging on the dorsal thalamus and TRN that would tend

to perturb the intrinsic and synaptic conductances underlying the waveform suggests a countervailing

source of spindle maintenance: this role could conceivably be filled by corticothalamics, which have been

demonstrated to synchronize spindle-wave activity over large distances in the brain (Bal et al., 1995;

Contreras and Steriade, 1996; Timofeev et al., 2001; Bonjean et al., 2011).

We finally wish to comment on the finding that intrareticular synapses, both GABAergic and electrical,

manifested relatively little effect on the signal propagation and oscillation we observed in this study,

despite having been copiously studied, simulated, and demonstrated to modulate certain waveforms

arising in both the isolated thalamus and broader thalamocortical system (Ahlsén and Lindström, 1982;

Steriade et al., 1990; Sanchez-Vives et al., 1997; Sohal et al., 2000; Landisman et al., 2002; Long et al.,

2004; Fuentealba et al., 2004; Deleuze and Huguenard, 2006; Lam et al., 2006). To the extent that

GABAergic intrareticular synapses disrupted efficient signal propagation along the length of our

networks, this could conceivably lend credence to the contention of Deleuze and Huguenard (2006), who,

based on their observation that TRN-TRN$_{GABA}$ synapses were more densely distributed along the

dorsoventral (vertical) axis of the TRN, posited that these synapses may selectively filter sensory input as

it traverses the thalamus to reach the cortex and play a less consequential role with respect to horizontal

(intrathalamic) signal propagation. Notwithstanding the potentially paradigm-shifting question as to

whether TRN-TRN$_{GABA}$ synapses are altogether absent in certain mammals or degenerate as a function of

increasing age (Landisman et al., 2002; Cruikshank et al., 2010; Hou et al., 2016), it has moreover been

shown in thalamic slice preparations derived from juvenile rodents, with supporting computational

models, that pharmacologically blocking or knocking out the expression of intrareticular GABAergic

synapses promotes large-scale synchronization of ~3-Hz spike-wave discharges in the thalamus that

resemble absence seizures; TRN-TRN$_{GABA}$ synapses are thus thought to desynchronize activity in a way

that preserves normal thalamic function while staving off epileptiform patterns in affected animals (von

Krosigk et al., 1993; Bal et al., 1995; Huntsman et al., 1999; Sohal et al., 2000; Sohal and Huguenard,

2003; Traub et al., 2005). Although intrareticular synapses were shown in the present study to modestly

dampen propagation through open-loop TC-TRN-TC architectures, both the relative lack of synaptic

divergence, as discussed above, and the limited scale of our model network would have likely precluded

any meaningful observations of neuronal synchrony or lack thereof as a function of TRN-TRN$_{GABA}$

weighting. It should additionally be stated that reticulothalamic signaling mediated through GABA$_B$

receptors is thought to contribute to the sustained hyperpolarization of thalamic relay neurons observed in

absence-like waveforms, and these receptors were not included in our model (von Krosigk et al., 1993;

Bal et al., 1995; Pinault et al., 1998; Crunelli and Leresche, 2002; Sohal and Huguenard, 2003).

As was the case with chemical synapses in the TRN, TRN-TRN$_{Elec}$ synapses did not mediate signal

propagation and supported comparatively weak forms of oscillation in a minority of network

permutations. With respect to the former property, our results were consistent with predictions that follow

from two sets of physiological observations made on these synapses: 1) across three separate studies,

electrical postsynaptic potentials or "spikelets," from which electrical coupling can be inferred, were

observed in less than 50% of observed TRN neuron pairs [Landisman et al., 2002; Deleuze and

Huguenard, 2006; Lam et al., 2006; Lee et al., 2014; note that Long et al. (2004) reported this figure at

71% but speculated that the discrepancy owed to recording from cell pairs with <5 μm edge-edge

separation]; and 2) mean electrical coupling coefficients between TRN neurons were small [0.032, as

reported by Landisman et al. (2002); ~0.02, Fuentealba et al. (2004); 0.11, as measured in closely apposed

cells by Long et al. (2004)]. These data suggest that 1) TRN-TRN$_{Elec}$ synapses are too sparsely distributed

to facilitate uninterrupted signal propagation within the TRN over large scales and 2) electrical coupling

between TRN neurons, even between those extremely closely situated to one another, is typically

insufficient to allow action potentials in one neuron to induce concurrent action potentials in its coupled

partner. Indeed, we only observed direct electrical induction of action potentials in those network

permutations exhibiting the highest degree of electrical coupling [0.36, close to the maximum coupling

strength observed by Landisman et al. (2002) and Long et al. (2004); see the middle panel of Fig. 4C for

an example of this phenomenon]. Nevertheless, gap junctions found between TRN neurons have been

demonstrated to play important roles vis-à-vis thalamic signaling: in keeping with their ability to act as

low-pass filters, TRN-TRN$_{Elec}$ synapses have been documented to promote thalamocortical

synchronization of subthreshold activity and low-frequency oscillations (<10 Hz), including components

of spindle waves (Long et al., 2004; Fuentealba et al., 2004; Traub et al., 2005), and can conceivably

facilitate the temporal discrimination of multiple near-coincident sensory stimuli by the cortex (Pham and

Haas, 2018). Furthermore, it has been suggested, based on the recent discovery of activity-dependent

plasticity of electrical coupling strength in brain regions including the TRN (Haas et al., 2011; Haas and

Landisman, 2012; Wang et al., 2015; Kohmann et al., 2016), that gap junction plasticity in the TRN might

itself underlie the generation of spindle waves (Pernelle et al., 2017). It remains an outstanding question

and one easily amenable to future investigation through computational modeling whether open-loop TC-

TRN-TC architectures, either independently of or in cooperation with intrareticular synapses, modulate

neuronal synchrony.

*The function and scale of open-loop thalamo-reticulo-thalamic synaptic motifs*

It is essential to contextualize the predictive potential of our model and the broader functions conceivably

served by open-loop TC-TRN-TC architectures. As discussed, the networks designed for this study did

not and were not intended to produce the gamut of phenomenology documented in the thalamocortical

system. Rather, they function as idealized network modules that could hypothetically exist within

surrounding neuronal populations. Open-loop TC-TRN-TC configurations have thus far been

observed both within and across individual thalamic nuclei (both first- and higher-order) and are thought

to serve as pathways for intra- and cross-modal modulation, respectively (Crabtree et al., 1998; Pinault

and Deschênes, 1998; Crabtree and Isaac, 2002; Lam and Sherman, 2005; Lee et al., 2010; Kimura et al.,

2007; Kimura, 2014; Lam and Sherman, 2015). However, despite available data on the distances over

which TRN neurons may project to synapse with TC cells (Cox et al., 1996), the proportion of recurrently

to laterally inhibitory TRN-TC synapses in any thalamic nucleus and the spatial homogeneity of these

connections remain unknown, and thus the density and distribution of open-loop TC-TRN-TC

architectures are questions as yet unanswered; the full extent of open-loop connections between

thalamic nuclei is similarly indeterminate. As such, our small network model could be easily rescaled to

approximate the physiological dimensions of these connections as such information becomes available,

adjusting synaptic delays accordingly.

Although the anatomical bases of open-loop TC-TRN-TC motifs have been partially characterized,

their functional significance in the brain lingers as a subject of continued speculation. As has been

previously surmised, these synaptic pathways could plausibly lend themselves to sensory enhancement,

multisensory integration, and attentional mechanisms (Crabtree and Isaac, 2002; Pinault, 2004; Willis et

al., 2015; Crabtree, 2018). Whereas a full elucidation of such phenomena vis-à-vis thalamoreticular

connectivity lies well beyond the scope of this study and will likely depend upon a far more nuanced

understanding of how perceptual processes are born out of interplay between the cortex, dorsal thalamus,

TRN, and likely other brain structures, our model can nevertheless offer more fundamental

predictions about the capacity of open-loop TC-TRN-TC configurations to accommodate basic signaling

properties within the thalamus. First, as inferred from physiological studies, open-loop pathways should

be fully capable of supporting signaling propagation from one thalamic relay neuron to another through a

limited number of intervening synapses (with a disynaptic pathway serving as the shortest such

configuration), whether or not the target neuron lies in the same thalamic nucleus. Secondly, a localized

thalamic region consisting exclusively of interconnected open-loop TC-TRN-TC motifs (homogeneously

synaptic) or a sequence of spatially separated, interconnected laterally inhibitory TRN-TC synapses,

should either exist, might be capable of supporting a non-periodic signal (e.g., a transient sensory-evoked

signal) propagating between multiple thalamic relay neurons, subject to the nature of the signal, the

input(s) generating it, and potentially limited by the degree of spontaneous (decohering) activity along the

signaling pathway. It is unclear how the fidelity of a transient signal in either scenario would compare to

that measured along a more densely interconnected segment of thalamus characterized by greater

thalamoreticular and reticulothalamic divergence, as traditionally conceived of in earlier modeling studies

(Destexhe et al., 1996a; Golomb et al., 1996; Bazhenov et al., 1998) and in which a signal might feed

back on itself; this is despite the fact that the latter architectures are capable of accommodating sustained,

periodic waveforms such as sleep spindles. Thirdly, discrete thalamic regions marked by synaptic

heterogeneity should be equally capable of propagating signals as those characterized by TC-TRN and

TRN-TC synaptic uniformity, after accounting for differences in the average weights or spatial densities of

laterally inhibitory TRN-TC synapses. Indeed, if the morphological, intrinsic, and synaptic heterogeneity

of TRN neurons are any indication (Schiebel and Schiebel, 1966; Jones, 1975; Spreafico et al., 1988;

Spreafico et al., 1991; Cox et al., 1996; Brunton and Charpak, 1997; Lee et al., 2007; Bickford, 2016), it

is reasonable to assume that both thalamorectiular and reticulothalamic synapses are typically distributed

in a heterogeneous manner across the thalamus. Lastly, unless driven by an external oscillatory source,

whether ascending or descending, oscillatory waveforms should not subsist in a localized thalamic region

populated exclusively by open-loop connections. Both forthcoming physiological investigation and future

modeling studies will undoubtedly be able to evaluate some of these predictions.

1025

**Acknowledgments:**

This work made use of the Illinois Campus Cluster, a computing resource that is operated by the Illinois

Campus Cluster Program (ICCP) in conjunction with the National Center for Supercomputing

Applications (NCSA) and which is supported by funds from the University of Illinois at Urbana-

Champaign. We additionally thank Profs. Thomas Anastasio and Ruoqing Zhu, Drs. Rama Ratnam and

Baher Ibrahim, and Mr. Weddie Jackson for their valuable analytical, statistical, and technical insights.

1032

1033 **References:**

1034    1.     Smith PH, Bartlett EL, Kowalkowski A (2006) Unique combination of anatomy and
1035               physiology in cells of the rat paralaminar thalamic nuclei adjacent to the medial geniculate
1036               body. *J Comp Neurol* 496(3):314-334.

1037
1038    2.     Lee SC, Cruikshank SJ, Connors BW (2010) Electrical and chemical synapses between relay
1039         neurons in developing thalamus. *J Physiol* 588(13):2403-2415.
1040
1041    3.     Guillery SW, Sherman SM (2002) Thalamic relay functions and their role in corticocortical
1042         communication: generalizations from the visual system. *Neuron* 33(2):163-175.
1043
1044    4.     Sherman SM (2004) Interneurons and triadic circuitry of the thalamus. *Trends Neurosci*
1045         27(11):670-675.
1046
1047    5.     Pinault D (2004) The thalamic reticular nucleus: structure, function and concept. *Brain Res*
1048         *Brain Res Rev* 46(1):1-31.
1049
1050    6.     Crick F (1984) Function of the thalamic reticular complex: the searchlight hypothesis. *Proc*
1051         *Natl Acad Sci USA* 81(14):4586-4590.
1052
1053    7.     Guillery RW, Feig SL, Lozsádi DA (1998) Paying attention to the thalamic reticular nucleus.
1054         *Trends Neurosci* 21(1):28-32.
1055
1056    8.     McAlonan K, Cavanaugh J, Wurtz RH (2006) Attentional modulation of thalamic reticular
1057         neurons. *J Neurosci* 26(16):4444-4450.
1058
1059    9.     Llinás RR, Paré D (1991) Of dreaming and wakefulness. *Neuroscience* 44(3):521-535.
1060
1061    10.    Steriade M, McCormick D, Sejnowski T (1993) Thalamocortical oscillations in the sleeping
1062         and aroused brain. *Science* 262(5134):679-685.
1063
1064    11.    von Krosigk, M, Bal T, McCormick DA (1993) Cellular mechanisms of a synchronized
1065         oscillation in the thalamus. *Science* 261(5119):361-364.
1066
1067    12.    Bal T, von Krosigk M, McCormick DA (1995) Role of the ferret perigeniculate nucleus in the
1068         generation of synchronized oscillations in vitro. *J Physiol* 483(3):665-685.
1069
1070    13.    Destexhe A, Bal T, McCormick DA, Sejnowski TJ (1996) Ionic mechanisms underlying
1071         synchronized oscillations and propagating waves in a model of ferret thalamic slices. *J*
1072         *Neurophysiol* 76(3):2049-2070.
1073
1074    14.    Huguenard JR (1998) Neuronal circuitry of thalamocortical epilepsy and mechanisms of
1075         antiabsence drug action. *Adv Neurol* 79:991-999.
1076
1077    15.    McCormick DA, Contreras D (2001) On the cellular and network bases of epileptic seizures.
1078         *Annu Rev Physiol* 63:815-846.
1079
1080    16.    Wells MF, Wimmer RD, Schmitt LI, Feng G, Halassa MM (2016) Thalamic reticular
1081         impairment underlies attention deficit in Ptchd1(Y/-) mice. *Nature* 532(7597):58-63.
1082
1083    17.    Krol A, Wimmer RD, Halassa MM, Feng G (2018) Thalamic reticular dysfunction as a
1084         circuit endophenotype in neurodevelopmental disorders. *Neuron* 98(2):282-295.
1085
1086    18.    Sherman SM, Guillery RW (2001) Exploring the thalamus (Academic Press, San Diego).
1087
1088    19.    Steriade M, Deschênes M (1984) The thalamus as a neuronal oscillator. *Brain Res* 320(1):1-
1089         63.

20. Steriade M, Domich L, Oakson G, Deschênes M (1987) The deafferented reticular thalamic nucleus generates spindle rhythmicity. *J Neurophysiol* 57(1):260-273.

21. Kim U, Bal T, McCormick DA (1995) Spindle waves are propagating synchronized oscillations in the ferret LGNd in vitro. *J Neurophysiol* 74(3):1301-1323.

22. Ulrich D, Huguenard, JR (1997) GABA(A)-receptor-mediated rebound burst firing and burst shunting in thalamus. *J Neurophysiol* 78(3):1748-1751.

23. Hale PT, Sefton AJ, Bauer LA, Cottee LJ (1982) Interrelations of the rat's thalamic reticular and dorsal lateral geniculate nuclei. *Exp Brain Res* 45(1-2):217-229.

24. Warren RA, Agmon A, Jones EG (1994) Oscillatory synaptic interactions between ventroposterior and reticular neurons in mouse thalamus in vitro. *J Neurophysiol* 72(4):1993-2003.

25. Sherman SM, Guillery RW (1996) Functional organization of thalamocortical relays. *J Neurophysiol* 76(3):1367-1395.

26. Shosaku A (1986) Cross-correlation analysis of a recurrent inhibitory circuit in the rat thalamus. *J Neurophysiol* 55(5):1030-1043.

27. Lo FS, Sherman SM (1994) Feedback inhibition in the cat's lateral geniculate nucleus. *Exp Brain Res* 100(2):365-368.

28. Gentet LJ, Ulrich D (2003) Strong, reliable and precise synaptic connections between thalamic relay cells and neurones of the nucleus reticularis in juvenile rats. *J Physiol* 546(3):801-811.

29. Pinault D, Deschênes M (1998) Anatomical evidence for a mechanism of lateral inhibition in the rat thalamus. *Eur J Neurosci* 10(11):3462-3469.

30. FitzGibbon T, Solomon SG, Goodchild AK (2000) Distribution of calbindin, parvalbumin, and calretinin immunoreactivity in the reticular thalamic nucleus of the marmoset: evidence for a medial leaflet of incertal neurons. *Exp Neurol* 164(2):371-383.

31. Crabtree JW, Collingridge GL, Isaac JT (1998) A new intrathalamic pathway linking modality-related nuclei in the dorsal thalamus. *Nat Neurosci* 1(5):389-394.

32. Crabtree JW, Isaac JT (2002) New intrathalamic pathways allowing modality-related and cross-modality switching in the dorsal thalamus. *J Neurosci* 22(19):8754-8761.

33. Lam YW, Sherman SM (2005) Mapping by laser photostimulation of connections between the thalamic reticular and ventral posterior lateral nuclei in the rat. *J Neurophysiol* 94(4):2472–2483.

34. Lam YW, Sherman SM (2015) Functional topographic organization of the motor reticulothalamic pathway. *J Neurophysiol* 113(9):3090-3097.

35. Kimura A, Imbe H, Donishi T, Tamai Y (2007) Axonal projections of single auditory neurons in the thalamic reticular nucleus: implications for tonotopy-related gating function and cross-modal modulation. *Eur J Neurosci* 26(12):3524-3535.

1143

1144 36.    Kimura A (2014) Diverse subthreshold cross-modal sensory interactions in the thalamic
1145 reticular nucleus: implications for new pathways of cross-modal attentional gating function.
1146 *Eur J Neurosci* 39(9):1405-1418.

1147

1148 37.    Asanuma C, Porter LL (1990) Light and electron microscopic evidence for a GABAergic
1149 projection from the caudal basal forebrain to the thalamic reticular nucleus in rats. *J Comp*
1150 *Neurol* 302(1):159-172.

1151

1152 38.    Bickford ME, Günlük AE, Van Horn SC, Sherman SM (1994) GABAergic projection from
1153 the basal forebrain to the visual sector of the thalamic reticular nucleus in the cat. *J Comp*
1154 *Neurol* 348(4):481-510.

1155

1156 39.    Zikopoulos B, Barbas H (2006) Prefrontal projections to the thalamic reticular nucleus form a
1157 unique circuit for attentional mechanisms. *J Neurosci* 26(28):7348-7361.

1158

1159 40.    Sun YG, Pita-Almenar JD, Wu CS, Renger JJ, Uebele VN, Lu HC, Beierlein M
1160 (2013) Biphasic cholinergic synaptic transmission controls action potential activity in
1161 thalamic reticular nucleus neurons. *J Neurosci* 33(5):2048–2059.

1162

1163 41.    Pita-Almenar JD, Yu D, Lu HC, Beierlein M (2014) Mechanisms underlying
1164 desynchronization of cholinergic-evoked thalamic network activity. *J*
1165 *Neurosci* 34(43):14463–14474.

1166

1167 42.    Destexhe A, McCormick DA, Sejnowski TJ (1993) A model for 8–10 Hz spindling in
1168 interconnected thalamic relay and reticularis neurons. *Biophys J* 65(6):2473–2477.

1169

1170 43.    Destexhe A, Contreras D, Sejnowski TJ, Steriade M (1994) A model of spindle rhythmicity
1171 in the isolated thalamic reticular nucleus. *J Neurophysiol* 72(2):803–818.

1172

1173 44.    Golomb D, Wang XJ, Rinzel J (1996) Propagation of spindle waves in a thalamic slice
1174 model. *J Neurophysiol* 75(2):750-769.

1175

1176 45.    Sohal VS, Huguenard JR (1998) Long-Range Connections Synchronize Rather Than Spread
1177 Intrathalamic Oscillations: Computational Modeling and In Vitro Electrophysiology. *J*
1178 *Neurophysiol* 80(4):1736-1751.

1179

1180 46.    Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ (1998) Computational models of
1181 thalamocortical augmenting responses. *J Neurosci* 18(16):6444-6465.

1182

1183 47.    Wasylenko TM, Cisternas JE, Laing CR, Kevrekidis IG (2010) Bifurcations of lurching
1184 waves in a thalamic neuronal network. *Biol Cybern* 103(6):447-62.

1185

1186 48.    Pham T, Haas JS (2018) Electrical synapses between inhibitory neurons shape the responses
1187 of principal neurons to transient inputs in the thalamus: a modeling study. *Sci Rep* 8(1):7763.

1188

1189 49.    Destexhe A, Contreras D, Steriade M (1998) Mechanisms underlying the synchronizing
1190 action of corticothalamic feedback through inhibition of thalamic relay cells. *J*
1191 *Neurophysiol* 79(2):999–1016.

1192

1193 50.    Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ (2002) Model of thalamocortical slow-
1194 wave sleep oscillations and transitions to activated states. *J Neurosci* 22(19):8691–8704.

1195

51. Traub RD, Contreras D, Cunningham MO, Murray H, LeBeau FE, Roopun A, Bibbig A,
Wilent WB, Higley MJ, Whittington MA (2005) Single-column thalamocortical network
model exhibiting gamma oscillations, sleep spindles, and epileptogenic bursts. *J*
*Neurophysiol* 93(4):2194-2232.

52. Izhikevich EM, Edelman GM (2008) Large-scale model of mammalian thalamocortical
systems. *Proc Natl Acad Sci USA* 105(9):3593-8.

53. Rogala J, Waleszczyk WJ, Lęski S, Wróbel A, Wójcik DK (2013) Reciprocal inhibition and
slow calcium decay in perigeniculate interneurons explain changes of spontaneous firing of
thalamic cells caused by cortical inactivation. *J Comput Neurosci* 34(3):461-476.

54. Willis AM, Slater BJ, Gribkova ED, Llano DA (2015) Open-loop organization of thalamic
reticular nucleus and dorsal thalamus: a computational model. *J Neurophysiol* 114(4):2353-
2367.

55. Sherman SM (2001) Tonic and burst firing: dual modes of thalamocortical relay. *Trends*
*Neurosci* 24(2):122–126.

56. Halassa MM, Siegle JH, Ritt JT, Ting JT, Feng G, Moore CI (2011) Selective optical drive of
thalamic reticular nucleus generates thalamic bursts and cortical spindles. *Nat*
*Neurosci* 14(9):1118–1120.

57. Ahlsén, G, Lindström, S (1982) Mutual inhibition between perigeniculate neurones. *Brain*
*Res* 236(2):482-486.

58. Steriade M, Jones EG, Llinás RR (1990) Thalamic Oscillations and Signaling (Wiley-
Interscience, New York).

59. Cox CL, Huguenard JR, Prince DA (1996) Heterogeneous axonal arborizations of rat
thalamic reticular neurons in the ventrobasal nucleus. *J Comp Neuro* 366(3):416-430.

60. Sanchez-Vives, MV, Bal T, McCormick DA (1997) Inhibitory interactions between
perigeniculate GABAergic neurons. *J Neurosci* 17(22):8894-8908.

61. Shu Y, McCormick DA (2002) Inhibitory interactions between ferret thalamic reticular
neurons. *J Neurophysiol* 87(5):2571-2576.

62. Deleuze C, Huguenard (2006) Distinct electrical and chemical connectivity maps in the
thalamic reticular nucleus: potential roles in synchronization and sensation. *J Neurosci*
26(33):8633-8645.

63. Lam, YW, Nelson CS, Sherman SM (2006) Mapping of the functional interconnections
between thalamic reticular neurons using photostimulation. *J Neurophysiol* 96(5):2593–2600.

64. Landisman CE, Long MA, Beierlein M, Deans MR, Paul DL, Connors BW (2002) Electrical
synapses in the thalamic reticular nucleus. *J Neurosci* 22(3):1002–1009.

65. Long MA, Landisman CE, Connors BW (2004) Small clusters of electrically coupled neurons
generate synchronous rhythms in the thalamic reticular nucleus. *J Neurosci* 24(2):341-349.

66.  Fuentealba P, Crochet S, Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M (2004)
Experimental evidence and modeling studies support a synchronizing role for electrical
coupling in the cat thalamic reticular neurons in vivo. *Eur J Neurosci* 20(1):111-9.
1250
1251  67.  Deleuze C, David F, Béhuret S, Sadoc G, Shin HS, Uebele VN, Renger JJ, Lambert RC,
1252       Leresche N, Bal T (2012) T-type calcium channels consolidate tonic action potential output
1253       of thalamic neurons to neocortex. *J Neurosci* 32(35):12228–12236.
1254
1255  68.  Pospischil M, Toledo-Rodriguez M, Monier C, Piwkowska Z, Bal T, Frégnac Y, Markram H,
1256       Destexhe A (2008) Minimal Hodgkin-Huxley type models for different classes of cortical and
1257       thalamic neurons. *Biol Cybern* 99(4-5):427–441.
1258
1259  69.  Yamada WM, Koch C, Adams PR (1989) Multiple channels and calcium dynamics. *Methods
1260       in Neuronal Modeling*, eds Koch C, Segev I (MIT Press, Cambridge, MA), pp 97–133.
1261
1262  70.  Tsodyks MV, Markram H (1997) The neural code between neocortical pyramidal neurons
1263       depends on neurotransmitter release probability. *Proc Natl Acad Sci USA* 94(2):719–723.
1264
1265  71.  Tsodyks MV, Pawelzik K, Markram H (1998) Neural networks with dynamic synapses.
1266       *Neural Comput* 10(4):821-835.
1267
1268  72.  Pinault D, Smith Y, Deschênes M (1997) Dendrodendritic and axoaxonic synapses in the
1269       thalamic reticular nucleus of the adult rat. *J Neurosci* 17(9):3215-3233.
1270
1271  73.  Cruikshank SJ, Urabe H, Nurmikko AV, Connors BW (2010) Pathway-specific feedforward
1272       circuits between thalamus and neocortex revealed by selective optical stimulation of axons.
1273       *Neuron* 65(2):230-45.
1274
1275  74.  Hou G, Smith AG, Zhang ZW (2016) Lack of Intrinsic GABAergic Connections in the
1276       Thalamic Reticular Nucleus of the Mouse. *J Neurosci* 36(27):7246-7252.
1277
1278  75.  Dayan P, Abbott LF (2005) Theoretical Neuroscience, revised edition (MIT Press,
1279       Cambridge, MA).
1280
1281  76.  Shimizu K, Stopfer M (2013) Gap junction. *Curr Biol* 23(23):R1026-1031.
1282
1283  77.  Chen C, Regehr WG (2003) Presynaptic modulation of the retinogeniculate synapse. *J
1284       Neurosci* 23(8):3130–3135.
1285
1286  78.  Siegel A, Sapru HN (2015) Essential Neuroscience, Third Edition (Lippincott Williams &
1287       Wilkins, Baltimore).
1288
1289  79.  Jones EG (1975) Some aspects of the organization of the thalamic reticular complex. *J Comp
1290       Neurol* 162(3):285-308.
1291
1292  80.  Sohal VS, Huntsman MM, Huguenard JR (2000) Reciprocal inhibitory connections regulate
1293       the spatiotemporal properties of intrathalamic oscillations. *J Neurosci* 20(5):1735-1745.
1294
1295  81.  Crabtree JW (1996) Organization in the somatosensory sector of the cat's thalamic reticular
1296       nucleus. *J Comp Neurol* 366(2):207-222.
1297
1298  82.  Cox CL, Huguenard JR, Prince DA (1997) Nucleus reticularis neurons mediate diverse
1299       inhibitory effects in thalamus. *Proc Natl Acad Sci USA* 94(16):8854-8859.

1300
1301   83.   Moore JW, Ramon F (1974) On numerical integration of the Hodgkin and Huxley equations
1302         for a membrane action potential. *J Theor Biol* 45(1):249–273.
1303
1304   84.   R Core Team (2013) R: A language and environment for statistical computing (R Foundation
1305         for Statistical Computing, Vienna). URL http://www.R-project.org/.
1306
1307   85.   Friedman J, Hastie J, Tibshirani R (2010) Regularization paths for generalized linear models
1308         via coordinate descent. *J Stat Softw* 33(1):1-22.
1309
1310   86.   Herz AV, Gollisch T, Machens CK, Jaeger D (2006) Modeling single-neuron dynamics and
1311         computations: a balance of detail and abstraction. *Science* 314(5796):80-5.
1312
1313   87.   McCormick DA, Pape HC (1980) Properties of a hyperpolarization-activated cation current
1314         and its role in rhythmic oscillation in thalamic relay neurones. *J Physiol (Lond)* 431:291-318.
1315
1316   88.   Williams SR, Stuart GJ (2000) Action potential backpropagation and somato-dendritic
1317         distribution of ion channels in thalamocortical neurons. *J Neurosci* 20(4):1307-17.
1318
1319   89.   Contreras D, Curró Dossi R, Steriade, M (1993) Electrophysiological properties of cat
1320         reticular thalamic neurones in vivo. *J Physiol* 470:273-94.
1321
1322   90.   Destexhe A, Contreras D, Steriade M, Sejnowski TJ, Huguenard JR (1996) In vivo, in vitro,
1323         and computational analysis of dendritic calcium currents in thalamic reticular neurons. *J*
1324         *Neurosci* 16(1):169-85.
1325
1326   91.   Crandall SR, Govindaiah G, Cox CL (2010) Low-threshold $Ca^{2+}$ current amplifies distal
1327         dendritic signaling in thalamic reticular neurons. *J Neurosci* 30(46):15419-29.
1328
1329   92.   Leresche N, Lightowler S, Soltesz I, Jassik-Gerschenfeld D, Crunelli V (1991) Low-
1330         frequency oscillatory activities intrinsic to rat and cat thalamocortical cells. *J*
1331         *Physiol* 441:155-74.
1332
1333   93.   Spreafico R, Battaglia G, Frassoni C (1991) The reticular thalamic nucleus (RTN) of the rat:
1334         cytoarchitectural, Golgi, immunocytochemical, and horseradish peroxidase study. *J Comp*
1335         *Neurol* 304(3):478-90.
1336
1337   94.   Lee SH, Govindaiah G, Cox CL (2007) Heterogeneity of firing properties among rat thalamic
1338         reticular nucleus neurons. *J Physiol (Lond)* 582(1):195-208.
1339
1340   95.   Landau, ID, Egger R, Dercksen VJ, Oberlaender M, Sompolinsky H (2016) The impact of
1341         structural heterogeneity on excitation-inhibition balance in cortical
1342         networks. *Neuron* 92(5):1106-1121.
1343
1344   96.   Cucchiaro JB, Bickford ME, Sherman SM (1991) A GABAergic projection from the
1345         pretectum to the dorsal lateral geniculate nucleus in the cat. *Neuroscience* 41(1):213-26.
1346
1347   97.   Winer JA, Saint Marie RL, Larue DT, Oliver DL (1996) GABAergic feedforward projections
1348         from the inferior colliculus to the medial geniculate body. *Proc Natl Acad Sci USA* 93:8005–
1349         10.
1350
1351   98.   Peruzzi D, Bartlett E, Smith PH, Oliver DL (1997) A monosynaptic GABAergic input from
1352         the inferior colliculus to the medial geniculate body in rat. *J Neurosci* 17:3766–77.

1353

99.    Llano DA, Slater BJ, Lesicko AM, Stebbings KA (2014) An auditory colliculo-
       thalamocortical brain slice preparation in mouse. *J Neurophysiol* 111:197–207.

100.   Steriade M, Wyzinski P, Apostol V (1972) Corticofugal projections governing rhythmic
       thalamic activity. Corticothalamic Projections and Sensorimotor Activities,
       eds Frigyesi TL, Rinvik E, Yahr MD (Raven, New York), pp 221–72.

101.   White EL, Hersch SM (1982) A quantitative study of thalamocortical and other synapses
       involving the apical dendrites of corticothalamic projection cells in mouse SmI cortex. *J
       Neurocytol* 11(1):137-57.

102.   De Curtis M, Spreafico R, Avanzini G (1989) Excitatory amino acids mediate responses
       elicited in vitro by stimulation of cortical afferents to reticularis thalami neurons of the rat.
       *Neuroscience* 33(2):275-83.

103.   Contreras D, Destexhe A, Sejnowski TJ, Steriade M (1996) Control of spatiotemporal
       coherence of a thalamic oscillation by corticothalamic feedback. *Science* 274(5288):771-4.

104.   Blumenfeld H, McCormick DA (2000) Corticothalamic inputs control the pattern of activity
       generated in thalamocortical networks. *J Neurosci* 20(13):5153-62.

105.   Zhang L, Jones EG (2004) Corticothalamic inhibition in the thalamic reticular nucleus. *J
       Neurophysiol* 91(2):759-66.

106.   Crandall SR, Cruikshank SJ, Connors BW (2015) A corticothalamic switch: controlling the
       thalamus with dynamic synapses. *Neuron* 86(3):768-82.

107.   Roy JP, Clercq M, Steriade M, Deschênes M (1984) Electrophysiology of neurons of lateral
       thalamic nuclei in cat: mechanisms of long-lasting hyperpolarizations. *J Neurophysiol*
       51(6):1220-35.

108.   Contreras D, Steriade M (1996) Spindle oscillation in cats: the role of corticothalamic
       feedback in a thalamically generated rhythm. *J Physiol* 490(1):159-179.

109.   Suga N, Ma X (2003) Multiparametric corticofugal modulation and plasticity in the auditory
       system. *Nat Rev Neurosci* 4(10):783-94.

110.   Sillito AM, Cudeiro J, Jones HE (2006) Always returning: feedback and sensory processing
       in visual cortex and thalamus. *Trends Neurosci* 29(6):307-16.

111.   Muller L, Destexhe A (2012) Propagating waves in thalamus, cortex and the thalamocortical
       system: Experiments and models. *J Physiol Paris* 106(5-6):222-38.

112.   Destexhe A, Sejnowski TJ (2003) Interactions between membrane conductances underlying
       thalamocortical slow-wave oscillations. *Physiol Rev* 83(4):1401-53.

113.   Andersen P, Andersson SA (1968) The physiological basis of the alpha rhythm (Appleton-
       Century-Crofts, New York).

114.   Jones EG (1985) The Thalamus (Plenum Press, New York).

115. Alonso JM, Usrey WM, Reid RC (2001) Rules of connectivity between geniculate cells and simple cells in cat primary visual cortex. *J Neurosci* 21(11):4002-15.

116. Miller LM, Escabí MA, Read HL, Schreiner CE (2001) Functional convergence of response properties in the auditory thalamocortical system. *Neuron* 32(1):151-60.

117. Rinzel J, Terman D, Wang X, Ermentrout B (1998) Propagating activity patterns in large-scale inhibitory neuronal networks. *Science* 279(5355):1351-5.

118. Steriade M, Llinás RR (1988) The functional states of the thalamus and the associated neuronal interplay. *Physiol Rev* 68(3):649-742.

119. Soltesz I, Lightowler S, Leresche N, Jassik-Gerschenfeld D, Pollard CE, Crunelli V (1991) Two inward currents and the transformation of low-frequency oscillations of rat and cat thalamocortical cells. *J Physiol (Lond)* 441:175-97.

120. Gibbs FA, Gibbs EL (1950) Atlas of Electroencephalography, volume one (Addison-Wesley, Cambridge, MA)

121. Bazhenov M, Timofeev I, Steriade M, Sejnowski TJ (1999) Self-sustained rhythmic activity in the thalamic reticular nucleus mediated by depolarizing GABA$_A$ receptor potentials. *Nat Neurosci* 2(2):168-74.

122. Bal T, Mccormick DA (1996) What stops synchronized thalamocortical oscillations? *Neuron* 17(2):297-308.

123. Lüthi A, McCormick DA (1998) Periodicity of thalamic synchronized oscillations: the role of Ca2+-mediated upregulation of Ih. *Neuron* 20(3):553-63.

124. Timofeev I, Bazhenov M, Sejnowski TJ, Steriade M (2001) Contribution of intrinsic and synaptic factors in the desynchronization of thalamic oscillatory activity. *Thalamus Relat Syst* 1(1):53-69.

125. Bonjean M, Baker T, Lemieux M, Timofeev I, Sejnowski T, Bazhenov M (2011) Corticothalamic feedback controls sleep spindle duration in vivo. *J Neurosci* 31(25):9124-34.

126. Huntsman MM, Porcello DM, Homanics GE, Delorey TM, Huguenard JR (1999) Reciprocal inhibitory connections and network synchrony in the mammalian thalamus. *Science* 283(5401):541-3.

127. Sohal VS, Huguenard JR (2003) Inhibitory interconnections control burst pattern and emergent network synchrony in reticular thalamus. *J Neurosci* 23(26):8978-88.

128. Crunelli V, Leresche N (2002) Childhood absence epilepsy: genes, channels, neurons and networks. *Nat Rev Neurosci* 3(5):371-82.

129. Lee SC, Patrick SL, Richardson, KA, Connors, BW (2014) Two functionally distinct networks of gap junction-coupled inhibitory neurons in the thalamic reticular nucleus. *J Neurosci* 34(39):13170-82.

130. Haas JS, Zavala B, Landisman CE (2011) Activity-dependent long-term depression of electrical synapses. *Science* 334(6054):389-93.

1458 131.  Haas JS, Landisman CE (2012) State-dependent modulation of gap junction signaling by the
1459       persistent sodium current. *Front Cell Neurosci* 5:31. doi:10.3389/fncel.2011.00031
1460
1461 132.  Wang Z, Neely R, Landisman CE (2015) Activation of group I and group II metabotropic
1462       glutamate receptors causes LTD and LTP of electrical synapses in the rat thalamic reticular
1463       nucleus. *J Neurosci* 35(19):7616-25.
1464
1465 133.  Kohmann D, Lüttjohann A, Seidenbecher T, Coulon P, Pape HC (2016) Short-term
1466       depression of gap junctional coupling in reticular thalamic neurons of absence epileptic rats. *J
1467       Physiol (Lond)* 594(19):5695-710.
1468
1469 134.  Pernelle G, Nicola W, Clopath C (2017) Gap junction plasticity can lead to spindle
1470       oscillations. arXiv:1710.03999 [q-bio.NC]
1471
1472 135.  Crabtree JW (2018) Functional Diversity of Thalamic Reticular Subnetworks. *Front Syst
1473       Neurosci* 12:41. doi:10.3389/fnsys.2018.00041
1474
1475 136.  Scheibel ME, Scheibel AB (1966) The organization of the nucleus reticularis thalami: a Golgi
1476       study. *Brain Res* 1(1):43-62.
1477
1478 137.  Spreafico R, De Curtis M, Frassoni C, Avanzini G (1988) Electrophysiological characteristics
1479       of morphologically identified reticular thalamic neurons from rat slices. *Neuroscience*
1480       27(2):629-38.
1481
1482 138.  Brunton J, Charpak S (1997) Heterogeneity of cell firing properties and opioid sensitivity in
1483       the thalamic reticular nucleus. *Neuroscience* 78(2):303-7.
1484
1485 139.  Bickford ME (2016) Thalamic Circuit Diversity: Modulation of the Driver/Modulator
1486       Framework. *Front Neural Circuits* 9:86. doi: 10.3389/fncir.2015.00086
1487

1488

1489

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499 

**A**

1500

**B**

1501

1502 **Figure 1. A:** Closed- vs. open-loop thalamo-reticulo-thalamic configurations. **B:** Three possible pathways

1503 through which a signal might propagate from one thalamic relay (TC) neuron to another via the thalamic

1504 reticular nucleus (TRN).

**A**

L4

TRN

TC

Excitatory    External
Inhibitory    Gap Junction

1505

**B**

Input to TC_A

Input to TC_B

Input to TC_C

120 mV

100 ms

1506

# C

1507

1508      **Figure 2. A:** Baseline three-layer, three-"column" thalamo-reticulo-cortical model network. Broken-line

1509      synapses were allowed to vary either as a class (homogeneously) or independently of one another

1510      (heterogeneously). **B,C:** Samples of the external input delivered to the TC neuron layer. $TC_A$ received a

1511      combination of Poisson-modulated and fixed input, either punctate (B) or sustained (C).

1512

1513

1514

1515

1516

1517

1518

1519

1520

1521

1522

**Figure 3.** Sample detrended L4 spike histograms in a network permutation responding to a fixed,

sustained stimulus delivered to $TC_A$ (yellow arrow). The propagation score assigned to any network

permutation was quantified as the amplitude of the initial stimulus-evoked response in the detrended $L4_c$

histogram; response propagation across the L4 subnetwork (orange arrow) was consistently linear, and

thus the initial response in $L4_C$ was observed at a fixed interval relative to the onset of stimulation.

Oscillation intrinsic to any network variant was quantified as the amplitude of the first off-center peak in

the normalized autocorrelogram (right) of post-stimulation activity in the detrended $L4_C$ histogram

(within broken black box). The initial 400 ms of activity preceding the fixed stimulus (in grey) is shown

here for each histogram but was not included in the calculations of either propagation or oscillation. Note

that the bin heights in the $L4_A$ histogram shown here were truncated in order to maintain identical vertical

scaling across all three L4 histograms.

1534

1535

1536

**A**

Propagation

Oscillation

TRN-TRN
GABA_A Conductance

TRN-TRN
Electrical Coupling

TC-TRN-TC Openness

a c    b

a  c  b

c    a,b

1537

**B**

Propagation Scores
(Openness Coeff. = 1.0)

GABA_A Cond. (nS)

Electrical Coupling Coeff.

Oscillation Scores
(Openness Coeff. = 0)

GABA_A Cond. (nS)

Electrical Coupling Coeff.

1538

1539

**Figure 4. A:** Ordinal heat maps ranking homogeneously synaptic network permutations according to the

extent of supported signal propagation and oscillation in response to a fixed, punctate stimulus. Warmer

colors correspond to higher values for each of the three synaptic class variables. The positions of three

selected networks are indicated on both heat maps according to their propagation and oscillation scores.

Each network is indexed by the synaptic variables ([TRN-TRN GABA$_A$ conductance (in nS), TRN-TRN

electrical coupling coefficient, TC-TRN-TC openness coefficient]): a, [0,0,1.0]; b, [450,0.36,1.0]; and c,

[50,0.18,0]. Networks with high degrees of TC-TRN-TC openness tended to best support signal

propagation, while GABAergic synapses and electrical coupling between TRN neurons modestly

impeded it. By contrast, networks with strong closed-loop reticulothalamic architectures were most

permissive of oscillation, with TRN-TRN synaptic strength negligibly correlated with this property. **B:**

Heat maps displaying propagation scores for the 70 fully open-loop homogeneously synaptic networks

permutations (left, openness coefficient=1.0) and oscillation scores for the 70 fully closed-loop network

permutations (right, openness coefficient=0), with the three selected networks indicated on the maps.

Propagation was typically stronger in network permutations with minimal or nonexistent TRN-TRN

connectivity (e.g., Network a, cf. Network b). **C:** Representative simulations and circuit diagrams

1555 depicting the normalized synaptic makeups for each of the three selected networks. The yellow arrow

1556 indicates when the fixed, punctate stimulus was delivered to $TC_A$ in each simulation. Orange highlighting

1557 indicates epochs of linear propagation, while circles are placed above spikes occurring during periods of

1558 oscillatory activity. Note that the episode of oscillatory activity depicted in the simulation of Network c

1559 changes frequency slightly following the third action potential.

1560

1561

1562

1563

1564

1565

1566

1567

1568

1569

1570

1571

1572

1573

1574

1575

1576

1577

1578

1579

1580

**A**

Propagation

Oscillation

TRN-TRN
GABA_A Conductance

d e    f

TRN-TRN
Electrical Coupling

d e    f

TC-TRN-TC Openness

fe  d

1581

**B**

Propagation Scores
(Openness Coeff. = 1.0)

GABA_A Cond. (nS)

1000

Electrical Coupling Coeff.

Oscillation Scores
(Openness Coeff. = 0.4)

GABA_A Cond. (nS)

0.5

0.2

Electrical Coupling Coeff.

Oscillation Scores
(Openness Coeff. = 0.2)

GABA_A Cond. (nS)

0.5

0.2

Electrical Coupling Coeff.

1582

**C**

**D**

**Figure 5. A:** Ordinal heat maps ranking homogeneously synaptic network permutations according to the extent of supported signal propagation and oscillation in response to a fixed, sustained stimulus. The positions of three additional selected networks are indicated on both heat maps, with each selected network indexed as in Figure 4: d, [0,0.06,1.0]; e, [100,0.12,0.4]; and f, [450,0.36,0.2]. As was the case

with fixed, punctate stimulation, network propagation was strongest in networks exhibiting strong open-

loop TC-TRN-TC connectivity, while the inverse was true with oscillation. Both varieties of TRN-TRN

synapse tended to diminish propagation while exerting a negligible effect on oscillation. **B:** Heat maps

displaying propagation and oscillation scores in TRN-TRN synaptic parameter space at different values of

TC-TRN-TC openness, with the three selected networks indicated on the maps. Propagation was strongest

in exclusively open-loop networks (left, openness coefficient=1.0), while oscillation was more likely to

occur in networks with stronger closed-loop connectivity (middle, openness coefficient=0.4; right,

openness coefficient=0.2). **C:** Representative simulations and circuit diagrams depicting the normalized

synaptic makeups for each of the three additional selected networks. The yellow arrow indicates when the

fixed, sustained stimulus was delivered to $TC_A$ in each simulation. Note the two different mechanisms

underlying oscillation in the simulations of Networks e and f. **D:** Mean oscillation scores for networks

varied nonlinearly as a function of their openness coefficients, with networks possessing openness

coefficients of 0 and 0.4 ($\alpha$) supporting oscillation to equal extents (one-way ANOVA with Tukey post-

hoc tests, $F(10,759)=137.8$, $p<0.0001$). Error bars indicate standard errors of the mean.

1603

1604

1605

1606

1607

1608

1609

1610

1611

1612

1613

1614

**A**

Positive Propagation Coeffs.          Negative Propagation Coeffs.

-1.00          0          1.00

Regression Coefficient (Normalized)

1615

**B**

Positive Oscillation Coeffs.          Negative Oscillation Coeffs.

-1.00          0          1.00

Regression Coefficient (Normalized)

1616

**C**

Network a'  Network b'  Network c'

L4 A B C
TRN A B C
TC A B C

200 mV
250 ms

Synaptic Weight (Normalized)
0  1

1617

**Figure 6. A,B:** Network regression models illustrating how propagation (A) and oscillation (B) varied as

a function of individual synaptic weights across simulated heterogeneously synaptic network

permutations responding to a fixed, sustained stimulus delivered to TC$_A$. Gray synapses are either non-

variable or associated with normalized regression coefficients with absolute values under 0.05. Synapses

with positive and negative coefficients in the regression models are depicted separately in the left- and

right-sided circuit diagrams, respectively. **C:** Representative simulations for three selected heterogeneous

networks, whose normalized synaptic weights are depicted in the circuit diagrams. The yellow arrow

indicates when the fixed, sustained stimulus was delivered to TC$_A$ in each simulation. Networks a', b',

and c' respectively illustrate propagation, self-contained generation of oscillation in network "Column C,"

and propagation of oscillation from Column A to Column C.

1628

1629

1630

1631

1632

**Figure 7**. Propagation, as measured in those network permutations scoring highest with respect to the property, was equally supported in networks where synaptic weights varied independently of one another (**heterogeneously**; red) as in networks where synaptic strength varied **homogeneously** (blue, yellow) by class [one-way ANOVA, $F(2,59)=0.84$, $p=0.437$]. By contrast, oscillation scores were significantly higher in top-performing heterogeneous networks than their homogeneous counterparts [one-way ANOVA with Tukey's HSD tests, $F(2,59)=166.14$, $p<0.0001$], while there was no significant difference in oscillation among top-performing homogeneous network variants when stimulated externally in a fixed, punctate (blue) or sustained (yellow) manner (Tukey's test, $p=0.153$). Each bar corresponds to a mean of the top 20 network propagation or oscillation scores within each synaptic architecture/fixed stimulation group; error bars indicate standard errors of the mean. ****=Tukey's test, $p<0.0001$; N.S.=not significant.

1644

1645

1646

1647

| Model Cellular Parameters | | | |
|---|---|---|---|
| Parameter | TC cell | TRN cell | L4 cell |
| Leak conductance, $g_L$ (nS) | 3.263 | 3.7928 | 4.8128 |
| Leak reversal potential, $E_L$ (mV) | -60.03 | -57 | -60.2354 |
| Transient sodium conductance, $g_{Na}$ (nS) | 1,500 | 3,000 | 3,000 |
| Sodium equilibrium potential, $E_{Na}$ (mV) | 50 | | |
| Delayed-rectifier potassium conductance, $g_K$ (nS) | 520 | 400 | 140 |
| M-type potassium conductance, $g_M$ (nS) | - | 3.5 | 1.5 |
| M-type potassium time constant, $\tau_M$ (ms) | - | 200 | 180 |
| Potassium equilibrium potential, $E_K$ (mV) | -100 | | -90 |
| T-type calcium conductance, $g_T$ (nS) | 45 | 21 | - |
| Calcium equilibrium potential, $E_T$ (mV) | 120 | | |
| H-current conductance, $g_H$ (nS) | 0.608 | 0.0192 | - |
| H-current reversal potential, $E_H$ (mV) | -33 | | - |
| Membrane capacitance, $C_m$ (pF) | 100.4 | 75.0 | 109.3865 |

1648

1649    **Table 1.** Intrinsic model cellular parameters.

1650

1651

1652

1653

1654

1655

1656

1657

1658

1659

1660

1661

1662

| Model Synaptic Parameters | | | | | | |
|---|---|---|---|---|---|---|
| Synapse | Neurotransmitter | Conductance (nS) | $\tau_{recov}$ (ms) | $\tau_{inact}$ (ms) | Reversal Potential (mV) | $U_{SE}$ |
| External synapse to TC cell | (Glutamate) | 32 | 125 | 2.64 | 0 | 0.76 |
| TC-to-TRN cell synapse (TC-TRN) | Glutamate | 150 | 500 | 2.64 | 0 | 0.76 |
| TC-to-L4 cell synapse (TC-L4) | Glutamate | 50 | 160 | 11.52 | 0 | 0.8113 |
| TRN-to-TC cell synapse (TRN-TC) | GABA$_A$ | Variable (0-80) | 167.29 | 16.62 | -80 | 0.62 |
| Chemical TRN-to-TRN cell synapse (TRN-TRN$_{GABA}$) | GABA$_A$ | Variable (0-450) | 225 | 15 | -75 | 0.62 |

1663

**Table 2.** Model synaptic parameters. The external synapse to TC neurons, through which both fixed and

stochastic inputs were delivered, was generically excitatory but explicitly modeled on glutamatergic

retinogeniculate synapse (Chen and Regehr, 2003). The $\tau_{recov}$, $\tau_{inact}$, and $U_{SE}$, parameters derived from the

synaptic depression model of Tsodyks and Markram (1997; Tsodyks et al., 1998), represent the time

constant for post-exocytotic synaptic recovery, the time constant characterizing synaptic inactivation

following the onset of presynaptic neurotransmitter release, and the fraction of available "resources"

(synaptic vesicles) available for release at the presynaptic terminal following action potential induction.

The conductances of both TRN-TC and TRN-TRN$_{GABA}$ synapses were allowed to vary within the

specified ranges.

1673

1674

1675

1676

1677

1678

1679

1680

1681

1682

1683

| Normalized Regression Coefficients (NRCs) | | | | |
|---|---|---|---|---|
| Synaptic Variable | Propagation Linear | Propagation 2° | Oscillation Linear | Oscillation 2° |
| TRN-TRN$_{GABA}$ | -0.137 | -0.413 | 0.058 | - |
| TRN-TRN$_{Elec}$ | -0.193 | -0.499 | 0.064 | 0.056 |
| TRN-TC | 1.000 | 1.000 | -1.000 | -1.000 |
| (TRN-TRN$_{GABA}$)$^2$ | - | 0.117 | - | - |
| (TRN-TRN$_{Elec}$)$^2$ | - | 0.266 | - | - |
| (TRN-TC)$^2$ | - | 0.213 | - | -0.010 |
| TRN-TRN$_{GABA}$ x TRN-TRN$_{Elec}$ | - | 0.226 | - | - |
| TRN-TRN$_{Elec}$ x TRN-TC | - | -0.239 | - | - |

1684

**Table 3.** Normalized linear and second-order regression coefficients for propagation and oscillation in

homogeneously synaptic networks excited by a fixed, punctate stimulus. The regressions include 1°, 2°,

and interaction terms corresponding to TRN-TRN$_{GABA}$, TRN-TRN$_{Elec}$, and open-loop TC-TRN-TC

synapses/pathways. Terms associated with regression coefficients of absolute values < 0.05 are omitted.

Positive and negative terms are highlighted in red and blue, respectively. Linear regression for

propagation, $R^2$=0.829, RMSE=0.042, $p$<0.0001; second-order regression for propagation, $R^2$=0.863,

RMSE=0.039, $p$<0.0001; linear regression for oscillation, $R^2$=0.661, RMSE=0.137, $p$<0.0001; second-

order regression for oscillation, $R^2$=0.687, RMSE=0.132, $p$<0.0001.

1693

1694

1695

1696

1697

1698

1699

1700

| Normalized Regression Coefficients (NRCs) | | | | |
|---|---|---|---|---|
| Synaptic Variable | Propagation Linear | Propagation 2° | Oscillation Linear | Oscillation 2° |
| TRN-TRN$_{GABA}$ | -0.173 | -0.670 | - | 0.060 |
| TRN-TRN$_{Elec}$ | -0.136 | -0.347 | - | - |
| TRN-TC | 1.000 | 1.000 | -1.000 | -0.052 |
| (TRN-TRN$_{GABA}$)$^2$ | - | 0.332 | - | - |
| (TRN-TRN$_{Elec}$)$^2$ | - | 0.164 | - | - |
| (TRN-TC)$^2$ | - | 0.594 | - | -1.000 |
| TRN-TRN$_{GABA}$ x TRN-TRN$_{Elec}$ | - | 0.262 | - | - |
| TRN-TRN$_{GABA}$ x TRN-TC | - | -0.152 | - | - |
| TRN-TRN$_{Elec}$ x TRN-TC | - | -0.365 | - | - |

**Table 4.** Normalized linear and second-order regression coefficients for propagation and oscillation in homogeneously synaptic networks excited by a fixed, sustained stimulus. Linear regression for propagation, $R^2$=0.793, RMSE=0.047, $p<0.0001$; second-order regression for propagation, $R^2$=0.842, RMSE=0.041, $p<0.0001$; linear regression for oscillation, $R^2$=0.526, RMSE=0.145, $p<0.0001$; second-order regression for oscillation, $R^2$=0.630, RMSE=0.128, $p<0.0001$.

| Normalized Regression Coefficients (NRCs) | | | | |
|---|---|---|---|---|
| Synaptic Variable | Propagation Linear | Propagation 2° | Oscillation Linear | Oscillation 2° |
| $TRN_A\text{-}TRN_C$ | - | - | 0.115 | - |
| $TRN_C\text{-}TRN_A$ | -0.088 | | | |
| $TRN_C\text{-}TRN_B$ | -0.084 | -0.073 | - | - |
| $TRN_A\text{=}TRN_B$ | -0.051 | -0.091 | - | - |
| $TRN_A\text{=}TRN_C$ | -0.072 | - | - | - |
| $TRN_B\text{=}TRN_C$ | - | -0.113 | 0.117 | - |
| $TRN_A\text{-}TC_A$ | -0.075 | - | 0.621 | 0.077 |
| $TRN_A\text{-}TC_B$ | 0.608 | 0.571 | -0.289 | -1.000 |
| $TRN_B\text{-}TC_B$ | -0.128 | -0.196 | 0.333 | 0.417 |
| $TRN_B\text{-}TC_C$ | 1.000 | 1.000 | -0.379 | -0.892 |
| $TRN_C\text{-}TC_C$ | -0.207 | -0.239 | 1.000 | 0.107 |
| $(TRN_C\text{-}TRN_B)^2$ | - | 0.079 | - | - |
| $(TRN_A\text{-}TC_B)^2$ | - | -0.245 | - | 0.189 |
| $(TRN_B\text{-}TC_B)^2$ | - | 0.174 | - | -0.093 |
| $(TRN_B\text{-}TC_C)^2$ | - | -0.472 | - | 0.278 |
| $(TRN_C\text{-}TC_C)^2$ | - | 0.187 | - | -0.146 |

| Normalized Regression Coefficients (NRCs), continued | | |
|---|---|---|
| Synaptic Variable | Propagation 2° | Oscillation 2° |
| $TRN_A\text{-}TRN_B$ x $TRN_A\text{-}TC_B$ | 0.070 | - |
| $TRN_A\text{-}TRN_C$ x $TRN_C\text{-}TC_C$ | - | 0.215 |
| $TRN_B\text{-}TRN_A$ x $TRN_A\text{=}TRN_B$ | - | 0.111 |
| $TRN_B\text{-}TRN_A$ x $TRN_A\text{-}TC_A$ | - | -0.186 |
| $TRN_C\text{-}TRN_A$ x $TRN_A\text{-}TC_A$ | - | -0.172 |
| $TRN_C\text{-}TRN_A$ x $TRN_A\text{-}TC_B$ | -0.119 | - |
| $TRN_C\text{-}TRN_A$ x $TRN_B\text{-}TC_C$ | -0.096 | - |
| $TRN_C\text{-}TRN_B$ x $TRN_B\text{-}TC_C$ | -0.153 | - |
| $TRN_A\text{=}TRN_B$ x $TRN_B\text{-}TC_C$ | - | -0.129 |
| $TRN_A\text{=}TRN_C$ x $TRN_A\text{-}TC_A$ | - | -0.114 |
| $TRN_A\text{=}TRN_C$ x $TRN_C\text{-}TC_C$ | -0.079 | - |
| $TRN_A\text{-}TC_A$ x $TRN_A\text{-}TC_B$ | - | 0.634 |
| $TRN_A\text{-}TC_A$ x $TRN_B\text{-}TC_C$ | - | 0.449 |
| $TRN_A\text{-}TC_B$ x $TRN_B\text{-}TC_B$ | -0.166 | 0.361 |
| $TRN_A\text{-}TC_B$ x $TRN_B\text{-}TC_C$ | 0.753 | -0.274 |
| $TRN_A\text{-}TC_B$ x $TRN_C\text{-}TC_C$ | -0.106 | 0.669 |
| $TRN_B\text{-}TC_B$ x $TRN_B\text{-}TC_C$ | - | 0.345 |
| $TRN_B\text{-}TC_B$ x $TRN_C\text{-}TC_C$ | - | -0.192 |
| $TRN_B\text{-}TC_C$ x $TRN_C\text{-}TC_C$ | -0.124 | 0.399 |

**Table 5.** Normalized linear and second-order regression coefficients for oscillation, propagation, and optimization in heterogeneously synaptic networks excited by a fixed, sustained stimulus. The regressions include 1°, 2°, and interaction terms corresponding to the 14 variable synapses in the networks. Equal signs denote gap junctions. Linear regression for propagation, $R^2=0.742$, RMSE=0.069, $p<0.0001$; second-order regression for propagation, $R^2=0.857$, RMSE=0.051, $p<0.0001$; linear regression for oscillation, $R^2=0.253$, RMSE=0.131, $p<0.0001$; second-order regression for oscillation, $R^2=0.388$, RMSE=0.118, $p<0.0001$.