The structure of species discrimination signals across a primate radiation

Sandra Winters[1], William L. Allen[1,2], & James P. Higham[1]

[1]Department of Anthropology, New York University, 25 Waverly Place, New York, NY 10003,

USA

[2]Department of Biosciences, Swansea University, Wallace Building, Singleton Park, Swansea

SA2 8PP, Wales, UK

Discriminating between conspecifics and heterospecifics potentially challenging for closely

related sympatric species. The guenons, a recent primate radiation, exhibit high degrees of

sympatry and form multi-species groups in which hybridization is possible but rare in most

populations. Guenons have species-specific colorful face patterns hypothesized to function in

species discrimination. Here, we apply a novel machine learning approach to identify the face

regions most essential for correct species classification across fifteen guenon species. We then

demonstrate the validity of these computational results using experiments with live guenons,

showing that facial traits identified as critical for accurate classification do indeed influence

selective attention toward con- and heterospecific faces. Our results suggest variability among

guenon species in reliance on single-trait-based versus holistic facial characteristics when

discriminating between species, and differences in behavioral responses to faces can be linked to

whether discrimination is based on a single trait or whole-face pattern. Our study supports the

hypothesis that guenon face patterns function to promote species discrimination and provides

novel insights into the relationship between species interactions and phenotypic diversity.

Keywords: species discrimination, visual signal form, complex signals, reproductive isolation,

Cercopithecini

Introduction

Closely related species living in sympatry face a potential challenge in discriminating between

conspecifics and heterospecifics. Such decision-making has important selective outcomes,

particularly in behaviors such as mate choice, with individuals choosing heterospecific mates

often incurring substantial fitness costs [1]. One mechanism for avoiding the costs if interacting

with heterospecifics is the use of species-specific signals that structure behavioral interactions

between species. For instance, mating signals and associated mating preferences that differ

between sympatric heterospecifics can function to maintain reproductive isolation across species

boundaries [2]. Such signals are predicted to be salient and distinctive [3], with sympatric species

under selective pressure to diversify. A pattern in which signal distinctiveness increases with

degree of sympatry, known as character displacement [4,5], has been observed in a wide variety

of animal groups [6–13]. Importantly, signals that function to maintain reproductive isolation via

mate choice should elicit increased mating interest from conspecifics compared to

heterospecifics [14].

Species in evolutionarily young animal radiations may be at particular risk of

hybridization and other costly interactions with heterospecifics due to behavioral similarities and

a lack of post-mating barriers to reproduction [15]. One such radiation is the guenons (tribe

Cercopithecini), a group of African primates consisting of 25-38 recognized species [16–18] that

diverged from papionin primates around 11.5 million years ago [19]. Guenons exhibit high

degrees of sympatry and often form polyspecific groups in which multiple species travel and

forage together [20]. Many guenons therefore interact with heterospecifics that share general

patterns of morphology (e.g. overall body size/shape) and behavior (e.g. activity patterns). In

such circumstances, discriminating between con- and heterospecifics may be particularly

important, especially in a mating context. Hybridization between sympatric guenon species is

possible but rare in natural circumstances [21], suggesting the existence of barriers to

heterospecific mating within mixed-species groups.

Guenons are among the most colorful and visually patterned groups of primates with

many species exhibiting extraordinary and unique face markings [10,23,25–27], which are

minimally variable between sexes across all guenon species [23,24]. Kingdon [23,26,27]

hypothesized that guenons use their divergent facial appearances to distinguish between species

and therefore select appropriate mates. This young and impressively diverse primate radiation

represents a fascinating test case of how visual signals are involved in species radiations and

mixed-species interactions [5,28–30]. Recent empirical work has begun to generate evidence for

their key role in guenon phenotypic and species diversification. Images of guenon faces can be

reliably classified by species using computer algorithms [10,24], demonstrating that guenon

faces contain species-specific identifying information. Guenon face patterns also exhibit

character displacement, with facial distinctiveness between species increasing with degree of

sympatry across the group [10]. Moreover, facial components common across species (nose

spots and eyebrow patches) alone can be used to computationally classify species [24]. This

suggests that guenon faces may be somewhat modular, with species information encoded in

particular face regions. Which face regions are most important, and the extent to which such

regions vary across species remains an open question that is of key importance to understanding

how complex signals involved in species discrimination evolve. Critically, it is unknown whether

variation across guenon species in purported species discrimination signals is perceived and

acted on by con- and heterospecific receivers.

Here, we use a machine learning approach to identify guenon face regions that are most

important for correct species classification by a computer. These results objectively identify the

signal components most likely to be useful to guenon receivers. We use them to determine which

signal properties to systematically investigate in behavioral experiments with guenon observers.

The machine-learning stage is critical, as many experiments that investigate behavioral responses

to complex signals select manipulations based on the perceptions of investigators, which

introduces anthropocentric bias [31]. Using the guenon face image database produced by Allen et

al. [10], we couple eigenface decomposition of the faces [32] with a novel occlude-reclassify

scheme in which we systematically block each part of the face and reclassify the image. This

allows us to document the spatial distribution of species-typical information across guenon faces

by identifying which face regions, when obscured, cause the break-down of correct species

classification. Eigenface decomposition was originally developed for individual face

discrimination in humans [32]; feature detection based on eigenfaces is also applicable to other

types of discrimination tasks involving complex animal signals [33–35] and has been used

previously to quantify guenon facial variation [10]. The perceptual face space generated by

eigenface decomposition parallels mammalian visual processing [36], lending biological

credibility.

After identifying the face regions that cause break-down in classification, and thus those

that should be important for correct species identification, we then present captive putty nosed

monkeys (*Cercopithecus nictitans*) and mona monkeys (*C. mona*) with images of con- and

heterospecific faces exhibiting variation in these regions and measure their resulting eye gaze to

assess their ability to distinguish between species based on face patterns. Ours is the first direct

measure of guenon responses to con- and heterospecific faces, which is crucial for clarifying the

biological relevance of guenon face patterns and for validating previous correlational results.

Differences in looking time between classes of stimuli can be difficult to interpret due to various

and often unpredictable novelty and familiarity effects [37], however primates reliably exhibit a

visual bias (i.e. greater looking time) toward images of conspecifics compared to those of

heterospecifics [38–42]. We follow the interpretation that longer looking time at a particular face

reflects level of interest. This is consistent with an interpretation that the face resembles a

conspecific face more closely, though other explanations are possible.

Our experimental trials involve the simultaneous presentation of paired con- and

heterospecific faces, focusing on a particular facial trait for each species. For putty nosed

monkeys we focus on nose spots and for mona monkeys on eyebrow patches, on the basis that

each of these features is within the region of the face identified by our machine learning

approach as being critical for that species. In each trial, heterospecific faces either do or do not

share a focal face trait with the subject, and conspecific faces are presented either naturally or

after being modified to remove the focal trait (for example stimuli, see Figure 1). This approach

allows us to assess generalized species biases in degree of interest as well as the extent to which

particular face regions influence these biases.

We predicted variability across species in the face regions identified by our occlude-

reclassify procedure, but made no predictions regarding which regions in particular would be

essential for each species. In looking time experiments, we predicted that putty nosed and mona

monkeys would exhibit visual biases toward face images of conspecifics, and that these biases

would be influenced by species-typical facial characteristics identified as important for correct

species classification. Such a pattern of results would support a role for species discrimination

signals likely used to facilitate inter-specific interactions such as maintaining reproductive

isolation via mate choice in generating and maintaining phenotypic variation in one of the most

speciose and diverse primate radiations. Ultimately, by examining how aspects of highly

complex signals encode species identity and influence receiver biases, this research increases

our understanding of how selection for species identity signaling generates phenotypic diversity.

Methods

Image collection & processing

Guenon face pattern analyses are based on an existing database of guenon face images from 22

guenon species [10]. Detailed methods of image collection and processing have been published

elsewhere [10]. Briefly, we used digital images of captive guenons collected using a color-

calibrated camera. Multiple images were taken of each subject while in a front-facing position

under indirect light. Images were transformed from camera RGB color space to guenon LMS

color space, defined by the peak spectral sensitivities of guenon long, medium, and short

wavelength photoreceptors. All images were then standardized with respect to illumination, size,

blur, and background. Each image was resized to be 392 by 297 by 3 pixels. All pixel values

were represented using double-level precision.

To avoid classifying species based on a very small number of exemplars, we restricted

our analyses to species represented by at least four individuals in our image database (i.e. all

classifications in a leave-one-out procedure are made based on at least three exemplars; see

below). Our analysis is therefore based on 599 total images of 133 individuals, collectively

representing fifteen guenon species (for species-specific sample sizes, see Figure 3).

Identification of face regions important for species classification

Guenon face images can be reliably classified by species based on eigenface features [10,24].

This approach relies on dimensionality reduction via principal component analysis (PCA) to

extract relevant features from face images; these features can then be used for the classification

of new faces [32]. In this procedure, each 'eigenface' (i.e. the eigenvectors resulting from PCA

of all face images) represents a different dimension of facial variability and each face image can

be represented by a series of weights associated with each eigenface. This creates a multi-

dimensional 'face space' in which faces are represented as points based on their eigenface

weights, and zero weights for all eigenfaces (i.e. the center of the space) represents the average

face across all images. Such face spaces have psychophysical parallels in primate face processing

centers in the visual cortex [36]. Multiple images of each subject were averaged to generate

average individual faces, which in turn were used to generate the average species faces that were

used in eigenface decomposition. We classified new images using a nearest-neighbor classifier

based on minimum Euclidean distance to each average species face in face space. This scheme

corresponds to an average face model of guenon face learning, which assumes that guenons

cognitively encode different species' face patterns as the mean of all encountered examples. In

previous work using similar methods, results were robust to the choice of learning model [10].

156   To avoid using the same individual guenons to both train and test our species classifier

we used a leave-one-out procedure for all analyses. For this procedure, we systematically

removed each individual from the image set, repeated the analysis procedure outlined above,

then classified each image of the excluded individual based on the features generated from all

other images. All species included in these analyses are represented by at least four individuals

(range: 4-23). We present results for all species, however results for species with samples sizes

in the lower end of this range should be considered less robust and interpreted with caution.

Eigenface-based features can be used to reliably classify guenons by species based on

axes of variation, however the extent to which specific facial characteristics are relevant for

correct classification of each species is difficult to determine. We used an occlude-reclassify

scheme developed to identify which image regions contribute most to correct classification in

computer vision classification tasks [43]. For each correctly classified image, we systematically

blocked each image region and re-classified the image; a correct re-classification indicates that

the occluded region of the face was unnecessary for correct classification, while an incorrect re-

classification indicates that the occluded region was essential. Occlusion of face regions was

accomplished by setting the relevant pixel as well as all those in a thirty-pixel radius to the mean

face color of that species. This procedure was repeated for every pixel in the image, effectively

sliding the occluded region across all face areas. A radius of thirty pixels occludes approximately

five percent of the image (Figure 2), with the specific region being occluded shifting by one

pixel at each iteration. Primate faces are broadly symmetrical, therefore to avoid the presence of

duplicate spatial information that may support species classification when part of the face is

occluded, we ran analyses on the left and right halves of the face separately. Results differed

little, so for clarity we report the results from the left hemi-face classification in the main text,

with right-side results summarized in the supplementary results. For more details on the

implementation of the occlude-reclassify procedure, see supplementary methods. Based on this

occlude-reclassify scheme, we generated a binary image for each image in our data set, with each

pixel being either zero (black) or one (white) base on whether the image was correctly classified

when that pixel and its neighbors was occluded. We then averaged these binary images across

individuals and species to generate species level heatmaps depicting face regions that are

essential for correct classification across species. For visualization, we converted greyscale

heatmaps to color using a color mapping function. To facilitate the identification of critical face

regions, occlusion results are presented as composite images combining heatmaps and a

greyscale version of the relevant species average face, with transparency set to 0.5.

Heatmaps vary across species in the extent to which face regions identified as essential

for correct species classification are spread across the face (i.e. ranging from small and isolated

face regions to large portions of the face identified as critical) as well as the relative import of

identified regions (i.e. the likelihood that identified regions caused misclassification, encoded as

how dark identified regions are in the heatmap). To quantify the spread and relative importance

of the identified face regions across species, we calculated the proportion of the face

misclassified and the mean classification error, respectively. The proportion of the face

misclassified was calculated as the number of heatmap pixels less than one (i.e. those that were

ever incorrectly classified) divided by the total number of pixels in the average face for each

species; higher values indicate that the face regions essential for correct species classification are

spread more widely across the face. The mean classification error was calculated as the mean

value of all heatmap pixels less than one; higher values indicate that the face regions identified

are particularly critical and more often lead to misclassification when occluded (i.e. the identified

regions are darker in the heatmaps). Computational analyses were conducted in MATLAB^{TM} and

run on the High Performance Computing cluster at New York University.

Looking time experiments

Looking time experiments were conducted at CERCOPAN sanctuary in Calabar, Nigeria, and

included 18 adult putty nosed monkeys (6 males, 12 females) and 16 adult mona monkeys (10

males, 6 females). Each species was divided into four experimental groups (based on socially

housed groups), with all individuals in the group viewing the same images in the same order. In

each species, two experimental groups were presented with male stimulus images and two with

female stimulus images across all trials. Experiments involved the simultaneous presentation of

two stimulus images to subjects, with their resulting eye gaze measured to determine visual

biases. Stimulus preparation and experimental procedures were carried out following the

recommendations of Winters et al. [37]. Briefly, we prepared stimulus images depicting guenon

faces which were presented approximately life-sized (image size on screen: 500 x 500 pixels,

11.96 x 11.96 cm), with accurate colors, and standardized for relevant characteristics. Stimulus

image pairs were presented to subjects side-by-side using a custom-designed experimental

apparatus. For more details regarding subjects, stimuli preparation, and experimental apparatus

design, see supplemental methods.

Each subject participated in three trials, with stimulus image pairs depicting the

following: (1) a conspecific and a heterospecific that shares a focal trait with the conspecific, (2)

a conspecific and a heterospecific that does not share a focal trait with the conspecific, and (3) a

conspecific for which the focal trait has been modified and a heterospecific that shares the focal

trait with the conspecific. Heterospecifics presented to putty nosed monkeys were Wolf's

guenons (*C. wolfi*, no nose spot) and red-tailed monkeys (*C. ascanius*, nose spot); heterospecifics

presented to mona monkeys were red-tailed monkeys (no eyebrow patches) and Diana monkeys

(*C. diana*, eyebrow patches). Heterospecific species were selected based on the presence/absence

of the relevant facial trait, a lack of range overlap with the subject species, and availability of

sufficient and appropriate images in our database. Image presentation locations (i.e. left verses

right) were counterbalanced across trials, and trial order was varied across subjects; both factors

were included in statistical analyses. For each trial, we placed the experimental apparatus

immediately outside the relevant enclosure and recorded the identities of participating subjects.

We waited a minimum of one week between trials of the same subject to minimize habituation or

trial order effects.

235   Videos of each trial were coded frame by frame to quantify the amount of time subjects

spent looking at each stimulus image. All coding was done blind to trial conditions and stimulus

image location. Reliability was assessed using approximately 10% of all trial videos, in which

we assessed agreement between two coders on the direction of jointly coded looks within these

trials as being in agreement in 94.46% of frames (Cohen's kappa $= 0.883$), which is well within

the range of acceptable reliability scores for this type of data [37,44]. Raw looking time data was

compiled to yield a total number of frames spent looking at each stimulus image for each subject

in each trial. Subjects varied widely in their level of interest in experiments, resulting in

considerable variation in overall looking time. We therefore used only the first five seconds of

looking for each subject in each trial, while allowing them to complete the current look at the

five second mark (i.e. we required at least one second of non-looking before terminating coding

for each subject). This resulted in a mean total looking time ($\pm$ standard deviation) of 3.89s ($\pm$

1.98s) for putty nosed monkeys and 4.58s ($\pm$ 2.52s) for mona monkeys, which is similar to

durations reported in previous looking time experiments in primates [37,44]. Because a direct

comparison is made between the species depicted in stimuli, each trial effectively serves as its

own control.

Statistical analyses

We analyzed differences in looking time elicited by subjects in experimental trials using

generalized linear mixed models (GLMMs). Models were fit using a binomial family

distribution, with the number of video frames spent looking at the targeted stimulus image and

the number of video frames spent looking at the paired image set as the binomial outcome

variable. This structure allowed us to assess looking biases while accounting for any differences

in total looking time across subjects. All models included group, subject, and unique trial (i.e. a

unique identifier for each subject in each trial, included to account for our analysis of the two

images presented in each trial as separate data 'rows') as nested random effects. Stimulus species

(conspecific v. heterospecific) and focal trait similarity (presence of nose spots for putty nosed

monkeys and eyebrow patches for mona monkeys), were included as fixed effects. We also

included the following additional factors as fixed effects: subject age (log transformed), sex, and

origin (captive v. wild born); stimulus image presentation spot (right v. left), eye contact (direct

eye contact with the camera or looking slightly away), sex, and degree of familiarity to the

subject; and trial order, apparatus pattern, and display ICC profile. For more details about these

variables see supplemental methods.

To determine which variables significantly influenced subject looking biases, we

compared models with different parameterizations using likelihood ratio tests (LRTs). A single

model including all fixed effects simultaneously would involve an excessive number of

predictors. We therefore first analyzed each variable separately via comparisons to a null model

including only random effects, and excluded non-significant predictors from subsequent

analyses. We generated an initial model composed of factors that were statistically significant

(alpha < 0.05) or exhibited a trend (alpha < 0.1) when tested alone. To determine the statistical

significance of these factors we then systematically excluded each factor from this model and

tested its contribution to the fit of the model to the data using LRTs. When species (conspecific

v. heterospecific) and focal trait (shared v. not shared) were both significant predictors in this

model we also tested a species*trait interaction. Within a final model composed of significant

predictors we compared across factor levels of fixed effects using z scores calculated using a

normal approximation. Adherence to model assumptions was verified based on plots of fitted

values and residuals. Trials from putty nosed and mona monkeys were analyzed separately.

GLMMs were run using the 'lme4' package version 1.0.12 [45] in R version 3.3.3 [46].

Results

Occlude-reclassify machine classification

We began by confirming that guenons could be reliably classified by species based on eigenface

decomposition [10]. Average subject images were correctly classified by species 99.31% of the

time, and distinct images were correctly classified 93.03% of the time. All correctly classified

images (n = 654) were used to identify face regions of critical importance to correct species

classification by the computer algorithm, using our occlude-reclassify scheme. We identified

essential face regions in all guenon species that, when occluded, led to incorrect species

classification (Figure 3; for full resolution images see Supplementary File 1). Species differed in

the importance of different face regions as well as the extent to which important regions were

concentrated in specific facial features or were more widely distributed across larger face areas

(Figure 4). For example, the nose spot of the putty nosed monkey was the most critical facial

feature identified across all species. The putty nosed monkey had the highest mean error rate for

misclassified face regions – indicating that the face regions identified had the highest likelihood

of causing misclassification when occluded – with the essential regions centered exclusively on

the nose. Thus, in the putty nosed monkey the nose is the only essential face feature; when the

nose is occluded species classification breaks down, whereas occluding any other face region has

no effect. In contrast, in other species our classifier relied on broader regions of the face, with

larger face regions identified as important for correct classification and the classifier relying less

exclusively on a single feature. The mona monkey is a good example of this, with disparate face

regions including the cheeks, eyebrows, and ear tufts all influencing correct classification of this

species. In some species negative space is important, suggesting that what makes the faces of

species distinctive may be the absence of certain facial traits. For instance, in *M. talapoin* the

absence of distinctive traits along the sides of the face – such as cheek and/or ear tufts observed

in other species – appears to be important.

Looking time experiments

Our experiments presenting subjects with pairs of con- and heterospecific faces revealed visual

biases in resulting eye gaze in both putty nosed and mona monkeys. In the subset of trials that

included a natural conspecific and a heterospecific without the relevant face trait (i.e. those

where the relevant facial traits are not spread across both con- and heterospecific faces), species

(and therefore also facial trait) was a significant predictor of looking behavior (putty nosed

monkeys: Chisq = 63.312, p < 0.001; mona monkeys: Chisq = 30.755, p < 0.001), with both

putty nosed and mona monkeys exhibiting a conspecific bias (respectively: z = 7.920, p < 0.001;

z = 5.536, p < 0.001; Figure 5).

Across all trials, in putty nosed monkeys model comparisons revealed that looking

behavior was significantly influenced by facial trait (nose spot v. no nose spot; Chisq = 11.511, p

< 0.001) and image location (right v. left; Chisq = 18.065, p < 0.001), but not by species

(conspecific v. heterospecific; Chisq = 3.051, p = 0.081). Overall, putty nosed monkeys looked

longer at stimulus faces that displayed a white nose patch (z = 3.343, p < 0.001; Figure 5), their

diagnostic species trait, regardless of species identity. Putty nosed monkeys also exhibited a

significant right gaze bias (z = 4.289, p < 0.001). None of the other variables relating to subject,

stimulus, or trial characteristics were statistically significant (all p > 0.1; Supplementary Table

1).

In mona monkeys, model comparisons revealed that looking behavior was significantly

influenced by species (conspecific v. heterospecific; Chisq = 177.480, p < 0.001), facial trait

(eyebrow patches v. no eyebrow patches; Chisq = 29.462, p < 0.001) and a species*trait

interaction (Chisq = 8.242, p = 0.004). Across all trials, mona monkeys looked longer at

conspecifics (z = 9.945, p < 0.001; Figure 5) and as a separate effect, faces without white

eyebrow patches, one component of their overall wider diagnostic discrimination area (z = 5.851,

p < 0.001). There was also an interaction between these two variables, with mona monkeys

looking longer at heterospecific faces with white eyebrow patches (z = 2.868, p = 0.004). None

of the other variables relating to subject, stimulus, or trial characteristics played a significant role

in mona monkey visual biases (all p > 0.1; Supplementary Table 2).

Discussion

Our experiments show that eye gaze in guenons is influenced by face regions identified as

critical to correct species classification by our machine classifier. This convergence of results

using disparate methods reinforces the validity of both, and ties computationally derived results

directly to guenon perception, demonstrating the utility of machine learning for identifying

biologically relevant signal components. To our knowledge, ours is the first analysis to use

machine classification combined with the systematic occlusion of image regions to characterize

the relevant signaling information encoded in an animal's appearance. This approach, based on

research in the field of computer vision designed to assess the contribution of image contents to

object classification [43], is useful for objectively quantifying the relative roles of different

signal components with respect to overall signal function. In closely-related sympatric species,

selection against mating or interacting with heterospecifics is often associated with the evolution

of species-typical traits used to maintain reproductive and behavioral isolation. The guenons, a

recent and diverse radiation that exhibit mixed species groups in which hybridization is rarely

observed, exemplify this phenomenon. By showing how species classification is dependent on

different aspects of face patterning and that this links with looking time toward con and

heterospecifics, our analyses support a role for guenon face patterns in species discrimination,

and identify specific face regions critical for this function. This parsing of critical signal

components is critical for understanding the phenotypic evolution of complex signals and

identifying relevant axes of signal variation for additional analyses.

Our occlude-reclassify analysis identified face regions critical to correct species

classification by a machine classifier in all guenon species included in our study. Critical regions

differed in both location and spread across the face, suggesting variation in potential use across

species. For some guenons, reliance on a single facial characteristic may be sufficient for species

discrimination. The best example of this in our data set is the putty nosed monkey, where our

machine classifier relied exclusively on the white nose spot to classify this species. That is,

occlusion of any other region resulted in correct classification, but when the nose spot was

occluded classification failed. This result is reinforced by our experiments, in which putty nosed

monkey visual attention was driven wholly by the presence of nose spots. Putty nosed monkeys

exhibited a conspecific bias when presented with natural con- and heterospecific faces, as is

typical in primates, however including stimuli depicting heterospecifics with nose spots and

conspecifics without nose spots completely obscured this conspecific bias. This combination of

results illustrates the importance of nose spots in this species. It is worth noting that putty nosed

monkey nose spots are the most straightforward facial trait documented in our analysis (i.e. putty

nosed monkeys were only misclassified when the nose spot was occluded and occluding the nose

spot led to a high rate of misclassification) and the relative simplicity of the face and related

visual biases in this species is likely exceptional. On the whole, species discrimination signals in

a large radiation with varying patterns of sympatry are expected to be complex and

multidimensional, and it is likely that only some species can exhibit single-trait-based signals

and visual biases without the system breaking down. This is supported by our results showing

that for most guenon species our classifier relied on multiple face regions for species

discrimination.

Not all guenons exhibited critical face regions restricted to a single facial trait, and our

machine classifier sometimes relied on disparate face regions. In our data set, the mona monkey

is a good example of such a species. Like in putty nosed monkeys, our experiments with mona

monkeys supported these computational results. Mona monkeys exhibited a conspecific bias

across all trials, regardless of single trait manipulations, as well as an additional bias based on the

presence of eyebrow patches. Thus, eyebrow patches alone do not appear to be the sole focus of

attention in mona monkeys. We predict that additional manipulation of other face regions would

be necessary to redirect their visual attention. Nonetheless, that mona monkey attention is still

influenced by this species-typical trait shows that it is important but not essential, a result

predicted by our computational analyses. It is unclear why mona monkeys would look longer at

stimuli without eyebrow patches, however it is possible that utilization of the whole face causes

increased attention to incongruency (e.g. conspecifics without eyebrow patches or

heterospecifics with them). Our results suggest that in mona monkeys, species discrimination

may be based on broader face information, and the perceptual processes involved in assessing

potential mates could be similar to generalized holistic face processing mechanisms observed in

other primates [47].

Our results suggest that guenons, while united by a general pattern of facial

diversification and the probable use of faces in mate choice, may vary across species in the

specific traits and processes that are involved in discriminating between conspecifics and

heterospecifics. Our pattern of results for putty nosed monkey nose spots and mona monkey

eyebrow patches is interesting because we know that both traits do contain sufficient information

to discriminate between species that share these features [24], yet they influence attention

differently in the two species. This disparity highlights the importance of testing receiver

perception directly. The fact that our experimental results with guenons line up with predictions

generated by our occlude-reclassify analysis implies that these computationally derived results

are biologically valid. Interestingly, we found no sex differences in visual biases for either

species, suggesting that selective pressures on species discrimination signaling and preference

traits are similar between sexes.

In guenons, an observed lack of hybrids in most polyspecific groups [21] is notable given

that hybridization is known to be possible between many guenon species [21,23,27], and

indicates the existence of pre-mating barriers to reproduction. Increased eye gaze is associated

with increased mating interest in humans [48] and non-human primates [44,49], suggesting that

our experimental results would generalize to mating contexts in guenons. Combined with

previous work [10,23,24,26,27], our results support the hypothesis that guenon face patterns play

a role in mate choice and reproductive isolation in this group. However, it remains possible that

the selection pressure for species discrimination traits in guenons arises partially or entirely from

other functions where behavioral coordination or avoidance between species is advantageous,

such as in foraging decisions [20,23]. Careful field observations would be needed to distinguish

between such possibilities.

Our occlude-reclassify approach is a novel method for identifying the distribution of

information in complex signals and can be used for any question that can be conceptualized as a

discrimination problem and analyzed using machine classification. This method therefore has

broad utility within sensory ecology and could help to better understand the link between form

and function in the complex signals that are common in many animal groups. The objectivity of

the approach is important, as it allows researchers to intelligently target specific signal

components for further analysis without reference to their own perceptions of their salience. This

is particularly important when studying species with sensory and perceptual systems very

different from our own [50,51]. Where possible, combining this approach with a biologically

realistic classification scheme, such as classification within a perceptual face space based on

eigenface scores [36] as used here, increases the biological validity of results.

Our research broadens our understanding of how morphology and social decision-making

can interact to structure interactions between species living in sympatry. In guenons, facial

features like white nose spots are highly salient, attention-grabbing, and distinctive, and our

combined results demonstrate the importance of these traits in species discrimination. Guenon

behavioral repertoires, such as nose-to-nose touching observed in wild putty nosed monkeys

(SW, personal observation) and red-tailed monkeys [52], further reflect the importance and

biological relevance of these traits. Primates preferentially attend to facial information [53,54],

making face patterns particularly suited to influencing behavior and decision-making in con- and

heterospecifics. The evolution of signals facilitating species discrimination may be a major

driver of biological diversity, and our work linking mating signal form and function in a recent

and diverse primate radiation highlights how such evolutionary processes can be important in

generating animal phenotypes.

References

1. Coyne JA, Orr HA. 2004 *Speciation*. Sunderland, MA: Sinauer Associates, Inc.
2. Andersson MB. 1994 *Sexual selection*. Princeton, NJ: Princeton University Press.
3. Dawkins MS, Guilford T. 1997 Conspicuousness and diversity in animal signals. In
Communication (eds DH Owings, MD Beecher, NS Thompson), pp. 55–75. Boston, MA:
Springer US.
4. Brown WL, Wilson EO. 1956 Character displacement. *Syst. Zool.* **5**, 49–64.
5. Pfennig KS, Pfennig DW. 2009 Character displacement: ecological and reproductive
responses to a common evolutionary problem. *Q. Rev. Biol.* **84**, 253–276.
(doi:10.1086/605079)
6. Higgie M, Chenoweth S, Blows MW. 2000 Natural selection and the reinforcement of mate
recognition. *Science* **290**, 519–521. (doi:10.1126/science.290.5491.519)
7. Lukhtanov VA, Kandul NP, Plotkin JB, Dantchenko A V, Haig D, Pierce NE. 2005
Reinforcement of pre-zygotic isolation and karyotype evolution in Agrodiaetus butterflies.
Nature **436**, 385–9. (doi:10.1038/nature03704)
8. McNaught MK, Owens IPF. 2002 Interspecific variation in plumage colour among birds:
species recognition or light environment? *J. Evol. Biol.* **15**, 505–514.
9. Kameda Y, Kawakita A, Kato M. 2009 Reproductive Character Displacement in Genital
Morphology in *Satsuma* Land Snails. *Am. Nat.* **173**, 689–697. (doi:10.1086/597607)
10. Allen WL, Stevens M, Higham JP. 2014 Character displacement of Cercopithecini primate
visual signals. *Nat. Commun.* **5**, 4266. (doi:10.1038/ncomms5266)
11. Pfennig KS, Rice AM. 2014 Reinforcement generates reproductive isolation between
neighbouring conspecific populations of spadefoot toads. *Proc. R. Soc. B Biol. Sci.* **281**,
20140949–20140949. (doi:10.1098/rspb.2014.0949)
12. Stanger-Hall KF, Lloyd JE. 2015 Flash signal evolution in *Photinus* fireflies: Character
displacement and signal exploitation in a visual communication system. *Evolution* **69**, 666–
682. (doi:10.1111/evo.12606)

13. Gordon NM, Ralph MZ, Stratman KD. 2017 Rapid character displacement of different call
472   parameters in closely related treefrogs (Hyla cinerea and H. gratiosa). *Behav. Ecol.*
473   *Sociobiol.* **71**. (doi:10.1007/s00265-017-2341-1)
474   14. Mendelson TC, Shaw KL. 2012 The (mis) concept of species recognition. *Trends Ecol. Evol.*
475   **27**, 421–427. (doi:10.1016/j.tree.2012.04.001)
476   15. Mallet J. 2005 Hybridization as an invasion of the genome. *Trends Ecol. Evol.* **20**, 229–237.
477   (doi:10.1016/j.tree.2005.02.010)
478   16. Groves CP. 2005 Order Primates. In *Mammal Species of the World: A Taxonomic and*
479   *Geographic Reference*, pp. 111–184. Baltimore, MD: Johns Hopkins University Press.
480   17. Grubb P, Butynski TM, Oates JF, Bearder SK, Disotell TR, Groves CP, Struhsaker TT. 2003
481   Assessment of the Diversity of African Primates. *Int. J. Primatol.* **24**, 1301–1357.
482   (doi:10.1023/B:IJOP.0000005994.86792.b9)
483   18. Hart JA *et al.* 2012 Lesula: a new species of Cercopithecus monkey endemic to the
484   Democratic Republic of Congo and implications for conservation of Congo's central basin.
485   *PloS One* **7**, e44271. (doi:10.1371/journal.pone.0044271)
486   19. Tosi AJ, Detwiler KM, Disotell TR. 2005 X-chromosomal window into the evolutionary
487   history of the guenons (Primates: Cercopithecini). *Mol. Phylogenet. Evol.* **36**, 58–66.
488   (doi:10.1016/j.ympev.2005.01.009)
489   20. Jaffe KE, Isbell LA. 2011 The guenons: polyspecific associations in socioecological
490   perspective. In *Primates in Perspective* (eds CJ Campbell, A Fuentes, KC MacKinnon, SK
491   Bearder, RM Stumpf), pp. 277–300. New York, NY: Oxford University Press.
492   21. Detwiler KM, Burrell AS, Jolly C. 2005 Conservation implications of hybridization in
493   African cercopithecine monkeys. *Int. J. Primatol.*
494   22. Cords M. 2012 The behavior, ecology, and social evolution of Cercopithecine monkeys. In
495   *The Evolution of Primate Societies* (eds JC Mitani, RA Palombit, JB Silk), pp. 91–112.
496   Chicago: University of Chicago Press.
497   23. Kingdon J. 1997 *The Kingdon guide to African mammals*. London: Academic Press.
498   24. Allen WL, Higham JP. 2015 Assessing the potential information content of multicomponent
499   visual signals: a machine learning approach. *Proc. R. Soc. B Biol. Sci.*
500   (doi:10.1098/rspb.2014.2284)
501   25. Bradley BJ, Mundy NI. 2008 The primate palette: The evolution of primate coloration. *Evol.*
502   *Anthropol.* **17**, 97–111. (doi:10.1002/evan.20164)
503   26. Kingdon J. 1988 What are face patterns and do they contribute to reproductive isolation in
504   guenons. In *A Primate Radiation: Evolutionary Biology of the African Guenons*, pp. 227–
505   245.
506   27. Kingdon J. 1980 The role of visual signals and face patterns in African forest monkeys
507   (guenons) of the genus Cercopithecus. *Trans. Zool. Soc. Lond.* **35**, 425–475.
508   (doi:10.1111/j.1096-3642.1980.tb00062.x)
509   28. Pfennig DW, Pfennig KS. 2010 Character Displacement and the Origins of Diversity. *Am.*
510   *Nat.* **176**, S26–S44. (doi:10.1086/657056)
511   29. Pfennig KS. 2016 Reinforcement as an initiator of population divergence and speciation.
512   *Curr. Zool.* **62**, 145–154. (doi:10.1093/cz/zow033)
513   30. Stuart YE, Inkpen SA, Hopkins R, Bolnick DI. 2017 Character displacement is a pattern: so,
514   what causes it? *Biol. J. Linn. Soc.* **121**, 711–715. (doi:10.1093/biolinnean/blx013)
515   31. Patricelli GL, Hebets EA. 2016 New dimensions in animal communication: the case for
516   complexity. *Curr. Opin. Behav. Sci.* **12**, 80–89. (doi:10.1016/j.cobeha.2016.09.011)

32. Turk M, Pentland A. 1991 Eigenfaces for Recognition. *J. Cogn. Neurosci.* **3**, 71–86.
(doi:10.1162/jocn.1991.3.1.71)
33. Perera TA, Collins J. 2015 Imaged based species recognition system. In *9th International*
Conference on Sensing Technology (ICST), pp. 195–199. IEEE.
34. Shiau Y-H, Lin S-I, Chen Y-H, Lo S-W, Chen C-C. 2012 Fish observation, detection,
recognition and verification in the real world. In *Proceedings of the International*
Conference on Image Processing, Computer Vision, and Pattern Recognition (IPCV), pp. 1–
6. The Steering Committee of The World Congress in Computer Science, Computer
Engineering and Applied Computing (WorldComp).
35. Weeks P, O'Neill M, Gaston K, Gauld I. 1999 Species–identification of wasps using
principal component associative memories. *Image Vis. Comput.* **17**, 861–866.
(doi:10.1016/S0262-8856(98)00161-9)
36. Chang L, Tsao DY. 2017 The code for facial identity in the primate brain. *Cell* **169**, 1013-
1028.e14. (doi:10.1016/j.cell.2017.05.011)
37. Winters S, Dubuc C, Higham JP. 2015 Perspectives: the looking time experimental paradigm
in studies of animal visual perception and cognition. *Ethology* **121**, 625–640.
(doi:10.1111/eth.12378)
38. Fujita K. 1987 Species recognition by five macaque monkeys. *Primates* **28**, 353–366.
(doi:10.1007/BF02381018)
39. Demaria C, Thierry B. 1988 Responses to animal stimulus photographs in stumptailed
macaques (Macaca arctoides). *Primates* **29**, 237–244. (doi:10.1007/BF02381125)
40. Dufour V, Pascalis O, Petit O. 2006 Face processing limitation to own species in primates: a
comparative study in brown capuchins, Tonkean macaques and humans. *Behav. Processes*
73, 107–13. (doi:10.1016/j.beproc.2006.04.006)
41. Méary D, Li Z, Li W, Guo K, Pascalis O. 2014 Seeing two faces together: preference
formation in humans and rhesus macaques. *Anim. Cogn.* **17**, 1107–1119.
(doi:10.1007/s10071-014-0742-3)
42. Rakotonirina H, Kappeler PM, Fichtel C. 2018 The role of facial pattern variation for species
recognition in red-fronted lemurs (Eulemur rufifrons). *BMC Evol. Biol.* **18**.
(doi:10.1186/s12862-018-1126-0)
43. Zeiler MD, Fergus R. 2014 Visualizing and Understanding Convolutional Networks. In
Computer Vision – ECCV 2014 (eds D Fleet, T Pajdla, B Schiele, T Tuytelaars), pp. 818–
833. Cham: Springer International Publishing. (doi:10.1007/978-3-319-10590-1_53)
44. Dubuc C, Allen WL, Cascio J, Lee DS, Maestripieri D, Petersdorf M, Winters S, Higham JP.
2016 Who cares? experimental attention biases provide new insights into a mammalian
sexual signal. *Behav. Ecol.* **27**, 68–74. (doi:10.1093/beheco/arv117)
45. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models using
lme4. *J. Stat. Softw.* **67**, 1–48. (doi:10.18637/jss.v067.i01)
46. R Core Team. 2018 *R: a language and environment for statistical computing, v3.5.2.* R
Foundation for Statistical Computing, Vienna, Austria. See https://www.R-project.org.
47. Dahl CD, Wallraven C, Bülthoff HH, Logothetis NK. 2009 Humans and macaques employ
similar face-processing strategies. *Curr. Biol.* **19**, 509–13. (doi:10.1016/j.cub.2009.01.061)
48. Shimojo S, Simion C, Shimojo E, Scheier C. 2003 Gaze bias both reflects and influences
preference. *Nat. Neurosci.* **6**, 1317–1322. (doi:10.1038/nn1150)

49. Dubuc C, Allen WL, Maestripieri D, Higham JP. 2014 Is male rhesus macaque red color
ornamentation attractive to females? *Behav. Ecol. Sociobiol.* **68**, 1215–1224.
(doi:10.1007/s00265-014-1732-9)
50. Endler JA. 1990 On the measurement and classification of colour in studies of animal colour
patterns. *Biol. J. Linn. Soc.* **41**, 315–352.
51. Bennett ATD, Cuthill IC, Norris K. 1994 Sexual selection and the mismeasure of color. *Am.*
Nat. , 848–860.
52. Estes R. 1991 *The Behavior Guide to African Mammals: Including Hoofed Mammals,*
Carnivores, Primates. Berkeley & Los Angeles, CA: University of California Press.
53. Tsao DY, Schweers N, Moeller S, Freiwald WA. 2008 Patches of face-selective cortex in the
macaque frontal lobe. *Nat. Neurosci.* **11**, 877–879. (doi:10.1038/nn.2158)
54. Kano F, Tomonaga M. 2009 How chimpanzees look at pictures: a comparative eye-tracking
study. *Proc. R. Soc. B Biol. Sci.* **276**, 1949–1955. (doi:10.1098/rspb.2008.1811)

Acknowledgements

We thank Allegra Depasquale and Laura Newman for assistance coding looking time videos, and

Kathryn Yee for assistance with stimuli preparation. Special thanks the directors and staff of

CERCOPAN sanctuary, particularly Claire Coulson and Isabelle Theyse, for access to the

facility and support during data collection.

Data accessibility

The guenon face image database, experimental looking time data, and all project code will be

uploaded to Dryad. During review, all files are available at:

http://www.dropbox.com/sh/1yf97kbhggzi5a8/AAAJ0dQMEqeFOMf9aHzGBMHNa?dl=0

Figure 1

Example experimental stimulus pairs. Subjects were shown a pair of stimulus images consisting

of a conspecific and a heterospecific. Facial traits (nose spots for putty nosed monkeys and

eyebrow patches for mona monkeys) were varied across trials, with conspecifics paired with a

heterospecific species that shares the facial trait (row 1) and one that does not (rows 2 and 3).

Conspecifics were displayed either naturally (rows 1 and 2) or with the facial trait removed (row

3). All subjects participated in all three trial types. Trial order and stimulus image side were

counterbalanced across subjects.

Figure 2

Average guenon face with an occluder shown in the top left. The occluder is depicted in black

for maximal visibility, but in analyses presented here is set to the mean face color of the relevant

species. During the occlude-reclassify analysis, the occluder is slid across the image and the

image re-classified; an incorrect classification at a given occluder location indicates the presence

of face information critical to correct classification. Analyses are run on hemi-faces to account

for facial symmetry. Image borders outside the radius of the occluder are not tested; the dashed

line encloses the region of the image analyzed using the occlude-reclassify procedure.

Figure 3

Likelihood of correct classification based on occlusion of different face regions. Species average faces are displayed on the left and

heatmaps identifying critical face regions on the right. Sample size is reported as n = number of individuals (number of total images).

Figure 4

Variation across species in face regions identified as essential for correct species classification.

The proportion of the face misclassified (y-axis) indicates the spread of essential regions across

the face; higher values signify broader spread and lower values more concentrated regions. The

mean classification error (x-axis) measures the relative importance of identified features; higher

values indicate higher rates of misclassification, suggesting identified regions are particularly

essential for correct species classification. Experimental results are presented for C. mona and C.

nictitans (Figure 5).

Figure 5

Species and trait biases observed during looking time tasks with (a) putty nosed monkeys and (b)

mona monkeys. Leftmost plots depict differences in looking time in trials consisting of

conspecifics and heterospecifics without the relevant facial trait. Center and right plots depict

looking time differences across all trials – which also include heterospecifics with the relevant

facial trait and conspecifics without it – with species biases depicted in the center and trait biases

on the right. Results are based on 18 putty nosed monkeys and 16 mona monkeys. Each subject

participated in three trials (see Figure 1 for example stimuli for each trial type).